


Z-Selective Synthesis of Vinyl Boronates through Fe-Catalyzed Alkyl Radical Addition

Guido Barzanò, Alexis Cheseaux, and Xile Hu*^{1b}

Laboratory of Inorganic Synthesis and Catalysis, Institute of Chemical Sciences and Engineering, Ecole Polytechnique Fédérale de Lausanne (EPFL), ISIC-LSCI, BCH 3305, Lausanne 1015, Switzerland

S Supporting Information

ABSTRACT: Z-Selective synthesis of vinyl boronates is challenging. This work describes Fe-catalyzed addition of alkyl radicals, formed by the corresponding alkyl halides, to ethynyl ethynylboronic acid pinacol ester that gives rise to Z-vinyl boronates in high stereoselectivity. The method works best for tertiary and secondary alkyl iodides. The Z-vinyl boronate products are easily converted to other Z-alkenes by transformation of the boronate group. The method is applied for the formal total synthesis of a 5-HT_{2c} receptor agonist.



Z-Selective synthesis of 1,2-disubstituted alkenes is an important yet challenging task in organic synthesis.^{1,2} Because the boronate group can be readily transformed into many common functional groups,³ Z-olefins might be conveniently accessed from Z-vinyl boronates. While hydroborylation of terminal alkynes is an efficient method to synthesize vinyl boronates, it typically leads to the thermodynamically more stable *E*-isomers (Figure 1a).^{4–11} Until now, there have been only several reports of Z-selective hydroboration of terminal

alkynes, with limited scope for alkyl alkynes.^{12–15} Here, we describe an alternative approach to the synthesis of Z-vinyl boronates. Our approach is based on Fe-catalyzed stereoselective addition of an alkyl radical, generated from an alkyl halide under reductive conditions, to ethynylboronic acid pinacol ester (**1**, Figure 1b). The approach is built upon our previous work,^{16–18} which showed that an alkyl radical was generated through the reaction of an alkyl halide with an Fe(I) species, produced by reduction of an Fe(II) catalyst with Zn (Figure 1c,d). The alkyl radical could be trapped by a terminal aryl alkyne to form a vinyl radical stabilized by an aryl group. Recombination of the vinyl radical with Fe(I) gave an Fe(II) vinyl intermediate. This step was stereoselective and favored a *cis*-arrangement of the aryl and alkyl groups. Transmetalation of the Fe(II) vinyl intermediate to Zn(II), followed by protonation, yielded the *cis*-alkene. In this chemistry, the aryl substituent of the alkyne was essential. Replacement of the aryl group by another electron-withdrawing group such as keto, ester, nitrile, or amide led to no formation of alkene. Here, we show that the boronic acid pinacol ester group (Bpin) can serve the same function of an aryl group in stabilizing the vinyl radical. An important advantage of the Bpin group over an aryl group is that the former could be readily converted to numerous other functional groups, making this method of Z-alkene synthesis truly general.

The reaction of **1** with *tert*-butyl iodide (**2a**) was chosen as the test reaction (Table 1). At first, the conditions previously developed for the reductive coupling of ethynylbenzene with **2a**, that is, 1 equiv of **1**, 3 equiv of **2a**, 20 mol % of FeBr₂, 3 equiv of Zn metal, and 2 mol % of I₂ activator, DMA as solvent, 80 °C, were employed (Table 1, entry 1). To our delight, a yield of 48% and a Z/E selectivity of 92:8 were





Figure 1. (a) Hydroborylation of alkyne to give vinyl boronates. (b) Synthesis of Z-vinyl boronates by Fe-catalyzed alkyl radical addition to **1**. (c) Synthesis of Z-styrenes by Fe-catalyzed alkyl radical addition to aryl alkynes. (d) Proposed mechanism for the reaction in c.

Received: November 26, 2018

Table 1. Optimization of Conditions for the Reaction of 1 with 2a^a



entry	cat.	n	aa	temp (°C)	yield (%)	Z/E ^b
1	FeBr ₂	3	I ₂	80	48	12:1
2	CuBr ₂	3	I ₂	80	33	13:1
3	NiI ₂	3	I ₂	80	0	nd
4	CoI ₂	3	I ₂	80	22	16:1
5	FeI ₂	3	I ₂	80	53	12:1
6	Fe(OTf) ₂	3	I ₂	50	82	6:1
7	Fe(OTf) ₂	3	TMS-I	50	95	44:1
8	Fe(OTf) ₂ ^c	3	TMS-I	50	42	53:1
9	Fe(OTf) ₂	1	TMS-I	50	2	nd
10	Fe(OTf) ₂	2	TMS-I	50	47	130:1

^aGeneral conditions: alkyne boronate ester (0.5 mmol), *tert*-butyl iodide (1.5 mmol), under N₂, in DMA (1 mL). aa = activating agent; temp = temperature; nd = not determined. ^bAccording to quantification by GC analysis. ^c10 mol % loading of catalyst.

obtained. Other metal salts such as CuBr₂, NiI₂, and CoI₂ were tested as catalysts. NiI₂ gave no yield, while CuBr₂ and CoI₂ gave lower yields but similar Z/E selectivity (Table 1, entries 2–4). FeI₂ had a similar efficiency to FeBr₂, but Fe(OTf)₂ (OTf = triflate) gave a much higher yield of 82% when the reaction was conducted at 50 °C (Table 1, entries 5 and 6). Unfortunately, the Z/E selectivity dropped to 6:1. Replacement of the I₂ activator by TMS-I (TMS = tetramethyl silyl) led to a yield of 95% and a Z/E selectivity of 44:1 (Table 1, entry 7). When the loading of Fe(OTf)₂ was decreased to 10 mol %, the yield dropped to 42% (Table 1, entry 8). When the loading of Zn was decreased to 2 or 1 equiv, the yields also dropped significantly (Table 1, entries 9 and 10).

The optimized conditions in Table 1 (entry 7) were applied for the coupling of 1 with various tertiary and secondary alkyl iodides, yielding the corresponding Z-alkenes (Figure 2). *tert*-Butyl (3a), 1-adamantyl (3b), and *tert*-pentyl (3c) groups were added with high yields. Secondary alkyl groups such as 2-octyl (3d), cyclohexyl (3e), cyclooctyl (3f), 2-butyl (3g), and 1-*tert*-butyl-4-cyclohexyl (3h) were added with good yields and selectivity as well. The Z-isomer of 3h was obtained with 14:1 trans/cis diastereoselectivity. Addition of 2-propylbenzene (3i) and an ester-functionalized alkyl group (3j), on the other hand, had modest yields of about 40%. More sensitive functional groups were not tolerated. Alkyl bromides were then tested as substrates, but lower yields were obtained compared with alkyl iodides (e.g., for 3a, 3c, 3d, 3f, and 3k). In some cases (3f, 3k), a difference in stereoselectivity was observed using either alkyl iodides or bromides as the substrates. This difference is likely due to experimental uncertainty as a low yield and small amount of products were obtained using alkyl bromides. Cyclohexyl chloride was also tested, but no coupling was found.

When the same conditions were applied for the coupling of 1 with primary alkyl iodides, very low yields were obtained. Upon further optimization, complex 4 was found to be a better catalyst than Fe(OTf)₂ (Figure 3). A new set of conditions was also identified: 1 equiv of 1, 3 equiv of 2a, 5 mol % of 4, 3 equiv of Zn metal, and 2 mol % of I₂ activator, DMA as

**Figure 2.** Scope of the coupling of 1 with tertiary and secondary alkyl halides. The Z/E ratio was determined by ¹H NMR after purification.

solvent, room temperature (rt). The scope of this coupling was, however, limited. While for addition of *n*-octyl (3l) and dec-1-ene (3m) groups the yields were good, the addition of other alkyl groups (3n–r) had yields in the range of 20–35%. The higher reactivity of primary alkyl radicals compared to secondary and tertiary radicals might be the origin of the decreased efficiency of the reactions with primary alkyl iodides.

The Z-vinyl boronate products obtained above could be converted to Z-alkenes bearing various functional groups by transformation of the Bpin group. For example, Pd-catalyzed Suzuki coupling of 3e with ethyl 2-bromobenzoate gave 5 in a 60% yield, bromination of 3b with CuBr₂ gave 6 in an 85% yield, and Cu-catalyzed azidation of 3b with NaN₃ gave 7 in an 86% yield (Scheme 1).

The Fe-catalyzed Z-alkene synthesis was then applied for the formal total synthesis of (Z)-7-chloro-6-*l*-[5-(2-cyclohexylvinyl)pyridin-2-ylmethyl]amino}-2,3,4,5-tetrahydro-1*H*-benzo[*d*]azepine (8, Scheme 2). This compound was reported as a 5-HT_{2c} receptor agonist, which might lead to treatment of various neurological disorders.¹⁹ The key intermediate to 8 was a Z-alkene 9, which could be produced by Pd-catalyzed heterogeneous hydrogenation of 1,2-disubstituted alkyne 10, but in a low yield (Scheme 2a). The latter was obtained by Pd-catalyzed Sonogashira coupling of ethynylcyclohexane with the corresponding pyridinyl bromide 11. In our synthesis, the nitrile group in the commercially available reagent 12 was reduced by NaBH₄ in the presence of





Figure 3. Scope studies for primary aliphatic halides. The *Z/E* ratio was determined by ¹H NMR after purification.


Scheme 1. Further Transformations of *Z*-Vinyl Boronates



NiCl₂·6H₂O (10 mol %) followed by protection with BOC₂O to give 13 in a yield of 57% (Scheme 2b). Pd-catalyzed Suzuki coupling of 13 with 3e, assembled by the Fe-catalyzed coupling of 1 with cyclohexyl iodide, gave the alkylated *Z*-olefin 14 in a yield of 72% (Scheme 2b). Removal of the Boc group by TFA (TFA = trifluoroacetic acid) gave the key intermediate 9 in a quantitative yield (Scheme 2b). In the previous route, a terminal alkyl alkyne was needed, while in the present route an alkyl iodide is used. For many of the secondary alkyl groups shown in Figure 2, alkyl iodides are easily accessible while the corresponding alkyl alkynes are not. Thus, our synthetic route can be advantageous for obtaining certain alkylated derivatives of 8. Additionally, the stereoselective step in our synthesis has a higher yield than that in the previous route.

In summary, Fe-catalyzed reductive coupling of ethynylboronic acid pinacol ester with alkyl halides has been developed

Scheme 2. Total Synthesis of (*Z*)-7-Chloro-6-[[5-(2-cyclohexylvinyl)pyridin-2-ylmethyl]amino]-2,3,4,5-tetrahydro-1*H*-benzo[*d*]azepine: (a) Previous Route (ref 17); (b) This Work



for the stereoselective synthesis of *Z*-vinyl boronates. A simple salt, Fe(OTf)₂, could be used for the coupling of secondary and tertiary alkyl halides, particularly alkyl iodides, with good yields and high *Z*-selectivity. The coupling of primary alkyl halides has a limited scope, even when using a defined Fe complex (4) as catalyst. The *Z*-vinyl boronates could be converted to other *Z*-alkenes by transformation of the Bpin group. The method was applied for a formal total synthesis of a 5-HT_{2c} receptor agonist (8).

■ ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/acs.orglett.8b03772.

Experimental procedures and ¹H and ¹³C NMR spectra (PDF)

■ AUTHOR INFORMATION

Corresponding Author

*E-mail: xile.hu@epfl.ch.

ORCID

Xile Hu: 0000-0001-8335-1196

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

This work is supported by the Swiss National Science Foundation.

■ REFERENCES

- (1) Williams, J. M. J. *Preparation of Alkenes: A Practical Approach*; Oxford University Press, 1996.
- (2) Siau, W.-Y.; Zhang, Y.; Zhao, Y. Stereoselective Synthesis of Z-Alkenes. *Top. Curr. Chem.* **2012**, *327*, 33–58.
- (3) Hall, D. G. Structure, Properties, and Preparation of Boronic Acid Derivatives. In *Boronic Acids*; Hall, D. G., Ed.; Wiley-VCH Verlag, 2011; pp 1–133, DOI: 10.1002/9783527639328.ch1.
- (4) Rawat, V. S.; Sreedhar, B. Iron-Catalyzed Borylation Reactions of Alkynes: An Efficient Synthesis of E-Vinyl Boronates. *Synlett* **2014**, *25* (08), 1132–1136.
- (5) PraveenGanesh, N.; d'Hond, S.; Chavant, P. Y. Methylpentane-diolborane: Easy Access to New Air- and Chromatography-Stable, Highly Functionalized Vinylboronates. *J. Org. Chem.* **2007**, *72* (12), 4510–4514.
- (6) Ho, H. E.; Asao, N.; Yamamoto, Y.; Jin, T. Carboxylic Acid-Catalyzed Highly Efficient and Selective Hydroboration of Alkynes with Pinacolborane. *Org. Lett.* **2014**, *16* (17), 4670–4673.
- (7) Tucker, C. E.; Davidson, J.; Knochel, P. Mild and Stereoselective Hydroborations of Functionalized Alkynes and Alkenes Using Pinacolborane. *J. Org. Chem.* **1992**, *57* (12), 3482–3485.
- (8) Pereira, S.; Srebnik, M. A Study of Hydroboration of Alkenes and Alkynes with Pinacolborane Catalyzed by Transition Metals. *Tetrahedron Lett.* **1996**, *37* (19), 3283–3286.
- (9) Shirakawa, K.; Arase, A.; Hoshi, M. Preparation of (E)-1-Alkenylboronic Acid Pinacol Esters via Transfer of Alkenyl Group from Boron to Boron. *Synthesis* **2004**, *2004* (11), 1814–1820.
- (10) Ben-Daat, H.; Rock, C. L.; Flores, M.; Groy, T. L.; Bowman, A. C.; Trovitch, R. J. Hydroboration of Alkynes and Nitriles Using an α -Diimine Cobalt Hydride Catalyst. *Chem. Commun.* **2017**, *53* (53), 7333–7336.
- (11) Obligacion, J. V.; Chirik, P. J. Earth-Abundant Transition Metal Catalysts for Alkene Hydrosilylation and Hydroboration. *Nature Reviews Chemistry* **2018**, *2* (5), 15.
- (12) Gunanathan, C.; Hölscher, M.; Pan, F.; Leitner, W. Ruthenium Catalyzed Hydroboration of Terminal Alkynes to Z-Vinylboronates. *J. Am. Chem. Soc.* **2012**, *134* (35), 14349–14352.
- (13) Ohmura, T.; Yamamoto, Y.; Miyaura, N. Rhodium- or Iridium-Catalyzed Trans-Hydroboration of Terminal Alkynes, Giving (Z)-1-Alkenylboron Compounds. *J. Am. Chem. Soc.* **2000**, *122* (20), 4990–4991.
- (14) Gorgas, N.; Alves, L. G.; Stöger, B.; Martins, A. M.; Veiros, L. F.; Kirchner, K. Stable, Yet Highly Reactive Nonclassical Iron(II) Polyhydride Pincer Complexes: Z-Selective Dimerization and Hydroboration of Terminal Alkynes. *J. Am. Chem. Soc.* **2017**, *139* (24), 8130–8133.
- (15) Obligacion, J. V.; Neely, J. M.; Yazdani, A. N.; Pappas, I.; Chirik, P. J. Cobalt Catalyzed Z-Selective Hydroboration of Terminal Alkynes and Elucidation of the Origin of Selectivity. *J. Am. Chem. Soc.* **2015**, *137* (18), 5855–5858.
- (16) Cheung, C. W.; Zhurkin, F. E.; Hu, X. Z-Selective Olefin Synthesis via Iron-Catalyzed Reductive Coupling of Alkyl Halides with Terminal Arylalkynes. *J. Am. Chem. Soc.* **2015**, *137* (15), 4932–4935.
- (17) Cheung, C. W.; Hu, X. Stereoselective Synthesis of Trisubstituted Alkenes through Sequential Iron-Catalyzed Reductive Anti-Carbozincation of Terminal Alkynes and Base-Metal-Catalyzed Negishi Cross-Coupling. *Chem. - Eur. J.* **2015**, *21* (50), 18439–18444.
- (18) Zhurkin, F. E.; Hu, X. γ -Selective Allylation of (E)-Alkenylzinc Iodides Prepared by Reductive Coupling of Arylacetylenes with Alkyl Iodides. *J. Org. Chem.* **2016**, *81* (13), 5795–5802.
- (19) Briner, K.; Adeva, B.; Cases-Thomas, M.; Galka, C.; Marcos, L.; Martinez-Grau, M.; Mazanetz, M.; O'toole, J.; Rathmell, R.; Reinhard,

M.; et al. 6-Arylalkylamino- 2,3,4,5-Tetrahydro-1H-Benzo[d]Azepines as 5-Ht2c Receptor Agonists. WO2007028083 (A2), March 8, 2007.