MESOIONIC 5-ALKYL-1,3-DITHIOLIUM-4-THIOLATES: SYNTHESIS AND BRINE SHRIMP TOXICITY

Paulo Afonso de Almeida, Tânia Maria Sarmento da Silva and Aurea Echevarria* Departamento de Quimica, ICE, Universidade Federal Rural do Rio de Janeiro, 23851-970, Seropedica (RJ) Brazil

Abstract. A series of twelve 1,3-dithiolium-4-thiolate mesoionic compounds were synthesized and characterized. The synthetical approach starting from α -bromoalkancic acids to obtain the corresponding 2-N-morpholino-dithiocarbamoyl-carboxylic acids that by on-pot reaction with carbon disulfide and acetic anhydride in triethylamine formed not isolate intermediates, 1,3-dithiolium-4-olates. After, the 2-N-morpholino-5-alkyl-1,3-dithiolium-4-thiolates were obtained by retro 1,3-dipolar addition reactions. The alkyl moiety linked to C-5 of heterocyclic ring permitted the increase of the hydrophobic character and this effect was evaluated on *Artemia salina* lethality. The results indicated a bell-shaped relationship between the number of carbon of side chain in mesoionic derivatives and LD₅₀ in brine shrimp toxicity assays.

Introduction

Mesoionic compounds have received much more attention and have been extensively studied because of their special structure, reaction behaviour, and wide-ranging biological activities (1-5). Recently, these compounds have been used in a broad range of new material applications including nonlinear optical effects (6), analytical reagents (7) (e. g. Nitron^{*}), and photographic materials (8).

The mesoionic compounds can be considered as belonging to a class of mesomeric heterocyclic betaines (9). These betaines are planar five-member heterocycles with at least one side-chain whose α -atom is also in that plane and with dipole moments of 5 D's order. In addition, they also have eight π -electrons in association with both five atoms comprising the ring and the side chain α -atom, and cannot be represented satisfactorily by any one covalent or polar structure, these compounds are called mesoionic (6,10,11). The general formula 1 where a - f represent suitably substituted by carbons or heteroatoms.



Previously we have studied several mesoionic systems such as 1,3,4-tiadiazolium-2-aminide (5,12), 1,3,4-triazolium-2thiolate (13), and 1,2,3-oxadiazolium-2-olate (14,15), and these studies were very useful in structural elucidation and antitumour evaluation. In this work a special attention for 1,3-dithiolium-4-thiolate class (2), particularly our goal is to introduce a lipophylic moiety to afford a new mesoionic series with significant hydrophobic character with ceramide-like analogy. Ceramide are important metabolites, with polar and apolar moleties, related to cellular signalization mechanisms (16). Thus, in this paper we report the synthesis of twelve 2-N-morpholino-5-alkyl-1,3-dithiolium-4-thiolates (2a - I), where the alkyl group (CH_2)_n-CH₃ with n = 1 - 8, 10, 12, 14 and 16 to permit the increase of hydrophobic character. In order to verify the preliminary effects on lipophylic increment we used the brine shrimp (*Artemia salina* Leach) bioassay (17). This bioassay determines the lethality of compounds toward brine shrimp larvae and doing predicts about the ability to kill cancer cell culture, and also a wide range of other pharmacological effects (17,18).

Experimental

Melting points uncorrected were determined on Kofler hot stage. ¹H and ¹³C NMR spectra were recorded on a Brucker AC-200 (200 MHz and 50.3 MHz, respectively). The ¹H and ¹³C chemical shifts (δ) in CDCl₃ are given in ppm relative to tetramethylsilane as internal standard, and J values are given in Hz. Infrared spectra were recorded in KBr pellets, using a Perkin-Elmer 1420 spectrophotometer. U.V. spectra (nm) were taken on Hitachi U2000 spectrometer in methanol. MS spectra were obtained on a Shimadzu QP5050A. Yields are based on isolated products with sufficient purity.

General procedure for the preparation of α -bromo-alkanoic acids (3a – I): In a bottom of 150 mL attached at condenser with gas apparatus containing 65 mmol of alkanoic acid, 68 mmol of bromine and 22 mmol of phosphorus trichloride was heating at 80°C by seven hours and at 100°C by one hour more. When used alkanoic acids with 10 or more carbons atoms were putted additional 5.0 mL of water and at final reaction the products were extracted with chloroform. ¹H NMR identified all compounds.

General procedure for the preparation of 2-N-morpholino-dithiocarbamoyl-alkanoic acids (5a - I): 25.1 mmol of Et₃N was added at 0°C with stirring to a solution of 23 mmol of morpholine in 10 mL of anhydrous benzene. After 10 minutes, 23.2 mmol of CS₂ was added to afford the intermediate **4** (not isolated), and was put on 22.9 mmol of α -bromo-alkanoic acids (3a - I) in 20 mL of anhydrous benzene. The mixture was maintained at room temperature and stirred for 16 hours. The solvent was removed under reduced pressure, washed in chloroform with three successive portions of 5N HCl, and distilled water until pH 5. The organic phase was drying over anhydrous sodium sulphate and removed under reduced pressure. The products were obtained as a clear brown wax and characterized by ¹H NMR, ¹³C NMR and IR.

2-*N*-morpholino-dithiocarbamoyl-propanoic acid (**5a**): (75% yield), oil. ¹H NMR δ 4.77 (1H, q, *J* = 7.49 Hz, H4), 440-3.80 (4H, m, CH₂N), 3.73 (4H, t, *J* = 4.66 Hz, CH₂O), 1.59 (d, *J* = 7.38 Hz, CH₃). ¹³C NMR δ 195.01 (C-2), 176.12 (CO₂H), 65.96 (CH₂O), 50.92 (CH₂N), 48.22 (C-4), 16.66 (CH₃).

2-*N*-morpholino-dithiocarbamoyl-butanoic acid (**5b**): (90% yield), oil. ¹H NMR δ 4.77 (1H, q, *J* = 7.49 Hz, H4), 4.40-3.90 (4H, m, CH₂N), 3.75 (4H, t, *J* = 4.71 Hz, CH₂O), 2.10-1.87 (2H, m, H-5), 1.05 (t, *J* = 7.36 Hz, CH₃). ¹³C NMR δ 194.98 (C-2), 175.81 (CO₂H), 65.82 (CH₂O), 55.04 (C-4), 50.74 (CH₂N), 24.39 (C-5), 17.01 (CH₃).

2-N-morpholino-dithiocarbamoyl-pentanoic acid (5c): (85% yield), oil. ¹H NMR δ 4.77 (1H, q, *J* = 7.49 Hz, H4), 4.40-3.85 (4H, m, CH₂N), 3.75 (4H, t, *J* = 4.44 Hz, CH₂O), 2.03-1.84 (2H, m, H-5), 1.48 (2H, m, H-6), 0.93 (t, *J* = 7.27 Hz, CH₃). ¹³C NMR δ 195.43 (C-2), 176.00 (CO₂H), 66.05 (CH₂O), 53.47 (C-4), 51.01 (CH₂N), 32.94 (C-5), 20.38 (C-6), 13.95 (CH₃).

2-*N*-morpholino-dithiocarbamoyl-hexanoic acid (**5d**): (93% yield), oil. ¹H NMR δ 7.51 (1H, sl, OH), 4.73 (1H, q, *J*=7.09 Hz, H4), 4.40-3.85 (4H, m, CH₂N), 3.74 (4H, t, *J*=4.65 Hz, CH₂O), 2.03-1.85 (2H, m, H-5), 1.42-1.26 (4H, m, H6 and H-7), 0.87 (t, *J*=7.00 Hz, CH₃). ¹³C NMR δ 195.28 (C-2), 175.85 (CO₂H), 65.96 (CH₂O), 53.74 (C-4), 50.89 (CH₂N), 30.69 (C-5), 29.09 (C-6), 22.12 (C-7), 13.62 (CH₃).

2-*N*-morpholino-dithiocarbamoyl-heptanoic acid (**5e**): (77% yield), oil. ¹H NMR δ 7.64 (1H, sl, OH), 4.74 (1H, q, *J*=7.14 Hz, H4), 4.40 - 3.90 (4H, m, CH₂N), 3.74 (4H, t, *J* = 4.82 Hz, CH₂O), 2.03-1.85 (2H, m, H-5), 1.30 (2H, s, H-6), 1.27 (4H, s, H-7 to H-8), 0.85 (t, *J* = 7.00 Hz, CH₃). ¹³C NMR δ 195.01 (C-2), 175.57 (CO₂H), 65.75 (CH₂O), 53.71 (C-4), 50.62 (CH₂N), 30.94 (C-5 and C-7), 26.45 (C-6), 21.93 (C-8), 13.56 (CH₃).

2-*N*-morpholino-dithiocarbamoyl-octanoic acid (**5f**): (97% yield), oil. ¹H NMR δ 7.76 (1H, sl, OH), 4.74 (1H, t, *J* = 7.14 Hz, H4), 4.40-3.85 (4H, m, CH₂N), 3.77 (4H, t, *J* = 3.95 Hz, CH₂O), 2.08-1.82 (2H, m, H-5), 1.45-125 (8H, m, H-6 to H-9), 0.84 (t, *J* = 5.72 Hz, CH₃). ¹³C NMR δ 195.25 (C-2), 176.36 (CO₂H), 65.99 (CH₂O), 53.83 (C-4), 50.81 (CH₂N), 31.30 (C-8), 30.97 (C-5) 28.81 (C-7), 26.98 (C-6), 22.36 (C-9), 13.90 (CH₃).

2-N-morpholino-dithiocarbamoyl-nonanoic acid (**5g**): (86% yield), oil. ¹H NMR δ 7.92 (1H, sl, OH), 4.75 (1H, t, J = 6.68 Hz, H4), 4.40-3.85 (4H, m, CH₂N), 3.76 (4H, s, CH₂O), 2.04-1.86 (2H, m, H-5), 1.43-1 24 (10H, m, H6 to H-10), 0.85 (t, J = 6.56 Hz, CH₃). ¹³C NMR δ 195.28 (C-2), 176.27 (CO₂H), 66.02 (CH₂O), 53.83 (C-4), 50.95 (CH₂N), 31.51 (C-9), 30.94 (C-5), 28.97 (C-8), 28.78 (C-7), 27.03 (C-6), 22.42 (C-10). 13.93 (CH₃).

2-N-morpholino-dithiocarbamoyl-decanoic acid (**5h**): (100% yield), oil. ¹H NMR δ 8.44 (1H, s, OH), 4.73 (1H, t, J = 7.05 Hz, H4), 4.40-3.85 (4H, m, CH₂N), 3.73 (4H, t, J = 3.88 Hz, CH₂O), 2.01-1.77 (2H, m, H-5), 1.40-121 (12H, m, H-6 to H-11), 0.83 (t, J = 5.86 Hz, CH₃). ¹³C NMR δ 195.13 (C-2), 175.80 (CO₂H), 65.99 (CH₂O), 53.83 (C-4), 50.92 (CH₂N), 31.60 (C-10), 30.97 (C-5), 29.06 (C-7, C-8, and C-9), 27.00 (C-6), 22.45 (C-11), 13.93 (CH₃).

2-*N*-morpholino-dithiocarbamoyl-dodecanoic acid (**5i**): (93% yield), oil. ¹H NMR δ 7.10 (1H, sl, OH), 4.74 (1H, t, *J* = 7.14 Hz, H4), 4.40-3.85 (4H, m, CH₂N), 3.76 (4H, t, *J* = 4.74 Hz, CH₂O), 2.05-1.78 (2H, m, H-5), 1.43-120 (16H, m, H-6 to H-13), 0.87 (t, *J* = 7.00 Hz, CH₃). ¹³C NMR δ 195.49 (C-2), 175.63 (CO₂H), 66.05 (CH₂O), 53.80 (C-4), 51.04 (CH₂N), 31.30 (C-12), 30.94 (C-5), 29.18 - 30.94 (C-7 to C-11), 27.09 (C-6), 22.54 (C-13), 13.99 (CH₃).

2-*N*-morpholino-dithiocarbamoyl-tetradecanoic acid (**5**): (98% yield), oll. ¹H NMR δ 6.30 (1H, sl, OH), 4.74 (1H, t, *J* = 7.15 Hz, H4), 4.40-3.85 (4H, m, CH₂N), 3.76 (4H, t, *J* = 4.70 Hz, CH₂O), 2.06-1.79 (2H, m, H-5), 1.43-1.30 (20H, m, H-6 to H-15), 0.85 (t, *J* = 6.45 Hz, CH₃). ¹³C NMR δ 195.28 (C-2), 175.99 (CO₂H), 66.02 (CH₂O), 53.86 (C-4), 50.92 (CH₂N), 31.73 (C-14), 31.00 (C-5), 29.18 (C-7 to C-13), 27.06 (C-6), 22.51 (C-15), 13.96 (CH₃).

2-N-morpholino-dithiocarbamoyl-hexadecanoic acid (**5**k): (66% yield), oil. ¹H NMR δ 6.10 (1H, sl, OH). 4.73 (1H, t, J=7.16 Hz, H4), 4.40-3.85 (4H, m, CH₂N), 3.76 (4H, t, J=4.10 Hz, CH₂O), 2.06-1.79 (2H, m, H-5), 1.45-120 (24H, m, H-6 to H-17), 0.85 (t, J=6.37 Hz, CH₃). ¹³C NMR δ 195.52 (C-2), 174.84 (CO₂H), 66.05 (CH₂O), 53.74 (C-4), 51.25 (CH₂N), 31.79 (C-16), 30.94 (C-5), 29.21 (C-7 to C-15), 27.09 (C-6), 22.54 (C-17), (13.99 CH₃).

2-N-morpholino-dithiocarbamoyl-octadecanoic acid (**5**I): (95% yield), oil. ¹H NMR δ 4.73 (1H, t J = 7.13 Hz, H4), 440-3.85 (4H, m, CH₂N), 3.76 (4H, t J = 4.45 Hz, CH₂O), 1.76-2.10 (2H, m, H-5), 1.22-1.44 (28H, m, H-6 to H-19), 0.85 (t J = 6.37 Hz, CH₃). ¹³C NMR δ 195.52 (C-2), 174.96 (CO₂H), 66.05 (CH₂O), 53.77 (C-4), 51.25 (CH₂N), 31.79 (C-18), 31.00 (C-5), 29.24 (C-7 to C-17), 27.09 (C-6), 22.57 (C-19), 13.99 (CH₃).

General procedure for the preparation of 2-N-morpholino-5-alkyl-1,3-dithiolium-4-thiolates (2a - I): 12.8 mmol of (5a - I) was dissolved in 60 mL of anhydrous benzene at 0°C and was added 215 mmol of CS₂ and 43 mmol of anhydrous Et₃N at same temperature. Then was added 132 mmol of acetic anhydride, when the reaction mixture became carmine-red, and allowed to attain ambient temperature with stirring for 20 hours. The products (2a - I), obtained as a yellow precipitate, were filtered off and purified by recrystallization from acetone.

2-*N*-morpholino-5-methyl-1,3-dithiolium-4-thiolate (2a): 85% yield; M.p. 194–196°C. IR 1538 (C=S⁺ and C=C), 1478 (C=C), 2908, 2870, 1255, 1109, 1033, 883 cm⁻¹. UV λ_{max} (log ε) 407 (4.27), 273 (4.88), and 205 (4.61). MS m/z (%) 233 (M⁺, 70), 86 (100), 130 (90). ¹H NMR δ 3.88 (4H, t, J=5.3 Hz CH₂O), 3.56 (4H, t, J=5.2 Hz CH₂N), 226 (3H, s, CH₃), 1.56 (H, s, SH). 2-*N*-morpholino-5-ethyl-1.3-dithiolium-4-thiolate (2b): 80% yield; M.p. 191–193°C. IR 1546 (C=S⁺ and C=C), 1483 (C=C), 2926, 2865, 1431, 1255, 1109, 1066, 883 cm⁻¹. UV λ_{max} (log ε) 407 (4.05), 273 (4.66), and 205 (4.66). MS *m/z* (%) 247 (M⁺, 65), 130 (100), 86 (95), 60 (20), 73 (12), 44 (28). ¹H NMR δ 3.88 (4H, t, J=4.74 Hz, CH₂O), 3.61 (4H, t, J=3.68 Hz, CH₂N), 2.75 (2H, q, J=7.46 Hz, H-12), 1.69 (H, s, SH); 1.18 (3H, t, J=7.40 Hz, CH₃). ¹³C NMR δ 183.06 (C-2), 117.56 (C-4), 64.58 (CH₂O), 52.03 (CH₂N), 21.68 (C-13), 13.87 (CH₃).

2-N-morpholino-5-propyl-1,3-dithiolium-4-thiolate (2c): 85% yield; M.p. 167–169°C. IR 1535 (C=S⁺ and C=C), 1473 (C=C), 2923, 2869, 1246, 1107, 1033, 887 cm⁻¹. UV λ_{max} (log ε) 409 (4.17), 275 (4.83) and 210 (4.61). MS m/z 261 (M⁺, 50), 130 (100), 86 (90). ¹H NMR δ 3.88 (4H,t J=5.01 Hz CH₂O), 3.60 (4H,t J=5.12 Hz CH₂N), 2.68 (2H,t J=7.68 Hz H-13), 1.69 (H,s, SH); 1.58 (2H, m, J=7.34 Hz, H-14), 0.96 (3H,t J=723 Hz CH₃). ¹³C NMR δ 184.51 (C-2), 150.88 (C-5), 119.51 (C-4), 64.92 (CH₂O), 51.71 (CH₂N), 30.64 (C-13), 22.71 (C-14), 13.62 (CH₃).

2-*N*-morpholino-5-butyl-1,3-dithiolium-4-thiolate (2d): 65% yield; M.p. 154–156°C. IR 1548 (C=S⁺ and C=C), 1473 (C=C), 2928, 2862,1249,1105,1023, 882 cm⁻¹. UV λ_{max} (log ε) 407 (4.27), 273 (4.88) and 205 (4.61). MS m/z 275 (M⁺, 40), 130 (100), 86 (90), 76 (30). ¹H NMR δ 3.88 (4H, t, *J* = 5.00 Hz, CH₂O), 3.61 (4H, t, *J* = 3.70 Hz, CH₂N), 2.72 (2H, t, *J* = 7.55 Hz, H-13), 1.65 (H, s, SH); 157 (2H, dd, *J* = 1426, 727 Hz, H-14), 144 (2H,dd, *J* = 1426, 7.13 Hz, H-15), 0.90 (3H, t, *J* = 7.13 Hz, CH₃). ¹³C NMR δ 184.42 (C-2), 150.50 (C-5), 119.67 (C-4), 64.92 (CH₂O), 51.72 (CH₂N), 30.64 (C-13), 28.44, (C-14), 22.12 (C-15), 13.70 (CH₃).

2-*N*-morpholino-5-pentyl-1,3-dithiolium-4-thiolate (**2e**): 60% yield; M.p. 147–149°C. IR 1558 (C=S⁺ and C=C), 1473 (C=C), 2926. 2859, 1253, 1111, 1028, 885 cm⁻¹. UV λ_{max} (log ϵ) 407 (4.27), 273 (4.88) and 205 (4.61). MS *m/z* (%) 289 (M⁺, 31), 246 (10), 130 (100), 86(80), 76 (60), 57 (20), 44(21). ¹H NMR δ 387 (4H,t J=5.11 Hz CH₂O), 3.58 (4H,t J=5.11 Hz CH₂N), 269 (2H, t J=7.75 Hz H-13), 1.69 (H,s, SH), 1.56 (2H,t J=727 Hz H-14), 1.30 (4H,t J=3.63 Hz H-15 and H-16), 0.88 (3H,t J=6.94 Hz CH₃), ¹³C NMR δ 183.83 (C-2), 150.08 (C-5), 119.03 (C-4), 64.26 (CH₂O), 51.07 (CH₂N), 30.51 (C-13), 28.47 (C-14), 28.08 (C-15), 21.62 (C-16), 13.16 (CH₃).

2-N-morpholino-5-hexyl-1,3-dithiolium-4-thiolate (2f): 51% yield; M.p. 151–153°C. IR 1550 (C=S^{*} and C=C), 1483 (C=C), 2929, 2861, 1257, 1114, 1030, 884 (CH₂) cm⁻¹. UV λ_{max} (log ϵ) 408 (4.00), 273 (4.60) and 209 (4.38). MS m/z 289 (M^{*},

60), 130 (100), 86 (90), 233 (20). ¹H NMR 3.88 (4H, t, J = 4.96 Hz, CH₂O), 3.59 (4 H, t, J = 5.08 Hz, CH₂N), 2.72 (2H, t, J = 7.68 Hz, H-13), 1.65 (H, s, SH), 1.57(2H, s, H-14), 1.40 (6H, s, H-15 to H-17), 0.85 (3H, s, CH₃). ¹³C NMR δ 184.43 (C-2), 150.44 (C-5), 119.74 (C-4), 64.58 (CH₂O), 52.03 (CH₂N), 31.31 (C-13), 29.36 (C-14), 28.72 (C-15,16), 22.26 (C-17), ⁻¹3.81 (CH₃).

2-*N*-morpholino-5-heptyl-1,3-dithiolium-4-thiolate (**2g**): 45% yield; M.p. 151–153°C. IR 1547 (C=S⁺ and C=C), 1481 (C=C), 2926, 2859, 1261, 1114, 1030, 886 cm⁻¹. UV λ_{max} (log ε) 407 (3.42), 272 (4.11) and 214 (3.82). MS m/z 319 (M⁺, 20), 130 (100), 86 (90). ¹H NMR δ 3.88 (4H, t J = 5.00 Hz CH₂O), 3.60 (4H, t J = 4.93 Hz CH₂N), 2.71 (2H, t J = 7.60 Hz H-13), 1.59 (H, s, SH), 1.56(2H, t, J = 6.95 Hz, H-14), 124 (8H, sl, H-15 to H-18), 0.85 (3H, t, J = 5.88 Hz, CH₃). ¹³C NMR δ 184.39 (C-2), 150.22 (C-5), 119.75 (C-4), 64.81 (CH₂O), 51.65 (CH₂N), 31.42 (C-13), 29.38 (C-14), 28.93 (C-15), 28.78 (C-16), 28.70 (C-17), 22.30 (C-18), 13.81 (CH₃).

2-*N*-morpholino-5-octyl-1,3-dithiolium-4-thiolate (**2h**): 62% yield; M.p. 151–153°C. IR 1546 (C=S⁺ and C=C), 1476 (C=C), 2925, 2853, 1255, 1115, 1030, 884 cm⁻¹. UV λ_{max} (log ϵ) 409 (4.05), 274 (4.63) and 210 (4.43). MS *m/z* (%) 331 (M⁺, 17), 233 (10), 130 (55), 97 (10), 86 (40), 76 (30), 57 (28), 44 (100). ¹H NMR δ 3.90 (4H, t *J* = 5.00 Hz, CH₂O), 3.59 (4H, t *J* = 5.00 Hz, CH₂N), 2.71 (2H, t *J* = 7.55 Hz, H-13), 1.66 (H, s, SH), 1.57 (2H, t *J* = 7.22 Hz, H-14), 124 (10H, sl, H-15 to H-19), 0.84 (3H, t *J* = 6.28 Hz, CH₃). ¹³C NMR δ 184.28 (C-2), 150.13 (C-5), 119.60 (C-4), 64.77 (CH₂O), 51.63 (CH₂N), 31.46 (C-13), 29.33 (C-14), 29.03 (C-15), 28.94 (C-16), 28.85 (C-17), 28.66 (C-18), 22.27 (C-19), 13.77 (CH₃).

2-*N*-morpholino-5-decyl-1,3-dithiolium-4-thiolate (2i): 39% yield; M.p. 147–149°C. IR 1558 (C=S⁺ and C=C), 1483 (C=C), 2922, 2854, 1252, 1113, 1030, 885 cm⁻¹. UV λ_{max} (log ϵ) 408 (3.49), 273 (4.11) and 230 (3.74). MS *m/z* (%) 359 (M⁺, 20), 233 (15), 130 (55), 86 (40), 76 (30), 44 (100). ¹H NMR δ 3.88 (4H, t *J* = 5.03 Hz, CH₂O), 3.59 (4 H, t *J* = 5.02 Hz, CH₂N), 2.72 (2H, t *J* = 7.54 Hz, H-13), 1.62 (H, s, SH), 1.59 (2H, t *J* = 7.22 Hz, H-14), 123 (12H, sl, H-15 to H-20), 0.84 (3H, t *J* = 6.28 Hz, CH₃). ¹³C NMR δ 184.43 (C-2), 150.11 (C-5), 119.85 (C-4), 64.87 (CH₂O), 51.73 (CH₂N), 31.59 (C-13), 29.42-28.75 (C-14 to C-19), 22.38 (C-20), 13.87 (CH₃).

2-*N*-morpholino-5-dodecyl-1,3-dithiolium-4-thiolate (2j): 35% yield; M.p. 139–141°C. IR, 1539 (C=S⁺ and C=C), 1470, 2919, 28501252, 1116, 1035, 872 cm. UV λ_{max} (log ϵ) 407 (3.98), 275 (4.52) and 213 (4.34). MS m/z 387 (M⁺, 10), 312 (50), 130 (30), 86 (40), 76 (100). ¹H NMR δ 3.88 (4H,t,*J*=500 Hz, CH₂O), 3.60 (4H,t,*J*=5.02 Hz, CH₂N), 2.69 (2H,t,*J*=7.68 Hz, H-13), 1.55 (2H, sl, H-14), 122 (18H, sl, H-15 to H-22), 0.84 (3H, t, *J* = 6.28 Hz, CH₃). ¹³C NMR δ 184.44 (C-2), 150.67 (C-5), 119.65 (C-4), 64.17 (CH₂O), 51.72 (CH₂N), 31.67 (C-13), 29.42-28.82 (C-14 to C-21), 22.44 (C-22), 13.90 (CH₃).

2-*N*-morpholino-5-tetradecyl-1,3-dithiolium-4-thiolate (**2k**): 62% yield; M.p.147-149°C. IR, 1554 (C=S⁺ and C=C), 1480 (C=C), 2921, 28521254, 1115, 1033, 885 cm¹. UV λ_{max} (log ε) 407 (4.16), 271 (4.71) and 212 (4.41). MS m/z 415 (M+, 10), 382 (10), 233 (10), 130 (100), 86 (80). ¹H NMR δ 3.88 (4H, t, *J* = 5.06 Hz, CH₂O), 3.57 (4H, t, *J* = 5.06 Hz, CH₂N), 2.73 (2H, t, *J* = 7.73 Hz, H-13).157 (2H, s, H-14), 122 (22H, s, H-15to H-24), 0.84 (3H, t, *J* = 6.28 Hz, CH₃). ¹³C NMR δ 184.58 (C-2), 150.65 (C-5), 119.85 (C-4), 65.01 (CH₂O), 51.83 (CH₂N), 31.78 (C-13), 29.54-28.92 (C-14 to C-23), 22.55 (C-22), 14.00 (CH₃).

2-N-morpholino-5-hexadecyl-1,3-dithiolium-4-thiolate (2I): 28% yield; M.p. 143-145°C. IR 1553 (C=S⁺ and C=C), 1479 (C=C), 2922, 2853, 1255, 1114, 1032, 884 cm⁻¹. UV λ_{max} (log ϵ) 407 (4.75), 275 (4.70), 207 (4.49). MS *m/z* (%) 443 (M⁺, 5), 247 (10), 130 (30), 86 (42), 76 (55), 44(100). ¹H NMR δ 3.88 (4H, t, J=5.00 Hz, CH₂O), 3.58 (4H, t, J=4.76 Hz, CH₂N), 2.72 (2H, t, J=7.63 Hz, H-13), 1.63 (H, s, SH), 1.59 (2H, s, H-14), 122 (26H, s, H-15 to H-26), 0.85 (3H, t, J=6.50 Hz, CH₂). ¹³C NMR δ 184.58 (C-2), 150.65 (C-5), 119.87 (C-4), 64.97 (CH₂O), 51.76 (CH₂N), 31.74 (C-13), 29.52-28.87 (C-14 to C-25), 22.51 (C-26), 13.96 (CH₃).

Biological assay

The brine shrimp lethality bioassay was performed following the reported procedure (17) with some modifications. The growth medium was prepared with a water solution 38g/L of sea salt and in small divided tank adds shrimp eggs to one cover side. Lamp above other side was putted to attract hatched shrimp through perforations in the dam. After 48h the shrimp are mature as nauplii and prepared to assay. Test compounds were dissolved in three drops of Cremophor®, 2mL DMSO and saline solution to complete 5 ml of total volume. After, appropriate volumes were added to tubes with 5 mL of saline solution containing 10 nauplii to afford final drug concentration of 1000, 100, 10 and 1 µg/mL, in quadruplicate for each concentration. The control samples containing Cremophor® and DMSO, in same conditions, do not cause significant brine shrimp mortality. After 24 h of incubation under light, the number of dead and survivors brine shrimp in each tube was counted. The LD₅₀ were calculated by graphics from drug concentration vs. lethality percentage using a Probit scale adjust. Data analysis was performed with Origin 6.0 software (Microcal Software).

10.00

Results and Discussion

The 2-N-morpholino-5-alkyl-1,3-dithiolium-4-thiolates (2a - I) were prepared by general procedure shown in Scheme 1. To start, the α -bromo-alkanoic acids (3a - I) were prepared by traditional procedure (19) with bromine and phosphorous trichloride. When the fatty acids containing 10 or more carbon atoms were used water addition at the final of reaction to avoid the intermediate α -bromoalkylacetyl bromide, reinforcing the Hell-Volhard-Zeilinski reaction, afforded the α -bromo-alkanoic acids in quantitative yields. The 2-N-morpholino-dithiocarbamoyl-carboxilic acids (5a - I) were obtained in one-pot procedure. The triethylammonium-2-N-morpholino-dithiocarbamate (4) was initially formed by nucleophilic addition of morpholine to carbon disulfide in the presence of triethylamine in benzene. After 10 minutes the appropriate α -bromo-alkanoic acids in benzene were added at 25°C temperature for 16 hours and by substitution nucleophilic reaction afforded the corresponding 5a - I, obtained in 66 to 100% yield as a clear brown wax.



n = 0 (2a), 1 (2b), 2 (2c), 3 (2d), 4 (2e), 5 (2f), 6 (2g), 7 (2h), 9 (2i), 11 (2j), 13 (2k) and 15 (2l)

Scheme 1

The mesoionic compounds (2a - I) were synthesized according to the modified procedure previously described (20) to the analogues with different and single side chain. The intermediates 2-N-morpholino-dithiocarbamoyl-carboxilic acids in anhydrous benzene, triethylamine and carbon disulfide, were added in acetic anhydride at 0°C. When the reactions started, a change of yellow to carmine-red colour was observed, and after 20h stirring the target products were obtained as yellow solids. The 2a - I were purified from acetone afforded orange crystals with yields into 45 to 85%. In the synthesis of 2i - I (with exception of 2k 62%), the increase of length of side-chain linked at C-5 to heterocyclic ring, caused a high hindered in cyclic intermediate (not isolated) affording the target mesoionic compounds by retro-1,3-dipolar addition reaction, eliminating carbon oxysulfide, with lower yield, 39 to 28% (Scheme 2).



Scheme 2

The structures proposed are consistent with the chemical shifts assignments in 1D ¹H and ¹³C NMR (HBBD and PEDANT) and 2D ¹H – ¹³C HMBC (¹H x ¹³C – COSY, ^{2,3}J (CH)). The chemical shifts of carbon atoms of heterocyclic ring in **2a** – I, presented C-2 and C-5 at high frequencies when compared with values of other carbon atoms (see Table I).

Mesoionic Compound	n ^a	LD ₅₀ ^b (µM)		
2a	1	870		
2b	2	372		
2c	3	479		
2d	4	182		
2e	5	309		
2f	6	195		
2g	7	105		
2h	8	12		
2i	10	112		
2j	12	123		
2k	14	195		
21	16	513		

Table I. Effects of 1,3-dithiolium-4-thiolate derivatives on survival of brine shrimp nauplii.

^a Number of carbon atoms in side chain attached at C-5 of heterocyclic ring.

^b Compound concentration required to kill brine shrimp by 50%. All assays were performed in quadruplicate.

We found δ 183.06 – 184.58 for C-2 and δ 150.08 – 150.88 for C-5. Thus, for the **2h** derivative, for example, the spin-spin interactions of H-7 (δ 3.59) with C-2 (δ 184.28, ³*J* (CH)) and C-8 (δ 64.77, ²*J* (CH)); and of H-13 (δ 2.71) with C-5 (δ 150.13, ²*J* (CH)) and C-4 (δ 119.60, ³*J* (CH)), were observed in the HMBC spectrum. The chemical shifts of C-4 atoms showing lower frequencies δ 119.03 – 119.87 due to the greater electron-donating effect caused by the sulphur atom negatively charged directly attached. The other carbon atoms showed the expected chemical shifts.

The brine shrimp nauplii have been used previously in a number of bioassay systems (18) and the compounds are tested at initial concentrations of 10, 100 and 1000 ppm (or µg/ml) in vials containing 5 ml of brine and ten shrimps in each of five replicates. Survivors are counted after 24h. These data are processed to estimate LD₅₀ values with 95%

confidence intervals for statistically significant comparisons of potencies. The literature report a positive correlation between brine shrimp toxicity and 9KB cells cytotoxicity and has been observed that ED_{50} values for general cytotoxicities are about one-tenth LD_{50} values in the brine shrimp test (21).

The Table II show the results obtained in brine shrimp assay. In order to verify a possible relationship between the chemical structure of 1,3-dithiolium derivatives and the biological response, was plotted the $-\log LD_{50}$ values against the number carbon atoms of side chain attached on mesoionic ring. The continuous growth of an alkyl chain or of methylene units increases the hydrophobic part of the molecule.

	C 2	6.4	C 5		C-3	C-4	C-5
	6-2	U-4	C-5		0-2	U-4	C-5
2a	n.o. ^b	n.o. ^b	n.o. ^b	2g	184.39	119.75	150.22
2b	183.06	117.56	n.o. ^B	2h	184.28	119.60	150.13
2c	184.51	119.51	150.88	2i	184.43	119.85	150.11
2d	184.42	119.67	150.50	2j	184.44	119.65	150.67
2e	183.83	119.03	150.08	2k	184.58	119.85	150.65
2f	184.43	119.74	150.44	21	184.58	119.87	150.67

Table II. NMR-chemical shifts for main carbon atoms of mesoionic derivatives.^a

 δ^{*} (ppm) in CDCl₃ solutions at 50.3 MHz. h n.o. = no observed peak

The Graphic 1 indicates a bell shaped curve with serrated variations (zigzag) on left side, where the **2h** (n = 8) has been the most active derivative. Zigzag variations are well known in homologous series for physical properties such as melting points and solubilities. Thus, sometimes is observed alternating variations of biological activities according to whether the number of carbon atoms is even or odd (22-24). Furthermore, the curves with an activity maximum are the most common, and it is presumed that they reflect the existence of an optimal partition coefficient associated with the easiest crossing of biological membranes. Bell-shaped curves are also seen when using isolated cells, for which it can be demonstrated that the receptor is in outside the membrane (25). Another possibility is a lipophylic pocket chains making up part of the bilayer, examples being leukotriene D_4 -agonist/antagonists (26). The study of the activities of some homologous compounds can identify which term is associated with the highest potency, using an easy bioassay.



Graphic 1. Effect of carbon chain length on brine shrimp lethality.

Acknowledgements – The authors thank the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) and Coordenação de Aperfeiçoamento de Pessoal Docente (CAPES) for the bursaries and research grant.

References

- (1) W. D. Ollis, C. A. Ramsden, Adv. Heterocycl. Chem. 19, 1 (1976).
- (2) C. G. Newton, C. A. Ramsden, Tetrahedron 38, 2965 (1982).
- (3) M. H. Osterhout, W. R. Nadler, A. Padwa, Synthesis 123 (1994).
- (4) N. F. Grynberg, A. C. S. Santos, A. Echevarria, Anticancer Drugs 8, 88 (1997).
- (5) S. Cadena, A. Echevarria, E. Carnieri, M. B. Oliveira, FEBS Lett. 440, 46 (1998).
- (6) G. L. C. Moura, A. M. Simas, J. Miller, Chem. Phys Lett. 257, 639 (1996).
- (7) W. Bocian, J. Jazwinski, L. Stefaniak, G. A. Webb, Mag. Reson. Chem. 33, 134 (1995).
- (8) J. C. Deaton, H. R. Luss, J. Chem. Soc., Dalton Trans. 3163 (1999).
- (9) W. D. Ollis, S. P. Stanforth, C. A. Ramsden, Tetrahedron 41, 2239 (1985).
- (10) K. K. Cheung, A. Echevarria, S. E. Galembeck, M. A. M. Maciel, J. Miller, V. M. Rumjanek, A. M. Simas, Acta Crystall, C 49, 1092 (1993).
- (11) K. K. Cheung, A. Echevarria, S. E. Galembeck, M. A. M. Maciel, J. Miller, V. M. Rumjanek, A. M. Simas, Acta Crystall. C 48, 1471 (1992).
- (12) A. C. S. Santos, A. Echevarria, Magn. Reson. Chem. 39, 182 (2001).
- (13) A. Echevarria, J. Miller, J. Chem. Soc. Perkin Trans. 2, 1425 (1989).
- (14) N. Grynberg, A. Echevarria, R. M. Gomes, T. Shinzato, J. Miller, Anticancer Research 12, 1025 (1992).
- (15) V. Rodrigues, A. Echevarria, J. Miller, Heterocycl. Commun. 6, 507 (1996).
- (16) Y. A. Hannun, Science 274 (1996).
- (17) B. N. Meyer, N. R. Ferrigni, J. E. Putnan, D. E. Jacobsen, D. E. Nichols, J. L. Mclaughlin, Planta Medica 45, 31 (1982).
- (18) G. Persoone, ed. Proceeding of the International Symposium on Brine Shrimp, Artemia salina, vols. 1-3, Universal Press, Witteren, Belgium, 1980.
- (19) A. I. Vogel, Vogel's Textbook of Practical Organic Chemistry, 5th. Ed; Logman Group-UK Ltd: London, 1989, pp 720-723.
- (20) M. B. Oliveira, J. Miller, A. B. Pereira, S. E. Galembeck, G. L. C. Moura, A. M. Simas, Phosphorus, Sulfur, and Silicon 108, 75 (1996).
- (21) J. C. McLaughlin, Brine shrimp and crown gall tumours: simple bioassays for the discovery of plant anti-tumour agents. Proceedings, NIH Workshop, Bioassays for Discovery of Anti-tumour and Antiviral Agents from Natural Sources, Bethesda, Oct. 18 – 19, 1988, p.22.
- (22) D. K. Herron, T. Goodson, N. G. Bollinger, D. Swauson-Bean, L. L. Froelich, W. T. Jackson, J. Med. Chem. 35, 1818 (1992).
- (23) C. G. Raison, O. D. Standen, Br. J. Pharmacol. 10, 191 (1995).
- (24) T. A. Fairlay, R. R. Tidwell, L. Donkor, N. A. Naiman, K. A. Ohemeng, R. J. Lombardy, J. A. Bentley, M. Corey, J. Med. Chem. 36, 1746 (1993).
- (25) C. G. Wermuth, "The Practice of Medicinal Chemistry", chapter 12 Molecular Variations in Homologous Series: Vinylogues and Benzologues. Academic Press-Ltd, 1996, pg 181 – 201.
- (26) J. J. Godfroid, C. Broquet, S. Jouquery, J. Med. Chem. 30, 792 (1987).

Received on September 20, 2002