

Organic Preparations and Procedures International: The New Journal for Organic Synthesis

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/uopp20>

DIATOMITE-SUPPORTED PERACETIC ACID OXIDATION OF 1,4-DIHYDROPYRIDINES

Bu-Lan Wu^a & Zhao-Bin Chen^a

^a School of Chemistry and Chemical Engineering, Shanxi University Taiyuan, 030006, PR CHINA Phone: 86351-7011492 Fax: 86351-7011492 E-mail:
Published online: 09 Feb 2009.

To cite this article: Bu-Lan Wu & Zhao-Bin Chen (2005) DIATOMITE-SUPPORTED PERACETIC ACID OXIDATION OF 1,4-DIHYDROPYRIDINES, Organic Preparations and Procedures International: The New Journal for Organic Synthesis, 37:1, 80-83, DOI: [10.1080/00304940509355404](https://doi.org/10.1080/00304940509355404)

To link to this article: <http://dx.doi.org/10.1080/00304940509355404>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

DIATOMITE-SUPPORTED PERACETIC ACID OXIDATION OF 1,4-DIHYDROPYRIDINES


Submitted by
(09/27/04)

Bu-Lan Wu and Zhao-Bin Chen*

*School of Chemistry and Chemical Engineering, Shanxi University
Taiyuan 030006, P. R. CHINA*

E-mail: zchen@sxu.edu.cn; Tel: 86351-7011492; Fax: 86351-7011688

Aspects of the oxidation of 1,4-dihydropyridines (1,4-DHP) continue to be the focus point of research activity as evidenced by the appearance of a large number of publications.¹⁻¹⁷ We investigated diatomite-supported peracetic acid as a new oxidation reagent for 1,4-dihydropyridines (1,4-DHP).



The reaction was carried out by stirring a mixture of 1,4-DHP and diatomite-supported peracetic acid in chloroform at 0-5°C for 5-40 min and the corresponding pyridine derivatives were obtained after simple filtration and further purification by chromatography. The results and reaction conditions are summarized in *Table 1*. The products were characterized by mp., IR and ¹H-NMR.

To check the efficiency of the reagent, the following experiments were performed. When the oxidation of 1,4-DHP was carried out by using peracetic acid without diatomite solid-support, the yield was low with a number of side-products, which is in accordance with the previous observation that reagents and catalysts on solid supports often have modified activity as improved product selectivity.^{18,19} Further studies showed that using diatomite solid-support was better than using silica gel. The effect of different solvents on the oxidation reaction was also studied. When the solvent was ethyl acetate or acetic acid, the reaction times were longer and the yields of the products were lower as compared with chloroform.

In conclusion, a novel and simple oxidation reagent for the conversion of 1,4-DHP to the corresponding pyridines has been presented. The ready availability of the reagent, its ease of manipulation and the mild reaction conditions recommend the diatomite-supported peracetic acid as a useful reagent.

Table 1. Oxidation of 1,4-Dihydropyridines by Diatomite-supported Peracetic Acid^a

Cmpd	Yield (%)	mp. (°C)	lit.mp. (°C)	Temp. (°C)	Time (min)	¹ H NMR (δ)
2a	85	72-73	72 ¹⁷	RT	40	8.67 (s, 1H), 4.34-4.41 (q, 4H, J = 7.07 Hz), 2.84 (s, 6H), 1.38-1.43 (t, 6H, J = 7.08).
2b	62	oil	oil ¹⁶	RT	30	4.34-4.41 (q, 4H, J = 7.09 Hz), 2.49 (s, 6H), 2.24 (s, 3H), 1.34-1.38 (t, 6H, J = 7.11).
2c	49	oil	oil	0-5	35	4.33-4.40 (q, 4H, J = 7.07 Hz), 2.48-2.53 (m, 2H), 2.46 (s, 6H), 1.47-1.59 (m, 2H), 1.32-1.37 (t, 6H, J = 7.08 Hz), 0.86-0.91 (t, 3H, J = 7.35 Hz).
2d	75	63-64	62-64 ¹⁶	0-5	5	7.36-7.37 (br s, 3H), 7.25-7.27 (br s, 2H), 3.97-4.04 (q, 4H, J = 7.10 Hz), 2.62 (s, 6H), 0.88-0.93 (t, 6H, J = 7.10 Hz).
2e	80	49-50	51-53 ¹⁷	0-5	10	7.17-7.19 (b, 2H, J = 8.11 Hz), 6.87-6.90 (b, 2H, J = 8.12 Hz), 4.00-4.08 (q, 4H, J = 7.05 Hz), 3.81 (s, 3H), 2.58 (s, 6H), 0.95-1.00 (t, 6H, J = 7.06 Hz).
2f	88	72-74	68 ¹⁷	0-5	15	7.35-7.38 (b, 2H, J = 7.71 Hz), 7.20-7.22 (b, 2H, J = 7.72 Hz), 4.02-4.08 (q, 4H, J = 6.55 Hz), 2.61 (s, 6H), 0.96-1.01 (t, 6H, J = 6.78 Hz).
2g	53	50-51	63 ¹⁷	0-5	5	8.24-8.27 (m, 1H), 8.18 (m, 1H), 7.55-7.60 (m, 2H), 4.01-4.08 (q, 4H, J = 6.95 Hz), 2.62 (s, 6H), 0.96-1.01 (t, 6H, J = 6.95 Hz).

a) All reactions were carried out in chloroform using a molar ratio of 1:1.1 for DHP:PAA; Diatomite:DHP (0.75 g:1 mmol).

EXPERIMENTAL SECTION

All mps were determined on a Sanyo Gallenkamp MPD350 Melting Point apparatus and are uncorrected. IR spectra were determined on a IR-408 Infrared Spectrometer and ¹H-NMR spectra were recorded at 300 MHz on a R-24B Hitachi NMR Spectrometer in CDCl₃ using TMS as an internal standard. Diatomite may be obtained from Aldrich Chemical Co. and 1,4-dihydropyridines were prepared according to described procedures.¹⁶

General Procedure for the Oxidation of Hantzsch 1,4-DHP.- The reaction must be carried out in an efficient fume hood. In a 50 mL round-bottomed flask, 0.465 g (1.1 mmol) of 18.0%

peracetic acid was added dropwise to 0.75 g of diatomite and the mixture was left to stand for appropriate time (10-15 min) at room temperature. Then the resulting solid diatomite-supported PAA was added in portions over 5 min to a stirred solution of 1,4-dihydropyridine **1d** (0.329 g, 1 mmol) in chloroform (20 mL) at 0-5°C. After completion of the reaction monitored by TLC, the reaction mixture was filtered and the residue was washed with chloroform (2 x 10 mL). The combined filtrates were washed with saturated aq. NaHSO₃ to eliminate residual peracetic acid and then with saturated aq. Na₂CO₃, followed by drying over anhydrous Na₂SO₄. The crude product obtained on removal of the solvent at reduced pressure was then purified by column chromatography over silica gel (eluent: ethyl acetate:petroleum ether 1:3) to give the oxidized product **2d** as a colorless solid (0.245 g, 75%).

The oxidations of the other 1,4-dihydropyridines were carried out using a similar procedure.

Acknowledgement.– We are grateful to the Science Foundation of Taiyuan City of China, Laboratory of Organic Solid, Institute of Chemistry, Chinese Academy of Sciences and the Returned Student Science Foundation of Shanxi Province of China, for financial support.

REFERENCES

1. S. H. Mashraqui and M. A. Karnik, *Tetrahedron Lett.*, **39**, 4895 (1998).
2. M. F. Gordeev, D. V. Patel and E. M. Gordon, *J. Org. Chem.*, **61**, 924 (1996).
3. J. J. Vanden Eynde, R. D'Orazio and Y. van Haverbeke, *Tetrahedron*, **50**, 2479 (1994).
4. J. J. Vanden Eynde, A. Mayence and A. Maquestiau, *Tetrahedron*, **48**, 463 (1992).
5. B. Khadilkar and S. Borkar, *Synth. Commun.*, **28**, 207 (1998).
6. T. Itoh, K. Nagata, Y. Matsuya, M. Miyazaki and A. Ohsawa, *J. Org. Chem.*, **62**, 3582 (1997).
7. J. J. Vanden Eynde, F. Delfosse, A. Mayence and Y. van Haverbeke, *Tetrahedron*, **51**, 6511 (1995).
8. S. H. Mashraqui and M. A. Karnik, *Synthesis*, 713 (1998).
9. N. Nakamichi, Y. Kawashita and M. Hayashi, *Synthesis*, 1015 (2004).
10. S. P. Chavan, S. W. Dantale, U. R. Kalkote, V. S. Jyothirmai and R. K. Kharul, *Synth. Commun.*, **28**, 2789 (1998).
11. A. Maquestiau, A. Mayence and J. J. Vanden Eynde, *Tetrahedron Lett.*, **32**, 3839 (1991).
12. O. Garcia, F. Delgado, A. C. Cano and C. Alvarez, *Tetrahedron Lett.*, **34**, 623 (1993).

13. R. S. Varma and D. Kumar, *Tetrahedron Lett.*, **40**, 21 (1999).
14. I. Mohammadpoor-Baltork, M. M. Sadeghi, H. R. Memarian and R. Pairow, *J. Chem. Research*, **40** (2000).
15. M. A. Zolfigol, K. B. Maryam, M. M. Sadeghi, I. Mohammadpoor-Baltork and H. R. Memarian, *Synth. Commun.*, **30**, 3919 (2000).
16. B. Love, and K. M. Snader, *J. Org. Chem.*, **30**, 1914 (1965).
17. D. P. Cheng and Z. C. Chen, *Synth. Commun.*, **32**, 793 (2002).
18. C. N. Rhodes, M. Franks, G. M. B. Parkes and D. R. Brown, *J. Chem. Soc., Chem. Commun.*, 804 (1991).
19. J. H. Clark, S. R. Cullen, S. J. Barlow and T. W. Bastock, *J. Chem. Soc. Perkin Trans. II*, **6**, 1117 (1994).

OXIDATION OF TOLUENES TO AROMATIC ALDEHYDES WITH MOLECULAR OXYGEN IN SUBCRITICAL WATER

Submitted by Berkant Kayan, Recep Özen, Ahmet M. Gizir* and N. Simsek Kus
(07/22/04)

Department of Chemistry, University of Mersin-33342 Mersin, TURKEY

* e-mail: agizir@mersin.edu.tr

Oxidations are important transformations in organic synthesis.¹ In general, many oxidations are carried out with a large excess of chromium or manganese reagents. The metal residues are environmentally undesirable and often cause problems during work-up.² In addition organic oxidations in the chemical industry have traditionally been performed in either aromatic or halogenated solvents.³ One of the most attractive oxidants is molecular oxygen,⁴ since oxygen is highly soluble in subcritical water. Molecular oxygen has been used as the reagent for the oxidation of alkyl aromatic compounds in the presence of different catalysts, such as MnBr_2 , CoBr_2 ,

