

Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for
authors and subscription information:

<http://www.tandfonline.com/loi/gpss20>

MICROWAVE SYNTHESIS OF TRICHLORFON AND ITS ANALOGUES

Abdollah Javidan^a & A. R. Banan^a

^a Department of Chemistry, Imam Hossein
University, Tehran, I. R. Iran

Published online: 16 Aug 2010.

To cite this article: Abdollah Javidan & A. R. Banan (2004) MICROWAVE SYNTHESIS OF TRICHLORFON AND ITS ANALOGUES, *Phosphorus, Sulfur, and Silicon and the Related Elements*, 179:11, 2253-2258, DOI: [10.1080/10426500490475184](https://doi.org/10.1080/10426500490475184)

To link to this article: <http://dx.doi.org/10.1080/10426500490475184>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

MICROWAVE SYNTHESIS OF TRICHLORFON AND ITS ANALOGUES

Abdollah Javidan and A. R. Banan
Department of Chemistry, Imam Hossein University,
Tehran, I. R. Iran

(Received August 26, 2003; accepted April 8, 2004)

The synthesis of dialkyl-(2,2,2-trichloro-1-Hydroxyethyl) phosphonates using dialkyl hydrogen phosphite and chloral hydrate under solvent-free condition by microwave irradiation is reported. The products were characterized using ^1H NMR, ^{13}C NMR, and ^{31}P NMR spectroscopy.

Keywords: Anticholinestrase; insecticide; microwave; organophosphorus compounds; trichlorfon

INTRODUCTION

Dialkyl-(2,2,2-trichloro-1-hydroxyethyl)phosphonates are important from the point of view of biologic activity, and the agrochemical and medicine industries, for example, trichlorfon (**3a**) or dimethyl-2,2,2-trichloro-1-hydroxyethyl phosphonate is a pesticide.^{1–4} Since this compound is less toxic than DDT, it is a good candidate to replace for DDT. Trichlorfon, which is well known as metrifonate in the medicine industry, is under investigation as an affecting drug for alzheimer's disease and anticholinestrase.⁵

In recent years, there has been growing interest in the application of microwave irradiation in chemical reaction enhancement because of its cleaner reactions, decreased reaction time, and easier workup.^{6–8} In continuation of our ongoing efforts in this area, for the above reasons, and in view of our general interest in microwave synthesis, we wish to report here the synthesis of these organophosphorus compounds using microwave irradiation. This procedure is a general method for


We thank the Basic Science Research Center of I.H.U for financial support of this work. We also thank Dr. A. Hajipour and Dr. A. Moghimi for their helpful discussions.

Address correspondence to Abdollah Javidan, Department of Chemistry, Imam Hossein University, P.O. Box 16575-347, Tehran, I. R. Iran. E-mail: ajavidan@ihu.ac.ir

synthesis of dialkyl-2,2,2-trichloro-1-hydroxyethyl phosphonates **3a–c** by microwave irradiation, which in previous preparations required heating starting materials for several hours in the reflux condition. The data obtained show that the reaction time can be reduced to minutes in good yields.

RESULTS AND DISCUSSION

For our study we worked with neat dialkylphosphites **1** and chloral hydrate **2** using a domestic microwave oven under solvent-free conditions in the presence of alumina or silica gel. The synthetic conventional method for these compounds was the same as previously reported procedures.^{9–12} The reactions are completed within minutes and in good-to-excellent yields (65–93%, Table I). To evaluate the effect of microwave irradiation, we tried several experiments on compounds **3a** (3, 6, 9, and 12 min). We observed that increasing the reaction time had no significant effect on the yields to product. In addition, when we exposed the reaction mixture to microwave irradiation for 30 min the product decomposed.



SCHEME 1

The structure of the products was assigned with ¹H NMR, ¹³C NMR, and ³¹P NMR spectroscopic analysis and comparison with authentic samples.^{13–15} The existence of signal at 18.84 ppm and disappearance of the signal at 11 ppm in the ³¹P-{¹H} NMR spectrum of crude product shows the presence of trichlorfon **3a**. The ¹H NMR spectrum showed two doublets at 3.89 and 3.93 ppm, confirming the presence of two diastereotopic methoxy groups.

The signal for methine proton of trichlorfon appeared as a doublet at 4.54 ppm with *J* = 15. ²*J*_{P–H} = 11.5 Hz. With respect to the relationship between the magnitude of ²*J*_{P–H} and dihedral angle observed in

TABLE I Preparation of **3a–c** Using Microwave Irradiation under Solvent-Free Conditions


Starting phosphite	(R)	Molar ratio of (phosphite/chloral)	Catalyst	Reaction time (min)	Conversion (%) [*]
1a	Methyl	1: 1.01	Silica gel	1.5 ^{**}	3a (60)
1a	Methyl	1: 1.01	Alumina	3 (240)	3a (70)
1a	Methyl	1: 1.01	Alumina	6 (240)	3a (74)
1a	Methyl	1: 1.01	Alumina	9 (240)	3a (71)
1a	Methyl	1: 1.01	Alumina	12 (240)	3a (67)
1b	Ethyl	1: 1.16	Alumina	4 (300)	3b (76)
1c	Isopropyl	1: 1.20	Alumina	3 (420)	3c (65)
1c	Isopropyl	1: 1.40	Alumina	7	3c (82.5)
1c	Isopropyl	1: 2.00	Alumina	8	3c (93)

^{*}Based on the ³¹P NMR of crude reaction mixture.

^{**}450 W.

organophosphorus compounds and the present data, the conformation of this compound has been assigned. The observed $^2J_{P-H} = 11.5$ Hz in trichlorfon **3a** suggests that the preferred conformation is *anti* (*a*) in which the hydrogen group is anti to phosphoryl group ($\theta = 180^\circ$) but the coupling constant of the alternative conformation (*b*) ($\theta = 60^\circ$) is around 24 Hz.¹⁶ Existence of hydrogen bonding between hydrogen and the oxygen of the phosphoryl group has been reported in the literature.¹⁶ The proton of the OH group appears at 5.48 ppm as a broad peak.

Further evidence for determination the structure of the products were obtained from ¹³C NMR analysis. In the ¹³C NMR, the CHOH carbon appeared at 79.98 ppm as a doublet with $^1J(PC) = 662.5$ Hz. This large value of $^1J(PC)$ is consistent with coupling of phosphorus with adjacent carbon.¹⁶ The most downfield shift of CCl₃ carbon at 97.7, which split with phosphorus, appears as a doublet and is in agreement with the phosphorus attached to three electronegative chlorine atoms. The presence of hydroxyl group was indicated by a broad absorption band at 3350 cm⁻¹ in the infrared spectrum.

**FIGURE 1**

CONCLUSION

In this article a straightforward method for formation of dialkyl-(2,2,2-trichloro-1-Hydroxyethyl) phosphonates using dialkyl hydrogen phosphate and chloral hydrate under solvent-free conditions using microwave irradiation has been reported. This method is superior to the reported method from the point of view of green chemistry, yields, and short reaction time.

EXPERIMENTAL

General

^1H , ^{13}C , and ^{31}P NMR spectra were obtained using a Bruker 250 spectrometer using CDCl_3 as solvent and TMS as internal standard or 85% H_3PO_4 as external standard. Melting points were determined on a Mettler 9100 electrothermal and are uncorrected. IR spectra were recorded on a Perkin-Elmer IR-738 spectrometer. All reagents and chemicals were obtained from Merck or Aldrich Chemical Company and were used as received unless otherwise noted. The starting materials, 1a, 1b, and 1c, were prepared as described in the literature.^{17,18} A domestic microwave oven operating at 2450 MHz was used at power 600 W for all the microwave experiments except that.

General Procedure

Method A (Conventional)

Reactions were performed under the same conditions using an oil bath at the reflux temperature as previously reported.⁹ The product was extracted with benzen, the organic phase was dry over MgSO_4 , filtered, and put away until the title compound was isolated as white crystal.

Method B (MW)

General method. In a mortar, dialkyl hydrogen phosphite **1a-c** (46 mmol) and alumina or (silica gel) was ground with a pestle, and 48 mmol of chloral hydrate was added. The mixture was transfer to a Teflon vessel and irradiated in microwave oven for the time specified in Table I. As this study was done only for the comparison, the products were not isolated, and after evaporation of the solvent the resulting mixture was analyzed by ^{31}P NMR.

(3a). m.p. 75–76°C (lit. 76–78°C). ^{31}P NMR (CDCl_3) δ : 18.83. ^1H NMR (CDCl_3) δ : 3.88 (d, $^3J_{\text{P-H}} = 6.5$ Hz, 3H, POCH_3), 3.93 (d, $^3J_{\text{P-H}} = 6.5$ Hz, 3H, POCH_3), 5.45 (d, $^2J_{\text{P-H}} = 11.75$, 1H, CHOH), 5.48 (s, 1H, OH). ^{13}C { ^1H } NMR (CDCl_3) δ 54.22 (d, $^2J_{\text{P-C}} = 27.5$ Hz, POCH_3), 54.80 (d, $^2J_{\text{P-C}} = 27.5$ Hz, POCH), 79.98 (d, $^1J_{\text{P-C}} = 665$ Hz, CHOH), 97.75 (d, $^2J_{\text{P-C}} = 6.8$ Hz, C-Cl). IR (KBr) cm^{-1} : 3350, 3042, 2980, 2940, 1270, 1200 (P=O).

(3b). m.p. 54–55°C (lit. 55–56°C). ^{31}P NMR (CDCl_3) δ : 16.37. ^1H NMR (CDCl_3) δ : 1.38 (t, $^4J_{\text{P-H}} = 7.0$, 6H, POCH_2CH_3), 4.28 (m, 4H, POCH_2CH_3), 4.49 (d, $J = 11.75$, 1H, CHOH), 5.86 (s, 1H, OH). ^{13}C { ^1H } NMR (CDCl_3) δ 16.36 (d, $^3J_{\text{P-C}} = 5$ Hz, POCH_2CH_3), 16.46 (d, $^3J_{\text{P-C}} = 5$ Hz, POCH_2CH), 64.28 (d, $^2J_{\text{P-C}} = 27.5$ Hz, POCH_2CH_3), 64.33 (d, $^2J_{\text{P-C}} = 27.5$ Hz, POCH_2CH_3), 78.99 (d, C(P)OH , $^1J_{\text{P-C}} = 657.5$ Hz), 98.04 (d, 1C, $^2J_{\text{P-C}} = 40$ Hz, C-Cl). IR (KBr) cm^{-1} : 3355, 3100, 3020, 2980, 1280, 1230, 1205 (P=O).

(3c). m.p. 108–109°C (lit. 105–106°C). ^{31}P NMR (CDCl_3) δ : 14.49. ^1H NMR (CDCl_3) δ : 1.39 (d, $^3J_{\text{H-H}} = 7.0$, 8.1, 6H, $\text{CH}(\text{CH}_3)_2$), 1.39 (d, $^3J_{\text{H-H}} = 7.0$, 6H, $\text{CH}(\text{CH}_3)_2$), 4.39 (d, 1H, $J = 11.75$, CHOH), 4.86 (m, 1H, $\text{CH}(\text{CH}_3)_2$), 5.49 (sb, 1H, OH). ^{13}C { ^1H } NMR (CDCl_3) δ : 23.74 (d, $^3J_{\text{P-C}} = 25$ Hz, $\text{CH}(\text{CH}_3)_2$), 24.34 (d, $^3J_{\text{P-C}} = 25$ Hz, $\text{CH}(\text{CH}_3)_2$), 24.44 (d, $^3J_{\text{P-C}} = 25$ Hz, $\text{CH}(\text{CH}_3)_2$), 73.17 (d, $^2J_{\text{P-C}} = 30$ Hz, $\text{CH}(\text{CH}_3)_2$), 73.83 (d, $^2J_{\text{P-C}} = 32.5$ Hz, $\text{CH}(\text{CH}_3)_2$), 80.20 (d, $^1J_{\text{P-C}} = 655$ Hz, C(P)OH), 98.27 (d, $^2J_{\text{P-C}} = 37.5$ Hz, C-Cl). IR (KBr film) cm^{-1} : 3350, 3065, 3010, 2950, 1290, 1270, 1200 (P=O).

REFERENCES

- [1] G. W. Ware, *Pesticides, Theory and Application* (1982), Chap. 5, p. 5.
- [2] Schrader, G. *Final report* No. 1808, Item No 22 (British Intelligence Objectives Committee, London, 1947).
- [3] *Farm Chemicals Handbook* (Meister Pro, U.S. 1999), C 394.
- [4] M. A. Kamrin, Ed., *Pesticide Profiles: Toxicity, Environment and Fate* (CRC Press, New York, 1997), p. 229.
- [5] H. Papp and Z. Rakonczay, *Neurochem. Int.*, **39**, 19 (2001).
- [6] R. S. Varma, *Green Chem.*, **43** (1997).
- [7] L. Perreux and A. Loupy, *Tetrahedron*, **57**, 9199 (2001).
- [8] (a) A. R. Hajipour, S. E. Mallakpour, and S. Khoei, *Chemistry Lett.*, 120 (2000); (b) A. R. Hajipour, S. E. Mallakpour, and Gh. Imanzadeh, *J. Chem. Res. (S)*, 228 (1998); (c) A. R. Hajipour, S. E. Mallakpour, and H. Adibi, *Chem. Lett.*, 460 (2000).
- [9] T. M. Christopher, et al., *U. S. Patent* 5,922,404, 1999.
- [10] W. F. Barthel, P. A. Giang, and S. A. Hall, *J. A. Chem. Soc.*, **76**, 4186 (1954).
- [11] B. J. Teichmann, *Prakt. Chem.*, **28**, 94 (1965).
- [12] W. Lorenz, G. Henglein, and G. Schrader, *J.C.S.*, **77**, 2554 (1954).

- [13] E. P. Trutneva and R. R. Shagidullin, *Izv. Akad. Nauk SSSR, Ser. Khim.*, 1760 (1974).
- [14] K. V. Nikonorov, E. A. Gurylev, T. A. Zyabilkova, and I. D. Temyachev, *Izv. Akad. Nauk SSS, Ser. Khim.*, 1398 (1976).
- [15] I. V. S. Varaprasad and D. K. Jaiswal, *Ind. J. Chem.*, 2 (1982).
- [16] L. C. Thomas, in *The Identification of Functional groups in Organophosphorus Compounds* (Academic Press, London, 1974), p. 73.
- [17] C. H. Campblee, D. H. Chadwick, and Kaufman, *Ind. Eng. Chem.*, **49**, 1871 (1957).
- [18] G. P. Christopher and J. Peter, *J. Chem. Edu.*, **57**, 740 (1980).