## Accepted Manuscript

Stable diazonium salts of weakly basic amines—convenient reagents for synthesis of disperse azo dyes

Jinjing Qiu, Bingtao Tang, Benzhi Ju, Yuanji Xu, Shufen Zhang

PII: S0143-7208(16)30430-2

DOI: 10.1016/j.dyepig.2016.08.026

Reference: DYPI 5406

To appear in: Dyes and Pigments

Received Date: 29 June 2016

Revised Date: 8 August 2016

Accepted Date: 9 August 2016

Please cite this article as: Qiu J, Tang B, Ju B, Xu Y, Zhang S, Stable diazonium salts of weakly basic amines—convenient reagents for synthesis of disperse azo dyes, *Dyes and Pigments* (2016), doi: 10.1016/j.dyepig.2016.08.026.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



## Stable Diazonium Salts of Weakly Basic Amines—Convenient Reagents for Synthesis of Disperse Azo Dyes

Jinjing Qiu, Bingtao Tang<sup>\*</sup>, Benzhi Ju, Yuanji Xu, Shufen Zhang

State Key Laboratory of Fine Chemicals, Dalian University of Technology, Dalian 116024, PR China

#### ABSTRACT

A new synthetic strategy for industrially important deep-shade disperse azo dyes was presented in this study. The key procedure is to prepare stable solid diazonium salts of weakly basic amines in the absence of concentrated sulfuric acid. Diazotization by *tert*-butyl nitrite in ethyl acetate was allowed to proceed in the presence of equivalent 1,5-naphthalenedisulfonic acid as stabilizer of diazonium salts and donor of hydrogen ion for the reaction. The separated solid diazonium salts exhibited good thermal stability. The corresponding disperse azo dyes were subsequently synthesized through the azo-coupling of the prepared solid diazonium salts with a range of aromatic tertiary amines. The azo dyes were produced in short reaction time, excellent yields, mild reaction conditions, simple experimental procedure and low energy consumption.

<sup>\*</sup> Corresponding author. Tel.: +86 411 84986267; fax: +86 411 84986264.

E-mail address: tbt1976@sohu.com (B. Tang).

Keywords: Stable solid diazonium salts; Weakly basic amines; Disperse azo dyes; Diazotization

#### **1. Introduction**

Azo dyes are compounds containing electron-donating (D) and electron-accepting (A) functional groups through an azo  $\pi$ -conjugated linker [1] and the D- $\pi$ -A system has various electronic structures which can bring a full range of colors [2, 3]. Hence, as industrially important chromophores, azo dyes are widely used in dying of natural and synthetic fibers [4, 5], and many high technology fields, such as electronic devices, linear and nonlinear optics, reprography, sensors [6–8]. To date, approximately 3,000 different azo dyes are used industrially which is the important guarantee for meeting people's demand for color [9, 10].

Diazonium salts are precursors of the synthesis of azo dyes [11] and they are also widely used in organic synthesis such as Sandmeyer reaction [12], Balz–Schiemann reaction [13], carbon–carbon coupling reaction [14], and modification of polymeric materials and nanomaterials [15–17]. However, solid diazonium salts are very dangerous and have the tendency to explode because of their very poor thermal stability [18]. Thus, the diazotization of amines with sodium nitrite is usually carried out in the presence of excess hydrochloric or sulfuric acid solution and the obtained diazonium salt solution is directly poured into the coupling system to synthesize azo dyes resulting in large amounts of waste water containing strong liquid acid. Considering the above constraints, strong

solid acids as replacements for liquid acid catalysts [10, 19], nitrite ionic liquid or nitrite-functionalized graphene quantum dots as effective nitrosonium sources [9, 20] were reported to synthesize the corresponding azo dyes in excellent yields. Although each of these methods has specific merits, the substrates are simple aromatic amines or their derivatives, such as aniline, aminobenzenesulfonic acid and nitroaniline. Clean synthesis of azo dyes based on weakly basic amines is still a great challenge because they are very difficult to be diazotized due to their extremely weak alkalinity.

Weakly basic amines with two or three strong electron-withdrawing groups are the main raw materials on the preparation of dark disperse azo dyes with excellent color fastnesses [21]. However, their diazotization reactions using sodium nitrite are difficult to be carried out in hydrochloric or sulfuric acid solution because of their low reaction activity and solubility. In industrial production, the diazotization of weakly basic amines is always conducted in more than 10 times the molar quantity of concentrated sulfuric acid [22, 23] and has to use nitrosyl sulfuric acid as nitrosating agent which has remarkable diazotization ability. Subsequently, the concentrated acid solution of diazonium salt is poured into the coupling system. Due to the strong dilution heat effect of concentrated sulfuric acid, large amounts of ice were used to maintain low temperature of the system to avoid the decomposition of the diazonium salt. Meanwhile, it is difficult to handle the high acid content waste water.

Herein, we present a practical preparation method of disperse azo dyes based on the

novel stable solid diazonium salts of weakly basic amines. After the reactions of weakly basic amines with *tert*-butyl nitrite carried out in organic solution without any sulfuric acid in the presence of equivalent 1,5-naphthalenedisulfonic acid, the stable solid diazonium salts were obtained by simple filtration because of their very low solubility in organic solvent. The related disperse azo dyes were synthesized through the well-known azo-coupling reaction with aniline derivatives in good yields. Compared with traditional methods, coupling process can be conducted at room temperature, thus providing high reaction efficiency, excellent yields, mild reaction conditions, simple experimental procedure, and low energy consumption.

#### 2. Experimental

#### 2.1. General information

2,6-Dichloro-4-nitroaniline, 2-chloro-4-nitroaniline, 2,6-dibromo-4-nitroaniline, 2-amino-5-nitrobenzonitrile, 2,4-dinitroaniline, N,N-diethylaniline, tert-butyl nitrite and 1,5-naphthalenedisulfonic acid tetrahydrate were purchased from Sinopharm Chemical Reagent Co., Ltd. 1,5-naphthalenedisulfonic acid tetrahydrate was dried at 130 °C under high vacuum (2.7)3 kPa) for h. 3-(N-Ethylanilino)propionitrile, 3,3'-(phenylimino)dipropionitrile N-[3-(diethylamino)phenyl]acetamide and were purchased from Jiangsu Jihua Chemical Co., Ltd. The solvents were commercial analytical grade and used as received.

The diazotization was monitored by thin-layerchromatography on precoated silica gel glass based plates purchased from Branch of Qingdao Haiyang Chemical Co., Ltd and the yields of the diazonium salts were obtained by titration with the standard solution of N,N-diethylaniline at 0 °C-5 °C. The purity of dyes was determined by external standard method through an Agilent 1260 Infinity HPLC equipment. The diazonium salts and the dyes were all characterized by spectroscopic data with specified conditions. Tandem mass spectrometry (MS/MS) was recorded on a Themo Scientific TSQ Quantum Ultra equipment with suitable collision energies. MS spectra and HRMS spectra were recorded on a Themo Scientific LTQ Orbitrap XL mass spectrometer. FT-IR spectra were recorded as Nujol mulls or KBr disks using a Thermo Scientific Nicolet 6700 FT-IR spectrometer. <sup>1</sup>H NMR spectra and <sup>13</sup>C NMR spectra were recorded on a Bruker Avance III 500 MHz spectrometer using  $D_2SO_4$  or DMSO-d<sub>6</sub> as the deuterated solvents. The absorbtion spectra were recorded on an Agilent HP 8453 UV-Vis spectrophotometer. Elemental analysis was performed on an Element Vario EL III Analyzer. Thermal decomposition temperature of the diazonium salts were obtained by thermogravimetric analysis (TGA) conducted on a NETZSCH TG 209 C TGA instrument at 5 °C/min from ambient temperature to 200 °C under air atmosphere.

#### 2.2. Typical synthesis of stable solid diazonium salt of 2,4-dinitroaniline

2,4-Dinitroaniline (10 mmol) and 1,5-naphthalenedisulfonic acid (10 mmol) were dissolved in 100 mL ethyl acetate at 50 °C and the mixture was stirred for 10 min.

*Tert*-butyl nitrite (15 mmol) was added by dropping to the mixture for about 2 min at 25 °C. The reaction solution was further stirred for 20 min with the temperature maintained at 25 °C. A salmon pink solid was obtained by suction filtration and air drying. The yield was 99.0%.

#### 2.3. Characterization of the stable diazonium salts

2-Chloro-4-nitrobenzene diazonium salt (Table 1, 2a): MS/MS (10 eV), m/z: calcd for diazonium cation, 183.99; found, 184.03. <sup>1</sup>H NMR (D<sub>2</sub>SO<sub>4</sub>)  $\delta$  (ppm): 9.22 (2.00H, d, J = 8.7 Hz), 9.14–9.06 (3.72H, m), 8.94 (1.86H, dd, J = 9.1, 2.2 Hz), 8.85–8.76 (2.00H, m), 8.25 (2.00H, dd, J = 8.7, 7.5 Hz). FT-IR (Nujol mull, cm<sup>-1</sup>): 3100, 3071, 2311, 2280, 1569, 1528, 1499, 1461, 1377, 1358, 1298, 1237, 1164, 1098, 1040. Anal. Calcd: C, 40.35; H, 1.88; N, 12.41%. Found: C, 39.63; H, 1.80; N, 12.54%.

2,6-Dibromo-4-nitrobenzene diazonium salt (Table 1, 2b): MS/MS (10 eV), m/z: calcd for diazonium cation, 307.85; found, 307.63. <sup>1</sup>H NMR (D<sub>2</sub>SO<sub>4</sub>) δ (ppm): 9.22 (2.00H, d, *J* = 8.8 Hz), 9.15 (1.82H, s), 8.79 (2.00H, dd, *J* = 7.4, 10 Hz), 8.23 (2.00H, dd, *J* = 8.8, 7.5 Hz). FT-IR (Nujol mull, cm<sup>-1</sup>): 3084, 2288, 1561, 1539, 1499, 1460, 1377, 1345, 1292, 1231, 1199, 1176, 1152, 1076, 1043. Anal. Calcd: C, 32.29; H, 1.52; N, 7.06%. Found: C, 32.34; H, 1.44; N, 7.22%.

2,6-Dichloro-4-nitrobenzene diazonium salt (Table 1, 2c): MS/MS (5 eV), m/z: calcd for diazonium cation, 217.95; found, 218.18. <sup>1</sup>H NMR (D<sub>2</sub>SO<sub>4</sub>)  $\delta$  (ppm): 9.21 (2.00H, d,

*J* = 8.8 Hz), 8.98 (2.72H, s), 8.80 (2.00H, d, *J* = 7.5 Hz), 8.23 (2.00H, t, *J* = 8.2 Hz). FT-IR (Nujol mull, cm<sup>-1</sup>): 3087, 2285, 1551, 1460, 1415, 1377, 1345, 1293, 1268, 1244, 1220, 1201, 1167, 1110, 1034, 1019. Anal. Calcd: C, 37.30; H, 1.61; N, 9.77%. Found: C, 36.33; H, 1.82; N, 9.11%.

2-Cyano-4-nitrobenzene diazonium salt (Table 1, 2d): MS/MS (8 eV), m/z: calcd for diazonium cation, 175.03; found, 175.28. <sup>1</sup>H NMR (D<sub>2</sub>SO<sub>4</sub>) δ (ppm): 9.65 (1.59H, d, *J* = 1.8 Hz), 9.48–9.41 (3.18H, m), 9.23 (2.00H, d, *J* = 8.8 Hz), 8.81 (2.00H, d, *J* = 7.5 Hz), 8.25 (2.00H, t, *J* = 8.2 Hz), 4.88 (1.58H, q, *J* = 7.1 Hz), 1.80 (1.05H, d, *J* = 14.2 Hz). FT-IR (Nujol mull, cm<sup>-1</sup>): 3473, 3099, 2309, 2245, 1613, 1549, 1498, 1460, 1377, 1356, 1315, 1236, 1189, 1157, 1101, 1040. Anal. Calcd: C, 43.77; H, 2.23; N, 15.36%. Found: C, 43.04; H, 2.76; N, 14.79%.

2,4-Dinitrobenzene diazonium salt (Table 1, 2e): MS/MS (10 eV), m/z: calcd for diazonium cation, 195.01; found, 195.02. <sup>1</sup>H NMR (D<sub>2</sub>SO<sub>4</sub>) δ (ppm): 9.78 (1.85H, q, *J* = 2.0 Hz), 9.43 (3.70H, qd, *J* = 8.8, 5.0 Hz), 9.23 (2.00H, dd, *J* = 8.8, 2.9 Hz), 8.82 (2.00H, dd, *J* = 7.7, 2.9 Hz), 8.26 (2.00H, td, *J* = 8.3, 3.0 Hz). FT-IR (Nujol mull, cm<sup>-1</sup>): 3099, 3076, 3043, 2312, 2287, 1612, 1591, 1546, 1498, 1462, 1377, 1348, 1331, 1246, 1227, 1215, 1201, 1163, 1142, 1111, 1090, 1036. Anal. Calcd: C, 39.11; H, 1.81; N, 16.00%. Found: C, 39.07; H, 1.82; N, 15.83%.

#### 2.4. Typical procedure for coupling

N,N-Diethylaniline (10 mmol) was dissolved in 80 mL water with 1.2 mL 37% hydrochloric acid and then the salmon pink diazonium salt of 2,4-dinitroaniline was quickly added within 2 min at 0 °C–5 °C followed by adjusting the pH to 4–5 with 2 M NaOH solution. The reaction endpoint was confirmed by color reaction using H-acid. The precipitated dye was filtered, washed with water and dried at 80 °C to afford the crude product. The crude yield was 95.0% and the degree of purity was 97.0%.

#### 2.5. Spectroscopic data of disperse azo dyes

Compound 4a:  $C_{16}H_{17}CIN_4O_2$ : MS, m/z: calcd for  $[M+H]^+$ , 333.11; found, 333.10. HRMS, m/z: calcd for  $[M+H]^+$ , 333.1113; found, 333.1115. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>)  $\delta$ (ppm): 8.44 (1H, q, J = 2.5 Hz), 8.25 (1H, dq, J = 9.0, 2.5 Hz), 7.90–7.83 (2H, m), 7.83–7.76 (1H, m), 6.89 (2H, dt, J = 9.0, 2.8 Hz), 3.52 (4H, q, J = 7.2 Hz), 1.18 (6H, t, J= 6.9 Hz). <sup>13</sup>C NMR (126 MHz, DMSO-d<sub>6</sub>)  $\delta$  (ppm): 152.49, 151.88, 146.59, 143.12, 132.11, 126.92, 125.67, 123.33, 117.95, 111.53, 44.38, 12.50. FT-IR (KBr, cm<sup>-1</sup>): 3099, 2972, 2903, 1599, 1579, 1513, 1449, 1407, 1355, 1329, 1307, 1137, 1121, 1074, 1045, 1007, 887, 831, 819. UV ( $\lambda_{max}$  in acetone): 513 nm.

Compound 4b:  $C_{16}H_{16}Br_2N_4O_2$ : MS, m/z: calcd for  $[M+H]^+$ , 456.97; found, 457.01. HRMS, m/z: calcd for  $[M+H]^+$ , 456.9692; found, 456.9697. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>)  $\delta$ (ppm): 8.56 (2H, s), 7.84–7.78 (2H, m), 6.90–6.83 (2H, m), 3.51 (4H, q, *J* = 7.1 Hz), 1.17 (6H, t, *J* = 7.0 Hz). <sup>13</sup>C NMR (126 MHz, DMSO-d<sub>6</sub>)  $\delta$  (ppm): 155.23, 151.85, 145.67, 141.65, 127.78, 126.22, 115.16, 111.19, 44.32, 12.44. FT-IR (KBr, cm<sup>-1</sup>): 3084, 2971, 1601, 1521, 1394, 1355, 1336, 1273, 1253, 1191, 1163, 1131, 1072, 1051, 1007, 888, 819. UV (λ<sub>max</sub> in acetone): 441 nm.

Compound 4c:  $C_{16}H_{16}Cl_2N_4O_2$ : MS, m/z: calcd for  $[M+H]^+$ , 367.07; found, 367.08. HRMS, m/z: calcd for  $[M+H]^+$ , 367.0723; found, 367.0724. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>)  $\delta$ (ppm): 8.43 (2H, s), 7.84–7.78 (2H, m), 6.90–6.83 (2H, m), 3.51 (4H, q, J = 7.0 Hz), 1.17 (6H, t, J = 7.0 Hz). <sup>13</sup>C NMR (126 MHz, DMSO-d<sub>6</sub>)  $\delta$  (ppm): 153.12, 151.93, 145.21, 142.09, 126.77, 126.28, 124.46, 111.20, 44.33, 12.43. FT-IR (KBr, cm<sup>-1</sup>): 3089, 2974, 1603, 1521, 1396, 1355, 1340, 1274, 1253, 1192, 1148, 1127, 1061, 1008, 887, 821. UV ( $\lambda_{max}$  in acetone): 443 nm.

Compound 4d:  $C_{17}H_{17}N_5O_2$ : MS, m/z: calcd for  $[M+H]^+$ , 324.15; found, 324.13. HRMS, m/z: calcd for  $[M+H]^+$ , 324.1455; found, 324.1457. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>)  $\delta$ (ppm): 8.83 (1H, s), 8.52 (1H, dd, J = 9.1, 2.6 Hz), 7.95 (1H, d, J = 9.1 Hz), 7.89 (2H, d, J = 8.9 Hz), 7.01–6.90 (2H, m), 3.56 (4H, q, J = 7.0 Hz), 1.19 (6H, t, J = 7.1 Hz). <sup>13</sup>C NMR (126 MHz, DMSO-d<sub>6</sub>)  $\delta$  (ppm): 157.22, 152.71, 145.79, 142.96, 129.50, 128.92, 127.54, 117.43, 115.62, 111.96, 110.29, 44.58, 12.53. FT-IR (KBr, cm<sup>-1</sup>): 3077, 2975, 2931, 2229, 1597, 1577, 1516, 1414, 1355, 1324, 1271, 1249, 1171, 1136, 1117, 1071, 1007, 907, 838, 823. UV ( $\lambda_{max}$  in acetone): 536 nm.

Compound 4e:  $C_{16}H_{17}N_5O_4$ : MS, m/z: calcd for  $[M+H]^+$ , 344.14; found, 344.14.

HRMS, m/z: calcd for  $[M+H]^+$ , 344.1353; found, 344.1354. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>)  $\delta$  (ppm): 8.89–8.83 (1H, m), 8.50 (1H, ddd, J = 9.2, 6.5, 2.6 Hz), 7.95–7.87 (1H, m), 7.77 (2H, ddd, J = 9.6, 5.0, 2.1 Hz), 6.90 (2H, td, J = 7.1, 5.4, 2.1 Hz), 3.58–3.49 (4H, m), 1.21–1.14 (6H, m). <sup>13</sup>C NMR (126 MHz, DMSO-d<sub>6</sub>)  $\delta$  (ppm): 152.48, 148.07, 145.99, 145.39, 143.00, 127.55, 127.37, 119.80, 119.18, 111.77, 44.52, 12.48. FT-IR (KBr, cm<sup>-1</sup>): 3100, 2976, 2931, 1592, 1516, 1411, 1353, 1328, 1271, 1196, 1134, 1113, 1066, 911, 831, 822. UV ( $\lambda_{max}$  in acetone): 529 nm.

Compound 4f:  $C_{17}H_{15}Br_2N_5O_2$ : MS, m/z: calcd for  $[M+H]^+$ , 481.96; found, 482.12. HRMS, m/z: calcd for  $[M+H]^+$ , 481.9645; found, 481.9649. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>)  $\delta$ (ppm): 8.82–8.33 (2H, m), 7.88–7.79 (2H, m), 7.01–6.94 (2H, m), 3.80 (2H, t, J = 6.8Hz), 3.57 (2H, q, J = 7.0 Hz), 2.85 (2H, t, J = 6.8 Hz), 1.18 (3H, t, J = 7.0 Hz). <sup>13</sup>C NMR (126 MHz, DMSO-d<sub>6</sub>)  $\delta$  (ppm): 155.12, 151.51, 145.82, 142.35, 127.77, 125.94, 119.15, 115.01, 111.80, 45.41, 44.66, 15.72, 12.04. FT-IR (KBr, cm<sup>-1</sup>): 3100, 3066, 2970, 2933, 2248, 1600, 1522, 1399, 1342, 1280, 1245, 1142, 1077, 1054, 881, 825. UV ( $\lambda_{max}$  in acetone): 421 nm.

Compound 4g:  $C_{17}H_{15}Cl_2N_5O_2$ : MS, m/z: calcd for  $[M+H]^+$ , 392.07; found, 392.20. HRMS, m/z: calcd for  $[M+H]^+$ , 392.0676; found, 392.0677. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>)  $\delta$  (ppm): 8.42 (2H, s), 7.83 (2H, d, J = 9.0 Hz), 6.97 (2H, d, J = 9.1 Hz), 3.81 (2H, t, J = 6.8 Hz), 3.58 (2H, q, J = 7.0 Hz), 2.85 (2H, t, J = 6.7 Hz), 1.18 (3H, t, J = 7.0 Hz). <sup>13</sup>C NMR (126 MHz, DMSO-d<sub>6</sub>)  $\delta$  (ppm): 152.99, 151.59, 145.40, 142.76, 126.66, 125.99, 124.46, 119.12, 111.80, 45.41, 44.67, 15.70, 12.03. FT-IR (KBr, cm<sup>-1</sup>): 3059, 2980, 2252, 1604, 1572, 1529, 1519, 1394, 1336, 1311, 1280, 1148, 1134, 1078, 1065, 884, 822. UV (λ<sub>max</sub> in acetone): 423 nm.

Compound 4h:  $C_{18}H_{16}N_6O_2$ : MS, m/z: calcd for  $[M+H]^+$ , 349.14; found, 349.25. HRMS, m/z: calcd for  $[M+H]^+$ , 349.1408; found, 349.1412. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>)  $\delta$ (ppm): 8.84 (1H, d, J = 2.6 Hz), 8.54 (1H, dd, J = 9.1, 2.6 Hz), 7.96 (1H, d, J = 9.1 Hz), 7.90 (2H, d, J = 8.9 Hz), 7.04 (2H, d, J = 9.3 Hz), 3.84 (2H, t, J = 6.8 Hz), 3.61 (2H, J =7.1 Hz), 2.87 (2H, t, J = 6.7 Hz), 1.24–1.15 (m, 3H). <sup>13</sup>C NMR (126 MHz, DMSO-d<sub>6</sub>)  $\delta$ (ppm): 156.82, 152.29, 146.18, 143.47, 129.41, 128.92, 127.09, 119.01, 117.54, 115.44, 112.32, 110.76, 45.48, 44.86, 15.74, 12.12. FT-IR (KBr, cm<sup>-1</sup>): 3080, 2919, 2248, 2231, 1600, 1595, 1577, 1525, 1518, 1411, 1340, 1325, 1309, 1290, 1173, 1133, 1121, 910, 842, 825. UV ( $\lambda_{max}$  in acetone): 511 nm.

Compound 4i:  $C_{18}H_{15}CIN_6O_2$ : MS, m/z: calcd for  $[M+H]^+$ , 383.10; found, 383.15. HRMS, m/z: calcd for  $[M+H]^+$ , 383.1018; found, 383.1021. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>)  $\delta$ (ppm): 8.45 (1H, ddd, J = 7.4, 4.6, 2.4 Hz), 8.26 (1H, ddt, J = 8.9, 6.0, 2.2 Hz), 7.88 (2H, ddd, J = 9.4, 4.5, 2.3 Hz), 7.81–7.73 (1H, m), 7.09 (2H, dt, J = 9.4, 2.2 Hz), 3.89 (4H, t, J= 7.0 Hz), 2.86 (4H, t, J = 6.9 Hz). <sup>13</sup>C NMR (126 MHz, DMSO-d<sub>6</sub>)  $\delta$  (ppm): 152.11, 150.78, 147.27, 144.25, 132.64, 126.29, 125.73, 123.37, 118.99, 118.24, 112.55, 45.68, 15.40. FT-IR (KBr, cm<sup>-1</sup>): 2958, 2250, 1600, 1514, 1404, 1362, 1339, 1314, 1151, 1121, 1046, 888, 832. UV ( $\lambda_{max}$  in acetone): 459 nm. Compound 4j:  $C_{18}H_{14}Cl_2N_6O_2$ : MS, m/z: calcd for  $[M+H]^+$ , 417.06; found, 417.06. HRMS, m/z: calcd for  $[M+Na]^+$ , 339.0448; found, 439.0447. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>)  $\delta$ (ppm): 8.43 (2H, dd, J = 6.4, 1.9 Hz), 7.85 (2H, d, J = 9.2 Hz), 7.09 (2H, d, J = 9.2 Hz), 3.89 (4H, t, J = 6.8 Hz), 2.86 (4H, t, J = 6.8 Hz). <sup>13</sup>C NMR (126 MHz, DMSO-d<sub>6</sub>)  $\delta$ (ppm): 152.88, 150.97, 145.59, 143.38, 126.55, 125.78, 124.49, 119.00, 112.38, 45.66, 15.36. FT-IR (KBr, cm<sup>-1</sup>): 3079, 2918, 2247, 1603, 1568, 1518, 1406, 1377, 1345, 1249, 1208, 1147, 1062, 1026, 900, 831. UV ( $\lambda_{max}$  in acetone): 405 nm.

Compound 4k:  $C_{18}H_{19}Br_2N_5O_3$ : MS, m/z: calcd for  $[M+H]^+$ , 513.99; found, 513.99. HRMS, m/z: calcd for  $[M+H]^+$ , 513.9907; found, 513.9913. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>)  $\delta$ (ppm): 10.23 (1H, s), 8.54 (2H, s), 7.93 (1H, d, J = 3.0 Hz), 7.72 (1H, d, J = 9.4 Hz), 6.67 (1H, dd, J = 9.5, 2.8 Hz), 3.51 (4H, q, J = 7.1 Hz), 2.15 (3H, s), 1.20 (6H, t, J = 7.0 Hz). <sup>13</sup>C NMR (126 MHz, DMSO-d<sub>6</sub>)  $\delta$  (ppm): 169.25, 153.78, 153.28, 145.15, 139.97, 130.28, 128.06, 123.12, 116.07, 107.94, 100.07, 44.76, 25.06, 12.51. FT-IR (KBr, cm<sup>-1</sup>): 3091, 2981, 2930, 1690, 1622, 1616, 1529, 1513, 1494, 1336, 1291, 1184, 1167, 1125, 1074, 1055, 1005, 890, 873, 858. UV ( $\lambda_{max}$  in acetone): 491 nm.

Compound 41:  $C_{18}H_{19}Cl_2N_5O_3$ : MS, m/z: calcd for  $[M+H]^+$ , 424.09; found, 424.23. HRMS, m/z: calcd for  $[M+H]^+$ , 424.0938; found, 424.0950. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>)  $\delta$ (ppm): 10.30 (1H, s), 8.41 (2H, d, J = 1.0 Hz), 7.91 (1H, d, J = 2.7 Hz), 7.71 (1H, d, J =9.5 Hz), 6.67 (1H, dd, J = 9.4, 2.7 Hz), 3.51 (4H, q, J = 7.1 Hz), 2.14 (3H, s), 1.20 (6H, t, J = 7.0 Hz). <sup>13</sup>C NMR (126 MHz, DMSO-d<sub>6</sub>)  $\delta$  (ppm): 169.26, 153.34, 151.91, 144.66, 139.95, 130.92, 127.42, 124.66, 123.25, 108.00, 100.16, 44.78, 24.94, 12.51. FT-IR (KBr, cm<sup>-1</sup>): 3353, 3103, 2980, 2935, 2901, 1704, 1607, 1552, 1531, 1495, 1419, 1350, 1330, 1256, 1172, 1111, 1064, 1005, 901, 886, 855. UV (λ<sub>max</sub> in acetone): 496 nm.

Compound 4m:  $C_{18}H_{20}N_6O_5$ : MS, m/z: calcd for  $[M+H]^+$ , 401.16; found, 401.22. HRMS, m/z: calcd for  $[M+H]^+$ , 401.1568; found, 401.1573. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>)  $\delta$ (ppm): 10.69 (1H, s), 8.83 (1H, d, J = 2.5 Hz), 8.47 (1H, dd, J = 9.1, 2.5 Hz), 8.18 (1H, d, J = 9.1 Hz), 7.95 (1H, d, J = 2.7 Hz), 7.75–7.63 (1H, m), 6.73 (1H, dd, J = 9.5, 2.8 Hz), 3.53 (4H, q, J = 7.1 Hz), 2.21 (3H, s), 1.20 (6H, t, J = 7.0 Hz). <sup>13</sup>C NMR (126 MHz, DMSO-d<sub>6</sub>)  $\delta$  (ppm): 169.70, 153.93, 148.27, 145.13, 144.54, 132.33, 127.39, 123.76, 119.95, 119.40, 109.02, 100.47, 9.48, 45.02, 24.88, 12.58. FT-IR (KBr, cm<sup>-1</sup>): 3448, 3095, 2975, 2935, 1693, 1624, 1593, 1567, 1520, 1488, 1419, 1323, 1279, 1254, 1193, 1172, 1137, 1117, 1078, 897, 858, 845, 831, UV ( $\lambda_{max}$  in acetone): 546 nm.

#### 3. Results and discussion

#### 3.1. Preparation and characterization of stable solid diazonium salts

The solid diazonium salts of aromatic amines were conveniently obtained by simple filtration after the reactions of weakly basic amines with *tert*-butyl nitrite which were carried out in organic solution in the presence of equivalent 1,5-naphthalenedisulfonic acid (Scheme 1). The structures of diazonium salts were confirmed by FT-IR and tandem mass spectrometry according to references [24, 25].

#### ---Insert Scheme 1---

Optimization of the diazotization conditions was investigated with different molar ratios of the aromatic amine, 1,5-naphthalenedisulfonic acid and *tert*-butyl nitrite and the optimal ratio was 1:1:1.5. The new diazonium salts of weakly basic amines were formed by the diazonium cation and 1,5-naphthalenedisulfonic anion. Diazonium salts exist in two forms, i.e., structure A and structure B (Scheme 1). The ratio of each structure form was further confirmed by manual integration in <sup>1</sup>H NMR (Fig. 1). The corresponding results are summarized in Table 1. For 2,4-dinitrobenzene diazonium salt, the integral ratio in the <sup>1</sup>H NMR at 8.25 and 9.42 was about 1:1.85, which are corresponding to the 1,5-naphthalenedisulfonic acid anion and 2,4-dinitrobenzene diazonium cation (Fig. 1, 2e). Accordingly, the ratio of structures A and B is 0.15:0.85; thus, structure B accounts for 85% of the whole diazonium salt structures (Table 1, entry 5). This result can also be verified by elemental analysis (see Experimental section).

#### ---Insert Fig. 1---

#### ---Insert Table 1---

Reaction temperatures were stipulated in accordance with different substrates (Table 2). Although diazotization proceeded without any concentrated sulfuric acid or nitrosyl sulfuric acid, the reaction efficiency was very high. As a result, the very low basic amine, 2,4-dinitroaniline, could be diazotized completely within a short time (22 min, Table 2, entry 5). The diazotization times of three substrates (Table 2, entries 3-5) were much less than that of the traditional methods [28–30]. Meanwhile, the yields of the diazonium salts were excellent (Table 2). The high reaction efficiency may be attributed to the formation of nitrosonium cation (<sup>+</sup>NO) [31] in the presence of 1,5-naphthalenedisulfonic acid and tert-butyl nitrite.

---Insert Table 2---

3.2 Thermal stability of solid diazonium salts

### ----Insert Fig. 2----

TGA was used to characterize the thermal stability of the diazonium salts. The initial thermal decomposition temperatures of 2a, 2b, 2c, 2d, and 2e are 164.4, 175.5, 125.7, 115.9, and 121.9 °C, respectively (Table 2 and Fig. 2). Thus, the prepared diazonium salts of weakly basic amines are thermostable and can be stored at room temperature in dried state without disintegration.

#### 3.3. Synthesis of disperse azo dyes

The disperse azo dyes were synthesized by the well-known azo-coupling reaction between the as-prepared stable diazonium salts and the tertiary anilines at 0 °C–5 °C. The pH of the coupling system was adjusted and maintained at 4–6 by 2 M NaOH solution. The crude yields and purities of the relevant disperse azo dyes are shown in Table 3. ---Insert Scheme 2---

---Insert Table 3---

The coupling time of the present method was all much shorter than those reported in the literatures (a few to dozens of minutes, Table 3, entries 1, 6, and 9). The reaction ability of the tertiary anilines is restricted by concentrated sulfuric acid in traditional methods. However, large amounts of concentrated acid were replaced in the present method by equivalent organic acid in contrast to amines during diazotization. Consequently, the coupling reaction could be conducted with short time at low concentration acid.

#### ---Insert Fig. 3---

Compared with traditional coupling technology, the present method does not employ the dilution process of concentrated sulfuric acid because of the usage of solid diazonium salts. Thus, the generation of a large amount of heat produced from the dilution of sulfuric acid in the traditional coupling process was avoided. Meanwhile, the thermal analysis results of solid diazonium salts show that their thermal stabilities are sufficient. Therefore, it is not needed to use a lot of ice to keep the system at low temperature or even the reaction can be carried out at normal temperature without any cooling process. Compounds 4a, 4c, and 4d were chosen as objects of study and prepared with different temperatures. The yields of the dyes are shown in Fig. 3, which indicate no significant reducing of the yield during temperature increase. These results show that this new technology is of great significance for the realization of energy saving and emission reduction in the production of disperse azo dyes based on weakly basic amines.

#### 4. Conclusion

A novel strategy for synthesis of industrial disperse azo dyes based on weakly basic amines is presented. The key advantage of this new procedure is the formation and simple separation of stable solid diazonium salts of weakly basic amines. This new procedure can avoid the strong heat effect coming from the dilution of concentrated sulfuric acid in the traditional coupling process. Large amounts of ice were saved and the generation of highly concentrated acidic waste water was prevented. For these reasons, the efficient, rapid and economical synthesis of disperse azo dyes employing stable solid diazonium salts has a very good prospect in the industrial application.

#### Acknowledgments

This work was supported by the National Natural Science Foundation of China (21276042, 21576039 and 21421005), Program for Innovative Research Team in University (IRT\_13R06). Fundamental Research Funds for the Central Universities (DUT14YQ209), Program for New Century Excellent Talents in University (NCET130080), Program for Liaoning Excellent Talents in University (LJQ2013006).

#### References

- Deshmukh MS, Sekar N. A combined experimental and TD-DFT investigation of three disperse azo dyes having the nitroterephthalate skeleton. Dyes Pigments 2014;103:25–33.
- [2] Sunthankar SV, Thanumoorthy V. Disperse dyes: Dyeing properties and fastness of azo dyes. Text Res J 1977;47:126–8.
- [3] Li M, Zhang K, Hou AQ. Crystallographic study of two monoazo disperse dyes with a D-π-A system.
  Color Technol 2015;131:38–42.
- [4] Neamtu M, Yediler A, Siminiceanu I, Macoveanu M, Kettrup A. Decolorization of disperse red 354 azo dye in water by several oxidation processes—a comparative study. Dyes Pigments 2004;60:61–8.
- [5] Tskhovrebov AG, Naested LCE, Solari E, Scopelliti R, Severin K. Synthesis of azoimidazolium dyes with nitrous oxide. Angew Chem Int Edit 2015;54:1289–92.
- [6] Valizadeh H, Shomali A. A new nitrite ionic liquid (IL-ONO) as a nitrosonium source for the efficient diazotization of aniline derivatives and in-situ synthesis of azo dyes. Dyes Pigments 2012;92:1138–43.
- [7] Matei A, Constantinescu C, Mitu B, Filipescu M, Ion V, Ionita I, et al. Laser printing of azo-derivative thin films for non-linear optical applications. Appl Surf Sci 2015;336: 200–5.
- [8] Shahab S, Hajikolaee FH, Filippovich L, Darroudi M, Loiko VA, Kumar R, et al. Molecular structure and UV–Vis spectral analysis of new synthesized azo dyes for application in polarizing films. Dyes Pigments 2016;129:9–17.

- [9] Valizadeh H, Shomali A, Nourshargh S, Mohammad-Rezaei R. Carboxyl and nitrite functionalized graphene quantum dots as a highly active reagent and catalyst for rapid diazotization reaction and synthesis of azo-dyes under solvent-free conditions. Dyes Pigments 2015;113:522–8.
- [10] Koukabi N, Otokesh S, Kolvari E, Amoozadeh A. Convenient and rapid diazotization and diazo coupling reaction via aryl diazonium nanomagnetic sulfate under solvent-free conditions at room temperature. Dyes Pigments 2016;124:12–7.
- [11] Velasco MI, Kinen CO, De Rossi RH, Rossi LI. A green alternative to synthetize azo compounds. Dyes Pigments 2011;90:259–64.
- [12] Nemati F, Elhampour A. Green and efficient diazotization-iodination of aryl amines using cellulose sulfuric acid as a biodegradable and recyclable proton source under solvent-free condition. Sci Iran 2012;19:1594–6.
- [13] Laali KK, Gettwert VJ. Fluorodediazoniation in ionic liquid solvents: new life for the Balz–Schiemann reaction. J Fluorine Chem 2001;107:31–4.
- [14] Christoforou IC, Koutentis PA. New regiospecific isothiazole C–C coupling chemistry. Org Biomol Chem 2006;4:3681–93.
- [15] Mahouche-Chergui S, Gam-Derouich S, Mangeney C, Chehimi MM. Aryl diazonium salts: a new class of coupling agents for bonding polymers, biomacromolecules and nanoparticles to surfaces. Chem Soc Rev 2011;40:4143–66.

- [16] Mooste M, Kibena E, Kozlova J, Marandi M, Matisen L, Niilisk A, et al. Electrografting and morphological studies of chemical vapour deposition grown graphene sheets modified by electroreduction of aryldiazonium salts. Electrochim Acta 2015;161:195–204.
- [17] Assresahegn BD, Brousse T, Bélanger D. Advances on the use of diazonium chemistry for functionalization of materials used in energy storage systems. Carbon 2015;92:362–81.
- [18] Garel L, Saint-Jalmes L. One-pot fluoro-de-diazoniation of anilines in organic medium. Tetrahedron Lett 2006;47:5705–8.
- [19] Zarei A, Hajipour AR, Khazdooz L, Mirjalili BF, Chermahini AN. Rapid and efficient diazotization and diazo coupling reactions on silica sulfuric acid under solvent-free conditions. Dyes Pigments 2009;81:240–4.
- [20] Valizadeh H, Amiri M, Hosseinzadeh F. Nanoparticles of organosilane-based nitrite ionic liquid immobilized on silica for the diazotization of aniline derivatives and subsequent synthesis of azo dyes. Dyes Pigments 2012;92:1308–13.
- [21] Desai P, Himeno K, Desai N, inventor; Colourtex Industries Limited (IN), assignee. Disperse azo dyes. United States patent US 8409300. 2013 Apr 2.
- [22] Hooker J, Hinks D, Montero G, Conlee C. Synthesis of N,N-diethyl-N-{4-[(E)-(4-nitrophenyl)diazenyl]phenyl}amine via in situ diazotisation and coupling in supercritical carbon dioxide. Color Technol 2002;118:273–6.

- [23] Schoutissen HAJ. The diazotization of very weakly basic amines. J Am Chem Soc 1933;55:4531-4.
- [24] Broxton TJ, Colton R, Traeger JC. Electrospray mass spectral studies of aromatic diazonium cations. Part 2. The surprising stability of some substituted 2-nitrobenzene diazonium cations to collisionally activated fragmentation. J Phys Org Chem 1995;8:351–5.
- [25] Ben-Efraim DA. Detection and determination of diazo and diazonium groups. In: Patai S, editor. The chemistry of diazonium and diazo groups. New York; John Wiley & Sons Ltd; 1978. P. 149–78.
- [26] He L, El-Shafei A, Freeman HS, Boyle P. X-ray and molecular modelling studies of 4-[N-alkylamino]azobenzene dyes. Dyes Pigments 2009;82:299–306.
- [27] Sun J, Zheng CL, Hökelek T. X-ray crystal structure analysis for CI Disperse Orange 61. Colorn Technol 2012;128:91–4.
- [28] Raditoiu V, Raditoiu A, Wagner L, Amariutei V, Raduly M, Nicolae CA. Disperse dyes functionalized with trialkoxysilane groups. Rev Chim-Bucharest 2011;62:409–14.
- [29] Wagner L, Raditoiu V, Raditoiu A, Ardeleanu P, Amariutei V, Raduly M. Functionalized azobenzenes chromogens with aromatic amino primary groups. Rev Chim-Bucharest 2009;60: 444–9.
- [30] Nicolescu FA, Jerca VV, Draghici C, Vuluga DM, Vasilescu DS. Synthesis and characterization of novel azobenzene methacrylate monomers. Des Monomers Polym 2009;12:553–63.
- [31] Kraska J, Boruszczak Z. Studies on the diazotization of aromatic amines with glycol nitrite. Dyes Pigments 1990;12:173–7.

[32] Mungmeechai T, Suesat J, Suwanruji P. Study of the relationships between the chemical structures of azo disperse dyes and their dyeing properties on polyester. Advanced Materials Research 2010;93–94:332–5.

#### ACCEPTED MANUSCRIPT

Scheme 1. Preparation of stable diazonium salts in organic solvent.

Scheme 2. Azo-coupling of stable diazonium salts with tertiary aniline derivatives.

Fig. 1. <sup>1</sup>H NMR spectra of the diazonium salts in D<sub>2</sub>SO<sub>4</sub>.

Fig. 2. Thermogravimetric curves of five stable solid diazonium salts.

Fig. 3. Yields of three representative dyes at different coupling temperatures (Yield = Crude yield  $\times$  purity).

Entry	Amine	Diazonium salt	Structure	Ratio of
Linu y			number	structures A and B
1	1a	$O_2N$ $\sim$ $N_2O_3S$ $-Ar$	2a	0.14:0.86
2	1b	$O_2N$ $ N_2O_3S$ $ Ar$ $Br$ $Br$	2b	1.00:0.00
3	1c	$O_2N \longrightarrow O_2N \longrightarrow O_2O_3S - Ar$	2c	0.64:0.36
4	1d	$O_2N \rightarrow N_2O_3S - Ar$	2d	0.41:0.59
5	1e	$O_2N - NO_2 - NO_2O_3S - Ar$	2e	0.15:0.85

# Table 1Structures of the stable solid diazonium salts

Table 2	
Preparation of stable diazonium salts	

Entry	Structure number	Temperature (°C)	Reaction time (min)		Yield	Thermal decomposition	
			Found	Reported [Ref.]	(%)	temperature (°C)	
1	2a <sup>a</sup>	0	60	60[26]	92.8	164.4	
2	2b	50	120	120[27]	93.1	175.5	
3	2c	22	17	120[28]	98.0	125.7	
4	2d	0	60	120[29]	89.7	115.9 <sup>b</sup>	
5	2e	25	22	120[30]	99.0	121.9	

<sup>a</sup> The 2-chloro-4-nitroaniline was added slowly to the reaction system after *tert*-butyl nitrite.

<sup>b</sup> There was a continuous lower decline of the TG curves before 115.9 °C.

#### Table 3

Synthesis of azo dyes using stable diazonium salts

Entry s	Diazonium salt	Coupling reagent	Product	Structure number	Reaction time (min)		Crude	D :
					Found	Reported [Ref.]	yield (%) <sup>a</sup>	Purity (%)
1	2a		$O_2N \rightarrow N \rightarrow N \rightarrow N \rightarrow N$	4a	5	120 [32]	92.6	95.9
2	2b		$O_2N \longrightarrow Br$ Br	4b	5		90.8	97.9
3	2c		$O_2N \rightarrow \bigvee_{Cl}^{Cl} N = N \rightarrow \bigvee_{N}^{N} N$	4c	5		88.5	97.7
4	2d		$O_2N \rightarrow N = N \rightarrow N$	4d	5		90.9	88.8
5	2e		0 <sub>2</sub> N-\-N=N-\-N-N	4e	5		95.0	97.0
6	2b	CH <sub>2</sub> CH <sub>2</sub> CN	$O_2N - \bigvee_{Br}^{Br} N = N - \bigvee_{-N}^{CH_2CH_2CN} O_2N - \bigvee_{-N}^{CH_2CH_2CN}$	4f	12	120 [27]	93.5	96.7
7	2c	CH <sub>2</sub> CH <sub>2</sub> CN	$O_2N - \left\{ \begin{array}{c} Cl \\ N = N - \left\{ \begin{array}{c} CH_2CH_2CN \\ Cl \end{array} \right\}$	4g	10		96.1	96.0
8	2d	CH <sub>2</sub> CH <sub>2</sub> CN	$O_2N \rightarrow N \rightarrow$	4h	16		94.7	87.3
9	2a	CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CN CH <sub>2</sub> CH <sub>2</sub> CN	O <sub>2</sub> N- N=N- N=N- CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CN CH <sub>2</sub> CH <sub>2</sub> CN	4i	40	300 [26]	94.8	96.6
10	2c	CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CN CH <sub>2</sub> CH <sub>2</sub> CN	$O_2N - \underbrace{\begin{pmatrix} CI \\ -N = N - \underbrace{\end{pmatrix}}_{CI} N = N - \underbrace{\begin{pmatrix} CH_2CH_2CN \\ CH_2CH_2CN \end{pmatrix}}_{CH_2CH_2CN}$	4j	19		94.0	90.9



<sup>a</sup> The crude yields of disperse azo dyes were calculated based on the initial weakly basic amines.

CHER MAR







REPART





- $\begin{array}{ll} 4a\text{-}e: & R_1=R_2=\mathrm{CH}_2\mathrm{CH}_3, R_3=\mathrm{H}\\ \\ 4f\text{-}h: & R_1=\mathrm{CH}_2\mathrm{CH}_2\mathrm{CN}, R_2=\mathrm{CH}_2\mathrm{CH}_3, R_3=\mathrm{H}\\ \\ 4i,j: & R_1=R_2=\mathrm{CH}_2\mathrm{CH}_2\mathrm{CN}, R_3=\mathrm{H} \end{array}$
- $4k\text{-}m: \ R_1 = R_2 = CH_2CH_3, R_3 = NHCOCH_3$

#### Highlights

•Diazotization of weakly basic amines occurred in the absence of concentrated sulfuric acid.

•Solid diazonium salts with good thermal stability were obtained.

•There is no dilution process of concentrated acid in the coupling reaction.

•Large amounts of ice are saved and energy consumption can be reduced.