Reduction of Nitric Oxide by Tetramesityliridium(IV) and Cobaltocene. Reactions of Hyponitrite Complexes and of the Ether $[Co(\eta^5-C_5H_5)(\eta^4-C_5H_5)](\mu-O-exo)$ with Nitroalkanes, Acids and Amines[†]

Robyn S. Hay-Motherwell,^a Geoffrey Wilkinson,^{*,a} Tracy K. N. Sweet^b and Michael B. Hursthouse^{*,b}

^a Johnson Matthey Laboratory, Chemistry Department, Imperial College, London SW7 2AY, UK ^b School of Chemistry and Applied Chemistry, University of Wales Cardiff, PO Box 912, Cardiff CF1 3TB, UK

The interactions of either Ir(mes)₄ (mes = $C_6H_2Me_3$ -2,4,6) or Co(cp)₂ (cp = η^5 - C_5H_5) with NO in light petroleum gave unstable solids the IR spectra of which allow them to be formulated as hyponitrite complexes, e.g. $[Co(cp)_2]_2(N_2O_2)$ 1; their reaction with MeNO₂ is described. The compound $Co(cp)_2$ with either NO or Ag, N,O, in toluene gave N,O and the very reactive ether $[Co(cp)(\eta^4-C_5H_5)]_{,}(\mu-O-exo) 2$. This reacts with nitroalkanes where the product depends on the nature of the nitroalkane. Nitromethane produced both $Co(cp)_2^+ CH_2 = NO_2^- H_2O_3$ and $[Co(cp)(\eta^4 - C_5H_5)]_2[(\mu - CH(NO_2) - exo]_4$, while $Me_2CH(NO_2)$ gave $Co(cp)[\eta^4-C_5H_5(CMe_2NO_2)-exo]$ and $EtNO_2$ corresponding species to 3 and 4 (*i.e.* 6 and $\overline{7}$). Reactions of 2 with alcohols or phenylacetylene gave mononuclear species of the same type as 4; treatment with NH₂Ph or NHPh₂ also gave similar exo complexes 8 and 9 respectively. All these exo species reacted with CHCI₃ to give the exo-CCI₃ compound by facile C-C bond cleavage. Mechanisms for the various reactions are discussed. The structures of compounds 2-4, 8 and 9 have been confirmed by X-ray crystallography: 2 and 4 contain $Co(cp)(\eta^4-C_{_5}H_{_5})$ moieties bridged by oxygen or CH(NO₂) groups while 8 and 9 are monomers with exo-NHPh or -NPh, groups. In all cases but one (2) there are near-eclipsed C_xCoC_x geometries; for one of the cobalts in 2 there is a twist of ca. 16° from the eclipsed configuration. Compound 3 has Co(cp),⁺ and CH,NO,⁻ ions the latter being involved in strong hydrogen bonding with the H₂O molecule also present in the lattice. In both 8 and 9 there is structural evidence for π interaction between p lone pairs on the planar nitrogen atom and the π system of a phenyl ring.

This study arose from the observation that the paramagnetic (1e) compound $Ir(mes)_4$, where $mes = C_6H_2Me_3-2,4,6,^1$ reacted with nitric oxide in light petroleum to give an unstable solid, which was soluble in nitromethane or acetonitrile giving a solution ¹H NMR spectrum identical to that of the iridium(v) cation ¹ [Ir(mes)₄]⁺.

iridium(v) cation ¹ $[Ir(mes)_4]^+$. Previous studies ² of the reactions of paramagnetic (1e) alkyls and aryls of transition metals with nitric oxide led to insertion of NO into the M-C bond followed by the formation of an M=O bond and RN=NR. However, it was evident that despite the existence of $(mes)_3Ir^V=O^3$ the reaction, equation (1), does

NO +
$$lr(mes)_4$$
 (mes)₃ $lr \lesssim \int_{0}^{N(mes)}$
(mes)₃ $lr = 0 + \frac{1}{2}(mes)N=N(mes)$ (1)

not occur. In view of the limited quantities of $Ir(mes)_4$ available we studied the reaction of NO with $Co(cp)_2$ ($cp = \eta^5-C_5H_5$) which also is paramagnetic (1e) and is readily oxidised to the cobaltocenium ion, $Co(cp)_2^{+,4}$

Studies by Bottomley and co-workers,⁵ of paramagnetic η^5 -cyclopentadienyl compounds of Ti^{III} and V^{II} showed that

the interaction with NO led to compounds with M–O–M bonds. Although the dioxodinitrate(N-N)(2–) (hyponitrite) ion, N₂O₂²⁻, was considered as a possible intermediate in the reactions, no direct evidence for the formation of this ion was obtained ^{5c} although N₂O was formed; this gas could be obtained by the reaction (2). Unstable hyponitrite complexes

$$N_2 O_2^{2-} \longrightarrow N_2 O_1 + O^{2-}$$
 (2)

of several transition metals have been obtained in reactions of NO with low-valent metal compounds or in thermal reactions of bis(nitrosyl) complexes.⁶ Sodium hyponitrite is made⁷ by sodium reduction of NO in reactions presumably involving two one-electron transfers, equations (3)–(5).

$$M' + 'NO \longrightarrow M^+ + NO^-$$
(3)

$$NO^- + 'NO \longrightarrow N_2O_2'^-$$
 (4)

$$N_2O_2^{\bullet-} + M^{\bullet} \longrightarrow 2M^+N_2O_2^{2-}$$
 (5)

Results and Discussion

Reaction of $Ir(mes)_4$ and $Co(cp)_2$ with NO.—The interaction of $Ir(mes)_4$ with NO in light petroleum gives a greenblack precipitate the IR spectrum of which, taken as quickly as

[†] Supplementary data available: see Instructions for Authors, J. Chem. Soc., Dalton Trans., 1994, Issue 1, pp. xxiii-xxviii.



Scheme 1 Main reactions of $Co(cp)_2$ involving NO and nitroalkanes. $L = (\eta^5 - C_5H_5)Co(\eta^4 - C_5H_5)$ unit. (*i*) NO in light petroleum; (*ii*) NO in MeNO₂; (*iii*) NO or Ag₂N₂O₂ in toluene; (*iv*) toluene; (*v*) MeNO₂ in Et₂O; (*ii*) Me₂CH(NO₂) in Et₂O; (*vii*) CHCl₃; (*viii*) CCl₄ [50% yield + Co(cp)₂Cl]. Nitrous oxide is eliminated in reactions of compound 1 and in (*iii*)



possible, has three bands (1097, 1022, 802 cm⁻¹) in the regions found ⁶ for co-ordinated hyponitrites; ionic hyponitrites⁸ have two bands around 1058 and 571 cm⁻¹, as we have confirmed also for Ag₂N₂O₂. Since three bands are also found for the cobalt compound discussed below we consider that the iridium compound is the oxygen-bridged species $[Ir^{V}(mes)_{4}]_{2}(\mu$ -ONNO- $\kappa^{2}O$) formed by two one-electron transfers from Ir(mes)₄ to NO as in equations (3)–(5).

The interaction of $Co(cp)_2$ (see Scheme 1) in hexane or light petroleum with NO leads rapidly to a beige precipitate. This compound 1 is unstable at room temperature, becoming redorange (see below), but the IR spectrum, again taken as rapidly as possible, also has three bands at 1088, 1020 and 800 cm⁻¹ in addition to bands for cp and η^4 -C₅H₅ rings; a structure of the type I seems reasonable.

It is surprising considering the extensive work on reactions of NO with $M(cp)_2$ type compounds and especially in view of the innumerable studies⁹ on $Co(cp)_2$ that the reaction of the latter with NO appears not to have been reported, perhaps because of the problems encountered in this study. The only comparable reaction is that of $Co(cp)_2$ and O_2 at low temperatures,¹⁰ which gives a very thermally unstable compound (decomp. > -40 °C), formulated only on the basis of its chemical reactions (*cf.* later discussion) as the *exo*-peroxobridged η^4 -cyclopentadiene cobalt(1) complex II. It may be noted here, in view of the similar compounds discussed below, that a variety of *exo*-bridged species are known,^{9,10} *e.g.* with μ -CF₂CF₂ and μ -CH₂C(O)CH₂; some are noted later.

On standing in toluene compound 1 decomposes with loss of N_2O , confirmed by collection of gas and subsequent gas chromatography-mass spectrometry (GS-MS) study, to give an orange solution containing a cobalt(1) complex 2 described below.

The iridium hyponitrite dissolves in nitromethane with evolution of N_2O as confirmed by mass spectral study but, unlike the cobalt system discussed below, no crystalline product could be obtained; the ¹H NMR spectrum in CDCl₃ of the solid residue after pumping off MeNO₂ showed only bands for



Fig. 1 Structure of the ether $[Co(cp)(\eta^4-C_5H_5)]_2(\mu$ -O-exo) 2



 $[Ir(mes)_4]^+$ as noted in the Introduction and the IR spectrum showed OH stretches.

Reaction of $Co(cp)_2$ and NO in Toluene.—By contrast with the reaction in hexane, in toluene NO reacts immediately to give an orange solution (N₂O is also formed and detected in off-gas by mass spectrometry) from which, on cooling, orange crystals of a compound 2 can be isolated. That a hyponitrite intermediate is involved is confirmed directly by vigorous stirring of a toluene solution of Co(cp)₂ with solid Ag₂N₂O₂ which again leads to 2 and N₂O.

The compound 2 although stable at room temperature for only ca. 1 h can be kept below ca. -20 °C and the crystal structure was determined at low temperature. This shows that 2 is the exo-µ-O ether III and similar to that proposed for the µ-peroxo species noted above. The structure is shown in Fig. 1 and selected bond lengths and angles are given in Table 1. The main feature is the unsymmetrical disposition of the η^4 -C₅H₅ groups about the ether linkage which is probably due to steric effects. As a result, the methine H atom on ring 2 [at C(201)] lies approximately over the η^4 ring 1 [C(102)–C(105)] and the C-O-C angle at the ether oxygen at 114.7(3)° is slightly larger than expected. Other points of note relate to the geometries of the distorted rings. Not unexpectedly the η^4 ring has 'slipped' as a result of the loss of bonding to the aliphatic carbons C(101) and C(201) by approximately 0.25 Å. For Co(1) the two rings have a closely eclipsed relative arrangement whereas for Co(2) they are twisted by approximately 16° from an eclipsed configuration. The Co–C bond lengths to the η^5 rings lie in the range 2.037(5)–2.082(5) Å whereas the distances to the η^4 rings are significantly shorter on average and specifically shorter to the central atoms of the 'butadiene' fragment [C(103), C(104) and C(203), C(204)] at 1.967(5)-1.974(5) Å, cf. 2.027(5)-2.042(5) Å to C(102), C(105), C(202) and C(205). The C-O distances in the ether link at 1.456(5), 1.475(5) Å are normal. In

Table 1 Selected bond lengths (Å) and angles (°) for compounds 2-4, 8 and 9

	•			<u>^</u>	~
	2	3	4	8	9
Co(1)-C(101)	2.497(4)	2.021(4)	2.554(5)	2.552(7)	2.534(3)
Co(1)-C(102)	2.038(5)	2.019(4)	2.027(5)	2.023(7)	2.046(3)
Co(1)-C(103)	1.974(5)	2.017(4)	1.980(5)	1.974(7)	1.985(4)
$C_{0}(1) - C(104)$	1.968(5)	2.010(4)	1.967(5)	1.981(7)	1.982(4)
$C_0(1) - C(105)$	2.042(5)	2.015(5)	2.008(5)	2.019(8)	2.031(3)
$C_0(1) = C(106)$	2.044(5)	2 010(4)	2 030(5)	2 053(7)	2 070(4)
$C_0(1) - C(107)$	2.011(5) 2.037(5)	2.015(4)	2.039(5)	2.053(7) 2.063(7)	2 102(4)
$C_{0}(1) - C(108)$	2.037(5)	2.003(4) 2.023(4)	2.057(5)	2.003(7)	2.102(4)
$C_{0}(1) = C(100)$	2.070(5)	2.023(4)	2.092(5)	2.093(7)	2.077(4)
$C_0(1) = C(109)$	2.062(5)	2.022(4)	2.030(3)	2.000(7)	2.040(3)
CO(1) = C(110)	2.040(3)	2.012(4)	2.042(3)	2.042(7)	2.000(4)
C-(2) C(201)	2 527(5)		2 549(5)		
$C_{0}(2) = C(201)$	2.327(3)	_	2.348(3)	_	
$C_0(2) = C(202)$	2.027(5)		2.027(5)	_	
$C_0(2) = C(203)$	1.967(5)	_	1.964(5)		_
Co(2) = C(204)	1.96/(5)		1.973(5)		
$C_0(2) - C(205)$	2.041(5)	_	2.008(4)	_	
Co(2) - C(206)	2.055(5)	_	2.030(5)	_	
Co(2)-C(207)	2.061(5)	—	2.052(5)		
Co(2)–C(208)	2.077(5)		2.100(5)		—
Co(2)–C(209)	2.078(5)	—	2.081(5)		
Co(2)C(210)	2.050(5)	_	2.028(6)	—	_
Solvent in lattice					
O(1)-N		1 303(4)			
$O(2) \ge N$		1.305(4) 1.276(4)		_	_
C(11) - N		1.270(4)	_	_	
		1.500(5)			
Bridging stoms					
bridging atoms					
O(1)-C(101)	1.475(5)		—		
O(1)-C(201)	1.456(5)		—		_
C(1)-C(101)			1.540(6)		
C(1)-C(201)			1.554(6)		_
Pendant groups					
N-C(1)				1 475(8)	1 508(4)
$\mathbf{N} - \mathbf{C}(11)$				1 387(8)	1 400(4)
N-C(21)				1.507(0)	1 450(4)
N_H				1.05(8)	1.450(4)
14 11				1.05(8)	
X(1) = C(101) = C(102)	117 3(4)		115 4(4)		
X(1) - C(101) - C(102)	117.0(4)		113.4(4)		
X(1) - C(201) - C(205)	117.0(4) 110.7(4)		115.0(4)		
X(1) = C(201) = C(203)	110.7(4)		113.3(4)		_
$C(201) \times (1) - C(202)$	114.7(4) 114.7(2)		114.7(4)		
C(104) = C(105) = C(101)	114.7(3)	_	112.8(4)		_
C(104) = C(103) = C(101)	109.4(4)	_	110.0(4)		
C(103) - C(102) - C(101)	109.9(4)		109.8(4)		_
C(204) - C(205) - C(201)	110.1(4)		109.7(4)	_	
C(203)-C(202)-C(201)	108.9(4)	_	109.3(4)		—
N(1)-C(1)-C(101)		_	110.0(4)	—	_
N(1)-C(1)-C(201)			106.1(3)	_	—
O(2)-N-O(1)	—	117.5(3)	123.4(4)	_	
O(2) - N - C(11)		122.6(4)	117.9(4)	_	
O(1)-N-C(11)		120.0(4)	118.8(4)		—
C(2)-C(1)-N				117.7(5)	116.5(2)
C(5)-C(1)-N			_	113.2(6)	116.4(3)
C(3)-C(2)-C(1)				110.2(6)	110.3(3)
C(4)-C(5)-C(1)	_	_		112.1(7)	110.4(3)
C(1) - N - C(11)		_		122.7(6)	119.0(2)
C(1) - N - C(21)	_				119.2(2)
C(1)–N–H	_	-	_	118(4)	
С(11)-N-Н				114(4)	_

For compounds 3, 8 and 9, C(101) to C(110) refer to actual atoms C(1) to C(10); for 2, 4, 8 and 9, C(101) and C(201) (2 and 4) are the tilted atoms in the η^4 rings; X is the bridging atom, oxygen in complex 2 and carbon in 4; for compound 4, C(11) refers to the actual atom C(1).

the folded η^4 rings the envelope folds are 27.6(3) and 30.2(5)° for rings 1 and 2 respectively; the angles between the η^5 rings and η^4 moieties are close to zero at 3.9(3) and 3.4(4)°. The IR and ¹H NMR spectra of compound 2 are in accord

with the ether structure and are closely similar to those of the cobalt and rhodium cyclopentadiene complexes $M(cp)(\eta^4-C_5-H_5R)$ [M = Co, R = H, Me, CHCl₂ or CCl₃, M = Rh, R = H]; the cobalt compounds were made by interaction of



Fig. 2 Asymmetric unit in the structure of Co(cp)_2 $^+ \mathrm{CH}_2 = \mathrm{NO}_2 \,^- \cdot \, \mathrm{H}_2 \mathrm{O} \, 3$



Fig. 3 Packing diagram for compound 3

alkyl halides with $Co(cp)_2$.^{*,11a} The only difference is that **2** has a band in the IR spectrum at 898 cm⁻¹ that can be assigned to the ether, C–O–C, linkage. The ether **2** is exceptionally reactive towards compounds with acidic hydrogens and this behaviour is in striking contrast to the usual low reactivity



Fig. 4 Hydrogen-bonded ribbons of $H_2O \cdot CH_2NO_2^-$ present in the structure of compound 3



of ethers, except special cases like alkyl tetrahydropyranyl ethers. $^{\rm 12}$

Reactions of Compound 2 with Nitroalkanes.—Nitromethane reacts rapidly with compound 2 in diethyl ether solutions {cf. the reaction of $[Ir(mes)_4]_2(N_2O_2)$ with MeNO₂ above} to give two products easily separated by crystallisation. The first isolated is a yellow crystalline solid 3 which is unstable at room temperature or on pumping in vacuum, losing MeNO₂. The residue according to IR spectra appears to be $Co(cp)_2^+OH^-$. Since the crystal structure of 3 discussed below indicates that it is $Co(cp)_2^+CH_2=NO_2^-H_2O$ this observation is not surprising. In solution in CDCl₃ the ¹H NMR spectrum shows bands for $Co(cp)_2^+$ and MeNO₂, the latter formed by reaction of $CH_2=NO_2^-$ and the water of crystallisation.

The structure of compound $\hat{3}$ determined by X-ray crystallography at low temperature is shown in Fig. 2; selected

^{*} The compound $Rh(cp)(C_5H_6)$ was made by direct interaction of $RhCl_3$ and Na(cp) in the solid state at *ca*. 120 °C in November 1955 at Harvard University by G. W. and characterised by IR and ¹H NMR spectroscopy. These species were first formulated as having *endo*-R groups but later reformulated as *exo* species.^{9b} The mechanism of reactions of $Co(cp)_2$ and RX is known to be free radical in nature.^{11b}



Fig. 5 Structure of $[Co(cp)(\eta^4-C_5H_5)]_2[\mu-CH(NO_2)-exo] 4$

bond lengths and angles are given in Table 1. The rings in the cobaltocenium ion have a near-eclipsed configuration with Co-C bond lengths shorter than in the Co(cp) moiety for compound 2 as expected, 2.005(4)-2.023(4) Å. The compound also contains the methylideneazinate ion $CH_2=NO_2^{-1}$ the structure of which has not, to our knowledge, previously been determined.¹³ The ion is planar to within limits of experimental error and the geometry agrees with the results of theoretical calculations,^{13a} except that our measured C-N bond, which clearly has multiple character, is some 0.04 Å shorter than that predicted using an extended basis set (MP2-HF/6-31G*), but agrees very well with that computed at the Hartree-Fock (HF) level. The two N-O bond lengths are 1.276(4) and 1.303(4) Å, with the former agreeing well with theory. In the crystal structure the CH2=NO2 - ion is involved in an extended ribbonlike hydrogen bonding system with the water molecule as can be seen in Figs. 3 and 4. The strongest interactions are $O(3)-H(32) \cdots O(1)-N$ 1.83 and $O(3)-H(31) \cdots O(1)-N$ 1.81 Å. The involvement only of O(1) of the NO₂ group in hydrogen bonding probably accounts for the greater length of O(1)-N.

The second compound 4 formed in the reaction of 2 with MeNO₂ is red-orange; this is the *exo*-CH(NO₂) cyclopentadiene cobalt(1) complex shown in IV. In addition to the bands characteristic for the cp and η^4 -diene rings the IR spectrum shows NO₂ stretches; the ¹H NMR spectrum in C₆D₆ has a triplet for the H of the *exo*- μ -CH(NO₂) group while the *endo*-H atoms of the rings appear as a doublet of triplets, the coupling constant (J = 8.25 Hz) being as expected.

The structure of compound 4 as determined by X-ray diffraction is shown in Fig. 5; selected bond lengths and angles are given in Table 1. It is analogous to that of 2 but with O replaced by a doubly deprotonated MeNO₂ group, *i.e.* CH(NO₂). The presence of the latter results in a different conformational arrangement of the η^4 -C₅H₅ groups about the C-C bonds of the bridge and it is noteworthy that the C-C-C angle is some 2° less than the C-O-C angle in 2. The geometries of the Co(cp) and η^4 -C₅H₅ fragments are very similar to those in 2 in terms both of bond lengths (Table 1) and folds in the η^4 rings [33.0(5), 32.5(4)°]. However, the angles between the η^5 ring and the η^4 fragment are now 6.4(5) and 6.8(3)°, the slight increase possibly reflecting the greater steric crowding present in 4. The rings are close to eclipsed in both fragments of the molecule.

Reaction Mechanisms of NO and RNO₂ Reactions.— Assuming, as suggested above, that the hyponitrite is co-



Scheme 2 Interaction of compound 2 with nitromethane. Overall stoichiometry, $2 \times 2 + 3$ MeNO $\longrightarrow 2 \times 3 + 4$



Scheme 3 Carbonium ion route for interaction of compound 2 and $MeNO_2$

ordinated in compound 1 it seems most likely that the N_2O_2 group is *exo* bound to a η^4 -cyclopentadiene ring as in V [equation (6)] since ready decomposition to the ether 2 and



 N_2O can then occur. Co-ordination of hyponitrite to cobalt seems highly unlikely. The reaction of 2 with MeNO₂ could then proceed as in Scheme 2. An alternative in Scheme 3 involves carbonium ion intermediates. Although the hydroxo species Co(cp)[η^4 -C₅H₅(OH)-*exo*] shown in Scheme 2 has previously been postulated in the interaction of Co(cp)₂⁺ salts

with concentrated NaOH solutions under forcing conditions to give azulene in low yield ¹⁴ there appears to be no other evidence for its existence. The *exo*-hydroxide would doubtless react immediately with MeNO₂ to give Co(cp)[η^4 -C₅H₅(CH₂NO₂)-*exo*] which, with a second molecule of the hydroxide, would produce the unique μ -CH(NO₂) bridge in compound 4. The water formed in the reactions is then found in the salt 3.

The formation of the monomeric exo-CH₂NO₂ species as an intermediate in Scheme 2 suggested that treatment of compound 2 with Me₂CH(NO₂) would give only a monomer, since further H-atom transfer to give μ -CH(NO₂) could not occur. This was found to be the case and the exo-CMe₂NO₂ complex 5 was isolated in essentially quantitative yield according to equation (7). It is of interest that in neat nitroalkanes as solvent

 $Co(cp)_2$ reacts with NO to give only the $Co(cp)_2^+$ salts of the appropriate anion.¹⁵ Presumably the protonation reactions coupled with the driving force of N₂O elimination are more rapid than the reactions of the ether **2** with nitroalkanes.

The ether 2 does, however, react with $EtNO_2$ to give both the $Co(cp)_2^+$ salt of $Me(H)C=NO_2^- 6$ as well as $Co(cp)\{\eta^4-C_5H_5[CH(Me)NO_2-exo]\}$ 7. The failure to form the analogue of compound 4 despite the presence of a CH group in 7 is presumably due to steric factors. These are perhaps also the reasons why $Me_2CH(NO_2)$ in Et_2O gives only 5 without the formation of the cobaltocenium salt.

Other Reactions of Compound 2.—There have been previous syntheses of $exo-\eta^4$ -diene cobalt species⁹ from CH acids. Thus acetylene reacts with Co(cp)₂ to give a dimeric species with an exo-CH=CH bridge¹⁶ while secondary phosphine oxides HP(O)R₂, *e.g.* R = Buⁿ or C₆H₁₁, give Co(cp){ η^4 -C₅H₅[P(O)R₂-*exo*]} and hydrogen at 100–140 °C.¹⁷ However, most species [other than those formed along with Co(cp)X by interaction of Co(cp)₂ with halogenoalkanes, see *e.g.* refs. 9(*a*), (*b*) and 11(*a*)] have been obtained by interaction of Co(cp)₂ in *the presence of oxygen* with a variety of compounds^{9,10} and evidently these reactions involved the labile μ -perox species noted earlier. *exo*-Compounds were obtained from CHCl₃, MeCN, Me₂HCCN, PhC=CH, Me₂CO [which gives a CH₂-C(O)CH₂ bridge] as well as C₅H₆,¹⁸ C₅H₅Me, indene and methanol.

In addition to confirming the nature of some of these products in reactions of compound 2 other new ones have been obtained. In all cases stoichiometric amounts of the reagent were added to the toluene solution of 2 made *in situ* from NO and Co(cp)₂. The compounds were characterised by ¹H NMR spectra in benzene solution. Monomeric species were obtained from PhC=CH, ^{10b} MeOH^{9c} and adamantan-1-ol. Acetic acid and C₆F₅OH however gave the Co(cp)₂⁺ salts of the anion while SiHPh₃ gave Co(cp)(C₅H₆) and (Ph₃Si)₂O. There was no reaction with RhH(CO)(PPh₃)₃.

While $Co(cp)_2$ can readily be obtained by interaction of $CoCl_2$ with C_5H_6 in the presence of NEt_3 and $NHEt_2$,¹⁹ in toluene compound 2 reacts with NH_2Ph , $NHPh_2$, $NHEt_2$, 1-aminonaphthalene and carbazole. Although the complex from carbazole is insoluble, the other species are moderately soluble in benzene and stable at room temperature. The structures of 8 and 9 (Figs. 6 and 7), the *exo* compounds from aniline and diphenylamine, respectively, have been determined by X-ray crystallography, selected bond lengths and angles are given in Table 1. The structures show that the geometries of the $Co(cp)(\eta^4-C_5H_5)$ moieties in 8 and 9 are similar to those of the other monomeric compounds of the types discussed earlier.



Fig. 6 Structure of $Co(cp)[\eta^4-C_5H_5-(NHPh)-exo]$ 8



Fig. 7 Structure of $Co(cp)[\eta^4-C_5H_5-(NPh_2)-exo]$ 9

The envelope folds in the η^4 rings are 31.7(6) and 29.3(3)° and the cp/ η^4 rings are eclipsed in both molecules. Perhaps the most significant feature in the structures is that in both molecules the geometry about the N atom is planar with the one phenyl ring in 8 and one of the rings in 9 coplanar with the plane of the sp² nitrogen. This feature and the shortness of the relevant N–C bond length (especially in 9 where it is 0.05 Å shorter than that to the other ring which is orthogonal to the N plane) indicates some interaction ²⁰ between the π system of the coplanar phenyl ring and the lone pair on the N atom which is in the p orbital perpendicular to the plane.

Since the reaction of compound 2 with an amine is likely to differ from that of CH acids, the attack is more likely to be nucleophilic on the η^4 -diene ring than on the ether oxygen atom. A possible pathway is shown in Scheme 4.

Facile C–X *Cleavage.*—Finally, all of the *exo* compounds both mono- and bi-nuclear except $Co(cp)(C_5H_6)$ react rapidly and quantitatively with CHCl₃ or CDCl₃ at room temperature to give the *exo*-CCl₃ compound ^{11a} as, *e.g.*, in equation (8).

$$Co(cp)(C_5H_5R) + CHCl_3 \longrightarrow Co(cp)[C_5H_5(CCl_3)] + RH \quad (8)$$

Possible mechanisms are of the type shown in equations (9) and (10). It is also of interest that the C-P bond in the





Overall reaction:



Scheme 4 Interaction of amines with compound 2. Reaction (iv) follows a path similar to those in (i)–(iii). Nu = Nucleophile



 $P(O)R_2$ species noted above is also cleaved by $CDCl_3^{17}$ but the products were formulated as having *exo*-CDCl₂ groups according to the ¹H NMR spectra, which of course would not be definitive. We consider them also to have *exo*-CCl₃ groups.

Experimental

Microanalyses for the compounds (with one exception) could not be obtained due to the air and thermal sensitivity. The general techniques used have been described.¹ Infrared spectra (cm⁻¹) are in Nujol mulls. Proton NMR spectra ($\delta vs.$ SiMe₄) were obtained on a JEOL-EX-270 spectrometer at 270 MHz, mass spectra of NO and N₂O gas samples on a JEOL JMS-AX 505W spectrometer by the University of London MS Service at King's College.

Commercial samples were from Aldrich. The light petroleum used had b.p. 40–60 °C. Nitric oxide was passed through a copper coil at -78 °C and the purity checked by mass spectrometry. Sodium and silver hyponitrites ^{7,8} and Co(cp)₂²¹ were made as described. Nitroalkanes were dried over CaSO₄ and distilled under N₂. All operations were carried out under purified N₂ or Ar or under vacuum.

Interaction of Co(cp)₂ and NO.—The gas was slowly passed through Co(cp)₂ (ca. 0.2 g) in light petroleum (30 cm³) for 5–10 min until the purple colour disappeared and a beige precipitate of compound 1 was formed. After rapid filtration, the IR spectrum showed bands due to hyponitrite (see text) in addition to η^5 - and η^4 -C₅H₅ bands; the N₂O₂ bands disappear when the solid turns red-orange to be replaced by the C–O–C band²² at ca. 900 cm⁻¹ due to the ether link in **2**. The gas above the solid after decomposition was N₂O.

When a similar reaction was carried out in toluene a small amount of yellow solid was formed and a red-orange solution from which, after filtration, reduction in volume to ca. 10 cm³ and cooling (-20 °C), X-ray-quality crystals of $[Co(cp)(\eta^4 C_5H_5$]₂(μ -O-exo) 2 were obtained in > 90% yield. The crystals are stable for weeks at -20 °C. IR (cm⁻¹): Co(cp)(η^4 -C₅H₅) bands ^{11*a*} and 898 (C–O–C). ¹H NMR (C₆D₆): δ 4.71 (t, J_{av} = 1.5, 4 H, H^{3.4} of η^4 ring), 4.44 (s, 10 H, cp), 4.03 (t, $J_{av} = 2.3$, 2 H, endo H¹ of η^4 ring), and 3.04 [q, J (inner pair) = 2.0, J (outer pair) = 2.3 Hz, 4 H, H^{2.5} of η^4 ring]. This spectrum is identical to that of the complex obtained from decomposition of 1 in toluene at room temperature. Compound 2 was also obtained essentially quantitatively when a mixture of $Ag_2N_2O_2$ (ca. 0.07 g) and $Co(cp)_2$ (ca. 0.09 g) in toluene (30 cm³) was stirred very vigorously for 1 h; N₂O was lost and the solution became red-orange with formation of a black precipitate of silver. After filtration and reduction in volume, compound 2 was obtained in > 80% yield on cooling to -20 °C.

Reactions with Nitroalkanes.—(a) Compounds 3 and 4. A fresh sample of compound 2 (ca. 0.06 g) was dissolved in MeNO₂ (ca. 2 cm³) and the orange solution layered with Et₂O. After ca. 48 h at -20 °C yellow crystals of 3 (ca. 0.04 g, yield ca. 98%) were collected. The filtrate was evaporated in vacuum and the residue dissolved in toluene; cooling at -20 °C gave orange-red crystals of 4 (0.03 g, yield ca. 98%). X-Ray-quality crystals of 3 were obtained by recrystallisation from MeNO₂-Et₂O and of 4 from toluene. Compound 3: IR (cm⁻¹) 3400 (br, H₂O), ca. 1200 (C=NO₂⁻)^{23a} and strong bands^{23b} at 1417, 1012 and 868 [Co(cp)₂⁺]; ¹H NMR (CDCl₃) δ 5.8 (s, cp) and 4.3 (MeNO₂ from reaction with H₂O, see text). Compound 4: IR (cm⁻¹) cp and η^4 -C₅H₅ bands; 1528, 1360 (NO₂);^{15 1}H NMR (C₆D₆) δ 4.77 (m, 4 H, H^{3.4} of η^4 ring), 4.38 (s, 10 H, cp), 2.99 (d of pseudo-t, 2 H, J_d = 8.25, J_t = 2.31, endo-H¹), 2.44 (m, 2 H, H² or H⁵), 2.38 (m, 2 H, H² or H⁵) and 2.36 [t, 1 H, J = 8.25 Hz, CH(NO₂)].

(b) Compound 5. As in (a) but compound 2 (ca. 0.07 g) was dissolved with stirring in Me₂CH(NO₂) (20 cm³); reduction in volume and cooling to -20 °C gave a quantitative yield of orange crystals of 5. IR (cm⁻¹) 1575, 1151 (NO₂); cp and η^4 -C₃H₅ bands. ¹H NMR (C₆D₆): δ 4.74 (t, br, 2 H, H^{3.4} of η^4 ring); 4.34 (s, 5 H, cp), 3.20 (t, br, 1 H, endo-H¹), 2.42 (q, br, 2 H, H^{2.5}) and 0.85 (s, 6 H, Me₂CNO₂).

(c) Interaction of NO and $Co(cp)_2$ in either neat MeNO₂ or Me₂CH(NO₂) gave a quantitative yield of the respective salts on work-up; both are unstable when solvent is removed. The characterisation was by IR and NMR spectra.

(d) The reaction of compound 2 and EtNO₂ was carried out as in (a) but 2 (0.16 g) was dissolved in EtNO₂ (3 cm³) and Et₂O was added to precipitate the yellow salt 6 in quantitative yield (0.1 g). Evaporation of the mother-liquor and dissolution of the residue in toluene gave a quantitative yield (0.11 g) of orange Co(cp){ η^4 -C₅H₅[CH(Me)NO₂-exo]} 7. Compound 6: IR (cm⁻¹) 1552, 1260, 1370 and Co(cp)₂ + bands; ¹H NMR (CDCl₃) δ 5.93 (cp), 3.24 (q, J = 7.26, CH₂) and 0.62 (t, J = 7.26, CH₃), both for EtNO₂. Compound 7: IR (cm⁻¹) 1543, 1325, 1174 plus cp and η^4 -C₅H₅ bands; ¹H NMR (C₆D₆) δ 4.75 (m, 1 H, H³ or H⁴ of η^4 ring), 4.70 (m, 1 H, H³ or H⁴), 4.34 (s, 5 H, cp), 3.05 [d of q, 1 H, J = 6.6 for q, 7.6 for d, CH(Me)NO₂], 2.85 (d of pseudo-t, 1 H, J = 7.59 for d, 2.64, 2.31 Hz for t), 2.50 and 2.19 (m, 1 H, H² or H⁵ of η^4 ring).

 Table 2
 Crystal data and structure refinement for compounds 2-4, 8 and 9

	2	3	4	8	9
Formula	C20H20C02O	C11H14CoNO3	C ₂₁ H ₂₁ C ₀₂ NO ₂	C16H16CoN	C ₂₂ H ₂₀ CoN
M _r	394.22	267.17	437.25	281.23	357.32
Crystal system	Monoclinic	Monoclinic	Triclinic	Triclinic	Triclinic
Space group	$P2_1/a$	$P2_1/a$	ΡĪ	PĨ	РĪ
a/Å	11.537(4)	7.520(1)	5.615(1)	5.756(9)	8.036(8)
b/Å	10.197(2)	17.904(4)	11.920(2)	9.43(2)	11.060(3)
c/Å	13.265(2)	8.433(1)	13.583(3)	12.23(2)	11.363(2)
α/°	90(0)	90(0)	104.05(1)	98.7(1)	118.97(6)
β/°	96.81(2)	108.13(2)	94.42(1)	88.7(1)	108.94(4)
γ/°.	90(0)	90(0)	91.12(1)	106.26(8)	88.34(4)
$U/Å^3$	1549.5(7)	1079.0(3)	878.6(3)	629.8(2)	825.6(9)
Ζ	4	4	2	2	2
$D_{\rm c}/{\rm Mg}{\rm m}^{-3}$	1.690	1.645	1.653	1.483	1.437
F(000)	808	552	448	292	372
$\mu(Mo-K\alpha)/mm^{-1}$	2.141	1.537	1.903	1.341	1.040
Reflections collected	5050	5473	2769	2096	2962
Independent reflections (R_{int})	2471 (0.0485)	2774 (0.0664)	2440 (0.0487)	1689 (0.0651)	2284 (0.0329)
Maximum, minimum absorption correction factors	0.775, 0.597	1.293, 1.059	0.834, 0.573		
Data, parameters	2465, 288	2756, 201	2428, 310	1687, 227	2284, 297
Goodness of fit, F^2	0.383	0.417	0.337	0.382	0.381
Final R indices R1, wR2					
$[I > 2\sigma(I)]$	0.0320, 0.0800	0.0392, 0.0747	0.0356, 0.0863	0.0403, 0.0945	0.0348, 0.0884
(all data)	0.0566, 0.1230	0.0848, 0.1334	0.0505, 0.1592	0.0698, 0.1538	0.0406, 0.0996
Largest difference peak and hole/e Å ⁻³	0.376, -0.511	0.552, -0.539	0.334, -0.390	0.543, -0.238	0.395, -0.266
Goodness of fit, $S = [\sum w(F_0^2 - F_c^2)^2/(n-p)]^{\frac{1}{2}}$ wh	here $n =$ number of	of reflections and p	= total number of	parameters; R1 =	$\Sigma F_{\rm o} - F_{\rm c} / \Sigma F_{\rm o};$

Goodness of m, $S = \lfloor 2w(F_o^2 - F_c^2)^2/(n-p) \rfloor^2$ where n = number of reflections and p = total number of parameters; $KI = \Sigma |F_o - F_c| \Sigma F_o$; $wR2 = [\Sigma w(F_o^2 - F_c^2)^2 / \Sigma w(F_o^2)^2]^{\frac{1}{2}}$ where $w = 1/[\sigma^2(F_o)^2 + (xP)^2]$ with x = 0 for compound 2 and 0.1 for 3, 4, 8 and 9 and $P = [\max(F_o^2, 0) + 2F_c^2]/3$.

Reactions of Compound 2 with Other Substrates.—All reactions were performed in the same manner from compound 2 prepared in situ: NO was passed slowly through a stirred toluene solution of $Co(cp)_2$ (0.1 g in ca. 20 cm³) for 5–10 min as previously described. After filtration, the required amount of purified substrate [corresponding to 90% yield of 2 from $Co(cp)_2$] was added at room temperature. The formation of product was immediate with yields quantitative according to NMR spectroscopy, except in the cases of adamantan-1-ol (80) and PhC=CH (23%). All products were crystallised from toluene; in the case of carbazole the product precipitated from the reaction mixture. Proton NMR data, in C_6D_6 unless otherwise indicated, for new compounds are as follows.

Compound 8: δ 6.3–7.1 (m, 5 H, aromatic), 4.70 (pseudo-t, br, 2 H, $J_t = 1.98$, H^{3.4}), 4.49 (s, 5 H, cp), 4.00 (d of pseudo-t, $J_d = 8.59$, $J_t = 1.98$ and 2.64, endo-H¹), 2.95 (q, 2 H, J = 1.98, H^{2.5}) and 2.93 (d, br, 1 H, J ca. 8 Hz, NH).

Compound 9 (in C₆D₅CD₃): δ 6.8–7.15 (m, 10 H, aromatic), 4.70 (s, br, 1 H, endo-H¹), 4.49 (s, br, 7 H, cp and H^{3.4}) and 2.91 (s, br, 2 H, H^{2.5}); at -15 to -65 °C, δ 7.16–6.8 (m, 10 H, aromatic), 4.68 (s, br, 1 H, endo-H¹), 4.45 (s, 5 H, cp), 4.41 (s, br, 2 H, H^{3.4}) and 2.87 (s, br, 2 H, H^{2.5}).

Co(cp)[η^4 -C₅H₅(NHC₁₀H₇)-*exo*]: δ 7.4–6.8 (m, 7 H, aromatic), 4.74 (pseudo-t, 2 H, *J* 1.98 and 1.65 Hz, H^{3.4}), 4.50 (s, 5 H, cp), 4.22 (d pseudo-t, 1 H, J_d = 7.9, J_t = 2.31, *endo*-H¹), 3.72 (d, br, 1 H, J_d = 7.9, NH), 3.06 (q, 2 H, *J* = 1.98 between inner pairs, 2.31 Hz between outer pairs, H^{2.5}).

Co(cp)[η^4 -C₅H₅(NEt₂)-exo]: δ 4.86 (pseudo-t, 2 H, J = 1.98 and 2.65 Hz, H^{3.4}), 4.49 (s, 5 H, cp), 3.74 (pseudo-t, 1 H, J = 2.31 and 1.98, endo-H¹), 2.80 (q, 2 H, J = 2.65 between inner pairs, 2.31 between outer pairs, H^{2.5}) 2.23 (q, 4 H, J = 7.26, CH₂CH₃) and 0.92 (t, 6 H, J = 7.26 Hz, CH₂CH₃).

Co(cp)[η^4 -C₅H₅(OC₁₀H₁₅)-exo]: δ 4.76 (t, 2 H, J = 1.65, H^{3,4}), 4.50 (s, 5 H, cp), 4.27 (pseudo-t, 1 H, J = 2.64 and 2.31, endo-H¹), 3.05 (q, 2 H, J = 1.65 between inner pairs, 2.31 between outer pairs, H^{2,5}), 1.91 (s, br, 3 H, bridgehead H of adamantyl), 1.74 (d, 6 H, J = 2.96, adamantyl) and 1.45 (pseudo-t, 6 H, J = 2.97 and 3.29 Hz, adamantyl).

Data not previously reported for known compounds: $Co(cp)[\eta^4-C_5H_5(OMe)-exo]$,^{9c} δ 4.75 (t, br, 2 H, H^{3.4} of η^4 ring), 4.44 (s, 5 H, cp), 4.05 (t, br, 1 H, endo-H¹), 3.06 (q, br, 2 H, H^{2.5}) and 2.94 (s, 3 H, OCH₃); $Co(cp)[\eta^4-C_5H_5(C\equiv CPh)-exo]$,^{10b} δ 7.33 (m, 2 H, Ph), 6.83 (m, 3 H, Ph), 5.03 (t, 2 H, J =1.98, H^{3.4}), 4.40 (s, 5 H, cp), 3.66 (pseudo-t, 1 H, J = 2.64 and 2.31, endo-H¹) and 2.68 (q, 2 H, J = 1.98 Hz, H^{2.5}).

1.30, 11 '), 4.30 (s, 5 II, cp), 5.00 (pseudo-t, 1 II, J = 2.04 and 2.31, endo-H¹) and 2.68 (q, 2 H, J = 1.98 Hz, H^{2.5}). With SiPh₃H to give Co(cp)(η^{4} -C₅H₆).^{11a} At 270 MHz: δ 5.23 (m, 2 H, H^{3.4}), 4.56 (s, 5 H, cp), 2.64 (d pseudo-t, 1 H, $J_d = 13.87$, $J_t = 1.98$, endo-H¹), 2.40 (q, br, 2 H, H^{2.5}) and 2.05 (d pseudo-t, 1 H, $J_d = 13.88$, $J_t = 1.99$ Hz, exo-H¹). Other product: (Ph₃Si)₂O; δ 7.0–7.7 (m, aromatic H).

With $MeCO_2H$ (in $CDCl_3$). δ 5.82 (s, cp), 2.11 (CO_2Me).

With C₆F₅OH (in CDCl₃). δ 5.81 (s, cp). ¹⁹F (referenced to CFCl₃): δ -171.0 (m, br, *o*-, *m*-F) and -188.4 (m, br, *p*-F).

With carbazole. The red solid was insoluble in C_6D_6 and $C_6D_5CD_3$ but in CDCl₃ the compound $Co(cp)[\eta^4-C_5H_5-(CCl_3)-exo]$ was formed (Found: C, 74.7; H, 4.8; N, 4.5. $C_{22}H_{18}CoN$ requires C, 74.4; H, 5.1; N, 4.0%). FAB mass spectrum in 3-nitrobenzyl alcohol: m/z 189, $Co(cp)_2^+$.

Reactions of exo-R Compounds with CHCl₃ or CDCl₃.—All the nitro compounds when dissolved in the minimum amount of chloroform at room temperature were converted quantitatively into the nitroalkane and Co(cp)[C₅H₅(CCl₃)-exo] the spectra of which were identical to those of authentic samples.^{11a} ¹H NMR (CDCl₃): δ 5.28 (t, 2 H, J = 1.98, H³ and H⁴), 4.82 (s, 5 H, cp), 3.76 (t, 1 H, J = 2.31, H¹) and 3.00 (q, 2 H, J =1.65 between inner pairs, 2.31 Hz between outer pairs, H² and H⁵).

All of the other compounds except $Co(cp)(\eta^4-C_5H_6)$ reacted similarly.

Reaction of $Ir(mes)_4$ and NO.—Nitric oxide was slowly passed into $Ir(mes)_4$ (ca. 0.15 g) in light petroleum (ca. 30 cm³) until the initial dark brown colour disappeared (ca. 20 min) and a dark green-black precipitate formed. This was collected rapidly and the IR spectrum showed broad bands at 1097, 1022 and 802 cm⁻¹. The solid was soluble in MeCN or MeNO₂ when

Table 3Fractional atomic coordinates ($\times 10^4$)

222	1
223	L

				. .			
Atom	x	У	Z	Atom	x	У	Z
Compound	12						
	4 656(1)	2 319(1)	5 907(1)	C(201)	2 715(4)	3 068(5)	2 810(3)
$C_0(2)$	1 966(1)	2 088(1)	1 107(1)	C(202)	1 738(4)	2 074(5)	2 599(4)
C(101)	3 311(4)	3 540(4)	4 622(3)	C(203)	794(5)	2 666(5)	1 975(4)
C(102)	4 598(4)	3 374(5)	4 597(3)	C(204)	1 264(5)	3 747(5)	1 486(4)
C(102)	4 962(5)	2 024(5)	4 377(3) A A67(A)	C(204)	2 469(5)	3 796(5)	1 834(4)
C(103)	4002(3)	203+(3)	4 740(2)	C(205)	2 407(5)	622(5)	7/8(/)
C(104)	3 914(4) 2 07((4)	1 293(3)	4 /49(3) 5 029(4)	C(200)	3077(3)	1 725(5)	(+)0+/
C(105)	3 0 / 6(4)	2 202(5)	5 U38(4)	C(207)	3137(3)	1 723(3)	99(4) 449(4)
C(106)	4 740(5)	3 312(6)	7 252(4)	C(208)	2 027(4)	1 946(3)	440(4)
C(107)	4 309(5)	2 035(6)	/ 361(4)	C(209)	1 272(4)	982(5)	-135(4)
C(108)	5 171(5)	1 115(5)	7 143(4)	C(210)	1 902(4)	186(5)	604(4)
C(109)	6 135(4)	1 835(5)	6 892(4)	0	2 607(3)	3 931(3)	3 668(2)
C(110)	5 874(5)	3 195(5)	6 938(4)				
a	1.0						
Compound	13						
Co	1 494(1)	6 113(1)	2 935(1)	C(8)	- 594(6)	6 477(2)	932(5)
C(1)	4 125(6)	5 979(3)	4 495(6)	C(9)	-1 334(6)	6 136(2)	2 108(6)
C(2)	4 103(6)	5 942(3)	2 815(7)	C(10)	-632(7)	6 524(2)	3 642(6)
C(3)	2 904(7)	5 347(2)	2 042(6)	C(11)	4 440(6)	6 357(3)	8 673(6)
C(4)	2 174(6)	5 025(2)	3 246(6)	OÙÚ	1 652(5)	6 930(2)	8 253(5)
CO	2.936(6)	5 416(2)	4 757(6)	$\vec{O}(2)$	1 736(5)	5 701(2)	8 115(4)
C(6)	519(7)	7 105(2)	3 414(6)	O(3)	-2178(5)	6 886(2)	6 767(4)
C(0)	552(6)	7 077(2)	1761(6)	N N	2674(5)	6 320(2)	8 346(4)
C(I)	552(0)	1011(2)	1 /01(0)	19	2 024(3)	0 520(2)	0 540(4)
Compound	d 4						
$C_{\alpha}(1)$	1.026(1)	1 771(1)	4 005(1)	C(109)	1 120(10)	1 222(5)	2 127(2)
$C_0(1)$	1.920(1)	17/1(1)	4003(1)	C(100)	2 599(0)	1 233(3)	2 + 27(3)
CO(2)	$\frac{5779(1)}{10(4(7))}$	2 / 10(1)	9 304(1)	C(109)	2 300(9)	19(4)	2 147(4)
0(1)	1 964(7)	5 /95(3)	/ 036(3)	C(110)	1 444(9)	18(4)	3 487(4)
O(2)	-1243(7)	4 /13(3)	6 892(3)	C(201)	2 175(9)	3 3/1(4)	7 797(3)
N	949(8)	4 851(3)	6 926(3)	C(202)	3 562(9)	4 105(4)	8 751(3)
C(1)	2 414(8)	3 805(4)	6 818(3)	C(203)	5 895(9)	3 708(4)	8 822(3)
C(101)	1 532(8)	2 894(4)	5 838(3)	C(204)	5 848(9)	2 551(4)	8 232(3)
C(102)	1 196(9)	3 340(4)	4 882(3)	C(205)	3 459(8)	2 272(4)	7 835(3)
C(103)	3 417(9)	3 351(4)	4 442(4)	C(206)	961(11)	2 333(5)	10 092(4)
C(104)	4 828(9)	2 540(5)	4 809(4)	C(207)	2 582(10)	3 147(5)	10 793(4)
C(105)	3 417(9)	2.069(4)	5 440(4)	C(208)	4 762(10)	2 617(5)	10 863(4)
C(106)	-778(9)	550(4)	3 603(3)	C(209)	4 575(11)	1 528(5)	10 227(4)
C(107)	- 960(9)	1 315(4)	2 964(4)	C(210)	2 235(12)	1 349(5)	9 750(4)
. ,				· ·	. ,		.,
Compound	d 8						
Со	3 842(1)	7 402(1)	2 370(1)	C(8)	4 028(12)	8 706(8)	1 112(5)
Ν	2 739(9)	6 499(6)	5 413(4)	C(9)	4 981(14)	7 482(8)	742(5)
C(1)	2 421(12)	6 708(7)	4 260(5)	C(10)	3 119(12)	6 144(8)	840(5)
C(2)	3 468(13)	8 261(7)	3 962(5)	càń	1 505(11)	7 055(6)	6 285(5)
CÓ	5 890(12)	8 477(8)	3 677(5)	$\hat{\mathbf{C}}(12)$	2 054(12)	6 952(7)	7 380(5)
C(4)	6218(11)	7 019(7)	3 351(5)	C(13)	805(11)	7470(7)	8 268(5)
C(5)	3 027(13)	6 025(9)	3 473(6)	C(13)	-1.021(13)	8 114(7)	8 083(5)
C(6)	1.060(13)	6 545(8)	1 225(5)	C(15)	1 560(11)	8 238(7)	7 000(5)
C(0)	1.592(11)	8 114(7)	1 423(5)	C(15)	-1309(11) -344(11)	7 726(7)	6 117(5)
$\mathbf{C}(I)$	1 372(11)	0114(7)	1 425(5)	0(10)	544(11)	/ /20(7)	0117(3)
Compound	19						
Co	110(1)	2 351(1)	4 889(1)	C(11)	-6225(4)	2.047(3)	2 230(3)
Ň	-5131(3)	1 304(3)	2 794(3)	CUD	-5766(4)	3 495(4)	2 828(4)
C(1)	-3260(4)	1 088(3)	3 832(1)	C(12)	-6851(A)	A 222(A)	2 220(4)
C	2140(4)	1 1 2 9 (2)	$\frac{5}{4}\frac{5}{11}$			- 255(4) 2564(4)	2 200(4)
C(2)	4 1-77(4)	201(2)	2 545(2)	C(14)	-0 440(4)	5 504(4) 5 144(4)	524(4)
C(3)	-1110(4)	391(3)	3 343(3)		-8921(4)	2 144(4)	534(4)
U(4)	-1 009(4)	1 080(3)	2 /84(3)	C(16)	- / 846(4)	1 384(4)	1 064(4)
C(S)	-19/0(4)	2 224(3)	3 214(3)	C(21)	-5614(4)	- 194(3)	2 133(3)
C(6)	1 441(4)	3 378(4)	7 094(4)	C(22)	-6 512(4)	-688(3)	2 715(4)
C(7)	2 594(4)	2 543(4)	6 397(4)	C(23)	-6 996(4)	-2 128(3)	2 109(4)
C(8)	2 825(4)	3 006(3)	5 492(4)	C(24)	-6 593(4)	-3 067(4)	902(4)
C(9)	1 801(4)	4 103(3)	5 615(4)	C(25)	-5 706(4)	-2 579(4)	320(4)
C(10)	976(4)	4 359(3)	6 629(4)	C(26)	-5 225(4)	-1 139(4)	938(4)

 N_2O was evolved; evaporation or dilution with Et₂O gave no crystalline product only a solid the ¹H NMR spectrum of which was identical to that observed previously ¹ for [Ir(mes)₄]⁺. ¹H NMR (CD₃CN): δ 6.73 (s, br, 2 H, aromatic), 2.43 (s, 3 H, *p*-Me) and 2.19 (s, br, 6 H, *o*-Me). This green-black product was also formed using cyclohexane or toluene as solvent.

Crystal Structure Determinations of Compounds 2-4, 8 and 9.—The X-ray measurements were made on crystals handled under standard Schlenk procedures and mounted on glass fibres using silicone oil as both a coating and adhesive medium. The unit-cell and intensity data for crystals 2-4 were obtained at 150 K, those for the thermally unstable crystals 8 and 9 at 120 K,

J. CHEM. SOC. DALTON TRANS. 1994

- 7 G. D. Mendenhall, J. Am. Chem. Soc., 1974, 96, 5000.
- 8 L. Kuhn and E. R. Lippincott, J. Am. Chem. Soc., 1956, 78, 1820; D. J. Millen, C. N. Polydoropoulis and D. Watson, J. Chem. Soc., 1960, 689.
- 9 (a) J. E. Sheats, J. Organomet. Chem. Library, 1979, 7, 461; (b)
 R. D. W. Kemmitt and D. R. Russell, in Comprehensive Organometallic Chemistry, eds. G. Wilkinson, F. G. A. Stone and E. W. Abel, Pergamon, Oxford, 1982, vol. 5, sect. 34.44; (c) H. Bönnemann, Angew. Chem., Int. Ed. Engl., 1985, 24, 248.
- 10 (a) H. Kojima, S. Takahashi and N. Hagihara, J. Chem. Soc., Chem. Commun., 1973, 230; Tetrahedron Lett., 1973, 1991; (b) H. Kojima, S. Takahashi, H. Yamazaki and N. Hagihara, Bull. Chem. Soc. Jpn., 1970, 43, 2272.
- 11 (a) M. L. H. Green, L. Pratt and G. Wilkinson, J. Chem. Soc., 1959, 3753; (b) G. E. Herberich and J. Schwarzer, Angew. Chem., Int. Ed. Engl., 1970, 9, 897.
- 12 R. T. Morrison and R. N. Boyd, *Organic Chemistry*, 4th edn., Allyn and Bacon, Boston, 1983.
- 13 See (a) K. Lammerstein and B. V. Prasad, J. Am. Chem. Soc., 1993, 115, 2348; (b) A. T. Nielsen, in The Chemistry of the Nitro and Nitroso Groups, ed. H. Feuer, Interscience, New York, 1969, Part 1; (c) G. Klebe, K. H. Böhn, M. Marsch and G. Boche, Angew. Chem., Int. Ed. Engl., 1987, 26, 78 and refs. therein; (d) J. A. Cook, M. G. B. Drew and D. A. Rice, J. Chem. Soc., Dalton Trans., 1975, 1973; B. N. Diel and H. Hope, Inorg. Chem., 1978, 17, 4448; (e) J. Yarwood and W. T. Orville-Thomas, J. Chem. Soc., 1963, 5991.
- 14 C. J. Attridge, S. J. Baker and A. W. Parkins, Organomet. Chem. Synth., 1971, 1, 183.
- 15 See M. T. Reetz, T. Knauf, U. Minet and C. Bingel, Angew. Chem., Int. Ed. Engl., 1988, 27, 1373.
- 16 Y. Wakatsuki and H. Yamazaki, Synthesis, 1976, 26.
- 17 W. Kläui and C. E. Song, Inorg. Chem., 1989, 28, 3845.
- 18 H. Bönnemann, M. Radermacher, C. Krüger and H. J. Klaus, *Helv. Chim. Acta*, 1983, **66**, 185.
- 19 J. M. Birmingham, D. Seyferth and G. Wilkinson, J. Am. Chem. Soc., 1954, 76, 4179.
- N. Allinger and E. Eliel, Top. Stereochem., 1971, 6, 19; see also A. N. Sobolev, V. K. Belsky, I. P. Romm, N. Yu Chernikova and E. N. Guryanova, Acta Crystallogr., Sect. C., 1985, 41, 967; H. Bock, I. Göbel, C. Näther, Z. Havlas, A. Gavezzotti and G. Filippini, Angew. Chem., Int. Ed. Engl., 1993, 32, 1755.
- 21 J. J. Eisch and R. B. King, Organometallic Syntheses, ed. R. B. King, Academic Press, New York, 1965, vol. 1.
- 22 L. J. Bellamy, *The Infrared Spectra of Complex Molecules*, Chapman and Hall, London, 1975.
- 23 (a) M. J. Brooks and N. Jonathan, Spectrochim. Acta, Part A, 1969,
 25, 187; C. N. R. Rao, ref. 13(b), ch. 2, p. 88; (b) D. Hartley and
 M. J. Ware, J. Chem. Soc. A, 1969, 138.
- 24 A. A. Danopoulos, G. Wilkinson, B. Hussain-Bates and M. B.
- Hursthouse, J. Chem. Soc., Dalton Trans., 1991, 1855.
- 25 G. M. Sheldrick, University of Göttingen, 1992.
- 26 N. P. C. Walker and D. Stuart, Acta Crystallogr., Sect. A., 1983, 39, 158.
- 27 A. Karaulov, University of Wales, Cardiff, 1991.

Received 28th January 1994; Paper 4/00539B

using a Delft-Instruments FAST TV area detector diffractometer and graphite-monochromated Mo-K α radiation (λ = 0.71069 Å) following previously described procedures.²⁴ Structures 2, 3, 8 and 9 were solved by direct methods whilst that for compound 4 was derived via a Patterson solution. All refinements were by full-matrix least squares on F^2 (SHELX 92).²⁵ The data for compounds 2-4 were corrected for absorption using the program DIFABS²⁶ adapted for FAST geometry.²⁷ The non-hydrogen atoms were refined anisotropically and all of the hydrogen atoms in compounds 2, 3, 8 and 9 were located from the difference map and freely refined. The cp rings in all of the compounds were refined without geometrical restraints. However, in compound 4, whilst most of the hydrogens in the structure were experimentally located from the difference map, and were freely refined, three hydrogens were placed in theoretical positions: H(1) (on the bridging group) was refined as a tertiary hydrogen in the riding mode and with a fixed thermal parameter: the cp hydrogens H(207) and H(208) were refined as riding model aromatic C-H with the H atoms on the external bisector of the C-C-C angles. The crystal data and refinement details are listed in Table 2, fractional atomic coordinates in Table 3.

Additional material available from the Cambridge Crystallographic Data Centre comprises H-atom coordinates, thermal parameters and remaining bond lengths and angles.

Acknowledgements

We thank Mr. Tye, King's College, London for mass spectra, Professors W. B. Motherwell and C. W. Rees for discussions, Johnson Matthey plc for a loan of iridium and the SERC (M. B. H.) for provision of X-ray facilities.

References

- 1 R. S. Hay-Motherwell, G. Wilkinson, B. Hussain-Bates and M. B. Hursthouse, J. Chem. Soc., Dalton Trans., 1992, 3477.
- 2 F. Bottomley, in *Reactions of Coordinated Ligands*, ed. P. S. Braterman, Plenum, New York, 1986, vol. 2, ch. 3, sect. 6; F. A. Cotton and G. Wilkinson, *Advanced Inorganic Chemistry*, 5th edn., Wiley, New York, 1988, p. 1311; A. Guttierrez, G. Wilkinson, B. Hussain-Bates and M. B. Hursthouse, *Polyhedron*, 1990, 9, 2081.
- 3 R. S. Hay-Motherwell, G. Wilkinson, B. Hussain-Bates and M. B. Hursthouse, *Polyhedron*, 1993, 12, 2009.
- 4 G. Wilkinson, J. Am. Chem. Soc., 1952, 74, 6146.
- 5 (a) F. Bottomley and I. J. B. Lin, J. Chem. Soc., Dalton Trans., 1981, 271; F. Bottomley, J. Darkwa and P. S. White, J. Chem. Soc., Dalton Trans., 1985, 1435; Organometallics, 1985, 4, 691; (b) F. Bottomley, personal communication.
- 6 D. A. Muccigrosso, F. Mares, S. E. Diamond and J. P. Solar, *Inorg. Chem.*, 1983, 22, 960 and refs. therein; S. Bhaduri, B. F. G. Johnson, A. Pickard, P. R. Raithby, G. M. Sheldrick and C. I. Zuccaro, *J. Chem. Soc., Chem. Commun.*, 1977, 354; S. Bhaduri, I. Bratt, B. F. G. Johnson, A. Khair, J. A. Siegel, R. Waller and C. I. Zuccaro, *J. Chem. Soc., Dalton Trans.*, 1981, 234.