CHEMISTRY A European Journal



Accepted Article Title: Terminal Uranium(V/VI)-Nitride Activation of Carbon Dioxide and Disulfide: Factors Governing Diverse and Well-Defined Cleavage and Redox Reactions Authors: Stephen Liddle, Peter Cleaves, Christos Kefalidis, Benedict Gardner, Floriana Tuna, Eric McInnes, William Lewis, and Laurent Maron This manuscript has been accepted after peer review and appears as an Accepted Article online prior to editing, proofing, and formal publication of the final Version of Record (VoR). This work is currently citable by using the Digital Object Identifier (DOI) given below. The VoR will be published online in Early View as soon as possible and may be different to this Accepted Article as a result of editing. Readers should obtain the VoR from the journal website shown below when it is published to ensure accuracy of information. The authors are responsible for the content of this Accepted Article. To be cited as: Chem. Eur. J. 10.1002/chem.201605620 Link to VoR: http://dx.doi.org/10.1002/chem.201605620

Supported by ACES



Terminal Uranium(V/VI)-Nitride Activation of Carbon Dioxide and Disulfide: Factors Governing Diverse and Well-Defined Cleavage and Redox Reactions

Peter A. Cleaves,^[a] Christos E. Kefalidis,^[b] Benedict M. Gardner,^[a] Floriana Tuna,^[c] Eric J. L. McInnes,^[c] William Lewis,^[d] Laurent Maron,^{*[b]} and Stephen T. Liddle^{*[a]}

^[a] School of Chemistry, The University of Manchester, Oxford Road, Manchester, M13 9PL, UK. *Email: steve.liddle@manchester.ac.uk.

^[b] LPCNO, CNRS & INSA, Université Paul Sabatier, 135 Avenue de Rangueil, Toulouse 31077, France. *Email: laurent.maron@irsamc.ups-tlse.fr.

^[c] School of Chemistry and Photon Science Institute, The University of Manchester, Oxford Road, Manchester, M13 9PL, UK.

^[d] School of Chemistry, The University of Nottingham, University Park, Nottingham, NG7 2RD, UK.

Abstract

We report the reactivity of terminal uranium(V/VI)-nitrides with CE_2 (E = O, S), where we observe well-defined C=E cleavage followed by zero-, one-, and two-electron redox events. The uranium(V)-nitride $[U(Tren^{TIPS})(N)][K(B15C5)_2]$ [1, $Tren^{TIPS} = N(CH_2CH_2NSiPr^i_3)_3$; B15C5 =benzo-15-crown-5] reacts with CO₂ to give $[U(Tren^{TIPS})(O)(NCO)][K(B15C5)_2]$ (3), whereas the uranium(VI)-nitride $[U(Tren^{TIPS})(N)]$ (2) reacts with CO₂ to give isolable $[U(Tren^{TIPS})(O)(NCO)]$ (4); complex 4 rapidly decomposes to known $[U(Tren^{TIPS})(O)]$ (5) with concomitant formation of N_2 and CO proposed, with the latter trapped as a vanadocene adduct. In contrast, 1 reacts with CS_2 to give $[U(Tren^{TIPS})(\kappa^2-CS_3)][K(B15C5)_2]$ (6), 2, and $[K(B15C5)_2][NCS]$ (7), whereas 2 reacts with CS_2 to give [U(Tren^{TIPS})(NCS)] (8) and "S", with the latter trapped as Ph₃PS. Calculated reaction profiles reveal outer-sphere reactivity for uranium(V) but inner-sphere mechanisms for uranium(VI); together the experimental and theoretical data reproduce the experimental outcomes and suggest that despite the wide divergence of products the initial activation of CE_2 follows mechanistically related pathways, providing insight into the factors of uranium oxidation state, chalcogen, and NCE groups that govern the subsequent divergent redox reactions that include common one-electron reactions and a less-common two-electron redox event. This work highlights that caution is warranted utilising CS_2 as a reactivity surrogate for CO_2 .

Introduction

The activation and cleavage of carbon dioxide and carbon disulfide, CE_2 (E = O, S), are burgeoning areas given current environmental and sustainability agendas.^[1] The former is an abundant, lowcost, and renewable C₁-source for fine chemicals and fuels production,^[2] and the latter is a versatile building block in organic synthesis that is often used to model CO₂ reactivity.^[3] In recent years felements,^[4] and especially uranium,^[5,6] have emerged as effective, appealing candidates for the reductive activation of CE_2 ,^[7] but studies of such activations remain in the shadow of those involving transition metals.^[1-3,8] Metal-nitride chemistry is now well-developed for transition metals with a wide range of reactivity types.^[9] However, although electron-rich terminal M=N triple bonds are primed for bond metathesis chemistry, and are therefore excellent candidates for nitrogen-atom transfer reactions with heteroallenes,^[10] there are few examples of transition metal-nitride reactivity with CE₂ molecules.^[8,9] Germane to this work, $[V{N(R)(Ar)}_3(N)Na]_2$ (R = Bu^t or Ad; Ar = 3,5-Me₂C₆H₃) reacts with CE₂ to give $[V{N(R)(Ar)}_3(NCE_2)][Na(THF)_2]$, which for E = O reacts no further but for E = S and R = Bu^t extrudes NaNCS following bond metathesis to give $[V{N(Bu^t)(Ar)}_3(S)]$.^[10d] In contrast, $[Nb{N(R)(Ar)}_3(N)Na]_2$ reacts with CO₂ to give $[Nb{N(R)(Ar)}_3(NCO_2)][Na(THF)_n]$, but C-O splitting only occurs with addition of external electrophiles (such as Ac₂O, O{C(O)CF₃}₂, or Bu^tCOCl); this opens the door to elimination of the CO₂-derived oxide as RCO₂⁻ and decarbonylation of the resulting niobium-cyanate to -nitride to close a synthetic cycle.^[10b]

Regarding uranium, a promising candidate in this arena given its redox chemistry,^[5,6] it was not until recently that terminal uranium-nitride complexes became available under ambient conditions,^[11] and so their inherent reactivity patterns are yet to be established. Indeed, there are no other classes of terminal f-block nitrides available for reactivity studies,^[12] and although a number of bridged diuranium-nitrides are known,^[6a,b,13] their reactivity is also almost unknown so determining and comparing the reactivity of terminal to bridging uranium-nitrides would present opportunities to better understand their intrinsic reactivities. Furthermore, uranium sometimes exhibits reactivity that parallels transition metal chemistry, but also often exhibits divergent reactivity, and so it is of interest to compare terminal uranium- and transition metal-nitride reactivity. Recently, we showed that the terminal uranium-nitrides [U(Tren^{TIPS})(N)][K(B15C5)₂] [1, Tren^{TIPS} = N(CH₂CH₂NSiPrⁱ₃)₃; B15C5 = benzo-15-crown-5]^[14] and [U(Tren^{TIPS})(N)] (2)^[11a] react with CO by reductive carbonylation,^[14,15] to give cyanate derivatives followed by complete denitrification and *N*-atom transfer.^[16] Whilst terminal M≡N triple bond reactivity with CE₂ has potential for C-N bond formation chemistry, since so little reactivity has been reported the factors that govern their reactivity are yet to be elucidated. With the reactivity of **1** and **2** towards CO suggesting a rich seam of *N*-atom transfer reactivity we turned our attention to the reactivity of **1** and **2** towards CO_2 and CS_2 .

Here, we report the first benchmarking study of the reactivity of terminal f-block-nitrides towards CO₂ and CS₂ with uranium(V) and (VI). In all cases we find facile CE₂ cleavage chemistry and that the split E and NCE products react further in zero-, one- and two-electron redox processes. In contrast to the d-block, one-electron redox chemistry generally dominates in the f-block, and although becoming more common two-electron redox process are unusual so understanding the factors that promote such reactivity is important given the fundamental role electron transfer plays in chemical reactivity.^[11,13a,b,c,e,g,17] Calculated reaction profiles reveal outer-sphere reactivity for uranium(V) but inner-sphere mechanisms for uranium(VI) and provide insight into the factors that govern the observed reactivity. Thus this work furthers our understanding of the nature and reactivity of terminal metal-nitride reactivity, permitting mechanistic similarities and differences between terminal d- and f-block and bridging and terminal f-block metal-nitride reactivity to be determined. Lastly, the divergent reactivities observed show that care should be taken extrapolating surrogate reactivity between CO₂ and CS₂.

Results and Discussion

Experimental terminal uranium(V)-nitride reactivity with CO_2



Scheme 1. a) Reactions of the terminal uranium(V)-nitride complex 1 with CO₂ and CS₂ to give the ate uranium(V)-oxo-cyanate 3 and the uranium(IV)-trithiocarbonate 6, uranium(VI)-nitride 2, and potassium bis(crown) thiocyanate 7, respectively. b) Reactions of the terminal uranium(VI)-nitride complex 2 with CO₂ and CS₂ to give the uranium(VI)-oxo-cyanate 4, which decomposes to the uranium(V)-oxo 5, N₂, and CO₂, and the uranium(IV)-thiocyanate 8 with expulsion of elemental "S" that can be trapped by Ph₃P as Ph₃PS. A control reaction between 8 and Ph₃P confirms that the S does not originate from the NCS group of 8.

Exposure of a degassed toluene solution of **1** to an atmosphere of CO₂ at -78 °C resulted in a colour change from yellow-brown to red-brown. After work-up and recrystallisation from benzene the orange complex formulated as the oxo-cyanate derivative [U(Tren^{TIPS})(O)(NCO)][K(B15C5)₂] (**3**) was isolated in 36% crystalline yield, Scheme 1a, which reflects the high solubility of this complex; analysis of the mother liquor by NMR spectroscopy suggests the formation of **3** is essentially quantitative (Figure S1).

The ¹H NMR spectrum of **3** spans 16 to -3 ppm, which suggests the presence of uranium(V) and the resonances are broad so inferences about the symmetry of the Tren^{TIPS} ligand (C_s - or C_3 -

coordination) could not be discerned; however, the ²⁹Si NMR spectrum exhibits two resonances (-27.1 and -27.8 ppm), suggesting that the anion component of **3** possesses approximate C_s rather than $C_{3\nu}$ symmetry.



Figure 1. Variable temperature SQUID magnetometry of the effective magnetic moment (μ_B) of 3, 6, and 8 in the temperature range 2-298 K. Lines are to guide the eye only and have no further significance.

The formation of cyanate and oxo functional groups is supported by inspection of the FTIR spectrum of **3**, which exhibits absorptions at 2194, 800, and 742 cm⁻¹, with the first attributed to cyanate and the latter two to oxo stretches, and these compare very well to calculated frequencies of 2195 and 799 and 747 cm⁻¹, respectively. The two stretches for the oxo result from coupling to the uranium-amine stretch to give asymmetric and symmetric combinations. The cyanate stretch compares well to a value of 2187 cm⁻¹ for [U(Tren^{TIPS})(NCO)] (**D**),^[14] prepared by reductive carbonylation of **2**, and those found in related examples of uranium cyanate complexes (ca 2122-2201 cm⁻¹).^[5a,17i]

The UV/Vis/NIR spectrum of **3** (Figure S8) exhibits a characteristic absorption at 6693 cm⁻¹ ($\epsilon = 43$ M⁻¹ cm⁻¹), consistent with uranium(V), as well as three other absorptions in that region resulting from splitting of the 7/2 excited state of ²F by the *O*_h crystal field into four levels (Γ_7 ', two Γ_8 ' and

6

 Γ_6 in double-group notation).^[18] The uranium(V) assignment is confirmed by variable temperature SQUID magnetometry of **3** (Figure 1), which reveals a magnetic moment of 1.86 μ_B at 298 K, that compares well to a solution magnetic moment of 1.69 μ_B at 298 K; this changes little until 50 K where it falls more rapidly to 1.20 μ_B at 2 K consistent with the presence of uranium(V) that is a magnetic doublet.^[19] Final confirmation of the presence of uranium(V) in **3** is provided by EPR spectroscopy (see Figure S15) which reveals a strong absorption at g = 2.85 at 5 K, which is consistent with a Kramers uranium(V) ion; uranium(IV) would be expected to be EPR-silent under these measurement conditions.^[20]



Figure 2. Molecular structure of **3** at 120 K with displacement ellipsoids at 50% and hydrogen atoms omitted for clarity. Only one of the two independent ate pairs in the crystallographic asymmetric unit is shown. The other pair is essentially identical with statistically indistinguishable bond lengths and angles. Selected bond lengths (Å) for the ate pair shown: U1-N1, 2.316(7); U1-N2, 2.302(7); U1-N3, 2.342(7); U1-N4, 2.517(7); U1-N5, 2.404(8); U1-O1, 1.848(7); N5-C34, 1.199(13); C34-O2, 1.163(13).

In order to confirm the structure of 3 we determined the molecular structure by single crystal X-ray diffraction (Figure 2). Complex 3 retains the separated ion pair motif from 1, and the structure reveals a pseudo-octahedral uranium centre where the oxo is *trans* to the Tren-amine and the cyanate group is *trans* to a Tren-amide and mutually *cis* to the oxo. Competitive refinement of the

diffraction data suggest that the cyanate is *N*- not *O*-bound, and this is supported by DFT calculations that show the former isomer to be more stable than the latter by 9 kcal/mol. The U-O, U-N_{cyanate}, and U-N_{amine} bond lengths of 1.848(7), 2.404(8), and 2.517(7) Å, respectively, are typical of such linkages,^[14,17d,21] though the latter is towards the lower end of the range of Tren-N_{amine}-U distances, which suggests there might be a weak inverse-*trans*-influence in the N \rightarrow U=O unit.^[22] The U-N_{amide} bond lengths span the range 2.302(7)-2.342(7) Å and can be considered long for uranium(V) when compared to Tren uranium complexes, which most likely reflects the electron rich nature of this anionic fragment of the compound. The O-U-N_{amine} and N_{amide}-U-N_{cyanate} angles deviate from the octahedral ideal at 168.0(3) and 168.3(3)°, respectively, reflecting the steric demands of the Tren^{TIPS} ligand.

Experimental terminal uranium(VI)-nitride reactivity with CO₂

When a degassed toluene solution of $[U(\text{Tren}^{\text{TIPS}})(N)]$ (2) is exposed to an atmosphere of CO₂ at -78 °C the brown solution turns wine-red (Scheme 1b). After quick work-up a red solid is obtained, and on the basis of the data presented below we formulate this red solid as $[U(\text{Tren}^{\text{TIPS}})(O)(\text{NCO})]$ (4), obtained in essentially quantitative yield. The ¹H NMR spectrum of the red solid retains the appearance of a diamagnetic complex, spanning 0-10 ppm, suggesting the retention of uranium(VI), but it is not particularly informative. We were unable to obtain the potentially more informative ²⁹Si NMR spectrum due to rapid decomposition of **4** in solution.

The FTIR spectrum of **4** exhibits strong absorptions at 2176 and 827 cm⁻¹, which are assigned as cyanate and oxo stretches, respectively. The former absorption compares well to the cyanate stretch for **3** and is supported by a calculated stretching frequency of 2198 cm⁻¹. Interestingly, the oxo stretch for **4**, unlike **3**, is now decoupled from the U-N_{amine} vibration resulting in only one absorption being observed; this is also reproduced by the frequency calculation, which yields a calculated oxo stretching frequency of 812 cm⁻¹. Although the formation of **4** seems certain on the

basis of **3** and computationally supported spectroscopy, despite exhaustive attempts we have been unable to obtain a crystal structure of **4** due to decomposition during recrystallisation attempts; however, this is consistent with the computed reaction profile (see below). Attempts to recrystallise **4** resulted in isolation of the known mono-oxo uranium(V) complex $[U(Tren^{TIPS})(O)]$ (**5**),^[17d] as evidenced by single crystal X-ray diffraction and FTIR data which are devoid of any absorptions in the cyanate region and that show the characteristic oxo stretch of **5** at 910 cm⁻¹. Monitoring the conversion of **4** to **5** by ¹H NMR spectroscopy (for a representative spectrum see Figure S2) reveals clean, quantitative conversion over 24 hours, with **4**:**5** ratios of 1:0 (<30 minutes), 2:1 (2 hours), 1:9 (5 hours), 0:1 (20 hours).

The isolation of 5 suggests that a cyanate radical has been ejected from the coordination sphere of uranium; this could react with solvent to abstract a proton to give HNCO, or dimerise to the putative molecule diisooxocyan followed by facile decomposition to N₂ and CO.^[23] Although the bond dissociation enthalpy of HNCO is high (~110 kcal/mol),^[24] suggesting that it could be formed - with toluene implicated as the H-source - we have not observed at any stage of the reaction any IR absorbances consistent with its formation;^[25] further, production of HNCO ($pK_a = 3.7$)^[26] would be anticipated to result in extensive decomposition of 5, which is not the case experimentally. To probe the possibility that diisooxocyan is formed, we reacted 50% ¹⁵N_{nitride}-labelled 2 with CO₂, but could not detect a ¹⁵N NMR resonance for dissolved N₂. Due to diamond absorbance from the probe-tip in the region 2000-2250 cm⁻¹ we could not observe CO formation in solution by ReactIR, and sampling of the gas headspace resulted in IR spectra dominated by solvent vapour. Unfortunately, sampling and MS-analysis of the headspace from the decomposition of 4 was inconclusive, likely due to concentration effects and the presence of N₂ even in high purity Ar gas. However, vacuum transfer of the headspace atmosphere from reactions that generate 4 onto a thawing toluene solution of vanadocene gave a weak absorption at 1895 cm⁻¹ in the resulting IR spectrum, which is consistent with the formation of a vanadocene carbonyl.^[5k] Attempts to trap the putative cyanate

radical were inconclusive (see SI). Attempts to oxidise **3** to **4** with numerous oxidants gave intractable product mixtures, but oxidation of $\mathbf{D}^{[14]}$ with trimethylamine-*N*-oxide or 4-methylmorpholine-*N*-oxide gave mixtures of **D**:**4**:**5** in 6:1:1 and 3:2:2 ratios, respectively, which is not inconsistent with oxidation of **D** to **4** followed by decomposition to **5** (Figures S6 and S7).

Experimental terminal uranium(V)-nitride reactivity with CS_2

The reactions of CS₂ with **1** and **2** proceed very differently, in terms of the final products, to the CO₂ reactivity (Scheme 1a). Addition of one equivalent of CS₂ to a solution of pentavalent **1** in toluene at -78 °C results in a colour change from yellow-brown to brown. After removal of solvent a brown oil is formed, which after washing with hexanes is isolated as a brown solid. Work-up of the hexane washings results in isolation of hexavalent **2**, suggesting that overall disproportionation has occurred. Extraction of the brown solid with aromatic solvent and recrystallisation resulted in the isolation of an orange complex formulated as the uranium(IV)-trithiocarbonate complex [U(Tren^{TIPS})(κ^2 -CS₃)][K(B15C5)₂] (**6**) in crystalline yields of ca 4%; this yield reflects the oily nature of this complex and although it is low it is reproducible. The formulation of **6** suggests the presence of a uranium(IV) ion, which would account for the remainder of the disproportion balance together with **2**. Inspection of crude reaction mixtures by ¹H NMR spectroscopy (Figure S4) suggests that **2** and **6** are formed in a 1:1 ratio, as would be expected, and together they constitute ~90% of the reaction mixture (ignoring the thiocyanate component, see below) and thus **2** and **6** constitute the major reaction products.

The ¹H NMR spectrum of **6** (Figure S3) spans the range 9 to -26 ppm in-line with the uranium(IV) assignment. Further confirmation that **6** is a uranium(IV) complex comes from solid-state SQUID magnetometry (Figure 1), which reveals a magnetic moment of 2.4 μ_B at 298 K, whose value decreases with decreasing the temperature to 0.6 μ_B at 2 K and tending to zero.^[19]

The solid that remains after the toluene or benzene extraction was examined by FTIR spectroscopy revealing strong absorbances at 2060 and 2052 cm⁻¹ which are assigned as thiocyanate stretchs.^[6c,f,j] For mass balance this is proposed to be $[K(B15C5)_2][NCS]$ (7), confirmed by comparison to an authentic sample prepared independently from KNCS and B15C5 that also exhibits two thiocyanate stretches in its FTIR spectrum.



Figure 3. Molecular structure of **6** at 120 K with displacement ellipsoids at 50% and hydrogen atoms omitted for clarity. Selected bond lengths (Å): U1-N1, 2.277(3); U1-N2, 2.290(3); U1-N3, 2.296(3); U1-N4, 2.670(3); U1-S1, 2.8520(10); U1-S2, 2.8415(8); S1-C34, 1.729(4); S2-C34, 1.731(4); S3-C34, 1.665(4).

The molecular structure of **6** was determined by single crystal X-ray diffraction to confirm its identity (Figure 3). The U-N_{amine} (2.670(3) Å) and U-N_{amide} (range: 2.277(3)-2.296(3) Å) distances are typical of U-N distances in Tren-uranium(IV) complexes.^[21] Complex **6** the first example where the $(CS_3)^{2^-}$ dianion is terminally bound to one f-element centre, and is only the third example of a f-block trithiocarbonate complex.^[6c,f,j] Befitting the unique coordination mode of the $(CS_3)^{2^-}$ dianion in f-block chemistry, the U-S bond lengths of 2.8415(8) and 2.8520(10) Å are shorter than in complexes where the $(CS_3)^{2^-}$ dianion bridges (range: 2.9488(19)-3.130(2) Å). The κ^2 -coordination mode of the $(CS_3)^{2^-}$ dianion appears to localise negative charge on the two coordinated sulfur-atoms, resulting in some localisation of a C=S double bond on the remaining uncoordinated sulfur-

carbon linkage; this is evidenced by a terminal C=S bond distance of 1.665(4) Å compared to the other two C-S bond lengths of 1.731(4) and 1.729(4) Å.

Experimental terminal uranium(VI)-nitride reactivity with CS₂

Since the uranium(V)-nitride **1** undergoes, overall, disproportion reactivity with CS_2 we were interested to determine how the analogous uranium(VI)-nitride **2** would react since it lacks the requisite valence electron to engage in disproportionation chemistry. When a toluene solution of **2** is treated with CS_2 over six days a new uranium-containing product is formed, Scheme 1b. Work-up and recrystallisation from pentane affords the green uranium(IV)-thiocyanate complex $[U(Tren^{TIPS})(NCS)]$ (**8**) in 51% crystalline yield.

The ¹H NMR spectrum of **8** (Figure S5) spans the range 9 to -37 ppm, consistent with the uranium(IV) formulation and the ²⁹Si NMR spectrum exhibits only one resonance at 2.4 ppm that is consistent with a $C_{3\nu}$ symmetric complex. The FTIR spectrum exhibits a strong absorbance at 2013 cm⁻¹ and this is typical of uranium(IV)-thiocyanate stretches.^[27] In support of the uranium(IV) assignment, variable temperature SQUID magnetometry data of **8** reveal a magnetic moment of 2.52 μ_B at 298 K (2.91 μ_B by Evans method in solution) that decreases monotonously down to 0.24 μ_B at 2 K and tending to zero in-line with the magnetic singlet ground state of uranium(IV) at low temperature (Figure 1).^[19]



Figure 4. Molecular structure of **8** at 120 K with displacement ellipsoids at 50% and hydrogen atoms omitted for clarity. Only one of the two parts of the disordered NCS unit are shown. Selected bond lengths (Å): U1-N1, 2.238(3); U1-N2, 2.240(3); U1-N3, 2.237(3); U1-N4, 2.626(4); U1-N5, 2.377(4); N5-C34, 1.181(11); C34-S1, 1.589(9).

In order to confirm the identity of **8** we prepared it independently from $[U(\text{Tren}^{\text{TIPS}})(\text{Cl})]$ and KSCN (isolated in 53% yield) and the characterisation data are in agreement. The formation of **8** suggests that elemental sulfur is the by-product of this reaction, presumably extruded from the putative complex $[U(\text{Tren}^{\text{TIPS}})(S)(\text{NCS})]$ analogously to **4**. Therefore, either post-reaction or all in one-pot, we treated the reactions with Ph₃P and observed the formation of Ph₃PS by ³¹P NMR spectroscopy (³¹P NMR: Ph₃P = -5.2; Ph₃PS = 42.8 ppm). In a control experiment to rule out the abstraction of sulfur from **8** to give the plausible cyanide product $[U(\text{Tren}^{\text{TIPS}})(\text{CN})]$, we treated **8** with Ph₃P and found that no reaction occurs, even on extended reflux in toluene.

Final confirmation of the identity of **8** was obtained by examination of its single crystal X-ray diffraction structure (Figure 4), which unambiguously reveals a complex of approximate $C_{3\nu}$ symmetry with an *N*-bound thiocyanate ligand at uranium. The metrical data for **8** are unexceptional.^[27]

Given this diverse range of reaction outcomes, we modelled the reaction profiles by using DFT calculations (at B3PW91 level of theory) to provide useful insights for the mechanisms and in order to understand why such a diverse range of reactivity occurs and what the controlling factors are. In all cases the computed energy pathways were found to be kinetically accessible and thermodynamically favourable and completely consistent with experimental outcomes.







Figure 5. Energy profile (at 298 K) for the gas phase reaction of CO_2 with the uranium(V)-nitride 1 computed at the B3PW91 level of theory. [U] = [U(Tren^{TIPS})].

The computed energy pathway for the reaction of complex **1** with CO₂ (Figure 5) is reminiscent of the outer-sphere CO₂ addition reported by Cummins.^[10a,d] In other words, the highly nucleophilic nitride attacks the incoming CO₂ molecule without any coordination to the poorly electrophilic U^V centre. Indeed, a very low-lying outer-sphere addition transition state (TS), **TS_{A-B}**, was located (enthalpy barrier of 1.4 kcal/mol⁻¹), where the CO₂ molecule bends to allow the π^* of CO₂, that is mainly located at the 2p-orbital at the carbon atom, to overlap with the σ -lone pair of the nitride. This outer-sphere TS is lower in energy than the classical [2 + 2]-cycloaddition route ($\Delta H^{\neq} = 5.1$ kcal/mol⁻¹) (Figure S20). **TS_{A-B}** leads to the formation of a η^1 -carbamate intermediate, that readily isomerises ($\Delta H^{\neq} = 0.2$ kcal/mol⁻¹) to a stable metal-based-four-membered ring (-23.1 kcal/mol⁻¹).

anuscri

cepte

Finally, following a low-lying transition state, TS_{I-3} , involving bond metathesis (-20.8 kcal/mol⁻¹), the cleavage of the C-O bond occurs to give the oxo-cyanate that is 57.1 kcal/mol⁻¹ more stable than the initial complex 1.



Figure 6. Energy profile (at 298 K) for the gas phase reaction of CO_2 with the uranium(VI)-nitride 2 computed at the B3PW91 level of theory. $[U] = [U(Tren^{TIPS})]$.

In the same way as for complex 1, the energy profile was computed for the reaction of complex 2 with CO₂ (Figure 6) at the same level of theory. Unlike 1, the outer-sphere mechanism is not found to be operative. Indeed, the U^{VI} centre is more electrophilic and at the same time the nitride is less nucleophilic compared to the U^V case, so that the CO₂ molecule needs to be activated through coordination to the Lewis acid centre to react. Hence, the reaction profile evolves firstly by a [2 + 2]-cycloaddition step, surmounting an activation barrier of 12.1 kcal/mol⁻¹ and leading to the formation of the intermediate **II**. The geometry of the latter is close to the intermediate **I** in uranium(V)-nitride reactivity (Fig. 5), but energetically more unstable by almost 22 kcal/mol⁻¹. The following step corresponds to the C-O bond-breaking which proceeds via a bond metathesis transition state **TS**_{1.3}- (Δ H[#] = 3.5 kcal/mol⁻¹) reminiscent of activated alkene metathesis^[28] to give the oxo-cyanate (-38.1 kcal/mol⁻¹). Although **4** is an enthalpic minimum with respect to the oxo **5** (-25.1 kcal/mol⁻¹), when the Gibbs free energy term is considered the formation of **5** from **4** becomes more favourable by 5.7 kcal/mol⁻¹ at -43.8 kcal/mol⁻¹ due to the entropic gain from N₂ and CO elimination. This is in-line with the strong electrophilic character of U^{VI} that prevents it from

15

tightly binding to the cyanate radical (U-N distance of 2.36 Å in 4 vs. 2.27 Å for 3^{-}) as it prefers a strongly anionic ligand. This finding is in-line with the initial experimental isolation of 4 but conversion to 5 on attempted recrystallisation of the former.

The initial reactivity of **2** with CO₂ to give **4** clearly follows the same pattern as **1** reacting to give **3**. However, although **3** is stable in the absence of O₂ or H₂O in solution or the solid state, **4** is stable for only ca 30 minutes in solution and the cyanate group is rapidly lost. The formation of **5** suggests that the cyanate group is ejected as a radical in a homolytic U-N_{cyanate} bond cleavage step which would provide the requisite electron for the one-electron reduction of the uranium(VI) centre. Comparing the computed barriers for the first step is informative regarding the nucleophilic nature of the nitride in **1** and **2**. The computed barrier for **2** (12.1 kcal/mol) although relatively low is > 30 kcal/mol higher relative to the equivalent point computed for **1**. This indicates that the nitride is of lower nucleophilic character in **2** than in **1**,^[29,30] which can be attributed to the shorter U-N distance in the former than in the latter, inducing a stronger U-N interaction. On the other hand, the barrier for the C-O bond-breaking in the carbamate intermediates are similar for both complexes, indicating that the oxophilicity of uranium is affected very little by the oxidation state in these steps. However, the final oxo-cyanate **4** further evolves by ejecting the coordinated cyanate group yielding **5**.

The question then arises as to the fate of the cyanate radical; H-abstraction to give HNCO can seemingly be ruled out, but the coupling of cyanate radicals to give the putative molecule diisooxocyan followed by facile decomposition to N_2 and CO has been computationally described,^[23] and seems a credible, if not conclusively proven, fate for the cyanate radical given the indirect supporting evidence by FTIR of the formation of vanadocene carbonyl as a CO-trap. This step is computed to be endothermic in terms of enthalpy, but is favoured by entropic effects (Figure 6) and indeed becomes exergonic overall when entropy is factored in (although it is known that the

computation of entropy is not very accurate).^[31] The difference between the reactivity of **3** and **4** is attributed to steric hindrance around the metal centre^[32] as well as electronic effects in the latter. Indeed, the shorter distances in the U^{VI} complexes makes the geometry more compact around the metal centre and also induces stronger interactions with the oxo ligand with respect to the Tren^{TIPS} ligand. Therefore, in 4 the uranium centre is less available for interacting with the radical cvanate that undergoes the entropically-driven radical coupling reaction to yield $N_2 + 2CO$.^[33] This is reflected in the calculations where 4 is more stable from its start point by 38.1 kcal/mol, the corresponding uranium(V) component of **3** is more stable than its start point by a much larger 57.1 kcal/mol, which must to a significant extent reflect the size differences between uranium(V) and uranium(VI) where the former is larger so an additional ligand in the coordination sphere of uranium brings greater stability through the formation of another U-N bond with only a minimal thermodynamic penalty to pay from steric aspects. We further suggest that this reactivity can be rationalised on the basis that uranium(VI) is more oxidising than uranium(V) and so although the uranium(V) component in 3 is stable to such homolytic extrusion of cyanate this is not the case for uranium(VI) in 4. A parallel can be drawn here to the chemistry of terminal uranium-nitrides under photolytic conditions,^[11a,13c] where the uranium(VI)- nitride activates ancillary ligand C-H bonds but uranium(V) analogues do not; this can be rationalised on the basis of the oxidising power of uranium(VI) compared to uranium(V). The fact that the cyanate linkage is lost in preference to the oxo group is testament to the highly favourable nature of the U=O bond, but also that one-electron redox events are far more common in f-element chemistry than two-electron redox events; if the uranium were to be reduced by two units to generate [U(Tren^{TIPS})(NCO)] instead this would require the elimination of oxidised 'O' which would be expected to be thermodynamically far uphill given the hard nature of electronegative oxide and its ability to stabilise high oxidation states at metals, i.e. resist oxidation. We note there is precedent for elimination of the cyanate radical from uranium,^[34] where in that report the reaction takes a different course because the cyanate radical can

be intercepted by low valent and electron rich uranium(III), which is not an available pathway in the reaction with hexavalent **2**.



Computed reaction profiles of terminal uranium-nitride reactivity with CS₂



Figure 7. Energy profile (at 298 K) for the gas phase reaction of CS_2 with the uranium(V)-nitride 1 computed at the B3PW91 level of theory. [U] = [U(Tren^{TIPS})]. Extrusion of NCS· from VI and subsequently the oxidation of 1 by NCS· to give 2 and 7 is highly favoured (-97.6 kcal/mol) (see Figures S17 and S19).

As for the reaction of 1 with CO₂, the reaction of 1 with CS₂ (Figure 7) begins by a low energy outer-sphere addition of CS₂, that after isomerisation yields the η^2 -N,S carbamate intermediate (-33.8 kcal/mol⁻¹). From this intermediate, rather than the C-S bond breaking, which was found to be higher in energy (Figure S21), a kinetically facile outer-sphere electrophilic attack of a CS₂ molecule to the pendant sulfur atom of the carbamate intermediates occurs (enthalpy barrier of only 2.0 kcal/mol⁻¹). Such process does not work for the reaction with CO₂ as every attempt to locate this kind of TS leads to the departure of the incoming CO₂ molecule. This yields, after an isomerisation process ($\Delta H^{\neq} = 11.6$ kcal/mol⁻¹), to a metal-based-six-membered ring intermediate (-36.5 kcal/mol⁻¹). In this intermediate, a C-S bond is strongly activated and thus easily broken ($\Delta H^{\neq} = 1.4$ kcal/mol⁻¹) to form an isothiocyanate adduct to the thiocarbonate complex. In the last step, the driving force

for the extrusion of NCS and the subsequent oxidation of 1 by NCS to give 2 and 7 is mainly due to the resulted high exothermicity of such sequence of events $(-97.6 \text{ kcal/mol}^{-1})$ as would be expected for two open shell radicals comproportionating to two closed shell species.



Figure 8. Energy profile (at 298 K) for the gas phase reaction of CS_2 with the uranium(VI)-nitride **2** computed at the B3PW91 level of theory. [U] = [U(Tren^{TIPS})].

Finally, the reactivity of complex **2** with CS₂ was investigated (Figure 8). Once again, like the analogous CO₂ reaction the outer-sphere mechanism does not operate since the formed product of this step is endothermic (Figure S22). Therefore, the two first steps of the reaction are similar to that found for the CO₂ reactivity previously described, that is a [2 + 2]-cycloaddition via TS_{2-VI} ($\Delta H^{\neq} = 23.4 \text{ kcal/mol}^{-1}$) yielding a thiocarbamate complex (-3.7 kcal/mol⁻¹) followed by a low-lying bond metathesis transition state TS_{VI-VII} ($\Delta H^{\neq} = 5.1 \text{ kcal/mol}^{-1}$) with C-S bond disruption to form a terminal sulfido-thiocyanate complex VII (-19.6 kcal/mol⁻¹). Extrusion of sulfur by the PPh₃, computed as Ph₃P=S, as indeed can be done experimentally, is enthapically highly favoured (-38.3 kcal/mol⁻¹) yielding eventually complex **8** (overall exothermicity of reaction –57.9 kcal/mol⁻¹).

Interestingly, for the reaction of **1** with CS_2 the thiocarbamate intermediate can easily be electrophilically attacked by another CS_2 molecule, accounting for the formation of the isothiocyanate adduct of terminal uranium(IV)-trithiocarbonate. However, the NCS ligand cannot

19

react as the cyanate ligand in a radical way. Indeed, a radical coupling of two NCS· would ultimately lead to the formation of two CS molecules that are kinetically highly unstable.^[35] Instead, the NCS· extrusion occurs in a different way, by employing a redox process. The NCS· radical can react with complex **1**, which acts as a single electron donor, yielding complex **2** by an esoteric but new way to prepare a uranium(VI)-nitride, to give NCS⁻ that coordinates the counter-cation of **1** to form **7**. This reaction is thermodynamically highly favourable (by 97.6 kcal/mol) and represents an alternative to the unfavourable radical coupling reaction for NCS·. Including a solvation model, this reaction is of the same order of magnitude, being -87.2 kcal/mol (Figure S18).

The uranium(VI)-nitride 2 reacts with CS₂ to ultimately give the uranium(IV)-thiocyanate 8 and elemental sulfur. However, the calculated reaction profile supports the idea that the nitride undergoes a nucleophilic attack of the CS₂ and by a subsequent bond-metathesis generates a $[U(Tren^{TIPS})(S)(NCS)]$ analogue of the CO₂ reaction with 2 to give 4. However, whereas for the analogous CO₂ reaction 4 extrudes cyanate, for the sulfur analogue the uranium(VI)-sulfidethiocyanate intermediate instead extrudes sulfur. Here, the former reaction is a more common oneelectron redox event, whereas the latter is a less frequently observed example of a two-electron reduction. We suggest that the different reaction outcomes can be attributed directly to the hard nature of oxide compared to softer sulfide; even highly oxidising uranium(VI) cannot oxidise oxide so the cyanate is sacrificed and ejected for 4 but the softer sulfide can be oxidised to elemental sulfur by uranium(VI) and thus 8 is formed. The barrier for the [2 + 2]-cycloaddition is the highest found (23.4 kcal/mol), that is 14.7 kcal/mol higher than for the outer-sphere reaction of CS₂ with 1, possibly due to three effects: (i) the smaller size of uranium(VI) compared to uranium(V); (ii) the difference of nucleophilicity of the nitride; and (iii) the bond polarisation in CS₂. Also, this relatively high computed energy barrier nicely explains the fact that the corresponding reaction takes considerable time to occur experimentally with respect to the others. The barrier for the C-S bond disruption from the thiocarbamate complex is again low (5.1 kcal/mol) and similar to that found for **1**. In the terminal sulfide-thiocyanate complex, the thiocyanate ligand is strongly bonded (calculated BDE = 25.6 kcal/mol). This is due to the relatively long U-S bond, which renders the uranium centre more available for coordination with an extra ligand. Indeed, we note that the uranium(VI)-sulfide-thiocyanate is the least stable (-19.6 kcal/mol); this compares to -57.1 kcal/mol for the uranium(V)-oxo-cyanate which is more stable than the uranium(VI)-oxo-cyanate (-38.1 kcal/mol).

Finally, the most striking feature of the CS₂ reactivity is the completely different final outcomes of the reactions compared to the CO₂ reactions, but as suggested by the calculated reaction profiles the reactions actually proceed initially through structurally related intermediates. The barriers for either the outer-sphere addition (16 kcal/mol) or the [2 + 2]-cycloaddition (23.4 kcal/mol) are systematically higher than for the analogous reactions with CO₂ (11-13 kcal/mol). Therefore, this cannot be attributed to a difference in nucleophilicity of the nitride, but to the difference of bond polarisation in CS₂ versus CO₂. Indeed, in the latter, the carbon atom carries a formal δ^+ charge whereas it is δ in the former, making the formation of an empty 2p-orbital at the carbon atom more complicated and thus increasing the barrier.

Summary and Conclusions

In summary, we report the first terminal f-block nitride reactivity study with CE_2 (E = O, S), where we observe cleavage at terminal uranium-nitrides followed by zero-, one-, and two-electron redox events to give four different, yet related reactions that are well-defined; this highlights that care should be taken when using either as a reactivity surrogate of the other. For CO₂ the uranium(V)nitride reacts to give a uranium(V)-oxo-cyanate by bond metathesis and cleavage of a C=O bond and the uranium(VI)-nitride reacts to give an analogous uranium(VI)-oxo-cyanate complex that subsequently decomposes to a uranium(V)-oxo with elimination of a cyanate radical; the latter is proposed to dimerise to diisooxocyan that is unstable with respect to decomposition to N₂ and CO

formation. In contrast, for CS_2 the uranium(V)-nitride reacts to give a uranium(IV)-trithiocarbonate along with uranium(VI)-nitride and potassium thiocyanate, with formal overall disproportionation, but the uranium(VI)-nitride reacts to give a uranium(IV)-thiocyanate complex and extrusion of elemental sulfur that can be trapped as Ph₃PS by a Ph₃P scavenger.

Calculated reaction profiles consistently reproduce the experimental outcomes, and together the experimental and theoretical data suggest that the initial activation of CE_2 follows outer sphere attack of CE_2 by the nitrides for uranium(V) or direct [2 + 2]-cycloaddition for uranium(VI). Irrespective, carbamate-type species are ultimately formed which then engage in multiple bond metathesis steps, which may or may not involve redox chemistry. By dissecting each individual calculated step with reference to the experimental reaction outcome it has proven possible to delineate the effects that uranium oxidation state and nature of the coordinated E and NCE groups have on the resulting divergent reaction pathways. In one instance there is no subsequent redox chemistry, but the more common outcome is a subsequent one-electron redox step, which is a common feature of f-block chemistry. However, in one instance an unusual two-electron reduction event is found.

In a wider context, the outer-sphere reactivity of **1** parallels that of related transition metal(V) complexes, but the inner-sphere reactivity of **2** stands in contrast, and can be related to the stronger, less reactive U=N triple bond in **2** compared to **1**. Cleavage of CE₂ by **1** and **2** here is similar to that reported for diuranium-nitride complexes,^[6a,b] but **1** and **2** react further than the bimetallics, most likely because the bimetallics generally trap the products whereas monometallic **1** and **2** do not. A fuller comparison of terminal and bridging uranium nitrides will require mechanistic studies of the latter to complement the former reported here.

Acknowledgements

We thank the Royal Society, the Engineering and Physical Sciences Research Council, the European Research Council, The University of Manchester, The University of Nottingham, the EPSRC UK National EPR Facility, COST Action CM1006, and the National Nuclear Laboratory for supporting this work. LM is a member of the Institut Universitaire de France. The Humboldt foundation is also acknowledged for a grant of experienced researcher. CalMip is thanked for a generous grant of computing time.

References

- a) W. B. Tolman, *Activation of Small Molecules*, Wiley-VCH, Weinheim, Germany, 2006; b)
 H. Arakawa, M. Aresta, J. N. Armor, M. A. Barteau, E. J. Beckman, A. T. Bell, J. E. Bercaw,
 C. Creutz, E. Dinjus, D. A. Dixon, K. Domen, D. L. DuBois, J. Eckert, E. Fujita, D. H.
 Gibson, W. A. Goddard, D. W. Goodman, J. Keller, G. J. Kubas, H. H. King, J. E. Lyons, L.
 E. Manzer, T. J. Marks, K. Morokuma, K. M. Nicholas, R. Periana, L. Que, J. Rostrup Nielson, W. M. H. Sachtler, L. D. Schmidt, A. Sen, G. A. Somorjai, P. C. Stair, B. R. Stultls and W. Tumas, *Chem. Rev.* 2001, *101*, 953.
- [2]. a) C. Costentin, M. Robert and J. -M. Savéant, *Chem. Soc. Rev.* 2013, *42*, 2423; b) E. E. Benson, C. P. Kubiak, A. J. Sathrum and M. Smieja, *Chem. Soc. Rev.* 2009, *38*, 89; c) D. J. Darensbourg, *Chem. Rev.* 2007, *107*, 2388; d) T. Sakakura, J. -C. Choi and H. Yasuda, *Chem. Rev.* 2007, *107*, 2365.
- [3]. a) B. Ochiai and T. Endo, Prog. Polym. Sci. 2005, 30, 183; b) K. K. Pandey, Coord. Chem.
 Rev. 1995, 140, 37.
- [4]. Representative examples: a) G. -C. Wang, Y. -M. So, K. -L. Wong, K. -C. Au-Yeung, H. H. -Y. Sung, I. D. Williams and W. -H. Leung, *Chem. Eur. J.* 2015, *21*, 16126; b) J. Andrez, J. Pécaut, P. -A. Bayle and M. Mazzanti, *Angew. Chem. Int. Ed.* 2014, *53*, 10488; c) Y. -M. So, G. -C. Wang, Y. Li, H. H. -Y. Sung, I. D. Williams, Z. Lin and W. -H. Leung, *Angew. Chem.*

Int. Ed. 2014, 53, 1626; d) K. Wang, G. Luo, J. Hong, X. Zhou, L. Weng, Y. Luo and L. Zhang, Angew. Chem. Int. Ed. 2014, 53, 1053; e) W. S. Ren, H. B. Song, G. F. Zi and M. D. Walter, Dalton Trans. 2012, 41, 5965; f) W. S. Ren, G. F. Zi, D. C. Fang and M. D. Walter, J. Am. Chem. Soc. 2011, 133, 13183; g) D. Heitmann, C. Jones, D. P. Mills and A. Stasch, Dalton Trans. 2010, 39, 1877; h) W. J. Evans, T. J. Mueller and J. W. Ziller, J. Am. Chem. Soc. 2009, 131, 2678; i) W. J. Evans, D. B. Rego, J. W. Ziller, A. G. DiPasquale and A. L. Rheingold, Organometallics 2007, 26, 4737; j) N. W. Davies, A. S. P. Frey, M. G. Gardiner and J. Wang, Chem. Commun. 2006, 4853; k) W. J. Evans, J. M. Perotti and J. W. Ziller, J. Am. Chem. Soc. 2005, 127, 3894; l) W. J. Evans, J. M. Perotti, J. C. Brady and J. W. Ziller, J. Am. Chem. Soc. 2003, 125, 5204; m) W. J. Evans, C. A. Seibel and J. W. Ziller, Inorg. Chem. 1998, 37, 770.

Uranium-mediated CO₂ reactivity: a) C. Camp, L. Chatelain, C. E. Kefalidis, J. Pécault, L. [5]. Maron and M. Mazzanti, Chem. Commun. 2015, 51, 15454; b) J. A. Higgins, A. S. P. Frey, F. G. N. Cloke and S. M. Roe, Organometallics 2015, 34, 2102; c) O. Cooper, C. Camp, J. Pécault, C. E. Kefalidis, L. Maron, S. Gambarelli and M. Mazzanti, J. Am. Chem. Soc. 2014, 136, 6716; d) A. -C. Schmidt, F. W. Heinemann, L. Maron and K. Meyer, Inorg. Chem. 2014, 53, 13142; e) C. L. Webster, J. W. Ziller and W. J. Evans, Organometallics 2014, 33, 433; f) A. -C. Schmidt, F. W. Heinemann, C. E. Kefalidis, L. Maron, P. W. Roesky and K. Meyer, Chem. Eur. J. 2014, 20, 13501; g) P. L. Arnold, Z. R. Turner, A. I. Germeroth, I. J. Casely, G. N. Nichol, R. Bellabarba and R. P. Tooze, Dalton Trans, 2013, 42, 1333; h) C. L. Webster, J. W. Ziller and W. J. Evans, Organometallics 2013, 32, 4820; i) J. A. Higgins, F. G. N. Cloke and S. M. Roe, Organometallics 2013, 32, 5244; j) O. P. Lam, S. M. Franke, F. W. Heinemann and K. Meyer, J. Am. Chem. Soc. 2012, 134, 16877; k) V. Mougel, C. Camp, J. Pécault, C. Copéret, L. Maron, C. E. Kefalidis and M. Mazzanti, Angew. Chem. Int. Ed. 2012, 51, 12280; 1) C. L. Webster, J. W. Ziller and W. J. Evans, Organometallics 2012, 31, 7191; m) A. -C. Schmidt, A. V. Nizovtsev, A. Scheuer, F. W. Heinemann and K. Meyer, Chem.

Commun. 2012, 48, 8634; n) E, M. Matson, W. P. Forrest, P. E. Fanwick, S. C. Bart, J. Am.
Chem. Soc. 2011, 133, 4948; o) S. M. Mansell, N. Kaltsoyannis and P. L. Arnold, J. Am.
Chem. Soc. 2011, 133, 9036; p) E. M. Matson, P. E. Fenwick and S. C. Bart, Organometallics
2011, 30, 5753; q) O. Bénaud, J. -C. Berthet, P. Thuéry and M. Ephritikhine, Inorg. Chem.
2010, 49, 8117; r) W. J. Evans, J. R. Walensky and J. W. Ziller, Organometallics 2010, 29,
945; s) O. P. Lam, S. C. Bart, H. Kameo, F. W. Heinemann and K. Meyer, Chem. Commun.
2010, 46, 3137; t) O. T. Summerscales, A. S. P. Frey, F. G. N. Cloke and P. B. Hitchcock,
Chem. Commun. 2009, 198; u) S. C. Bart, C. F. Anthon, W. Heinemann, E. Bill, N. M.
Edelstein and K. Meyer, J. Am. Chem. Soc. 2008, 130, 12536; v) I. Castro-Rodriguez and K.
Meyer, J. Am. Chem. Soc. 2005, 127, 11242; w) I. Castro-Rodriguez, H. Nakai, L. N.
Zhakharov, A. L. Rheingold and K. Meyer, Science 2004, 305, 1757; x) K. G. Moloy and T.
J. Marks, Inorg. Chim. Acta, 1985, 110, 127; y) A. L. Arduini, J. D. Jamerson and J. Takats,

[6]. Uranium-mediated CS₂ reactivity: a) L. Chatelain, R. Scopelliti and M. Mazzanti, J. Am. Chem. Soc. 2016, 138, 1784; b) M. Falcone, L. Chatelain and M. Mazzanti, Angew. Chem. Int. Ed. 2016, 55, 4074; c) C. Camp, O. Cooper, J. Andrez, J. Pécault and M. Mazzanti, Dalton Trans. 2015, 44, 2650; d) E. M. Matson, A. T. Breshears, J. J. Kiernicki, B. S. Newell, P. E. Fanwick, M. P. Shores, J. R. Walensky and S. C. Bart, Inorg. Chem. 2014, 53, 12977; e) O. P. Lam, S. M. Franke, F. W. Heinemann and K. Meyer, J. Am. Chem. Soc. 2012, 134, 16877; f) O. P. Lam, L. Castro, B. Kosog, F. W. Heinemann, L. Maron and K. Meyer, Inorg. Chem. 2012, 51, 781; g) N. A. Siladke, J. LeDuc, J. W. Ziller and W. J. Evans, Chem. Int. Ed. 2011, 50, 5965; i) E. M. Matson, W. P. Forrest, P. E. Fanwick and S. C. Bart, J. Am. Chem. Soc. 2011, 133, 4948; j) C. Lescop, T. Arliguie, M. Lance, M. Nierlich and M. Ephritikhine, J. Organomet. Chem. 1999, 580, 137; k) J. G. Brennan, R. A. Andersen and A. Zalkin, Inorg.

Chem. 1986, 25, 1756; l) A. L. Arduini, J. D. Jamerson and J. Takats, J. Inorg. Chem. 1981, 20, 2474.

- [7]. a) S. T. Liddle, *Angew. Chem. Int. Ed.* 2015, *54*, 8604; b) H. S. La Pierre and K. Meyer, *Prog. Inorg. Chem.* 2014, *58*, 303; c) B. M. Gardner and S. T. Liddle, *Eur. J. Inorg. Chem.* 2013, 3753; d) A. R. Fox, S. C. Bart, K. Meyer, and C. C. Cummins, *Nature* 2008, *455*, 341.
- [8]. a) J. P. Krogman, B. M. Foxman and C. M. Thomas, J. Am. Chem. Soc. 2011, 133, 14582; b)
 A. R. Sadique, W. W. Brennessel and P. L. Holland, Inorg. Chem. 2008, 47, 784; c) C. H.
 Lee, D. S. Laitar, P. Müller and J. P. Sadighi, J. Am. Chem. Soc. 2007, 129, 13802; d) C. C.
 Lu, C. T. Saouma, M. W. Day and J. C. Peters, J. Am. Chem. Soc. 2007, 129, 4; e) D. S.
 Laitar, P. Müller and J. P. Sadighi, J. Am. Chem. Soc. 2005, 127, 17196; f) G. Fachinetti, C.
 Floriani, A. Chiesi-Villa and C. Guastini, J. Am. Chem. Soc. 1979, 101, 1767.
- [9]. a) J. M. Smith, Prog. Inorg. Chem. 2014, 58, 417 b) C. T. Saouma and J. C. Peters, Coord. Chem. Rev. 2011, 255, 920; c) R. A. Eikey and M. M. Abu-Omar, Coord. Chem. Rev. 2003, 243, 83; d) K. Dehnicke and J. Strähle, Angew. Chem. Int. Ed. Engl. 1992, 31, 955; e) K. Dehnicke and J. Strähle, Angew. Chem. Int. Ed. Engl. 1981, 20, 413.
- [10]. a) S. P. Semproni and P. J. Chirik, J. Am. Chem. Soc. 2013, 135, 11373; b) J. S. Silva and C. C. Cummins, J. Am. Chem. Soc. 2010, 132, 2169; c) U. J. Kilgore, F. Basuli, J. C. Huffman and D. J. Mindiola, Inorg. Chm., 2006, 45, 487; d) J. K. Brask, V. Durà-Villà, P. L. Diaconescu and C. C. Cummins, Chem. Commun. 2002, 902.
- [11]. a) D. M. King, F. Tuna, E. J. L. McInnes, J. McMaster, W. Lewis, A. J. Blake and S. T. Liddle, *Nat. Chem.* 2013, 5, 482; b) D. M. King, F. Tuna, E. J. L. McInnes, J. McMaster, W. Lewis, A. J. Blake and S. T. Liddle, *Science* 2012, 337, 717.
- [12]. D. M. King and S. T. Liddle, Coord. Chem. Rev. 2014, 266-267, 2.
- [13]. a) N. Tsoureas, A. F. R. Kilpatrick, C. J. Inman, and F. G. N. Cloke, *Chem. Sci.* 2016, 7, 4624; b) C. Camp, J. Pécaut and M. Mazzanti, *J. Am. Chem. Soc.* 2013, 135, 12101; c) R. K. Thomson, T. Cantat, B. L. Scott, D. E. Morris, E. R. Batista and J. L. Kiplinger, *Nat. Chem.*

2010, 2, 723; d) T. K. Todorova, L. Gagliardi, J. R. Walensky, K. A. Miller and W. J. Evans, J. Am. Chem. Soc. 2010, 132, 12397; e) S. Fortier, G. Wu and T. W. Hayton, J. Am. Chem. Soc. 2010, 132, 6888; f) A. R. Fox, P. L. Arnold and C. C. Cummins, J. Am. Chem. Soc. 2010, 132, 3250; g) A. R. Fox and C. C. Cummins, J. Am. Chem. Soc. 2009, 131, 5716; h) G. Nocton, J. Pécaut and M. Mazzanti, Angew. Chem. Int. Ed. 2008, 47, 3040; i) W. J. Evans, K. A. Miller, J. W. Ziller and J. Greaves, Inorg. Chem. 2007, 46, 8008; j) W. J. Evans, S. A. Kozimor and J. W. Ziller, Science 2005, 309, 1835; k) I. Korobkov, S. Gambarotta and G. P. A. Yap, Angew. Chem. Int. Ed. 2002, 41, 3433.

- [14]. P. A. Cleaves, D. M. King, C. E.; Kefalidis, L. Maron, F. Tuna, E. J. L. McInnes, J. McMaster, W. Lewis, A. J. Blake and S. T. Liddle, *Angew. Chem. Int. Ed.* 2014, 53, 10412.
- [15]. a) A. F. Cozzolino, J. S. Silvia, N. Lopez and C. C. Cummins, *Dalton Trans.* 2014, 43, 4639;
 b) B. L. Tran, M. Pink, X. Gao, H. Park and D. J. Mindiola, *J. Am. Chem. Sos.* 2010, 132, 1458; c) J. S. Silvia and C. C. Cummins, *J. Am. Chem. Soc.* 2009, 131, 446.
- [16]. a) F. Dielmann, O. Back, M. Henry-Ellinger, P. Jerabek, G. Frenking and G. Bertrand, *Science* 2012, 337, 1526; b) J. J. Scepaniak, R. P. Bontchev, D. L. Johnson and J. M. Smith, *Angew. Chem. Int. Ed.* 2011, 50, 6630.
- [17]. a) D. P. Halter, F. W. Heinemann, J. Bachmann and K. Meyer, *Nature* 2016, *530*, 317; b) C. Camp, M. A. Antunes, G. Garcia, I. Ciofini, I. C. Santos, J. Pecault, M. Almeida, J. Marcalo and M. Mazzanti, *Chem. Sci.* 2014, *5*, 841; c) E. Lu, O. J. Cooper, J. McMaster, F. Tuna, E. J. L. McInnes, W. Lewis, A. J. Blake and S. T. Liddle, *Angew. Chem. Int. Ed.* 2014, *53*, 6696; d) D. M. King, F. Tuna, J. McMaster, W. Lewis, A. J. Blake, E. J. L. McInnes and S. T. Liddle, *Angew. Chem. Int. Ed.* 2013, *52*, 4921; e) J. L. Brown, S. Fortier, G. Wu, N. Kaltsoyannis and T. W. Hayton, *J. Am. Chem. Soc.* 2013, *135*, 5352; f) D. P. Mills, O. J. Cooper, F. Tuna, E. J. L. McInnes, E. S. Davies, J. McMaster, F. Moro, W. Lewis, A. J. Blake and S. T. Liddle, *J. Am. Chem. Soc.* 2012, *134*, 10047; g) S. Fortier, J. L. Brown, N. Kaltsoyannis, G. Wu and T. W. Hayton, *Inorg. Chem.* 2012, *51*, 1625; h) C. R. Graves and J.

L. Kiplinger, *Chem. Commun.* 2009, 3831; i) I. Castro-Rodríguez, H. Nakai and K. Meyer, *Angew. Chem. Int. Ed.* 2006, 45, 2389; j) D. S. J. Arney and C. J. Burns, *J. Am. Chem. Soc.* 1995, *117*, 9448; k) D. S. J. Arney and C. J. Burns, *J. Am. Chem. Soc.* 1993, *115*, 9840; l) D. S. J. Arney, C. J. Burns and D. C. Smith, *J. Am. Chem. Soc.* 1992, *114*, 10068; m) J. G. Brennan and R. A. Andersen, *J. Am. Chem. Soc.* 1985, *107*, 514.

- [18]. J. Selbin and J. D. Ortego, Chem. Rev. 1969, 69, 657.
- [19]. D. R. Kindra and W. J. Evans, Chem. Rev. 2014, 114, 8865.
- [20]. D. Patel, F. Tuna, E. J. L. McInnes, J. McMaster, W. Lewis, A. J. Blake and S. T. Liddle, *Dalton Trans.* 2013, 42, 5224.
- [21]. B. M. Gardner and S. T. Liddle, Chem. Commun. 2015, 51, 10589.
- [22]. a) H. S. La Pierre, M. Rosenzweig, B. Kosog, C. Hauser, F. W. Heinemann, S. T. Liddle and K. Meyer, *Chem. Commun.* 2015, *51*, 16671; b) H. S. La Pierre and K. Meyer, *Inorg. Chem.* 2013, *52*, 529; c) A. J. Lewis, P. J. Carroll and E. J. Schelter, *J. Am. Chem. Soc.* 2013, *135*, 13185; d) B. Kosog, H. S. La Pierre, F. W. Heinemann, S. T. Liddle and K. Meyer, *J. Am. Chem. Soc.* 2012, *134*, 5284; e) E. O'Grady and N. Kaltosyannis, *J. Chem. Soc. Dalton Trans.* 2002, 1233; f) R. G. Denning, *J. Phys. Chem A* 2007, *111*, 4125; (g) R. G. Denning, *Struct. Bonding*, Springer, Berlin 1992, *79*, 215.
- [23]. A. Schulz and T. M. Klapötke, Inorg. Chem. 1996, 35, 4791.
- [24]. B. Ruscic and J. Berkowitz, Chem. Sci. 1994, 100, 4498.
- [25]. Recorded at 145 K (b.p. 296 K), the N-H IR frequency of HNCO is 2254 cm⁻¹ (v₂). See, M. S. Lowenthal, R. K. Khanna and M. H. Moore, *Spectrochim. Acta A* 2002, *58*, 73.
- [26]. A. R. Amell, J. Am. Chem. Soc. 1956, 78, 6234.
- [27]. a) C. Camp, N. Settineri, J. Lefèvre, A. R. Jupp, J. M. Goicoechea, L. Maron and J. Arnold, *Chem. Sci.* 2015, *6*, 6379; b) E. Hashem, J. A. Platts, F. Hartl, G. Lorusso, M. Evangelisti, C. Schulzke and R. J. Baker, *Inorg. Chem.* 2014, *53*, 8624; c) A. J. Lewis, E. Nakamaru-Ogiso, J. M. Kikkawa, P. J. Carroll and E. J. Schelter, *Chem. Commun.* 2012, *48*, 4977.

- [28]. See for instance P. Liu, B. L. H. Taylor, J. Garcia-Lopez, and K. N. Houk, Computational Studies of Ruthenium-Catalyzed Olefin Metathesis, in *Handbook of Metathesis, Set*, ed R. H. Grubbs, A. G. Wenzel, D. J. O'Leary and E. Khosravi, Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim, Germany, 2015, doi: 10.1002/9783527674107.ch7.
- [29]. C. E. Kefalidis, A. Stasch, C. Jones and L. Maron, Chem. Commun. 2014, 50, 12318.
- [30]. C. E. Kefalidis and L. Maron, Chem. Eur. J. 2016, 22, 4743.
- [31]. L. Watson and O. Eisenstein, J. Chem. Ed. 2002, 79, 1269.
- [32]. a) B. M. Gardner, G. Balázs, M. Scheer, A. J. Wooles, F. Tuna, E. J. L. McInnes, J. McMaster, W. Lewis, A. J. Blake and S. T. Liddle, *Angew. Chem. Int. Ed.* 2015, 54, 15250;
 b) B. M. Gardner, G. Balázs, M. Scheer, F. Tuna, E. J. L. McInnes, J. McMaster, W. Lewis, A. J. Blake and S. T. Liddle, *Nat. Chem.* 2015, 7, 582; c) B. M. Gardner, F. Tuna, E. J. L. McInnes, J. McMaster, W. Lewis, A. J. Blake and S. T. Liddle, *Nat. Chem.* 2015, 7, 582; c) B. M. Gardner, F. Tuna, E. J. L. McInnes, J. McMaster, W. Lewis, A. J. Blake and S. T. Liddle, *Angew. Chem. Int. Ed.* 2015, 54, 7068; d) B. M. Gardner, P. A. Cleaves, C. E. Kefalidis, J. Fang, L. Maron, W. Lewis, A. J. Blake and S. T. Liddle, *Chem. Sci.* 2014, 5, 2489; e) B. M. Gardner, G. Balázs, M. Scheer, F. Tuna, E. J. L. McInnes, J. McMaster, W. Lewis, A. J. Blake and S. T. Liddle, *Angew. Chem. Int. Ed.* 2014, 53, 4484; f) D. M. King, J. McMaster, F. Tuna, E. J. L. McInnes, W. Lewis, A. J. Blake and S. T. Liddle, *J. Am. Chem. Soc.* 2014, *136*, 5619.
- [33]. For examples of sterically compressed complexes see: a) N. A. Siladke, K. R. Meihaus, J. W. Ziller, M. Fang, F. Furche, J. R. Long and W. J. Evans, J. Am. Chem. Soc. 2012, 134, 1243; b)
 W. J. Evans, T. J. Mueller and J. W. Ziller, J. Am. Chem. Soc. 2009, 131, 2678; c) W. J. Evans, S. A. Kozimor and J. W. Ziller, J. Am. Chem. Soc. 2003, 125, 14264; d) W. J. Evans, S. A. Kozimor, G. W. Nyce and J. W. Ziller, J. Am. Chem. Soc. 2003, 125, 13831; e) W. J. Evans, and B. L. Davis, Chem. Rev. 2002, 102, 2119; f) W. J. Evans, G. W. Nyce, M. A. Johnston and J. W. Ziller, J. Am. Chem. Soc. 2000, 122, 12019.
- [34]. C. E. Kefalidis, A. S. P. Frey, S. M. Roe, F. G. N. Cloke and L. Maron, *Dalton Trans.* 2014, 43, 11202.

[35]. a) L. Castro, S. Labouille, D. R. Kindra, J. W. Ziller, F. Nief, W. J. Evans and L. Maron,

Chem. Eur. J. 2012, 18, 7886. b) L. Castro and L. Maron, Chem. Eur. J. 2012, 18, 6610.

Keywords

Uranium, nitride, carbon dioxide, carbon disulfide, density functional theory

ToC

