This article was downloaded by: [University of Sydney] On: 12 August 2013, At: 22:07 Publisher: Taylor & Francis Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information: <u>http://www.tandfonline.com/loi/gpss20</u>

Microwave-Induced Stereoselective Conversion of Dialkyl 2-(1,1,3-Trioxo-1,3-dihydro-2 H -1,2-Benzisothiazol-2-yl)-3-(triphenylphosphoranylidene)succi to Dialkyl 2-(1,1,3-Trioxo-1,3-dihydro-2 H -1,2-benzisothiazol-2-yl)-2butendioates in the Presence of Silica-Gel Powder in Solvent-Free Conditions Ali Ramazani^a, Akram Abbasi Motejadded^a & Ebrahim Ahmadi^a

^a Chemistry Department, Zanjan University, Zanjan, Iran

Published online: 01 Feb 2007.

To cite this article: Ali Ramazani , Akram Abbasi Motejadded & Ebrahim Ahmadi (2006) Microwave-Induced Stereoselective Conversion of Dialkyl 2-(1,1,3-Trioxo-1,3-dihydro-2 H -1,2-Benzisothiazol-2-yl)-3-(triphenylphosphoranylidene)succinates to Dialkyl 2-(1,1,3-Trioxo-1,3-dihydro-2 H -1,2-benzisothiazol-2-yl)-2-butendioates in the Presence of Silica-Gel Powder in Solvent-Free Conditions, Phosphorus, Sulfur, and Silicon and the Related Elements, 181:1, 233-236, DOI: <u>10.1080/104265090969838</u>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <u>http://www.tandfonline.com/page/terms-and-conditions</u>



Microwave-Induced Stereoselective Conversion of Dialkyl 2-(1,1,3-Trioxo-1,3-dihydro-2*H*-1,2-Benzisothiazol-2-yl)-3-(triphenylphosphoranylidene)succinates to Dialkyl 2-(1,1,3-Trioxo-1,3-dihydro-2*H*-1,2-benzisothiazol-2-yl)-2butendioates in the Presence of Silica-Gel Powder in Solvent-Free Conditions

Ali Ramazani Akram Abbasi Motejadded Ebrahim Ahmadi Chemistry Department, Zanjan University, Zanjan, Iran

Protonation of the highly reactive 1:1 intermediates produced in the reaction between triphenylphosphine and dialkyl acetylenedicarboxylates by saccharin leads to vinyltriphenylphosphonium salts, which undergo an addition reaction with a counter anion in CH_2Cl_2 at r.t. to produce the corresponding stabilized phosphorus ylides. Silica-gel powder was found to catalyze the stereoselective conversion of the stabilized phosphorus ylides to the corresponding electron-poor N-vinylated isothiazoles in solvent-free conditions under thermal and microwave irradiation in fairly good yields.

Keywords Microwave; phosphorus ylide; saccharin; silica gel; solvent-free conditions

INTRODUCTION

 β -Additions of nucleophiles to the vinyl group of vinylic phosphonium salts leading to the formation of new alkylidenephosphoranes has attracted much attention as a very convenient and synthetically useful method in organic synthesis.¹ Organophosphorus compounds have been used extensively in organic synthesis as useful reagents as well as ligands of a number of transition metal catalysts.² Silica gel as an additive promotes the Wittig reactions of phosphorus ylides with aldehydes, including sterically hindered aldehydes to increase the rate and yields of alkenes.^{3,4} In the past we have established a convenient, one-pot method for preparing stabilized phosphorus ylides

Received February 1, 2005; in final form February 3, 2005.

This work was supported by the Zanjan University.

Address correspondence to Ali Ramazani, Chemistry Department, Zanjan University, PO Box 45195-313, Zanjan, Iran. E-mail: aliramazani@yahoo.com

utilizing the in situ generation of the phosphonium salts.⁵ In this article, we report on catalytic role of silica-gel powder in the stereoselective conversion of dialkyl 2-(1,1,3-trioxo-1,3-dihydro-2*H*-1,2-benzisothiazol-2-yl)-3-(triphenylphosphoranylidene)succinates (**5**) to dialkyl 2-(1,1,3-trioxo-1,3-dihydro-2*H*-1,2-benzisothiazol-2-yl)-2-butendioates (**6**) in solvent-free conditions⁶ under thermal and microwave irradiation in fairly good yields (Scheme 1).



SCHEME 1

RESULTS AND DISCUSSION

Reactions are known in which an α,β -unsaturated carbonyl compound is produced from phosphonium salts.⁵ Thus, compounds **6** may result from an initial addition of triphenylphosphine **1** to the acetylenic ester **2** and concomitant protonation of the 1:1 adduct by the saccharin (**3**) to form the corresponding triphenylphosphonium salts **4**. A conjugate addition of the saccharin (**3**) anion to the vinyltriphenylphosphonium cation **4** leads to the formation of the stabilized phosphorus ylide **5**. Silica-gel powder was found to catalyze the stereoselective conversion of the stabilized phosphorus ylides 5 to the corresponding electron-poor N-vinylated isothiazoles 6 in solvent-free conditions under thermal and microwave irradiation in fairly good yields (Scheme 1). TLC indicated that the reaction was completed after 45 min at $105^{\circ}C$ (3 min at microwave power 0.5 KW). The reaction proceeded smoothly and cleanly under solvent-free conditions⁶ (in all cases the reaction worked efficiently with high coversions) and no side reactions were observed. We also have used KHSO₄, MgSO₄, Al₂O₃, MgO, ZnO, K₂HPO₄, KH₂PO₄, $NaHCO_3$, and $MnSO_4$ in this reaction, but the yields of the corresponnding products 6 in cases of MgO, $MgSO_4$, K_2HPO_4 , and KH_2PO_4 were low, and in the others cases, no products were observed; in all cases, decomposition was observed. In the absence of the silica-gel powder, the reactions were not completed at reflux temperature (toluene as solvent) after 24 h and decomposition of the starting materials and product were observed. The structures **6a-b** were deduced from their IR, ¹H NMR, and ¹³C NMR spectra.

In summary, we have found that silica-gel powder is able to catalyze stereoselective conversion of ylide **5** compound **6** in solvent-free conditions (Scheme 1). Other aspects of this process are under investigation.

EXPERIMENTAL

Commerical-oven Butane M245 was used for microwave irradiation. Melting points were measured on an Electrothermal 9100 apparatus and are uncorrected. IR spectra were recorded on a Shimadzu IR-460 spectromer. ¹H and ¹³C NMR spectra were measured with a BRUKER DRX-500 AVANCE spectrometer at 500 and 125 MHz, respectively.

General Procedure for the Preparation of Ylides 5 and Compounds 6a–b

To a magnetically stirred solution of triphenylphosphine 1 (1 mmol) and saccharin 3 (1 mmol) in CH_2Cl_2 , 6 mL was added dropwise a mixture of 2 (1 mmol) in CH_2Cl_2 (4 mL) at $-10^{\circ}C$ over 15 min. The mixture was allowed to warm up to r.t. Silica-gel power (2 g) was added and the solvent was evaporated. Dry silica gel and the residue were heated for 45 min at $105^{\circ}C$ (or irradiated at microwave power 0.5 KW for 3 min) and then placed over a column of silica gel (10 g). The column chromatography was washed using ethyl acetate-light petroleum ether (1:9) as an eluent. The solvent was removed under reduced pressure and products (**6a–b**) were obtained. The following are the characterization data of the compounds (**6a–b**).

Dimethyl 2-(1,1,3-Trioxo-1,3-dihydro-2H-1,2-benzisothiazol-2yl)-2-butendioates (6a)

White crystals, m.p. 130–137°C; Yield: 47%. IR (KBr) (ν_{max} , Cm⁻¹): 3085, 2923, 2854, 1751,1727, and 1635. ¹H NMR (CDCl₃), *Z*-isomer: 51%, *E*-isomer: 49%, $\delta_{\rm H}$ for *Z*-isomer: 3.74 and 3.90 (6 H, 2 s, 2 OCH₃); 7.47 (1 H, s CH=), 7.8–8.2 (4 H, m, arom.). $\delta_{\rm H}$ for *E*-isomer: 3.84 and 3.90 (6 H, 2 s, 2 OCH₃); 6.74 (1 H, s, CH=), 7.8–8.2 (4 H, m, arom.). ¹³C NMR (CDCl₃) $\delta_{\rm c}$ for *Z*-isomer: 53.55 and 53.45 (2 OCH₃); 121.12 (CH=); 137.89 (N-C=), 121.36, 125.85, 128.40, and 134.61 (4 CH, arom.), 125.93 and 135.23 (2 C, arom.), 161.35 and 162.53 (2C=O of esters); 157.18 (C=O, imide). $\delta_{\rm C}$ for *E*-isomer: 52.70 and 53.74 (2 OCH₃); 135.83 (CH=); 138.46 (N-C=), 121.47, 126.87, 131.18, and 134.95 (4 CH, arom.), 126.03 and 135.27 (2C, arom.), 161.80 and 163.75 (2C=O of esters); 158.15 (C=O, imide).

Diethyl 2-(1,1,3-Trioxo-1,3-dihydro-2H-1,2-benzisothiazol-2-yl)-2-butendioates (6b)

Viscous colorless oil; Yield: 40%. IR (CCl₄) (ν_{max} , Cm⁻¹): 3084, 2937, 2868, 1742, and 1641. ¹H NMR (CDCl₃), Z-isomer: 69.5%, E-isomer: 30.5%, $\delta_{\rm H}$ for Z-isomer: 1.15 (3H, t, ${}^{3}J_{\rm HH} = 7.1$ Hz, CH₃ of Et); 1.33 (3H, t, ${}^{3}J_{\rm HH} = 7.1$ Hz, CH₃ of Et), 4.17 (2 H, q, ${}^{3}J_{\rm HH} = 7.1$ Hz, OCH₂ of Et); $4.34 (2H, q, {}^{3}J_{HH} = 7.1 \text{ Hz}, \text{OCH}_{2} \text{ of Et}); 7.45 (1 \text{ H}, \text{s}, \text{CH}=), 7.8-8.2 (4 \text{ H}, \text{s})$ m, arom.). $\delta_{\rm H}$ for *E*-isomer: 1.25 (3H, t, ${}^{3}J_{\rm HH} = 7.1$ Hz, CH₃ of Et); 1.34 $(3H, t, {}^{3}J_{HH} = 7.1 \text{ Hz}, \text{CH}_{3} \text{ of Et}), 4.29 (2 \text{ H}, q, {}^{3}J_{HH} = 7.1 \text{ Hz}, \text{OCH}_{2}$ of Et); 4.35 (2 H, q, ${}^{3}J_{HH} = 7.1$ Hz, OCH₂ of Et); 6.73 (1 H, s, CH=), 7.8–8.2 (4 H, m, arom.). ^{13}C NMR (CDCl₃), δ_C for Z-isomer: 13.75 and 13.99 (2 CH₃ of 2 Et); 61.90 and 63.14 (2 OCH₂); 121.31 (CH=); 138.48 (N-C=), 122.25, 125.83, 134.57, and 135.24 (4 CH, arom.), 128.40 and 135.49 (2 C, arom.), 161.32 and 162.14 (2C=O of esters); 158.16 (C=O, imide). $\delta_{\rm C}$ for *E*-isomer: 13.75 and 14.04 (2 CH₃ of 2 Et); 61.74 and 62.85 (2 OCH₂); 121.45 (CH=); 137.94 (N-C=), 125.91, 125.95, 134.91, and 135.77 (4 CH, arom.), 126.91 and 135.49 (2 C, arom.), 160.77 and 163.29 (2C=O of esters); 157.27 (C=O, imide).

REFERENCES

- J. I. G. Cadogan, Organophosphorus Reagents in Organic Synthesis. J. I. G. Cadogon, Ed. (Academic Press, New York, 1979).
- [2] I. Yavari and A. Ramazani, Synth. Commun., 27, 1449 (1997).
- [3] V. J. Patil and U. Mavers, Tetrahedron Lett., 37, 1281 (1996).
- [4] C. Xu, G. Chen, C. Fu, and X. Huang, Synth. Commun., 25, 2229 (1995).
- [5] A. Ramazani, L. Yousefi, E. Ahmadi, and A. Souldozi, *Phosphorus, Sulfur, and Silicon*, 179, 1459 (2004) and references cited therein.
- [6] K. Tanaka and F. Toda, Chem. Rev., 100, 1025 (2000).