

Reactions of cyclomanganated complexes with carbon disulfide: routes to η^2 -aryldithiocarboxylate- $\text{Mn}(\text{CO})_4$ complexes and to the trithiocarbonate complex $(\mu_3\text{-CS}_3)_2\text{Mn}_4(\text{CO})_{16}$

Wade J. Mace, Lyndsay Main, Brian K. Nicholson*, Marion Hagyard

Department of Chemistry, University of Waikato, Private Bag 3105, Hamilton, New Zealand

Received 22 August 2002; accepted 29 September 2002

Abstract

Reaction of cyclomanganated aryl ketones with CS_2 proceeds with insertion into the Mn-C bond to give η^2 -dithiocarboxylato- $\text{Mn}(\text{CO})_4$ compounds. With other cyclomanganated substrates such as that from $\text{Ph}_3\text{P=S}$ and also with $\text{Mn}_2(\text{CO})_{10}$, CS_2 gives $(\mu_3\text{-CS}_3)_2\text{Mn}_4(\text{CO})_{16}$ with bridging trithiocarbonate ligands.

© 2002 Elsevier Science B.V. All rights reserved.


Keywords: Manganese; Cyclomanganates; Carbon disulfide; Insertion; Aryl dithiocarboxylate; Trithiocarbonate

1. Introduction

There is now a well-developed chemistry of cyclomanganated complexes [1]. These compounds are readily formed from aromatic and heteroaromatic ketones [2], from substrates with imine or amine donor groups [3], from triphenyl-phosphine or -arsine chalcogenides [4], and from chalcones [5] to give, for example, compounds 1–3.

The Mn-C bond of these species is a site of reactivity, and combination with alkenes [6], alkynes [7], organoiso-cyanates [8], sulfur dioxide [9] or mercuric chloride [10] have all been shown to generate novel derivatives. For many of these reactions the first step in the process is undoubtedly insertion of the substrate into the Mn-C bond, followed by rearrangement with or without demanganation.

Another molecule that is known to insert into M-C




* Corresponding author. Fax: +64-7-838-4219

E-mail address: b.nicholson@waikato.ac.nz (B.K. Nicholson).

bonds (as well as into M-H , M-O , M-P or M-S bonds) is CS_2 [11]. This generally leads to the formation of bidentate dithiocarboxylate complexes (e.g.

$\text{MeCS}_2\text{Mn}(\text{CO})_4$ from $\text{MeMn}(\text{CO})_5$ [12], although monodentate species sometimes form (e.g. $\text{PtCl}(\text{SC}(\text{S})\text{H})(\text{PPh}_3)_2$ from $\text{Pt}(\text{H})(\text{Cl})(\text{PPh}_3)_2$ [13]. The only previous investigation involving cyclometallated substrates that we are aware of is the reaction of a nickel complex to give, initially, a *gem*-dithiolate complex which reacted further to give a η^2 -trithiocarbonate complex and a thioketone (Eq. (1)) [14].



We now report the reactions of CS_2 with a range of cyclomanganated complexes.

2. Experimental

2.1. General

All manipulations were carried out in an oxygen-free N_2 atmosphere with dried solvents in standard Schlenk equipment or on a standard vacuum line. Orthomanganated substrates were prepared by published methods [2–5], while CS_2 was purified by trap-to-trap distillation on the vacuum line.

2.2. Instrumentation

Infrared spectra were recorded on a Digilab FTS-40 FTIR spectrophotometer. NMR spectroscopy was performed using a Bruker DRX400 Avance in CDCl_3 ; assignments were based on standard 2D experiments. Routine electrospray mass spectra (ESMS) were obtained on a VG Platform II spectrometer operating under standard conditions in MeOH. $\text{Na}[\text{OMe}]$ was added as an ionisation aid [15]. Elemental analysis was performed by the Campbell Microanalytical Laboratory, University of Otago. Melting points were measured on a Reichart Thermopan melting point apparatus and are uncorrected.

2.3. Reactions

2.3.1. Reaction of η^2 -(2-acetyl-5-methoxyphenyl)tetracarbonylmanganese with CS_2

The cyclomanganated ketone **1** (133 mg, 0.42 mmol) was placed in a thick-walled glass ampoule (ca. 20 ml capacity). This was attached to a vacuum line, evacuated, and CS_2 (ca. 10 ml) was distilled in. The ampoule

was sealed under vacuum and transferred to a Carius tube where it was heated at 85°C for 24 h. When the cooled ampoule was stored at -20°C for 72 h orange crystals formed. The ampoule was opened and the crystals were collected, and characterised as η^2 -(2-acetyl-5-methoxyphenyldithiocarboxylato)tetracarbonylmanganese, **4**, (91 mg, 55%). M.p. 116°C . Anal. Found: C, 44.2; H, 2.3. Calc. for $\text{C}_{14}\text{H}_9\text{MnO}_6\text{S}_2$: C, 42.9; H, 2.3%. IR: $\nu(\text{CO})$ (CH_2Cl_2) 2093 (m), 2016 (vs), 2002 (vs), 1963 (s) cm^{-1} ; ^1H -NMR: δ 7.59 (1H, d, $3J_{\text{H-H}} = 8.6$ Hz, H3), 7.02 (1H, dd, $3J_{\text{H-H}} = 8.5$ Hz, $4J_{\text{H-H}} = 2.2$ Hz, H4), 6.95 (1H, d, $4J_{\text{H-H}} = 2.1$ Hz, H6), 3.92 (3H, s, OCH_3), 2.59 (3H, s, $\text{C}(\text{O})\text{CH}_3$). ^{13}C -NMR: δ 199.2 (C7), 161.9 (C5), 149.2 (C1), 131.2 (C3), 129.6 (C2), 116.1 (C4), 113.0 (C6), 56.2 (OCH_3), 29.6 ($\text{C}(\text{O})\text{CH}_3$); ESMS: (MeOH with added $\text{Na}[\text{OMe}]$, cone voltage 20 V), m/z 423 (33%) $[\text{M} + \text{OMe}]^-$; 395 (100%) $[\text{M} + \text{OMe} - \text{CO}]^-$; 363 (48%) $[\text{M} - \text{H} - \text{CO}]^-$; 589 (28%) $[\text{2M} - \text{Mn}(\text{CO})_5]^-$. This compound was also characterised by an X-ray crystal structure determination; see below.

2.3.2. Reaction of η^2 -(2-benzoylphenyl)tetracarbonylmanganese with CS_2

Cyclomanganated benzophenone (190 mg, 0.55 mmol) was reacted with CS_2 (ca. 10 ml) in a sealed ampoule at 85°C for 24 h, following the same procedure as above. The cooled ampoule was opened and the excess CS_2 removed under vacuum. The residue was chromatographed on an alumina column, with diethyl ether: petroleum spirits (1:1). The first fraction eluted

gave a dark orange oil which did not crystallise. This was characterised as η^2 -(2-benzoyl-phenyldithiocarboxylato)tetracarbonylmanganese, **5**, (156 mg, 67%). IR: $\nu(\text{CO})$ (CH_2Cl_2) 2093 (m), 2019 (vs), 2007 (vs), 1964 (s) cm^{-1} ; $^1\text{H-NMR}$: δ 7.86–7.43 (m, Ar–H). $^{13}\text{C NMR}$: δ 196.9 (–COPh), 145.4 (C–CS₂), 137.6–125.7 (other Ar); ESMS: (MeOH with added Na[OMe], cone voltage 10 V), m/z 455 (40%) [$\text{M} + \text{OMe}$][–]; 427 (100%) [$\text{M} + \text{OMe-CO}$][–]; 653 (78%) [$2\text{M} - \text{Mn}(\text{CO})_5$][–].

2.3.3. Reaction of η^2 -(2-acetylthien-3-yl)tetracarbonylmanganese with CS₂

The cyclomanganated thiophene derivative (112 mg, 0.38 mmol) was reacted with CS₂ (ca. 10 ml) in a sealed ampoule at 85 °C for 24 h, as above. Excess CS₂ was removed and the residue recrystallised from dichloromethane: petroleum spirits at –20 °C. Dark yellow rods of η^2 -(2-acetyl-thien-3-yl-dithiocarboxylato)tetracarbonylmanganese, **6**, were obtained (114 mg, 80%). M.p. 108 °C (dec.). Anal. Found: C, 36.1; H, 1.4. Calc. for C₁₁H₅MnO₅S₃: C, 35.9; H, 1.4%. IR: $\nu(\text{CO})$ (CH_2Cl_2) 2093 (m), 2016 (vs, br), 1966 (s) cm^{-1} ; $^1\text{H NMR}$: δ 7.45 (1H, d, $^3J_{\text{H-H}} = 5.0$ Hz, Ar–H), 7.24 (1H, d, $^3J_{\text{H-H}} = 5.0$ Hz, Ar–H), 2.66 (3H, s, –C(O)CH₃). $^{13}\text{C NMR}$: δ 191.4 (–C(O)CH₃), 150.4 (Ar), 140.0 (Ar), 129.8 (Ar–H), 128.4 (Ar–H), 30.0 (–C(O)CH₃). ESMS: (MeOH with added Na[OMe], cone voltage 10 V), m/z 399 (68%) [$\text{M} + \text{OMe}$][–]; 371 (74%) [$\text{M} + \text{OMe-CO}$][–]; 339, (16%) [M-CO-H][–]; 541 (100%) [$2\text{M} - \text{Mn}(\text{CO})_5$][–].

2.3.4. Reaction of η^2 -[(2-diphenylthiophosphinyl)-phenyl]tetracarbonylmanganese with CS₂

Cyclomanganated triphenylphosphine sulfide (**2b**) (125 mg, 0.31 mmol) was reacted with CS₂ as above. $^{31}\text{P-NMR}$ and ESMS on the crude reaction mixture showed the presence of the starting material **2b** and Ph₃PS. Excess CS₂ was removed and the residue recrystallised from CH₂Cl₂: Et₂O at –20 °C. A small yield of orange blocks were obtained, identified as (μ_3 -CS₃)₂Mn₄(CO)₁₆ (**7**). M.p. > 115 °C (dec.). Anal. Found: C, 23.5; H, 0.0. Calc. for C₁₈Mn₄O₁₆S₆: C, 24.5; H, 0.0%. IR: $\nu(\text{CO})$ (CH_2Cl_2) 2091 (s), 2049 (m), 2009 (vs, br), 1979 (m), 1965 (m) cm^{-1} ; ESMS: (MeCN, cone voltage 10 V), m/z 717 (100%) [$\text{M} - \text{Mn}(\text{CO})_4$][–]; 689 (27%) [$\text{M} - \text{Mn}(\text{CO})_4 - \text{CO}$][–]; 661, (8%) [$\text{M} - \text{Mn}(\text{CO})_4 - 2\text{CO}$][–]. The characterisation was confirmed by an X-ray crystal structure analysis; see below.

Similar reactions of orthomanganated triphenylphosphine selenide (**2c**), orthomanganated triphenylphosphine phenylimine (**2a**) and the orthomanganated chalcone (**3a**), also gave low yields of (μ_3 -CS₃)₂Mn₄(CO)₁₆ as the only carbonyl-containing product.

2.3.5. Reaction of dimanganese decacarbonyl with CS₂

Mn₂(CO)₁₀ (517 mg, 1.33 mmol) was reacted with CS₂ in a sealed ampoule at 110 °C for 24 h. The unopened ampoule was stored at –20 °C for 48 h, to give crystals of the crude product. These were collected by filtration and chromatographed on silica gel plates, eluting with CH₂Cl₂:petroleum spirits (1:1) to give as the major product (μ_3 -CS₃)₂Mn₄(CO)₁₆, **7**, (217 mg, 39%). Spectroscopic data was identical to that given in Section 2.3.4.

2.4. X-ray crystallography

For compound **7** unit cell parameters and intensity data were collected using a Siemens SMART CCD diffractometer, using standard collection procedures, with monochromatic Mo–K α X-rays (0.71073 Å), while for **4** a Siemens P4 four-circle diffractometer was used. Corrections for absorption and other effects were carried out with SADABS [16]. All other calculations used the SHELX-97 programs [17]. The structures were solved by direct methods, and developed routinely with refinement based on F^2 . All non-hydrogen atoms were assigned anisotropic temperature factors, and hydrogen atoms were included in calculated positions.

2.4.1. Crystal data for η^2 -(2-acetyl-5-methoxyphenyl-dithiocarboxylato)tetracarbonylmanganese (**4**)

C₁₄H₉MnO₆S₂, M_r 392.27, orthorhombic, $Pnma$, $a = 22.878(8)$, $b = 7.005(3)$, $c = 9.973(6)$ Å, $V = 1598.1(14)$ Å³, $D_{\text{calc}} = 1.630$ g cm^{–3}, $Z = 4$, $F(000) = 792$, $\mu(\text{Mo-K}\alpha) 1.113$ mm^{–1}, $T_{\text{max}} 0.262$, $T_{\text{min}} 0.238$, crystal size $0.66 \times 0.17 \times 0.04$ mm³. T 168 K.

A total of 1999 reflections, 1364 unique ($R_{\text{int}} 0.0407$) was collected $2 < \theta < 24^\circ$. Final $R_1 0.0415$ [data with $I > 2\sigma(I)$], 0.0767 (all data), $wR_2 0.0863$, GoF 1.014, final $\Delta e +0.37/-0.35$.

The structure is illustrated in Fig. 1, while selected bond parameters are listed in the caption.

2.4.2. Crystal data for (μ_3 -CS₃)₂Mn₄(CO)₁₆ (**7**)

C₁₈Mn₄O₁₆S₆, M_r 884.30, triclinic, $P\bar{1}$, $a = 7.010(3)$, $b = 8.410(4)$, $c = 13.674(6)$ Å, $\alpha = 83.40(1)$, $\beta = 80.04(1)$, $\gamma = 74.21(1)^\circ$, $V = 762.1(5)$ Å³, $D_{\text{calc}} = 1.927$ g cm^{–3}, $Z = 1$, $F(000) = 432$, $\mu(\text{Mo-K}\alpha) 2.099$ mm^{–1}, $T_{\text{max}} 1.00$, $T_{\text{min}} 0.76$, crystal size $0.26 \times 0.16 \times 0.05$ mm³. T 168 K.

A total of 9436 reflections, 3016 unique ($R_{\text{int}} 0.0333$) was collected $2 < \theta < 26.5^\circ$. Final $R_1 0.0309$ [data with $I > 2\sigma(I)$], $wR_2 0.0702$, GoF 0.954, final $\Delta e +0.63/-0.50$. The structure is illustrated in Fig. 2, with selected bond parameters in the caption.




Fig. 1. The structure of η^2 -(2-acetyl-5-methoxyphenyldithiocarboxylato)tetracarbonylmanganese (**4**). Bond parameters include: Mn(1)–S(1) 2.381(1), S(1)–C(1) 1.676(3), Mn(1)–C(11) 1.815(5), Mn(1)–C(12) 1.869(7) Å; S(1)–Mn(1)–S(1') 72.73(6)°, Mn(1)–S(1)–C(1) 86.2(2)°, S(1)–C(1)–S(1') 114.7(3)°.

3. Results and discussion

The reaction of the cyclometallated acetophenone **1** with CS₂ at 85 °C for 24 h gave a reasonable yield of the dithiocarboxylate complex (η^2 -MeOC₆H₄CS₂)Mn(CO)₄ (**4**), which crystallised on cooling the reaction mixture. The temperature of reaction was critical; below 80 °C little change took place while above 90 °C extensive decomposition occurred.

The new complex **4** showed in the IR spectrum a characteristic ν CO pattern for a *cis*-L₂Mn(CO)₄ species, with a shift of around 10 cm^{−1} to higher frequencies compared with the starting complex **1**. The ¹H- and ¹³C-

NMR spectra were consistent with the dithiocarboxylate structure. Similarly the electrospray mass spectrum, with Na[OMe] added as an ionisation aid [15], gave expected peaks arising from both OMe[−] addition and H⁺ abstraction from **4**. However, the spectroscopic data did not completely distinguish between the η^2 -RCS₂ bonding pattern as in **4**, and an alternative which has an η^1 -RCS₂ with the >C=O of the acetyl group still coordinated, as in **8**. Therefore, a single crystal X-ray determination was carried out to distinguish between the two possibilities.

The structure is illustrated in Fig. 1. It shows that structure **4** is adopted, with the Mn(CO)₄ group attached to the organic ligand by a symmetrical η^2 -RCS₂ linkage. The CS₂ group is orthogonal to the aryl plane of the rest of the ligand, and the C=O group is uncoordinated. As expected, the Mn–CO distances of the two CO ligands *trans* to each other are significantly longer than the Mn–CO bonds *trans* to the dithiocarboxylate ligand.

The corresponding reactions of orthomanganated benzophenone, and the orthomanganated 2-acetylthiophene complex, with CS₂ under the same conditions produced the equivalent dithiocarboxylate complexes **5** and **6** respectively. This suggests that for manganated aryl ketones the conversion of the C,O-bonded R ligand to a RCS₂ ligand is general.

A suggested pathway to **4–6** in these reactions involves initial insertion of CS₂ into the Mn–C bond to give a S,O-bonded ligand with a seven-membered chelate ring, as in **8**. Displacement of the O-donor by the second sulfur atom would give the final product. This pathway is consistent with reactions of orthomanganated complexes involving other unsaturated molecules.

Although compounds **4–6** are new examples, other dithiocarboxylate complexes of Mn(CO)₄ are established, having been prepared previously by insertion of CS₂ into the Mn–C bond of RMn(CO)₅ compounds





Fig. 2. The structure of (μ_3 -CS₃)₂Mn₄(CO)₁₆ (**7**). Bond parameters include: Mn(1)–S(1) 2.3866(11), Mn(1)–S(2) 2.3971(13), Mn(2)–S(3) 2.4035(13), C(1)–S(1) 1.691(3), C(1)–S(2) 1.699(3), C(1)–S(3) 1.763(3) Å; S(1)–Mn(1)–S(2) 73.18(3)°, Mn(2)–S(3)–Mn(2') 100.18(3)°, S(3)–Mn(2)–S(3') 79.82(3)°, Mn(1)–S(1)–C(1) 86.41(10)°, S(1)–C(1)–S(2) 114.52(16)°, S(1)–C(1)–S(3) 122.84(16)°, S(2)–C(1)–S(3) 122.53(16)°. The inset is a side view emphasising the Z-shape of the molecule.



[12]. This earlier method, however, gave much lower yields (1–17%) than those found in the present study for the cyclomanganated substrates (55–80%). There appear to be only two instances of the direct reaction of dithiocarboxylate anions with either $\text{BrMn}(\text{CO})_5$ or $[\text{Mn}(\text{CO})_5(\text{MeCN})]^+$ to give related species [18]. Therefore the reaction of CS_2 with orthomanganated aryl ketones may provide a useful route to dithiocarboxylates, especially those with functional groups that would interfere with the other syntheses.

The reaction of CS_2 with orthomanganated triphenyl phosphine sulfide, **2b**, was also investigated. This gave no evidence for the formation of a dithiocarboxylate, with the only new carbonyl-containing product being dark-yellow crystals isolated in low yield. Spectroscopic and analytical data were inconclusive, so an X-ray structure determination was carried out. This showed the product to be **7** as illustrated in Fig. 2. The centrosymmetric molecule consists of two η^3 -trithiocarbonate anions, each with two of the sulfur atoms chelating to a $\text{Mn}(\text{CO})_4$ group, while the third acts as a bridging atom between two other $\text{Mn}(\text{CO})_4$ groups. The CS_3^{2-} groups are planar, with shorter C–S bonds to the chelating S atoms (1.695 Å) than to the bridging one (1.763 Å). The pyramidity at the bridging S atom induces a Z shape to the molecule (see inset to Fig. 2). The overall molecule is a combination of two common motifs — $(\eta^2\text{-RCS}_2)\text{Mn}(\text{CO})_4$ as in the dithiocarboxylates discussed above [12,18] and related species [19], and a $(\mu\text{-SR})_2\text{Mn}_2(\text{CO})_8$ unit which is well-known for manganese carbonyl thiolates [20].

Also related to compound **7** are the $[(\eta^2\text{-CS}_3)\text{Mn}(\text{CO})_4]^-$ and $(\text{CO})_5\text{Mn}(\eta^3\text{-CS}_3)\text{Mn}(\text{CO})_4$ complexes found by Benson et al in reactions of $[\text{Mn}(\text{CO})_5]^-$ with CS_2 [21].

Compound **7** does not appear to have been reported before, but the isomorphous Re analogue was isolated from the reaction of $\text{F}_3\text{CRe}(\text{CO})_5$ with CS_2 [12,22].

The route to **7** is not clear. It seemed possible that the third S atom might have come from the orthomanganated $\text{Ph}_3\text{P}=\text{S}$, but this was discounted when **7** was also the product (albeit in low yield) when using the corresponding orthomanganated $\text{Ph}_3\text{P}=\text{Se}$ (**2c**); ESMS of the crude reaction mixture showed no traces of a dithioselenocarbonate analogue of **7** which would have been expected if atom transfer from the phosphine chalcogenide was involved.

Similarly $\text{Ph}_3\text{P}=\text{NPh}$ (**2a**) and chalcone (**3**) gave low yields of **7** when they were reacted with CS_2 under the same conditions.

To find a more rational synthesis of **7**, $\text{Mn}_2(\text{CO})_{10}$ was reacted with CS_2 . Although a higher temperature was needed (110 °C) the isolated yield of **7** was reasonable, 39%. It seems that formation of CS_3^{2-} ligands can readily arise from a formal redox disproportionation of CS_2 in reactions of a variety of manganese [21] (or rhenium [22] or nickel [14]) compounds.

4. Supplementary material

Crystallographic data for the structural analysis have been deposited with the Cambridge Crystallographic Data Centre, CCDC nos. 194981 and 194982 for compounds **4** and **7**, respectively. Copies of this information may be obtained free of charge from The Director, CCDC, 12 Union Road, Cambridge CB2 1EZ, UK (Fax: +44-1223-336033; e-mail: deposit@ccdc.cam.ac.uk or www: <http://www.ccdc.cam.ac.uk>).

Acknowledgements

We thank the University of Waikato for financial support and for a postgraduate scholarship (W.M.). We thank Associate Professor C.E.F. Rickard, University of Auckland, and Professor W.T. Robinson, University of Canterbury, for X-ray data sets.

References

- [1] (a) M.I. Bruce, *Angew. Chem. Int. Ed. Eng.* 16 (1977) 73;
(b) L. Main, B.K. Nicholson, *Adv. Metal-Org. Chem.* 3 (1994) 1;
(c) J.M. Vila, M. Gayosos, M.T. Pereira, A. Suarez, J.J. Fernandez, J.M. Ortigueira, A. Fernandez, M.L. Torres, *Trends Organometal. Chem.* 2 (1997) 21.
- [2] (a) R.J. McKinney, G. Firestein, H.D. Kaesz, *Inorg. Chem.* 14 (1975) 3066;
(b) J.M. Cooney, L.H.P. Gommans, L. Main, B.K. Nicholson, *J. Organometal. Chem.* 349 (1988) 197;
(c) N.P. Robinson, L. Main, B.K. Nicholson, *J. Organometal. Chem.* 349 (1988) 209;
(d) J.M. Cooney, L.H.P. Gommans, L. Main, B.K. Nicholson, *J. Organometal. Chem.* 634 (2001) 157.
- [3] (a) R.L. Bennett, M.I. Bruce, B.L. Goodall, M.Z. Iqbal, F.G.A. Stone, *J. Chem. Soc. Dalton Trans.* (1972) 1787.;
(b) R.L. Bennett, M.I. Bruce, I. Matsuda, *Aust. J. Chem.* 28 (1975) 1265;
(c) A. Suarez, J.M. Vila, M.T. Pereira, E. Gayoso, M. Gayoso, *J. Organometal. Chem.* 335 (1987) 359;
(d) R.M. Ceder, J. Sales, X. Solans, M. Font-Altaba, *J. Chem. Soc. Dalton Trans.* (1986) 1351.;
(e) J.M. Cooney, L. Main, B.K. Nicholson, *J. Organometal. Chem.* 516 (1996) 191;
(f) M.B. Dinger, L. Main, B.K. Nicholson, *J. Organometal. Chem.* 565 (1998) 125.
- [4] (a) G.J. Depree, N.D. Childerhouse, B.K. Nicholson, *J. Organometal. Chem.* 533 (1997) 143;
(b) M.A. Leeson, B.K. Nicholson, M.R. Olsen, *J. Organometal. Chem.* 579 (1999) 243.
- [5] W. Tully, L. Main, B.K. Nicholson, *J. Organometal. Chem.* 503 (1995) 75.
- [6] (a) L.H.P. Gommans, L. Main, B.K. Nicholson, *J. Chem. Soc. Chem. Commun.* (1987) 761.;
(b) R.C. Cambie, M.R. Metzler, P.S. Rutledge, P.D. Woodgate, *J. Organometal. Chem.* 429 (1992) 59;
(c) R.C. Cambie, M.R. Metzler, C.E.F. Rickard, P.S. Rutledge, P.D. Woodgate, *J. Organometal. Chem.* 426 (1992) 213;
(d) W. Tully, L. Main, B.K. Nicholson, *J. Organometal. Chem.* 503 (1995) 75;
(e) W.J. Grigsby, L. Main, B.K. Nicholson, *J. Organometal. Chem.* 540 (1997) 185.
- [7] (a) L.S. Liebeskind, J.R. Gasdaska, J.S. McCallum, S.J. Tremont, *J. Org. Chem.* 54 (1989) 669;
(b) N.P. Robinson, L. Main, B.K. Nicholson, *J. Organometal. Chem.* 364 (1989) C37;
(c) R.C. Cambie, M.R. Metzler, P.S. Rutledge, P.D. Woodgate, *J. Organometal. Chem.* 429 (1992) 41;
(d) W.J. Grigsby, L. Main, B.K. Nicholson, *Organometallics* 12 (1993) 397;
(e) W. Tully, L. Main, B.K. Nicholson, *J. Organometal. Chem.* 507 (1996) 103;
(f) W. Tully, L. Main, B.K. Nicholson, *J. Organometal. Chem.* 633 (2001) 162–172.
- [8] L.S. Liebeskind, S.A. Johnson, J.S. McCallum, *Tetrahedron Lett.* 31 (1990) 4397.
- [9] J.M. Cooney, C.V. Depree, L. Main, B.K. Nicholson, *J. Organometal. Chem.* 515 (1996) 109.
- [10] J.M. Cooney, L.H.P. Gommans, L. Main, B.K. Nicholson, *J. Organometal. Chem.* 336 (1987) 293.
- [11] (a) A. Wojcicki, *Adv. Organometal. Chem.* 12 (1974) 31;
(b) K.K. Pandey, *Coord. Chem. Rev.* 140 (1995) 37 (and references therein).
- [12] E. Lindner, R. Grimmer, *J. Organometal. Chem.* 25 (1970) 493.
- [13] A. Palazzi, L. Busetto, M. Graziana, *J. Organometal. Chem.* 30 (1971) 273.
- [14] J. Campora, E. Gutierrez, A. Monge, P. Palma, M.L. Poveda, *Organometallics* 13 (1994) 1728.
- [15] W. Henderson, J.S. McIndoe, B.K. Nicholson, P.J. Dyson, *J. Chem. Soc. Dalton Trans.* (1998) 519.
- [16] R.H. Blessing, *Acta Crystallogr. Sect. A* 51 (1995) 33.
- [17] G.M. Sheldrick, *SHELX-97*, Programs for X-ray Crystallography, University of Göttingen, Germany, 1997.
- [18] (a) U. Mikloweit, R. Mattes, *Z. Anorg. Allg. Chem.* 532 (1986) 145;
(b) M. Herberhold, J. Ott, L. Haumaier, *Chem. Ber.* 118 (1985) 3143.
- [19] D. Rehder, R. Kramolowsky, K.G. Steinhäuser, U. Kunze, A. Antoniadis, *Inorg. Chim. Acta* 73 (1983) 243.
- [20] (a) B.F.G. Johnson, P.J. Pollick, I.G. Williams, A. Wojcicki, *Inorg. Chem.* 7 (1968) 831;
(b) P.M. Trierchel, in: G. Wilkinson, F.G.A. Stone, E.W. Abel (Eds.), *Comprehensive Organometallic Chemistry*, vol. 4, Pergamon, Oxford, 1982, p. 103ff.
- [21] I.B. Benson, J. Hunt, S.A.R. Knox, V. Oliphant, *J. Chem. Soc. Dalton Trans.* (1978) 1240.
- [22] G. Thiele, G. Liehr, E. Lindner, *J. Organometal. Chem.* 70 (1974) 427.