

# Role of Trace Amine in the Metathesis of Imines by $\text{CpTa(=NR)Cl}_2$

Matthew C. Burland, Timothy W. Pontz, and Tara Y. Meyer\*

Department of Chemistry, University of Pittsburgh, Pittsburgh, Pennsylvania 15260


Received December 21, 2001

Tantalum imido compound  $\text{CpTa(=NBu}^t\text{)Cl}_2$  (**1**) reacts with imines to give NR exchange (metathesis) products. Addition of <5 mol % of compound **1** to a 1:1 mixture of two different imines resulted in catalytic cross metathesis of the two imines to give an equilibrium mixture of imines. Kinetic studies of the reaction of **1** with excess imine  $(\text{Bu}^t\text{)HC=N}(p\text{-tolyl})$  (**2a**) gave an average  $k_{\text{obs}}$  of  $1.29 \times 10^{-4} \text{ min}^{-1}$  ( $\sigma$  0.28). Pseudo-first-order behavior was observed for **1**. The dependence of  $k_{\text{obs}}$  on the concentration of **2a** was found to be nonlinear and dependent on catalytic amounts of contaminant amine  $p$ -tolylamine (**9**). Reaction of **1** with *tert*-butylamine (**6**) results in the formation of the imide/amide  $\text{CpTa(=NBu}^t\text{)(NHBu}^t\text{)Cl}$  (**7**) and HCl. Compound **7** did not react with imines to give exchange products, so it is proposed that the reaction of **1** with imines is significantly accelerated by the presence of HCl.

## Introduction

We have discovered that compounds of the general formula  $\text{CpTa(=NR)Cl}_2$  represent a new class of imine metathesis catalysts. The impetus for these studies originated in our previous investigation of the imine reactivity of the group VI bis(imide) complexes  $(\text{DME})\text{Cl}_2\text{-Mo(=NR)}_2$ .<sup>1–3</sup> Although we had gained a significant understanding of the overall catalytic cycle, the complexity inherent in the dual-active site catalysts precluded the detailed mechanistic studies that are necessary to understand the system and enable the eventual application of the imine metathesis strategy to the catalytic synthesis of small molecules and polymers.

Interestingly, the two other examples of imine/imide exchange for which mechanistic data were reported apparently proceed by different pathways. The viability of a Chauvin-type<sup>4</sup> mechanism (Figure 1) was established by Bergman and co-workers in their observation of isolable diazametallacyclic intermediates in the reaction of  $\text{CpCp}'\text{Zr(=NR)(THF)}$  ( $\text{Cp}' = \text{Cp}, \text{Cp}^*$ ) with imines.<sup>5–7</sup> In contrast, Mountford and co-workers observed neither diazametallacycles nor kinetics consistent with a [2+2] mechanism and proposed an amine-catalyzed process to explain the reaction of  $(\text{py})_3\text{Cl}_2\text{Ti(=NR)}$  with imines.<sup>8,9</sup> Other examples of imine/imide and imine/carbodiimide metathesis are known, but little mechanistic data has been reported for these sys-



**Figure 1.** Chauvin-type mechanism for imine metathesis at an electrophilic metal center.

tems.<sup>10–13</sup> Given the variance in precedents, the determination of mechanism in our system is of particular importance to us as well as others.

Since the dual-site molybdenum catalysts proved unsuitable for kinetic studies, we sought to discover a new single-site catalyst. Metal imides of the type  $\text{CpTa(=NR)Cl}_2$  were selected because they possess a single imide functional group while being approximately isolobal with the molybdenum bis(imide) complex *sans* DME. Herein, we describe both the stoichiometric and catalytic metathetic reactions of these tantalum mono(imide) complexes with imines and, on the basis of kinetic analyses and <sup>15</sup>N-labeling studies, propose an acid-catalyzed mechanism.

## Results

**Imide/Imine Metathesis.**  $\text{CpTa(=NBu}^t\text{)Cl}_2$ , **1**, reacted with 15 equiv of  $(\text{Bu}^t\text{)HC=N}(p\text{-tolyl})$ , **2a**, in  $\text{C}_6\text{D}_6$  at 70 °C to yield imine  $(\text{Bu}^t\text{)HC=N}(\text{Bu}^t)$ , **4a**, as shown by comparison of the <sup>1</sup>H NMR spectroscopic resonances with those of an authentic sample (Scheme 1). A new Cp-containing compound, assigned as  $p$ -tolyl imide **3a**, was also produced, as evidenced by distinctive <sup>1</sup>H NMR resonances at  $\delta$  5.72 (Cp) and 2.19 (tolyl  $\text{CH}_3$ ). No other species were observed. Within 3–4 days nearly all of the starting imide **1** was converted to the new product **3a**. No precipitates formed and integration versus an

(1) Cantrell, G. K.; Meyer, T. Y. *J. Chem. Soc., Chem. Commun.* **1997**, 1551–1552.

(2) Cantrell, G. K.; Meyer, T. Y. *Organometallics* **1997**, *16*, 5381–5383.

(3) Cantrell, G. K.; Meyer, T. Y. *J. Am. Chem. Soc.* **1998**, *120*, 8035–8042.

(4) Hérisson, J.-L.; Chauvin, Y. *Macromol. Chem.* **1970**, *141*, 161.

(5) Krska, S. W.; Zuckerman, R. L.; Bergman, R. G. *J. Am. Chem. Soc.* **1998**, *120*, 11828–11829.

(6) Meyer, K. E.; Walsh, P. J.; Bergman, R. G. *J. Am. Chem. Soc.* **1994**, *116*, 2669–2670.

(7) Meyer, K. E.; Walsh, P. J.; Bergman, R. G. *J. Am. Chem. Soc.* **1995**, *117*, 974–985.

(8) McInnes, J. M.; Mountford, P. *J. Chem. Soc., Chem. Commun.* **1998**, 1669–1670.

(9) Mountford, P. *J. Chem. Soc., Chem. Commun.* **1997**, 2127–2134.


(10) Bruno, J. W.; Liu, X. J. *Organometallics* **2000**, *19*, 4672–4674.

(11) Royo, P.; Sánchez-Nieves, J. *J. Organomet. Chem.* **2000**, *597*, 61–68.

(12) Birdwhistell, K. R.; Lanza, J.; Pasos, J. *J. Organomet. Chem.* **1999**, *584*, 200–205.

(13) Weiss, K.; Kindl, P. *Angew. Chem., Int. Ed. Engl.* **1984**, *23*, 629–631.

Scheme 1




internal standard confirmed that all species remained soluble.

Tantalum imide **1** also reacted metathetically in C<sub>6</sub>D<sub>6</sub> with excess (Ph)HC=N(Pr<sup>n</sup>), **2b**, to give imine (Ph)HC=N(Bu<sup>t</sup>), **4b**, and a product assigned as CpTa(=NPr<sup>n</sup>)Cl<sub>2</sub>, **3b**. Imine **4b** was identified in the reaction mixture by comparison with independently prepared compound. The assignment of the exchange product was made on the basis of the Cp resonance at  $\delta$  5.74 and a multiplet at  $\delta$  4.23 that is characteristic of the  $\alpha$ -methylene of an N-alkyl group. The resonances for the hydrogens on the  $\beta$ - and  $\gamma$ -carbons of this new species appear to overlap with those of the starting material imide. After 1 day at 55 °C a third species, **5**, appeared. Compound **5** was identifiable only from a new Cp resonance at  $\delta$  6.10. Either this compound has no N-alkyl group or its resonances overlap with those of **1** and/or **3b**. We speculate that **5** may be the product of  $\beta$ -hydride elimination, which is unable to occur in the case of the *tert*-butyl imide, **1**, and the *p*-tolyl imide **3a**. After 10 days of heating there were no new species present by <sup>1</sup>H NMR spectroscopy, but precipitate was evident and the integration of all species against an internal standard showed a confirming decrease in the concentration of Cp-containing complexes.


**Catalytic Imine Metathesis.** CpTa(=NBu<sup>t</sup>)Cl<sub>2</sub>, **1**, acts as an imine metathesis catalyst. Addition of <5 mol % of **1** to a 1:1 mixture of imines **2a** and **2b** produces the metathesis products (Bu<sup>t</sup>)HC=N(Pr<sup>n</sup>) and (Ph)HC=N(*p*-tolyl) within hours at 70 °C (Scheme 2). The reaction is slow, reaching equilibrium after >1 week. The imines produced by imide/imine exchange and the imide products, **3a** and **3b**, were also observed in the reaction mixture. At longer reaction times (>2 weeks), precipitate was observed.

**Kinetic Studies.** The reaction of **1** with *N*-*p*-tolyl imine **2a** was monitored kinetically by <sup>1</sup>H NMR spectroscopy (Scheme 3). In the presence of excess imine, pseudo-first-order behavior was observed for **1**. The validity of a first-order analysis was confirmed by following several reactions for more than 5 half-lives and by the independence of the rate constant on the concentration of **1**. The observed pseudo-first-order rate constants,  $k_{\text{obs}}$ , were obtained by fitting the  $[\text{Imide } \mathbf{1}]_t - [\text{Imide } \mathbf{1}]_{\text{eq}}$  to a single-exponential function (Figure 2). An average  $K_{\text{eq}}$ , defined as  $\{[\text{Imide } \mathbf{3a}][\text{Imine } \mathbf{4a}]\}/\{[\text{Imide } \mathbf{1}][\text{Imine } \mathbf{2a}]\}$ , of 1.29 ( $\sigma = 0.28$ ) was calculated from the data for experiments 2–9 (experiment 1 omitted because only a 6.3:1 excess was used).

To determine the order of the imine in the reaction, the dependence of  $k_{\text{obs}}$  on imine was examined. The plot of  $k_{\text{obs}}$  versus [Imine] for experiments 1–9 did not show a simple linear dependence (Figure 3). Moreover, two control experiments established that the reaction depended, at least partially, on a catalytic contaminant




**Figure 2.** Pseudo-first-order disappearance of CpTa(=NBu<sup>t</sup>)Cl<sub>2</sub>, **1**, in the presence of 15 equiv of (Bu<sup>t</sup>)HC=NTol, **2a** ( $k_{\text{obs}} = 2.6 \times 10^{-4} \text{ min}^{-1}$ ). Corresponds to experiment 5 in Table 1.



**Figure 3.** Dependence of the pseudo-first-order rate constant ( $k_{\text{obs}}$ ) for the disappearance of **1** on imine concentration: Experiments 1–9 were performed with dried and distilled imine (Imine1x); experiment 10 used doubly purified imine (Imine2x). The imine1x data have been fitted with a polynomial to highlight the nonlinear trend. The curve does not represent a specific rate law.

present in the imine. In the first control experiment, the original kinetic experiment (Scheme 3) was repeated with *doubly purified imine*; the  $k_{\text{obs}}$  obtained was significantly smaller than the rate constant that would have been predicted on the basis of data obtained from the singly purified imine (Table 1, experiment 10, Figure 2). In the second control experiment, the relative ratios of imine to imide were reversed such that there was an excess of imide **1**. The resulting reaction was extremely slow, with only trace amounts of product visible after 1 week at 70 °C (not included in table).


Scheme 2



Scheme 3



Scheme 4



Since the most likely catalytically active contaminant of the imine is amine, the reaction of **1** with excess *p*-tolylimine **2a** was repeated in the presence of added *p*-tolylamine **9** (0.1–0.6 equiv with respect to **1**, Scheme 4). A plot of the  $k_{\text{obs}}$  obtained from these experiments showed a direct dependence on amine concentration (Table 1, experiments 11–13, Figure 4). Although the trend appears linear, implying a first-order dependence on amine, the experiments covered a relatively small range of concentrations and the relationship is likely not one of simple proportionality. Experimental difficulties precluded studies of significantly lower or higher concentrations.

Several attempts to remove the amine from the reaction mixture failed. Any reagent sufficiently acidic or basic enough to completely sequester the amine also reacted in some fashion with the imide complex or with the imine. The amine must be accepted as a necessary component of any reaction mixture containing excess imine.

**Amine/Imide Reactions.** To probe the role of amine in the imide/imine metatheses, the direct reaction of *tert*-butyl imide **1** with *tert*-butylamine, **6**, was examined (Scheme 5). By  $^1\text{H}$  NMR spectroscopy of the reaction mixtures, the instantaneous formation of a new imide/amide complex,  $\text{CpTa(=NBu}^t\text{)(NHBu}^t\text{)Cl}$  (**7**), was observed. Formation of the ammonium salt  $\text{NH}_3\text{Bu}^t\text{Cl}$  (**8**) was also noted during the course of the reaction. The imide/amide was the only soluble product observed, independent of the ratio of amine to imide; addition of a single equivalent of amine gave 0.5 equiv of compound **7**. Although we could not isolate imide/amide **7** in analytically pure form nor could we obtain X-ray quality crystals, its formulation was consistent with all spectroscopic data and an analogous  $\text{Cp}^*$  derivative is known.<sup>11</sup> A 2D heteronuclear multiple-bond correlation (HMBC) NMR spectrum showed the expected long-range coupling between the amide proton and the *ipso* carbon of the attached *tert*-butyl group. Also, the  $^{15}\text{N}$


Table 1. Kinetic Data for Imide/Imine Exchange Reactions

	[Imide <b>1</b> ]/M	[Imine <b>2a</b> ] <sub>init</sub> /M	[Amine <b>9</b> ] <sub>init</sub> /M	$k_{\text{obs}}$ ( $10^{-4}/\text{min}^{-1}$ )
Imide + Excess Imine <sup>d</sup>				
1	0.027	0.17	<0.002 <sup>b</sup>	1.2 <sup>a</sup>
2	0.022	0.22	<0.002 <sup>b</sup>	1.2
3	0.026	0.25	<0.002 <sup>b</sup>	1.7
4	0.022	0.32	<0.002 <sup>b</sup>	3.3
5	0.022	0.32	<0.002 <sup>b</sup>	2.6
6	0.022	0.36	<0.002 <sup>b</sup>	3.7
7	0.028	0.42	<0.004 <sup>b</sup>	4.5
8	0.039	0.58	<0.004 <sup>b</sup>	5.7
9	0.022	0.43	<0.004 <sup>b</sup>	4.7
Imide + Excess Doubly Purified Imine <sup>d</sup>				
10	0.028	0.42 <sup>c</sup>	<0.002 <sup>b</sup>	1.2
Imide + Excess Doubly Purified Imine + Amine <sup>e</sup>				
11	0.018	0.27 <sup>c</sup>	0.0018	9.6
12	0.018	0.27 <sup>c</sup>	0.0037	10.5
13	0.019	0.28 <sup>c</sup>	0.0077	12.6
14	0.018	0.27 <sup>c</sup>	0.011	14.5

<sup>a</sup>  $k_{\text{obs}}$  for this experiment is less certain due to an insufficient excess of imine to imide (6.3:1). <sup>b</sup> Upper limit based on NMR sensitivity and/or integration of observed baseline signals that are associated with trace amine. <sup>c</sup> Doubly purified imine. <sup>d</sup> Estimated uncertainty in  $k_{\text{obs}} = \pm 0.6 \times 10^{-4}$  based on uncertainties in concentration, integration, and [Imine **2a**]<sub>eq</sub>. <sup>e</sup> Estimated uncertainty in  $k_{\text{obs}} = \pm 2.5 \times 10^{-4}$  based on uncertainties in concentration, integration, and [Imine **2a**]<sub>eq</sub>. Amine **9** = *p*-tolylamine.


NMR spectra of the partially labeled and fully labeled derivatives  $\text{CpTa(=NBu}^t\text{)(}^{15}\text{NHBu}^t\text{)Cl}$  (**7N1a**) and  $\text{CpTa(}^{15}\text{NBu}^t\text{)(}^{15}\text{NHBu}^t\text{)Cl}$  (**7N2**) were consistent (vide supra).

The reaction of *tert*-butyl imide **1** with *p*-tolylamine **9** was also examined by  $^1\text{H}$  NMR (Scheme 6). The reaction mixture was significantly more complex than the reaction of **1** with *tert*-butylamine **6**. At equilibrium, at least five species were present including starting imide (20%) and the amine/imide exchange product,  $\text{CpTa(=NTol)Cl}_2$  **3a** (15%). Significant precipitate, along with a broad resonance at  $\delta$  8.0, was consistent with the formation of ammonium salts. Although we were



**Figure 4.** Dependence of the pseudo-first-order rate constant ( $k_{\text{obs}}$ ) for the disappearance of **1** vs amine concentration for a ternary reaction of **1**, 15 equiv of imine **2a**, and *p*-tolylamine **9**.

**Scheme 5**



unable to identify all species in solution, by analogy to the *tert*-butylamine reactions, some or all of the imide/amide complexes are likely to be present: CpTa(=NBu<sup>t</sup>)(NHTol)Cl (**10a**), CpTa(=NTol)(NHBu<sup>t</sup>)Cl (**10b**), CpTa(=NBu<sup>t</sup>)(NHBu<sup>t</sup>)Cl (**7**), and CpTa(=NTol)(NHTol)Cl (**10c**). The ratios of all species were found to vary reversibly as a function of temperature.

The ternary reaction of tantalum imide **1**, *p*-tolylamine **9** (0.5 equiv), and *N*-tolylimine **2a** (15 equiv) was also studied by <sup>1</sup>H NMR and, initially, was found to contain all of the species seen in the binary imide/amine reaction (Scheme 7). After 6 days at 70 °C, the reaction had reached equilibrium and the spectrum was much simplified. Only three Cp-containing species remained: the product **3a** (87%), starting *tert*-butylimide **1** (4%), and a compound whose Cp resonance corresponded exactly with the one observed in the binary imide/tolylamine reaction (Scheme 6) for the most abundant of the putative imide/amide compounds **10a–c**. Imide/amide **7** can be ruled out on the basis of the known shift.

**Reaction of Imide/Amide (7) with Imine.** To determine if imide/amide **7** generated in situ might be responsible for the amine-accelerated rates of imine/imide exchange, we examined the reaction of *isolated* imide/amide **7** with 15 equiv of imine (Bu<sup>t</sup>)CH=N(*p*-tolyl) (**2a**) (Scheme 8). Interestingly, no new imine product was detected after 2 days at 70 °C. The control, consisting of imide **1** and 15 equiv of the same imine, had proceeded to completion in the same period.

**<sup>15</sup>N-Labeling Studies.** Studies involving <sup>15</sup>N-labeled substrates provided further insight into these complex reaction systems. We prepared <sup>15</sup>NH<sub>2</sub>Bu<sup>t</sup> (**6N**) and (Ph)HC=N(Bu<sup>t</sup>) (**4bN**) using standard methods. The reaction of the labeled amine **6N** with imide **1** was studied over time by <sup>1</sup>H NMR and <sup>15</sup>N NMR spectroscopies (Scheme 9). Within a minute of the addition of a single

equivalent amine **6N** to the imide in C<sub>6</sub>D<sub>6</sub>, a <sup>1</sup>H NMR spectrum showed that half of the starting material had been converted to the partially labeled imide/amide **7N1a**. The amine was completely consumed, and precipitate, presumably ammonium salt, was visible. Long-range <sup>3</sup>J<sub>NH</sub> coupling of 2.2 Hz was observed from the amide nitrogen to the methyl peaks of the attached *tert*-butyl group. No such coupling was observed to the imide *tert*-butyl of **7N1a** or in the remaining starting material **1**. After heating for 2 h at 80 °C, the label began to wash into the imide positions, as evidenced by the broadening and splitting of the *tert*-butyl peaks. In the <sup>15</sup>N NMR spectra, parallel behavior was observed. Immediately after addition, only the amide resonance at δ 116.7 was easily detectable. *No resonance for the labeled free amine 6N was observed*, indicating that it had been completely consumed. After heating, the resonances for the imide groups for both **7N2** and imide complex **1N** were easily visible. Presumably the monolabeled compound CpTa(<sup>15</sup>NBu<sup>t</sup>)(NHBu<sup>t</sup>)Cl (**7N1b**) is also present, but it cannot be spectroscopically differentiated in mixtures containing **7N2**.


The ternary experiment involving stoichiometric imide **1**, (Ph)HC=N(Bu<sup>t</sup>) (**4b**), and labeled amine **6N** was also followed by NMR. Initial <sup>15</sup>N NMR spectra showed only the partially labeled imide/amide **7N1a**. With heating, resonances for the imide functionalities of **1N** and **7N2** became visible, as did the imine nitrogen for **4bN**. Again, no free amine was observed. The <sup>15</sup>N spectra were collected with an NOE-suppressing pulse sequence in order to make it possible to integrate the peaks. Although the differences were not dramatic, it appeared as if the label washed into the imide positions faster than into the imine (Scheme 10). A control experiment with the imine **4b** and the labeled amine **6N** showed no scrambling of the isotopic label, even after heating at 80 °C for 1 h.

Reaction of the labeled imine **4bN** with imide **1** and unlabeled amine **6** was also followed by <sup>15</sup>N and <sup>1</sup>H NMR spectroscopy (Scheme 11). After initial formation of imide/amide **7**, the resonances associated with the imide **1N** and the imide/amide **7N1a** and **7N2** were seen to grow in, while the imine resonance for **4bN** was reduced. It appeared that the <sup>15</sup>N label was incorporated into the imide **1N** at a faster rate than into the imide/amide **7N2**. No free amine was seen at any point during the reaction.


To determine if the precipitated ammonium salt played a role in our reaction, or whether the nonpolar nature of the solvent excluded its participation, we conducted experiments to determine if NR exchange between solution phase species and the precipitated ammonium salt was possible (Scheme 12). We heated a sample of imide **1** in C<sub>6</sub>D<sub>6</sub> with 2 equiv of solid, <sup>15</sup>N-labeled ammonium salt <sup>15</sup>BuNH<sub>3</sub>Cl, **8N**. Although the solid did not appear to dissolve after several days of heating at 70 °C, a <sup>15</sup>N NMR spectrum of the sample clearly showed the presence of labeled imide **1N**. A similar experiment with the *tert*-butylamine **6** and the labeled ammonium salt **8N** also showed incorporation of the label despite the nonpolar nature of the solvent. As a comparison, a sample of unlabeled amine showed no detectable signal in the <sup>15</sup>N NMR spectrum after a similar period of data collection.




Scheme 6




Scheme 7



Scheme 8



Scheme 9



### Discussion

$\text{CpTa(=NBu}^t\text{)Cl}_2$ , **1**, undergoes imide/imine exchange and catalyzes the metathesis of imines, but the primary mechanism appears to involve trace amine. The fact that the reaction is first order in **1** and that the rate depends on amine suggests that the reaction of these two components may be the rate-determining step of the reaction. These data rule out a Chauvin-type mechanism, which would be expected to be zero order in amine.


It is, at this point, relevant to discuss in more detail the mechanism proposed by Philip Mountford to explain the zero-order dependence on  $(\text{py})_3\text{Cl}_2\text{Ti(=NBu}^t\text{)}$  of the  $\text{=NR}$  exchange reaction with imine.<sup>8</sup> In the first step, Mountford postulates that a small amount of *tert*-butylamine reacts with the tolylimine to produce free *p*-tolylamine. Mountford and co-workers had previously established that *p*-tolylamine reacts with the butylimide titanium complex to regenerate the *tert*-butylamine and the product *p*-tolylimide complex (Scheme 13). If these two steps are combined, the overall reaction is catalytic in amine, and if the first step is rate determining, the zero-order dependence on imide complex is explained. In addition, the titanium complexes are available to act as Lewis acid catalysts for the amine/imine reaction.

An analogous mechanism can be proposed for the reaction of **1** with imine (Scheme 14). In this case, however, the relative rates of the two steps must be reversed in order to explain the first-order dependence of the reaction rate on **1**. It is also necessary to postulate that, in contrast with the Mountford system, step 2, the amine/imide exchange, is reversible.


In addition to the amine-catalyzed pathway, two alternative mechanisms can be postulated if, instead of assuming that the amine acts as catalyst as depicted in Scheme 14, it is hypothesized that one of the products of the amine reaction with imide **1** is the mediator (Scheme 15). The two products from the reaction of *tert*-butylamine **6** with imide **1** are the imide/amide **7** and the HCl that is sequestered as the ammonium salt,  $\text{NH}_3\text{-Bu}^t\text{Cl}$  **8**. The amide functional group of **7** could conceivably act as the mediator for the  $\text{NR}$  exchange reaction as shown in Scheme 16. The 1,3-proton transfer required could be intermolecular or intramolecular.

HCl as catalyst is also intriguing. Although it is mostly consumed by reaction with the basic *tert*-butylamine **6**, it is possible that some remains present in the broad reaction mixture. The observed exchange of  $^{15}\text{N}$


Scheme 10



Scheme 11



Scheme 12



label from ammonium salt into soluble imide and imine suggests that the amine-bound HCl can interact with the other species in solution. Moreover, HCl could theoretically act as a catalyst for virtually any reaction combination in the mixture: imide/imide, imide/amine,

imide/amine, amine/imine, imide/imine, amide/amine, and imine/imine (Scheme 17).

In considering these three mechanisms, we currently favor the third, HCl catalysis, as the most significant. The first mechanism, the amine-mediated pathway postulated by Mountford, is likely to occur at some rate in the reaction mixture. Both of the constituent reaction steps have been observed. However, this mechanism alone cannot explain some of our observations. First, the reactions of imide **1** with amines are not straightforward imide/amine exchanges. In contrast with the titanium complex studied by Mountford and co-workers, which gives simple amine/imide exchange, the reaction of amine with  $\text{CpTa}(\text{=NBu}^t)\text{Cl}_2$ , **1**, gives multiple products, including at least one stable imide/amide. Second, the strongly Lewis acidic **1** appears to nearly consume added amine. The concentration, therefore, of amine in solution when it originates as a trace contaminant must be very low. It is worth noting that the pyridine ligands coordinated to the titanium would be expected to diminish the Lewis acidity of the Mountford complex



$^{13}\text{C}$ , and  $^{15}\text{N}$  NMR spectra were recorded with a Bruker AF500 or AF300 NMR spectrometer. Chemical shifts for  $^1\text{H}$  and  $^{13}\text{C}$  are referenced to the residual protio impurity in the deuterated solvent.  $^{15}\text{N}$  chemical shifts are referenced to formamide used as an external standard.  $^{14}\text{N}$  NMR spectra were recorded by Dr. M. Minelli, Grinnell College, Iowa. They were measured on a Bruker AC 300 MHz NMR spectrometer with a 10 mm broadband probehead (109Ag-31P) with digital tuning. Nitromethane neat was used as external reference (0 ppm).

**CpTa(=NBu $^t$ )Cl $_2$  (1)** was prepared according to the method of Gibson for CpCl $_2$ Nb(=NBu $^t$ ) with slight modification.<sup>18</sup> To a stirred, room-temperature suspension of CpTaCl $_4$  (1.38 g, 3.57 mmol) in 1,2-dichloroethane (50 mL) in a 100 mL Schlenk flask was slowly added (TMS)NBu $^t$  (1.04 g, 7.17 mmol). The flask was fitted with a reflux condenser, and the suspension was refluxed under nitrogen overnight. The solvent was removed under vacuum, and the resulting yellow-brown solid was redissolved in hexane. The solution was filtered to remove an insoluble tan powder, and the filtrate was stored at  $-35^\circ\text{C}$  overnight. The product crystals were isolated as golden yellow needles by filtration in 72% yield (0.99 g, 2.55 mmol).  $^1\text{H}$  NMR (300.13 MHz, C $_6$ D $_6$ ,  $25^\circ\text{C}$ ):  $\delta$  1.10 (s, 9H, C(CH $_3$ ) $_3$ ), 5.93 (s, 5H, C $_5$ H $_5$ ).  $^{13}\text{C}\{^1\text{H}\}$  NMR (75.4 MHz, CDCl $_3$ ,  $25^\circ\text{C}$ ):  $\delta$  32.6 (C(CH $_3$ ) $_3$ ), 66.7 (C(CH $_3$ ) $_3$ ), 111.6 (C $_5$ H $_5$ ).  $^{14}\text{N}\{^1\text{H}\}$  NMR (21.69 MHz, CH $_2$ Cl $_2$ ,  $25^\circ\text{C}$ ):  $\delta$  30 (NC(CH $_3$ ) $_3$ ).  $^{15}\text{N}$  NMR of CpTa(=NBu $^t$ )Cl $_2$  (**1N**) (50.68 MHz, C $_6$ D $_6$ ,  $25^\circ\text{C}$ ):  $\delta$  292.4 (NC(CH $_3$ ) $_3$ ).

**CpTa(=NBu $^t$ )(NHBu $^t$ )Cl (7).** To a solution of CpTa(=NBu $^t$ )Cl $_2$  (0.49 g, 1.27 mmol) in toluene (5 mL) was added H $_2$ NBu $^t$  (300  $\mu\text{L}$ , 2.85 mmol). Precipitate formed immediately, and the solution was filtered through Celite to remove the ammonium salt. The solvent was removed from the filtrate under vacuum to give the product as an off-white powder in 86% yield (0.46 g, 1.09 mmol). Repeated recrystallizations did not give material suitable for chemical analysis nor for X-ray diffraction.  $^1\text{H}$  NMR (300.13 MHz, C $_6$ D $_6$ ,  $25^\circ\text{C}$ ):  $\delta$  1.05 (s, 9H, NC(CH $_3$ ) $_3$ ), 1.13 (s, 9H, NHC(CH $_3$ ) $_3$ ), 5.88 (s, 5H, C $_5$ H $_5$ ), 6.40 (s, 1H, NHC(CH $_3$ ) $_3$ ).  $^{13}\text{C}\{^1\text{H}\}$  NMR (75.4 MHz, CDCl $_3$ ,  $25^\circ\text{C}$ ):  $\delta$  33.6 (NC(CH $_3$ ) $_3$ ), 34.5 (NHC(CH $_3$ ) $_3$ ), 56.5 (NHC(CH $_3$ ) $_3$ ), 65.5 (NC(CH $_3$ ) $_3$ ), 108.5 (C $_5$ H $_5$ ).  $^{15}\text{N}$  NMR of CpTa(=NBu $^t$ )(NHBu $^t$ )Cl (**7N2**) (50.68 MHz, C $_6$ D $_6$ ,  $25^\circ\text{C}$ ):  $\delta$  116.7 (NHC(CH $_3$ ) $_3$ ), 267.0 (NC(CH $_3$ ) $_3$ ).

**$^{15}\text{NH}_2\text{Bu}^t$  (6N).** To a slurry  $^{15}\text{NH}_3\text{Bu}^t\text{Cl}$  (0.30 g, 2.78 mmol) in 1.75 mL of C $_6$ D $_6$  in a three-necked round-bottom flask under N $_2$  was added solid NaOH (0.2 g, 5 mmol, 1.8 equiv). Deionized water (250  $\mu\text{L}$ ) was added via syringe to help dissolve the solids. Some solid remained undissolved. The contents of the flask were stirred for 2 h and then vacuum transferred into a round-bottom flask containing CaH $_2$ . After stirring for 24 h, the solution was vacuum transferred again into a clean flask containing more CaH $_2$  and stirred overnight before being vacuum transferred into a clean, dry flask. The dry C $_6$ D $_6$  solution of the labeled amine was diluted to a total volume of 3 mL by addition of C $_6$ D $_6$ . An 80  $\mu\text{L}$  aliquot was added to a solution of 50  $\mu\text{L}$  of anisole (0.73 mmol) in C $_6$ D $_6$  in an NMR tube. Integration of the  $^1\text{H}$  NMR spectrum established that the concentration of the bulk solution was 0.626 M (68% yield).  $^{15}\text{N}$  NMR spectroscopy determined that the label had been incorporated into the product.  $^1\text{H}$  NMR (300.13 MHz, C $_6$ D $_6$ ,  $25^\circ\text{C}$ ):  $\delta$  0.98 (br, 2H,  $^{15}\text{NH}_2$ ), 3.29 (d, 9H, CH $_3$ ,  $^3J_{\text{NH}} = 2.3$  Hz).  $^{15}\text{N}$  NMR (50.68 MHz, C $_6$ D $_6$ ,  $25^\circ\text{C}$ ):  $\delta$   $-53.6$  (NH $_2$ ).

(15) TMS amines were synthesized by either (a) the addition of TMSCl to the parent lithium amine or (b) the reaction of TMSCl with the parent amine in basic solution.

(16) Glueck, D. S.; Wu, J.; Hollander, F. J.; Bergman, R. G. *J. Am. Chem. Soc.* **1991**, *113*, 2041–2054.

(17) Imines were prepared by stirring a benzene solution of the corresponding amine and aldehyde over molecular sieves, followed by filtration, removal of solvent in vacuo, and vacuum distillation.

(18) Williams, D. N.; Mitchell, J. P.; Poole, A. D.; Siemeling, U.; Clegg, W.; Hockless, D. C. R.; O'Neil, P. A.; Gibson, V. C. *J. Chem. Soc., Dalton Trans.* **1992**, 739–751.

**(Ph)HC=NBu $^t$  (4bN)** was prepared by standard methods.<sup>17</sup>  $^1\text{H}$  NMR (300.13 MHz, C $_6$ D $_6$ ,  $25^\circ\text{C}$ ):  $\delta$  8.12 (d, 1 H,  $^{15}\text{N}=\text{CHPh}$ ,  $J_{\text{NH}} = 3.8$  Hz), 7.80 (d, 2 H, CHPh), 7.06–7.26 (m, 3 H, CHPh), 1.23 (d, 9 H,  $^{15}\text{NC}(\text{CH}_3)_3$ ,  $J_{\text{NH}} = 1.3$  Hz).  $^{15}\text{N}$  NMR (50.68 MHz, C $_6$ D $_6$ ,  $25^\circ\text{C}$ ):  $\delta$  244.1.

**General Procedure for NMR Experiments.** Imine and amine standard solutions were prepared by weighing the imine into a volumetric flask and diluting with C $_6$ D $_6$ . Likewise, standard solutions of the internal standards were prepared using volumetric flasks. The necessary amounts of these solutions were then transferred into an NMR tube equipped with a Teflon stopcock. The catalyst was either weighed into a vial, dissolved in 100  $\mu\text{L}$  of C $_6$ D $_6$ , and transferred into the NMR tube or taken from a standard solution prepared in a volumetric flask. If needed, additional C $_6$ D $_6$  was added to the NMR tube so that concentration remained constant between analogous experiments. When two imines were used, relative stoichiometry was checked by integration prior to heating.

**Reaction of 1 and (Bu $^t$ )HC=N(*p*-tolyl), 2a.** Compound **1** (0.0066 g, 0.017 mmol, 1 equiv), (Bu $^t$ )HC=N(*p*-tolyl) (0.171 mol, 10 equiv), and anisole internal standard were added to an NMR tube. The reaction was monitored at  $55^\circ\text{C}$  for 45 days. The progress of the reaction was monitored by  $^1\text{H}$  NMR spectroscopy. Resonances for **3a** and **4a** were observed.

**Reaction of 1 and (Ph)HC=N(Pr $^n$ ), 2b.** Compound **1** (0.0066 g, 0.017 mmol, 1 equiv), (Ph)HC=N(Pr $^n$ ) (0.171 mol, 10 equiv), and hexamethylbenzene internal standard were added to an NMR tube. The reaction was monitored at  $55^\circ\text{C}$  for 45 days. The progress of the reaction was monitored by  $^1\text{H}$  NMR spectroscopy. Resonances for **3b** and **4b** were observed initially. After 1 day resonances for compound **5** were observed.

**Reaction of 1, (Bu $^t$ )HC=N(*p*-tolyl), 2a, and (Ph)HC=N(Pr $^n$ ), 2b.** Compound **1** (0.0066 g, 0.017 mmol, 1 equiv), (Ph)HC=N(Pr $^n$ ) (0.171 mmol, 10 equiv), (Bu $^t$ )HC=N(*p*-tolyl) (0.171 mmol, 10 equiv), and hexamethylbenzene internal standard were added to an NMR tube. The reaction was monitored at  $25^\circ\text{C}$  for 25 days, then  $55^\circ\text{C}$  for 35 days. The progress of the reaction was monitored by  $^1\text{H}$  NMR spectroscopy. Resonances for the metathesis products (Ph)HC=N(*p*-tolyl) and (Bu $^t$ )CH=N(Bu $^t$ ) were observed initially. After  $>2$  weeks, precipitate was observed.

**Reaction of Imide 1 and Labeled Amine 6N.** A 20  $\mu\text{L}$  portion of a 0.626 M (12.5  $\mu\text{mol}$ ) standard solution of labeled amine **6N**, in C $_6$ D $_6$ , was added to 100  $\mu\text{L}$  of a 0.0912 M (9.12  $\mu\text{mol}$ ) standard solution of imide **1** in C $_6$ D $_6$ , in a screw top NMR tube. A fine precipitate of ammonium salt formed immediately. The progress of the reaction was monitored by  $^1\text{H}$  and  $^{15}\text{N}$  NMR spectroscopy at  $80^\circ\text{C}$  for 3 days. Resonances for the labeled imide/amide **7N1a** were observed initially. Over the course of the reaction, resonances for **1N** and **7N2** were observed.

**Reaction of Imide 1, Imine 4b, and Labeled Amine 6N.** A 100  $\mu\text{L}$  portion of a 0.626 M (62.5  $\mu\text{mol}$ ) standard solution of labeled amine **6N**, in C $_6$ D $_6$ , was added to a screw top NMR tube containing 400  $\mu\text{L}$  of a 0.0912 M (36.5  $\mu\text{mol}$ ) standard solution of imide **1** and 100  $\mu\text{L}$  of a 0.33 M (33.0  $\mu\text{mol}$ ) standard solution of imine **4b**. The progress of the reaction was monitored by  $^1\text{H}$  and  $^{15}\text{N}$  NMR spectroscopy at  $80^\circ\text{C}$  for 3 days. Resonances for the labeled imide/amide **7N1a** were observed initially. Over the course of the reaction, resonance for **1N**, **4bN**, and **7N2** were observed.

**Reaction of Imide 1, Labeled Imine 4bN, and Amine 6.** A 200  $\mu\text{L}$  portion of a 0.472 M (94.4  $\mu\text{mol}$ ) standard solution of amine in C $_6$ D $_6$  was added to a screw top NMR tube containing 97.4 mg (251  $\mu\text{mol}$ ) of imide **1** and 200  $\mu\text{L}$  of a 1.24 M (248  $\mu\text{mol}$ ) standard solution of imine **4bN**. The progress of the reaction was monitored by  $^1\text{H}$  and  $^{15}\text{N}$  NMR spectroscopy at  $68^\circ\text{C}$  for 2 days. Resonances for the labeled imide **1N** were observed initially. Over the course of the reaction, resonances for **7N1a** and **7N2** were observed.



**Reaction of Imide 1 with Labeled Ammonium Salt 8N.** A 74.4 mg (0.19 mmol) portion of imide **1** and 42.9 mg (0.39 mmol) of labeled ammonium salt **8N** were added to a screw top NMR tube. Approximately 0.5 mL of  $\text{C}_6\text{D}_6$  was added to give a yellow solution with a white precipitate. The sample was heated at 70 °C for 2 weeks.  $^{15}\text{N}$  NMR spectroscopy revealed a resonance for labeled imide **1N**.

**Reaction of Amine 6 with Labeled Ammonium Salt 8N.** A 55.8 mg (0.50 mmol) portion of labeled ammonium salt was added to 0.5 mL of a 0.5 M (0.25 mmol) solution of amine

**6.** The sample was heated at 70 °C for 2 weeks.  $^{15}\text{N}$  NMR spectroscopy revealed a resonance for labeled amine **6N**.

**Acknowledgment.** This research was supported by NSF award CHE-0091400. We thank Dr. Martin Minelli and Dr. Fu-Tyan Lin for help in acquiring  $^{14}\text{N}$  and  $^{15}\text{N}$  NMR spectral data.

OM011085V