

Synthesis of bis(indolyl)methanes Catalyzed by Triethylborane

J. Pablo García Merinos^{a,b,*}, Heraclio López Ruíz^a, Yliana López^b and Susana Rojas Lima^a

^aÁrea Académica de Química, Universidad Autónoma del Estado de Hidalgo, Carretera Pachuca-Tulancingo Km. 4.5, Mineral de La Reforma, Hidalgo, CP 42076, Mexico, ^bInstituto de Investigaciones Químico-Biológicas, Universidad Michoacana de San Nicolás de Hidalgo, Ed.B-1, C.U., Morelia, Michoacán, 58030, México

Received November 20, 2014; Revised January 13, 2015; Accepted February 19, 2015

Abstract: Triethylborane (TEB) was found to be a mild, efficient, and acid catalyst in electrophilic substitution reaction of indoles with aldehydes compounds to afford the corresponding bis(indolyl)methanes. Vibrindole A (**5**) and bis(indolyl)methanes derivatives **16** and **18** were synthesized using this methodology. Compound **16** is an intermediary in the synthesis of the natural bisindoles arsindoline B (**2**) and streptindole (**6**). The structure of vibrindole A (**5**) was unequivocally confirmed by a single crystal X-ray diffraction analysis.



J. P. G. Merinos

Keywords: Arsindoline B, bis(indolyl)methanes, indole, streptindole, triethylborane.

INTRODUCTION

The indole nucleus, referred to as a “privileged structure” by some authors, [1] is very important in pharmaceutical products, biological systems, and in the field of materials science. Moreover, the indole nucleus is present in many natural products isolated from marine and terrestrial organisms. These products are a rich source of antitumor agents.

There are several reports related to the isolation, characterization, and biological evaluation of indole. Among them,

arsindoline A (**1**) and B (**2**) were isolated from a marine-derived bacterium strain CB101 identified as *Aeromonas sp* (Fig. 1) [2,3]. 2,2-Bis(6-bromo-3-indolyl)ethylamine (**3**) was isolated from the Californian tunicate *Didemnum candidum* [4]. However, there are no reported biological studies on the promising antitumor potential of these indole alkaloids (Fig. 1). On the other hand, there are several natural bioactive products that share a common bis(indolyl)methane molecular unit. Arundine (**4**), isolated from the root of *Arundo donax*, which exhibits potent carcinogenicity [5]; vibrindole A (**5**), a





Fig. (1). BIMs from the natural source.

*Address correspondence to these authors at the Área Académica de Química, Universidad Autónoma del Estado de Hidalgo, Carretera Pachuca-Tulancingo Km. 4.5, Mineral de La Reforma, Hidalgo, CP 42076, Mexico and Instituto de Investigaciones Químico-Biológicas, Universidad Michoacana de San Nicolás de Hidalgo, Ed.B-1, C.U., Morelia, Michoacán, 58030, México; Tel: (52)443 3 22 35 00 Ext. 4207; Fax: (52) 443 3 26 57 90 Ext. 103; E-mail: jpgarciam@gmail.com



Scheme 1.

metabolite of the marine *Vibrio parahaemolyticus*, shows antibacterial activity against *Staphylococcus aureus*, *Staphylococcus albus* and *Bacillus subtilis* [6], and streptindole, (6) isolated from the intestinal bacterium *Streptococcus faecium* IB 37, demonstrate DNA-damaging activity and genotoxicity (Fig. 1) [7].

In this context, bis(indolyl)methanes (BIMs) have attracted considerable interest in recent years [8]. An increasing number of naturally occurring bioactive BIMs have shown cytotoxic activity against several cancer lines. Consequently, numerous protocols for the synthesis of BIMs derivatives have been reported [9,10].


Common methodologies for the synthesis of BIMs involve the condensation of indoles with several aldehydes or ketones in the presence of protic acids or Lewis acids [11–15]. These methodologies are well documented as useful routes to a large variety of BIMs. As a continuation of our

research on the synthesis of heterocyclic compounds, herein are described preliminary results of the Friedel-Crafts alkylations using TEB as a catalyst. Through this methodology, the synthesis of vibrindole A (5) and BIM derivatives 16 and 18 was carried out in short time reaction. Compound 16 is an intermediary in the synthesis of the natural bisindoles arsin-doline B (2) and streptindole (6).

RESULTS AND DISCUSSION





First, we carried out the reaction of indole with acetaldehyde and triethylborane in 1,2-dichloroethane at room temperature. Corresponding bis(indolyl)methane was formed (TLC) in a short reaction time (Scheme 1).

To prove the generality of the protocol, the reaction was then extended towards a variety of aldehydes and the results are summarized in Table 1.


Table 1. Et₃B catalyzed synthesis of bis(indolyl)methanes.^a

Entry	Aldehyde	Product	Yield (%) ^c	References
1	(9)	10	32	[16,17]
2	(11)	12	28	[18,19]
3	(13)	14	12	[20]

Table 1. contd...

Entry	Aldehyde	Product	Yield (%) ^c	References
4	 (15)	16	92	[19,7]
5	 (17)	18	52	[17]
6	 (19)	20	12	[13,21]
7	 (21)	22	10	[18,19]

^a All reactions were carried out with 1 equiv. of aldehyde and 2 equiv. of indole in 1,2-DCE at room temperature for 20 min. ^b Reaction was carried out with (0.05 mL) of Et₃B. ^c Yields refer to pure, isolated products.






Fig. (3). The crystal packing of the title compound, viewed along the *c* axis. The dashed lines indicate the X-H... π interactions.

the natural bisindoles arsinoline B (**2**) and streptindole (**6**). The structure of vibrindole A (**5**) was unequivocally established by single-crystal X-ray diffraction. The reaction is simple, non-toxic, easy to handle, and with shorter reaction times. The triethylborane catalyzed C3-selective Friedel-Crafts alkylation of indoles with aldehydes. Work along these lines is underway.

EXPERIMENTAL

IR spectra were acquired on a Perkin Elmer TF-IR System 2000 using KBr pellets ($\bar{\nu}$, cm^{-1}). NMR spectra (^1H , ^{13}C , HETCOR and COSY) were determined on JEOL Eclipse +400 spectrometer, and chemical shifts are stated in ppm (δ) and are referred to the residual ^1H signal ($\delta = 7.27$) or to the central ^{13}C triplet signal ($\delta = 77.0$) for CDCl_3 . The products were separated by chromatography over silica gel (70-230 mesh). Crystals of **5**, suitable for X-ray analysis, were obtained from hexane-ethyl acetate 7:3 by slow evaporation of the solvent at room temperature. The X-ray measurement was performed at 293 K on a Bruker SMART CCD diffractometer with Mo $\text{K}\alpha$ -radiation, $\gamma = 0.71073 \text{ \AA}$. The structure was solved by direct methods (SHELXS-86) and refined using SHELXL-97. All non-hydrogen atoms were refined anisotropically, and the hydrogen atoms were included at calculated positions with the use of a riding model. CCDC 1050231 contained crystallographic data for this publication. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif.

Crystal Data for Compound 5

Crystal data for compound **5**: $\text{C}_{18}\text{H}_{16}\text{N}_2$, colorless prisms, $0.2 \text{ mm} \times 0.18 \text{ mm} \times 0.19 \text{ mm}$, formula weight $M = 260.33$, triclinic, $P1$, $a = 6.562 \text{ \AA}$, $b = 9.977 \text{ \AA}$, $c = 21.357 \text{ \AA}$,

$\alpha = \beta = \gamma = 90^\circ$, $V = 1398.2 \text{ \AA}^3$, $Z = 4$, $D_x = 1.237 \text{ Mg m}^{-3}$, $\mu = 0.073 \text{ mm}^{-1}$, $F(000) = 552$. Collected reflections: 10165 within a theta range of 1.91 – 27.07° . Unique reflections: 1794. Observed reflections: 1794 with $F > 4\sigma(F)$ the absorption correction was not applied. Refinement: (refinement on F^2), final $R = 0.0494$, $wR^2 = 0.1302$, goodness-of-fit = 0.976 , 1794 reflections, 190 parameters, maximum and minimum difference electron densities were $0.183 \text{ e \AA}^{-3}$ and $-0.118 \text{ e \AA}^{-3}$, respectively.

Typical Procedure for the Preparation of bis(indolyl) methanes

A solution of indole (0.2 g, 3.4 mmol) and aldehyde (0.07 g, 1.7 mmol) in $\text{ClCH}_2\text{CH}_2\text{Cl}$ (0.5 mL) was treated with Et_3B (0.05 mL) and the resulting reaction mixture was stirred at room temperature for 20 min. The solvent was removed *in vacuo* and the resulting mixture was purified by flash column chromatography eluting with hexane-ethyl acetate 7:3 or 9:1. The spectral data of some of the bis(indolyl)methanes are summarized below.

3,3'-Bis(indolyl)-2-(hydroxyphenyl)methane (Entry 3, 14): IR (ν , cm^{-1} KBr): 3382, 3226, 2918, 2849, 1735, 1664, 1619, 1459, 1376, 1268, 1195, 1108, 749. NMR ^1H 400 MHz (CDCl_3), δ : 7.80 (1H, s, NH), 7.40 (1H, d, H-5, $J = 7.7 \text{ Hz}$), 7.31 (1H, d, H-8, $J = 8.1 \text{ Hz}$), 7.20 (3H, m, H-12, H-13 y H-14), 7.04 (1H, td, H-6, $J = 0.7, 7.3 \text{ Hz}$), 6.87 (2H, d, H-2, H-7, H-15, $J = 2.2 \text{ Hz}$), 6.63 (1H, s, H-2), 5.9 (1H, s, H-3'), 5.48 (1H, s broad, OH). NMR ^{13}C 100 MHz (CDCl_3), δ : 154.5, (C-10), 136.9 (C-3), 130.1 (C-12), 129.2 (C-10), 128.1 (C-13), 126.8 (C-9), 123.8 (C-2), 122.4 (C-14), 120.9 (C-15), 120.0 (C-5), 119.9 (C-4), 119.6 (C-6), 117.1 (C-11), 116.6 (C-7), 111.4 (C-8), 35.9 (C-3').

3,3'-Bis(indolyl)-4-(bromophenyl)methane (Entry 6, 20): IR (ν , cm^{-1} KBr): 3417, 2918, 2849, 1618, 1482, 1455, 1414, 1199, 1033, 742, 541. NMR ^1H 400 MHz (CDCl_3), δ : 7.78 (1H, s, NH), 7.39 (2H, m, H-5 y H-11), 7.33 (1H, d, $J = 8.8 \text{ Hz}$, H-8), 7.20 (1H, t, H-6, H-12, $J = 8.2 \text{ Hz}$), 7.04 (1H, d, H-7, $J = 8.2 \text{ Hz}$), 6.56 (1H, s, H-2), 5.85 (1H, s, H-3'). NMR ^{13}C 100 MHz (CDCl_3), δ : 143.22 (C-10), 136.79 (C-14), 131.41 (C-3), 130.59 (C-4), 126.98 (C-13), 123.70 (C-6), 123.01 (C-9), 122.19 (C-2), 120.02 (C-7), 119.90 (C-5), 119.48 (C-12), 119.16 (C-11), 111.25 (C-8), 39.81 (C-10).

CONFLICT OF INTEREST

The authors confirm that this article content has no conflict of interest.

ACKNOWLEDGEMENTS

Financial support from CONACYT (project No. 183980) and CIC-UMSNH is gratefully acknowledged. The authors thank Ms. Carolina López Ortiz for revision of the manuscript. One of the authors, J.P.G.M. is thankful to CONACYT for a scholarship (grant: 186053).

REFERENCES

- [1] Bartoli, G.; Dalpozzo, R.; Nardi, M. Applications of Bartoli indole synthesis. *Chem. Soc. Rev.* **2014**, *43*, 4728–4750.

- [2] Cai, S.-X.; Li, D.-H.; Zhu, T.-J.; Wang, F.-P.; Xiao, X.; Gu, Q.-Q. Two new indole alkaloids from the marine-derived bacterium *Aeromonas* sp. CB101. *Helv. Chim. Acta* **2010**, *93*, 791-795.
- [3] Blunt, J. W.; Copp, B. R.; Keyzers, R. A.; Munro, M. H. G.; Prinsep, M. R. Marine natural products. *Nat. Prod. Rep.* **2012**, *29*, 144-222.
- [4] Fahy, E.; Potts, B. C. M.; Faulkner, D. J.; Smith, K. 6-Bromotryptamine derivatives from the gulf of California Tunicate *Didemnum candidum*. *J. Nat. Prod.* **1991**, *54*, 564-569.
- [5] Khuzhaev, B. U.; Aripova, S. F.; Shakirov, R. Sh. Arundine-A new dimeric alkaloid from the roots of *Arundo donax*. *Chem. Nat. Compd.* **1994**, *30*, 635-636.
- [6] Bell, R.; Carmelli, S. Vibrindole A, a Metabolite of the marine bacterium, *Vibrio parahaemolyticus*, isolated from the toxic mucus of the boxfish *Ostracion cubicus*. *J. Nat. Prod.* **1994**, *57*, 1587-1590.
- [7] Osawa, T.; Namiki M. Structure elucidation of streptindole, a novel genotoxic metabolite isolated from intestinal bacteria. *Tetrahedron Lett.* **1983**, *24*, 4719-4722.
- [8] Shiri, M.; Zolfigol, M. A.; Kruger, H. G.; Tanbakouchian, Z. Bis- and trisindolylmethanes (BIMs and TIMs). *Chem. Rev.* **2010**, *110*, 2250-2293.
- [9] Gibbs, T. J. K.; Tomkinson, N. C. O. Aminocatalytic preparation of bisindolylalkanes, *Org. Biomol. Chem.* **2005**, *3*, 4043-4045.
- [10] Abe, T.; Nakamura, S.; Yanada, R.; Choshi, T.; Hibino, S.; Ishikura, M. One-pot construction of 3,3'-bisindolylmethanes through Bartoli indole synthesis. *Org. Lett.* **2013**, *15*, 3622-3625.
- [11] Nagarajan, R.; Perumal, P. T. InCl_3 and $\text{In}(\text{OTf})_3$ catalyzed reactions: synthesis of 3-acetyl indoles, bis-indolylmethane and indolylquinoline derivatives. *Tetrahedron* **2002**, *58*, 1229-1232.
- [12] Jella, R. R.; Nagarajan, R. Synthesis of indole alkaloids arsendoline A, arsendoline B and their analogues in low melting mixture. *Tetrahedron Lett.* **2013**, *69*, 10249-10253.
- [13] Handy, S. N.; Westbrook, M. A mild synthesis of bis(indolyl)methanes using a deep eutectic solvent. *Tetrahedron Lett.* **2014**, *55*, 4969-4971.
- [14] Mishra, S.; Ghosh, R. Ecofriendly and sustainable efficient synthesis of bis(indolyl)methanes based on recyclable Brønsted (CSA) or Lewis ($\text{ZrOCl}_2 \cdot 8\text{H}_2\text{O}$) acid catalysts. *Indian J. Chem.* **2011**, *50B*, 1630-1636.
- [15] Govindarajulu, B.; Nimmagadda, S.; Paramasivan, T.; Perumal, A. Convenient method of synthesis of bis-indolylmethanes: Indium trichloride catalyzed reactions of indole with aldehydes and Schiff bases. *Synth. Comm.* **2000**, *30*, 1609-1614.
- [16] Pore, D. M.; Desai, U. V.; Thopate, T. S.; Wadgaonkar, P. P. A mild, expedient, solventless synthesis of bis(indolyl)alkanes using silica sulfuric acid as a reusable catalyst. *ARKIVOC* **2006**, (xii), 75-80.
- [17] Bandgar, B. P.; Patil, A. V.; Kamble, V. T. Fluoroboric acid adsorbed on silica gel catalyzed synthesis of bisindolyl alkanes under mild and solvent-free conditions. *ARKIVOC* **2007**, (xiv), 252-259.
- [18] Denis, J.-N.; Mauger, H.; Vallée, Y. The reaction of nitrones with indoles. Synthesis of asymmetrical diindolylalkanes. *Tetrahedron Lett.* **1997**, *38*, 8515-8518.
- [19] Giuseppe, B.; Marcella, B.; Gioia, F.; Giuliani, A.; Marcantoni, E.; Sambri, L. Solvent-Free indoles addition to carbonyl compounds promoted by $\text{CeCl}_3 \cdot 7\text{H}_2\text{O}$ -NaI- SiO_2 : An efficient method for the synthesis of streptindole. *Synthesis* **2004**, 895-900.
- [20] Antule, A. A. R.; Kazi, S. H.; Shaikh, A.; Ahmed, M. Z. An efficient and eco-friendly synthesis of bis (indolyl) methanes in aqueous medium. *Elixir Org. Chem.* **2014**, *71*, 25135-25137.
- [21] Jadhav, V. B. 5-Sulphosalicylic acid mediated expeditious synthesis of bis (indolyl) methanes. *Elixir Appl. Chem.* **2014**, *70*, 24010-24014.