



Synthesis of Protected Derivatives of 3-Pyrrolylalanine

Jody E. Beecher^a and David A. Tirrell^{b*}

^aAffymetrix, Santa Clara, California 95051

^bDepartment of Polymer Science and Engineering, University of Massachusetts, Amherst, MA 01003 USA

Received 27 February 1998; revised 24 March 1998; accepted 1 April 1998

Abstract: Protected derivatives of 3-pyrrolylalanine (3-PA) have been synthesized starting with *N*-tri(isopropyl)silylpyrrole. Following formylation and Boc protection, treatment with (±)-*N*-(benzyloxycarbonyl)-α-phosphonoglycine trimethyl ester provided a fully protected dehydro-3-pyrrolylalanine. Reduction of the double bond and removal of the methyl ester groups yielded 3-PA protected at the α- and pyrrole nitrogens. 3-PA is only transiently stable following *N*-deprotection. © 1998 Elsevier Science Ltd. All rights reserved.

Genetic engineering combined with *in vivo* bacterial protein synthesis offers a novel approach to the design and preparation of macromolecular materials, in that precise control of chain length, sequence, and stereochemistry can be achieved.¹ Since materials properties are closely tied to polymer microstructure, this synthetic strategy offers unique advantages in the creation of functionally novel biopolymers.^{2–4} In addition, protein-based materials can be engineered to combine natural and artificial domains to create polymers with unique structural and biological properties.⁵

The monomers available for *in vivo* protein synthesis include the 20 natural amino acids, as well as some analogues,^{6,7} which can be used to introduce novel functional groups into engineered proteins.^{8–10} One of our specific goals is the synthesis of proteins containing electroactive residues, which are of interest in the development of materials for the control of cell growth,¹¹ drug delivery,¹² and biosensors.¹³ While no translational studies of pyrrolylalanine have been reported, the ability of 3-thienylalanine¹⁰ to be incorporated into *E. coli* proteins in place of phenylalanine suggests that 3-pyrrolylalanine (3-PA) might be similarly incorporated, and the structural similarity of 3-PA and histidine indicates that 3-PA might also act as a histidine surrogate. Regardless of its translational activity, 3-PA and its derivatives should be useful building blocks for solution- and solid-phase peptide synthesis. After incorporation into target peptide or protein, 3-PA residues should be susceptible to oxidative polymerization through the 2- and 5- positions of the heterocycle to create materials with electrochemical properties analogous to those of the polypyrroles.¹¹ In addition, artificial heterocyclic amino acids, such as pyrrolylalanine, are useful in the design of potential therapeutic agents.¹⁴


Efficient polymerization of pyrroles requires use of derivatives free of substitution in the 2- and 5- positions; 3-PA (1) must be used rather than the known 2-isomer.^{14,15}

The initial reaction in the synthesis of 3-PA derivatives (Scheme) is the introduction of an aldehyde moiety at the 3-position of the pyrrolyl ring. While pyrrole is activated to electrophilic attack in the 2- and 5-positions, substitution can be directed to the 3-position by the introduction of a large protecting group, such as tri(isopropyl)silyl (TIPS), on the pyrrole nitrogen.¹⁶ Reaction of *N*-tri(isopropyl)silylpyrrole with the Vilsmeier-Haack reagent, followed by aqueous workup under alkaline conditions afforded pyrrolyl-3-carboxaldehyde (2) in 72% yield.¹⁷ After protection with (*tert*-butoxycarbonyl) (Boc),¹⁸ 3 was treated with (±)-*N*-

(benzyloxycarbonyl)- α -phosphonoglycine trimethyl ester in a Horner Emmons reaction to provide the dehydro-3-pyrrolylalanine derivative (**4**) in 85% yield.¹⁹ Reduction of the double bond with sodium borohydride in the presence of catalytic nickel(II) chloride hexahydrate produced the fully protected 3-PA (**5**) in 69% yield.²⁰ The

methyl ester protecting group was removed in aqueous sodium hydroxide to give the free carboxylic acid derivative (**6**).²¹ Finally, 3-PA (**1**) was liberated from the remaining protecting groups by reaction with trimethylsilyl iodide and pyridine in 77% yield.^{22,23} 3-PA decomposes rapidly in aqueous acid, less rapidly in neutral water, and negligibly in aqueous base.

Acknowledgment. This work was supported by a grant from the Polymers and Genetics Programs of the National Science Foundation. NMR spectra were recorded in the University of Massachusetts NMR Laboratory, which is supported by the NSF Materials Research Laboratory at the University. Jody Beecher thanks the National Institutes of Health for financial support of her postdoctoral research.



References and Notes

1. Tirrell, D. A.; Fournier, M. J.; Mason, T. L. *Curr. Opin. Struct. Biol.* **1991**, *1*, 638.
2. Anderson, J. P.; Cappello, J.; Martin, D. C. *Biopolymers* **1994**, *34*, 1049.
3. Krejchi, M. T.; Atkins, E. D. T.; Waddon, A. J.; Fournier, M. J.; Mason, T. L.; Tirrell, D. A. *Science* **1994**, *265*, 1427.
4. Yu, S. M.; Conticello, V.; Kayser, C.; Fournier, M. J.; Mason, T. L.; Tirrell, D. A. *Nature* **1997**, *389*, 167.
5. McPherson, D. T.; Morrow, C.; Minehah, D. S.; Wu, J.; Hunter, E.; Urry, D. W. *Biotechnol. Prog.* **1992**, *8*, 347.
6. Richmond, M. H. *Bacteriol. Rev.* **1962**, *26*, 398.
7. Wilson, M. J.; Hatfield, D. L. *Biochim. Biophys. Acta* **1984**, *781*, 205.
8. Ibba, M.; Hennecke, H. *Bio/Technology* **1994**, *12*, 678.
9. Yoshikawa, E.; Fournier, M. J.; Mason, T. L.; Tirrell, D. A. *Macromolecules* **1994**, *27*, 5471.
10. Kothakota, K.; Mason, T. L.; Tirrell, D. A.; Fournier, M. J. *J. Am. Chem. Soc.* **1995**, *117*, 536.
11. Wong, J. Y.; Langer, R.; Ingber, D. E. *Proc. Natl. Acad. Sci. USA* **1994**, *91*, 3201.
12. Pyo, M.; Maeder, G.; Kennedy, R. T.; Reynolds, J. R. *J. Electroanal. Chem.* **1994**, *368*, 329.
13. Bartlett, P. N.; Cooper, J. M. *J. Electroanal. Chem.* **1993**, *362*, 1.
14. Bladon, C. M. *J. Chem. Soc. Perkin Trans 1* **1990**, 1151.
15. Masquelin, T.; Broger, E.; Mueller, K.; Schmid, R.; Obrecht, D. *Helv. Chim. Acta* **1994**, *77*, 1395.
16. Anderson, H. J.; Loader, C. E. *Synthesis* **1985**, 353.

17. Bray, B. L.; Mathies, P. H.; Naef, R.; Solas, D. R.; Tidwell, T. T.; Artis, D. R.; Muchowski, J. M. *J. Org. Chem.* **1990**, *55*, 6317.
18. A solution of di-*tert*-butyl dicarbonate (3.80 g, 17.4 mmol) in dry CH_2Cl_2 (25 mL) was added dropwise to a solution of 1*H*-pyrrole-3-carboxaldehyde (1.52 g, 16.0 mmol) in dry CH_2Cl_2 (30 mL) at 0 °C. After 15 min, a solution of diisopropylethyl amine (3.0 mL, 17.2 mmol) and catalytic dimethylaminopyridine in dry CH_2Cl_2 (20 mL) was added dropwise over 30 min. The reaction was stirred for 3 h at 0 °C and then stirred for 1 h at room temperature. Aqueous acidic workup and purification by flash chromatography (hexanes/ethyl acetate 9:1 v/v) yielded 2.87 g (93%) of **3** as a colorless solid. ^1H NMR (CDCl_3): δ 1.59 (s, 9H, $\text{C}(\text{CH}_3)_3$); 6.64 (dd, $J=3.4$, 1.6 1H, pyr. H); 7.26 (dd, $J=3.4$, 2.0, 1H, pyr. H); 7.84 (dd, $J=2.0$, 1.6, 1H, pyr. H); 9.83 (s, 1H, CHO). ^{13}C NMR (CDCl_3): δ 27.9 ($\text{C}(\text{CH}_3)_3$); 85.5 ($\text{C}(\text{CH}_3)_3$); 109.3, 122.1, 128.2, 128.8 (4 pyr C); 147.9 (pyr-NCOO); 185.6 (CHO). mp 35-36 °C. Anal. calcd for $\text{C}_{10}\text{H}_{13}\text{NO}_3$ (195.22): C, 61.53; H, 6.71; N, 7.18; found: C, 61.44; H, 6.66; N, 7.20.
19. Tetramethylguanidine (1.41 mL, 11.2 mmol) was added dropwise to a solution of (\pm -*N*-(benzyloxycarbonyl)- α -phosphonoglycine trimethyl ester (3.73 g, 11.3 mmol) in dry CH_2Cl_2 (125 mL) and stirred for 20 min at room temperature. The mixture was then cooled to -41 °C (dry ice/ CCl_4 bath) and a solution of **3** (2.00 g, 10.2 mmol) in dry CH_2Cl_2 (50 mL) was added dropwise. The reaction was stirred for 1 h at -41 °C, then warmed slowly to room temperature (3 h), stirred for 18 h at room temperature, and finally refluxed for an additional 12 h. Aqueous workup and purification by flash chromatography (hexanes/ethyl acetate 8:2 v/v) yielded 3.48 g (85%) of **4** as a light yellow solid. ^1H NMR (CDCl_3): δ 1.59 (s, 9H, $\text{C}(\text{CH}_3)_3$); 3.77 (s, 3H COOCH_3); 5.16 (s, 2H, CH_2Ph); 6.15-6.34 (br s, 1H, NH); 6.39-7.01 (m, 1H, pyr. H); 7.18 (dd, $J=3.2$, 2.2, 1H, pyr. H); 7.26-7.41 (m, 6H, 5 arom H, $\text{C}=\text{CHR}$); 7.45-7.49 (m, 1H, pyr. H). ^{13}C NMR (CDCl_3): δ 27.9 ($\text{C}(\text{CH}_3)_3$); 52.3 (OCH_3); 67.4 (CH_2Ph); 84.5 ($\text{C}(\text{CH}_3)_3$); 112.0, 121.2, 121.3 (3 pyr C); 121.9 ($\text{CH}=\text{C}$); 123.4 (pyr C); 127.9, 128.2, 128.5, 136.0 (4 arom. C); 148.0 ($\text{COO}t\text{-Bu}$), 154.2 (COOCH_2Ph), 165.7 (COOCH_3). mp 86-88 °C. Anal. calcd for $\text{C}_{21}\text{H}_{24}\text{N}_2\text{O}_6$ (400.43): C, 62.99; H, 6.04; N, 7.00; found: C, 63.03; H, 6.16; N, 7.08.
20. Sodium borohydride (1.17 g, 30.9 mmol) was added in portions to a solution of **4** (2.50 g, 6.24 mmol) and nickel(II) chloride hexahydrate (0.26 g, 1.1 mmol) in MeOH (150 mL) at 0 °C. The reaction was warmed to room temperature and stirred for 16 h. After removal of the solvent *in vacuo*, the residue was redissolved in CH_2Cl_2 , washed with water and dried over MgSO_4 . Purification by flash chromatography (hexanes/ethyl acetate 7:3 v/v) yielded 1.74 g (69%) of **5** as a colorless wax. ^1H NMR (CDCl_3): δ 1.57 (s, 9H, $\text{C}(\text{CH}_3)_3$); 2.96 (d, $J=5.0$, 2H, CH_2CH); 3.75 (s, 3H, COOCH_3); 4.53-4.63 (m, 1H, CH_2CH), 5.11 (s, 2H, PhCH_2); 5.28 (d, $J=8.1$, 1H, NH); 5.96 (dd, $J=3.1$, $J=1.7$, 1H, pyr. H); 7.00 (dd, $J=2.0$, 1.7, 1H, pyr. H); 7.14 (dd, $J=3.1$, $J=2.0$, 1H, pyr. H); 7.34 (m, 5H, arom H). ^{13}C NMR (CDCl_3): δ 27.9 ($\text{C}(\text{CH}_3)_3$); 29.6 (CH_2CH); 52.3 (OCH_3); 54.1 (CH_2CH); 66.9 (CH_2Ph); 83.6 ($\text{C}(\text{CH}_3)_3$); 112.7, 118.4, 120.5, 120.7 (4 pyr. C); 128.0, 128.1, 128.5, 136.2 (4 arom. C) 148.6 ($\text{COO}t\text{-Bu}$), 155.7 (COOCH_2Ph); 172.1 (COOCH_3). Anal. calcd for $\text{C}_{21}\text{H}_{26}\text{N}_2\text{O}_6$ (402.44): C, 62.67; H, 6.51; N, 6.96; found: C, 62.75; H, 6.45; N, 6.72.
21. Aqueous NaOH (10% w/w, 20 mL) was added dropwise to a solution of **5** (1.27 g, 3.16 mmol) in MeOH (40 mL) at 0 °C. The solution was stirred for 15 min and adjusted to pH 6 with dilute HCl. After removing the solvent *in vacuo*, the residue was redissolved in CH_2Cl_2 , washed with water and dilute acid, and dried over MgSO_4 . Recrystallization from ether/hexanes gave 1.04 g (85%) of **6** as a colorless solid. ^1H NMR (CDCl_3): δ 1.58 (s, 9H, $\text{C}(\text{CH}_3)_3$); 2.97-3.02 (m, 2H, CH_2CH); 4.61 (m, 1H, CH_2CH); 5.12 (s, 2H, PhCH_2); 5.23 (br d, $J=8.1$, 1H, NH); 6.03 (dd, $J=3.2$, $J=1.7$, 1H, pyr. H); 7.04-7.07 (m, 1H, pyr. H);

- 7.14 (dd, $J=3.2$, $J=2.0$, 1H, pyr. H); 7.33-7.36 (m, 5H, arom. H). ^{13}C NMR (CDCl_3): δ 27.9 ($\text{C}(\text{CH}_3)_3$); 29.3 (CH_2CH); 53.8 (CH_2CH); 67.2 (PhCH_2); 83.8 ($\text{C}(\text{CH}_3)_3$); 112.8, 118.6, 120.5, 120.7 (4 pyr. C); 128.1, 128.3, 128.5, 136.0 (4 arom. C); 148.6 ($\text{COO}t\text{-Bu}$), 156.0 (COOCH_2Ph); 175.2 (COOH). mp 136-137 °C. Anal. calcd for $\text{C}_{20}\text{H}_{24}\text{N}_2\text{O}_6$ (388.42): C, 61.85; H, 6.23; N, 7.21; found: C, 61.90; H, 6.25; N, 7.29.
22. Lott, R. S.; Chauhan, V. S.; Stammer, C. H. *J. Chem. Soc., Chem. Comm.* **1979**, 495.
23. Trimethylsilyl iodide (0.24 mL, 1.7 mmol) was added dropwise to a solution of **6** (0.205 g, 0.53 mmol) and pyridine (0.046 mL, 0.57 mmol) in CH_2Cl_2 (8 mL) at 0 °C. The reaction was warmed to room temperature and stirred for 18 h. The soluble fraction of the reaction mixture was separated from the pyridine salt by-product by filtration and the solvent and other reaction by-products were removed *in vacuo*. Finally, the residue was treated with excess MeOH (10 mL) to yield 63 mg (77%) of **1** as a white precipitate. Purification (>98% as determined by ^1H NMR) was achieved by passing a cold solution of **1** in H_2O or D_2O through a plug of decolorizing carbon. ^1H NMR (D_2O): δ 2.82 (dd, $J=15.2$, $J=7.2$, 1H, CH_2CH); 2.85 (dd, $J=15.2$, $J=4.9$, 1H, CH_2CH); 3.70 (dd, 1H, $J=7.2$, $J=4.9$, CH_2CH); 5.94 (dd, $J=3.1$, $J=1.8$, 1H, pyr. H); 6.96 (dd, $J=2.0$, $J=1.8$, 1H, pyr. H); 7.02 (dd, $J=3.1$, $J=2.0$, 1H, pyr. H).