

One-Pot Synthesis of Functionalized 3-Aryl-1-phenyl-1*H*-pyrazoles from Hydrazonoyl Chlorides and Acetylenic Esters in the Presence of Ph₃P

by Issa Yavari*, Gholamhossein Khalili, and Anvar Mirzaei

Chemistry Department, Tarbiat Modares University, P. O. Box 14115-175, Tehran, Iran
(phone: + 98-21-82883465; fax: + 98-21-82883455; e-mail: yavarisa@modares.ac.ir)

The zwitterionic 1:1 intermediates generated by addition of Ph₃P to acetylenic esters is trapped by 1-[(aryl)chloromethylene]-2-phenylhydrazines (= *N*-phenylarenecarbohydrazonoyl chlorides) to yield functionalized 3-aryl-1-phenyl-1*H*-pyrazoles in good yields.


Introduction. – Pyrazoles have occupied a unique position in the design and synthesis of novel biologically active agents that exert remarkable anticancer activities. In fact, pyrazoles have been studied for over a century as an important class of heterocyclic compounds and still continue to attract considerable attention due to the broad range of biological activities they possess [1–4]. Among these, *N*-arylpyrazoles are an interesting class of heterocycles with remarkable pharmacological activities such as antibacterial-antifungal [5], tumor-necrosis inhibitor [6], antimicrobial [7], hypoglycemic [8], hypolipidemic [9], and anti-inflammatory [10][11].

Results and Discussion. – As a part of our current studies on the development of new routes in heterocyclic synthesis [12–15], we report the results of our studies involving the reaction of the zwitterionic intermediates derived from triphenylphosphine (Ph₃P) and acetylenic esters **1** with 1-[(aryl)chloromethylene]-2-phenylhydrazines (= *N*-phenylarenecarbohydrazonoyl chlorides) **2** [16–18], which constitutes a synthesis of functionalized 3-aryl-1-phenyl-1*H*-pyrazoles (**3**) in good yields (*Scheme 1*).


The structures of compounds **3a–3j** were deduced from their IR, ¹H-NMR, and ¹³C-NMR spectra. For example, the ¹H-NMR spectrum of **3a** exhibited two *ss* (δ 3.87 and 3.90) for the MeO groups, along with *ms* for the Ph groups. The ¹H-decoupled ¹³C-NMR spectrum of **3a** showed 15 distinct resonances that confirm the proposed structure. The IR spectrum of **3a** displayed characteristic C=O bands (1726 and 1725 cm^{–1}). The NMR spectra of **3b–3j** were similar to those for **3a**, except for the aryl and ester moieties, which exhibited characteristic resonances in appropriate regions of the spectra. The ¹H-NMR spectra of **3e–3j** exhibited a *s* at δ *ca.* 8.50 for the pyrazole CH group, which confirms the proposed structures.

Although the mechanistic details of the reaction are not known, a plausible rationalization may be advanced to explain the product formation. Presumably, the zwitterionic intermediate [19–21] **4**, formed from Ph₃P and **1**, adds to the hydrazonoyl chloride **2** to produce **5** (*Scheme 2*). This intermediate undergoes a cyclization reaction to afford **6**, which is converted to **3** by elimination of Ph₃P.

Scheme 1



Scheme 2



The formation of a mixture of two regioisomers of 1,3,4-trisubstituted and 1,3,5-trisubstituted pyrazoles, in a ratio of 1:4, has been reported [22] from the reaction of 1-[(aryl)chloromethylene]-2-phenylhydrazines with alkyl prop-2-ynolates **1c** and **1d** in the presence of Et₃N. When we carried out this reaction in the presence of Ph₃P, only one regioisomer, namely the 1,3,4-trisubstituted products **3e–3j**, were obtained in 83–92% yields. Structures **3e–3j** were distinguished from the corresponding 1,3,5-trisubstituted pyrazoles by their NMR spectra. Thus, the methine H-atom of **3e–3j** showed a *s* at $\delta > 8.5$, whereas the methine signal of the 1,3,5-trisubstituted pyrazoles is expected at δ ca. 7 ppm. Similarly, the ¹³C-NMR signal of the pyrazole CH group in compounds **3e–3j** appeared at $\delta > 130$, which is a deshielding by more than 10 ppm compared to the expected resonance for the 1,3,5-trisubstituted isomer.

In conclusion, the zwitterionic 1:1 intermediates generated by addition of Ph₃P to acetylenic esters is trapped by 1-[(aryl)chloromethylene]-2-phenylhydrazines to yield functionalized 3-aryl-1-phenyl-1*H*-pyrazoles in good yields. The present method may be considered as a practical route for the synthesis of pyrazole-ring systems.

Experimental Part

General. Compounds **1** and Ph₃P were obtained from Merck and used without further purification. Compounds **2** were prepared according to [16]. M.p.: *Electrothermal-9100* apparatus; uncorrected. IR Spectra: *Shimadzu-IR-460* spectrometer; $\tilde{\nu}$ in cm⁻¹. ¹H- and ¹³C-NMR Spectra: *Bruker-DRX-500-Avance* instrument; in CDCl₃ at 500.1 and 125.7 MHz, resp.; δ in ppm, *J* in Hz. MS: *Finnigan-MAT-8430* mass spectrometer, at 70 eV; in *m/z* (rel. %). Elemental analyses (C, H, N): *Heraeus-CHN-O-Rapid* analyzer.

Compounds 3: General Procedure. To a stirred soln. of Ph_3P (0.52 g, 2 mmol) and 1-[(aryl)chloromethylene]-2-phenylhydrazine **2** (2 mmol) in MeCN (10 ml) was added the acetylenic ester **1** (2 mmol) at -5° . The mixture was allowed to reach r.t. After completion of the reaction (2–4 h; TLC (AcOEt/hexane 2:1) monitoring), the solvent was evaporated, and the residue was purified by column chromatography (silica gel (230–240 mesh; Merck), hexane/AcOEt 4:1): pure product.

Dimethyl 1,3-Diphenyl-1H-pyrazole-4,5-dicarboxylate (3a): Yield 0.31 g (93%). Colorless crystals. M.p. $153-155^\circ$. IR (KBr): 1726 (C=O), 1725 (C=O), 1521, 1487, 1442, 1265, 1227, 1140, 1098, 1015, 786, 752, 688. $^1\text{H-NMR}$: 3.87 (s, MeO); 3.90 (s, MeO); 7.45–7.55 (m, 6 CH); 7.57–7.59 (m, 2 CH); 7.78–7.80 (m, 2 CH). $^{13}\text{C-NMR}$: 52.1 (MeO); 53.1 (MeO); 114.4 (C); 124.6 (2 CH); 128.2 (2 CH); 128.8 (2 CH); 129.0 (CH); 129.1 (CH); 129.2 (2 CH); 131.4 (C); 136.9 (C); 139.1 (C); 152.0 (C); 160.7 (COO); 163.4 (COO). EI-MS: 336 (75, M^+), 305 (90), 77 (100), 51 (60). Anal. calc. for $\text{C}_{19}\text{H}_{16}\text{N}_2\text{O}_4$ (336.34): C 67.85, H 4.79, N 8.33; found: C 67.72, H 4.85, N 8.20.

Dimethyl 3-(4-Methylphenyl)-1-phenyl-1H-pyrazole-4,5-dicarboxylate (3b): Yield 0.31 g (90%). Colorless crystals. M.p. $65-67^\circ$. IR (KBr): 1729 (C=O), 1706 (C=O), 1585, 1527, 1490, 1432, 1292, 1261, 1230, 1179, 1110, 1003, 817, 750, 688. $^1\text{H-NMR}$: 2.42 (s, Me); 3.85 (s, MeO); 3.88 (s, MeO); 7.26 (d, $^3J = 8.0$, 2 CH); 7.43–7.49 (m, CH); 7.55 (d, $^3J = 7.8$, 2 CH); 7.67 (d, $^3J = 8.0$, 2 CH). $^{13}\text{C-NMR}$: 21.3 (Me); 52.0 (MeO); 53.0 (MeO); 114.1 (C); 124.6 (2 CH); 128.5 (C); 128.7 (2 CH); 128.9 (2 CH); 129.1 (CH); 129.3 (2 CH); 136.8 (C); 138.3 (C); 139.2 (C); 152.0 (C); 160.7 (COO); 163.5 (COO). EI-MS: 350 (75, M^+), 319 (84), 77 (100), 51 (65). Anal. calc. for $\text{C}_{20}\text{H}_{18}\text{N}_2\text{O}_4$ (350.37): C 68.56, H 5.18, N 8.00; found: C 68.78, H 5.25, N 8.10.

Diethyl 1,3-Diphenyl-1H-pyrazole-4,5-dicarboxylate (3c): Yield 0.32 g (90%). Pale yellow crystals. M.p. $52-53^\circ$. IR (KBr): 1716 (C=O), 1702 (C=O), 1523, 1489, 1440, 1261, 1227, 1143, 1099, 1011, 756, 691. $^1\text{H-NMR}$: 1.26 (t, $^3J = 6.3$, Me); 1.31 (t, $^3J = 6.3$, Me); 4.33–4.35 (m, 2 CH_2O); 7.44–7.48 (m, 4 CH); 7.55–7.57 (m, CH); 7.78–7.80 (m, CH). $^{13}\text{C-NMR}$: 13.7 (Me); 14.0 (Me); 61.1 (CH_2O); 62.4 (CH_2O); 114.4 (C); 124.8 (2 CH); 128.1 (2 CH); 128.7 (CH); 128.8 (2 CH); 128.9 (CH); 129.1 (2 CH); 131.6 (C); 137.1 (C); 139.2 (C); 151.9 (C); 160.2 (COO); 162.9 (COO). EI-MS: 364 (69, M^+), 319 (85), 77 (100), 51 (55). Anal. calc. for $\text{C}_{21}\text{H}_{20}\text{N}_2\text{O}_4$ (364.40): C 69.22, H 5.33, N 7.69; found: C 69.30, H 5.40, N 7.80.

Diethyl 3-(4-Methylphenyl)-1-phenyl-1H-pyrazole-4,5-dicarboxylate (3d): Yield 0.33 g (89%). Pale yellow crystals. M.p. $210-212^\circ$. IR (KBr): 1728 (C=O), 1706 (C=O), 1525, 1488, 1432, 1261, 1229, 1170, 1153, 1109, 1002, 816, 749. $^1\text{H-NMR}$: 1.24 (t, $^3J = 7.1$, Me); 1.30 (t, $^3J = 7.1$, Me); 2.40 (s, Me); 4.28–4.33 (m, 2 CH_2O); 7.24 (d, $^3J = 7.9$, 2 CH); 7.45–7.50 (m, CH); 7.53–7.55 (m, 2 CH); 7.66 (d, $^3J = 7.9$, 2 CH). $^{13}\text{C-NMR}$: 13.8 (Me); 14.0 (Me); 21.3 (Me); 61.0 (CH_2O); 62.3 (CH_2O); 114.3 (C); 124.8 (2 CH); 128.6 (C); 128.7 (2 CH); 128.8 (2 CH); 128.9 (CH); 129.0 (2 CH); 136.8 (C); 138.7 (C); 139.3 (C); 152.0 (C); 160.1 (COO); 163.0 (COO). EI-MS: 378 (75, M^+), 333 (70), 77 (100), 51 (59). Anal. calc. for $\text{C}_{22}\text{H}_{22}\text{N}_2\text{O}_4$ (378.42): C 69.83, H 5.86, N 7.40; found: C 69.68, H 5.80, N 7.31.

Methyl 1,3-Diphenyl-1H-pyrazole-4-carboxylate (3e): Yield 0.25 g (92%). Pale yellow crystals. M.p. $110-112^\circ$. IR (KBr): 1692 (C=O), 1589, 1525, 1433, 1360, 1274, 1221, 1133, 1053, 1007, 753, 680. $^1\text{H-NMR}$: 3.85 (s, MeO); 7.39 (t, $^3J = 6.4$, CH); 7.42–7.53 (m, 5 CH); 7.81 (dd, $^4J = 1.0$, $^3J = 8.5$, 2 CH); 7.91 (dd, $^4J = 1.4$, $^3J = 8.0$, 2 CH); 8.53 (s, CH). $^{13}\text{C-NMR}$: 50.8 (MeO); 112.8 (C); 119 (2 CH); 126.9 (CH); 127.4 (2 CH); 128.2 (CH); 128.8 (2 CH); 129.0 (2 CH); 131.6 (C); 131.7 (CH); 138.7 (C); 153.5 (C); 162.8 (COO). EI-MS: 278 (70, M^+), 247 (63), 77 (100), 51 (66). Anal. calc. for $\text{C}_{17}\text{H}_{14}\text{N}_2\text{O}_2$ (278.31): C 73.37, H 5.07, N 10.07; found: C 73.50, H 5.10, N 10.15.

Methyl 3-(4-Methylphenyl)-1-phenyl-1H-pyrazole-4-carboxylate (3f): Yield 0.25 g (86%). Pale yellow crystals. M.p. $103-105^\circ$. IR (KBr): 1723 (C=O), 1587, 1517, 1492, 1443, 1275, 1201, 1108, 1024, 823, 750, 687. $^1\text{H-NMR}$: 2.44 (s, Me); 3.86 (s, MeO); 7.27 (t, $^3J = 7.8$, 2 CH); 7.37 (t, $^3J = 7.5$, CH); 7.48–7.51 (m, 2 CH); 7.78–7.80 (m, 4 CH); 8.51 (s, CH). $^{13}\text{C-NMR}$: 21.4 (Me); 51.3 (MeO); 113.3 (C); 119.5 (2 CH); 127.4 (CH); 126.6 (2 CH); 128.0 (C); 129.2 (2 CH); 129.5 (2 CH); 132.2 (CH); 138.6 (C); 139.3 (C); 154.1 (C); 163.4 (COO). EI-MS: 292 (72, M^+), 261 (68), 77 (100), 51 (69). Anal. calc. for $\text{C}_{18}\text{H}_{16}\text{N}_2\text{O}_2$ (292.33): C 73.96, H 5.52, N 9.58; found: C 73.90, H 5.60, N 9.64.

Ethyl 1,3-Diphenyl-1H-pyrazole-4-carboxylate (3g): Yield 0.25 g (87%). Pale yellow crystals. M.p. $63-64^\circ$. IR (KBr): 1679 (C=O), 1525, 1272, 1224, 1143, 1029, 744, 681. $^1\text{H-NMR}$: 1.34 (t, $^3J = 7.2$, Me); 4.33 (q, $^3J = 7.2$, CH_2O); 7.38 (t, $^3J = 7.4$, CH); 7.42–7.53 (m, 5 CH); 7.81 (d, $^3J = 8.3$, 2 CH); 7.89 (d, $^3J = 8.3$, 2 CH); 8.53 (s, CH). $^{13}\text{C-NMR}$: 14.2 (Me); 60.3 (CH_2O); 113.8 (C); 119.5 (2 CH); 127.4 (CH); 127.9

(2 CH); 128.6 (CH); 129.4 (2 CH); 129.6 (2 CH); 132.3 (CH); 138.5 (C); 139.3 (C); 154.0 (C); 162.9 (COO). EI-MS: 292 (74, M^+), 247 (65), 77 (100), 51 (66). Anal. calc. for $C_{18}H_{16}N_2O_2$ (292.33): C 73.96, H 5.52, N 9.58; found: C 73.90, H 5.58, N 9.71.

Ethyl 3-(4-Methylphenyl)-1-phenyl-1H-pyrazole-4-carboxylate (3h): Yield 0.26 g (85%). Pale yellow crystals. M.p. 98–100°. IR (KBr): 1682 (C=O), 1522, 1272, 1220, 1133, 1052, 765, 677. 1H -NMR: 1.34 (t , $^3J = 7.1$, Me); 2.41 (s , Me); 4.30 (q , $^3J = 7.1$, CH_2O); 7.26 (t , $^3J = 7.9$, 2 CH); 7.35 (t , $^3J = 7.4$, CH); 7.47–7.50 (m , 2 CH); 7.77–7.80 (m , 4 CH); 8.50 (s , CH). ^{13}C -NMR: 14.3 (Me); 21.3 (Me); 60.3 (CH_2O); 113.7 (C); 119.5 (2 CH); 127.3 (CH); 128.4 (C); 128.5 (2 CH); 129.3 (2 CH); 129.5 (2 CH); 132.3 (CH); 138.5 (C); 139.3 (C); 154.0 (C); 163.0 (COO). EI-MS: 306 (77, M^+), 261 (62), 77 (100), 51 (60). Anal. calc. for $C_{19}H_{18}N_2O_2$ (306.36): C 74.49, H 5.92, N 9.14; found: C 74.60, H 5.85, N 9.25.

Methyl 3-(4-Chlorophenyl)-1-phenyl-1H-pyrazole-4-carboxylate (3i): Yield 0.25 g (83%). Pale yellow crystals. M.p. 149°. IR (KBr): 1692 (C=O), 1597, 1527, 1501, 1436, 1277, 1222, 1138, 1090, 1061, 1091, 959, 838, 771, 702. 1H -NMR: 3.84 (s , MeO); 7.37 (t , $^3J = 7.4$, CH); 7.42 (d , $^3J = 8.5$, 2 CH); 7.50 (t , $^3J = 7.8$, 2 CH); 7.77 (d , $^3J = 7.8$, 2 CH); 7.87 (d , $^3J = 8.5$, 2 CH); 8.50 (s , CH). ^{13}C -NMR: 51.4 (MeO); 113.3 (C); 119.5 (2 CH); 127.6 (CH); 128.1 (C); 129.5 (2 CH); 130.5 (2 CH); 130.6 (2 CH); 132.4 (CH); 134.7 (C); 139.1 (C); 152.8 (C); 163.1 (COO). EI-MS: 312 (70, M^+), 281 (72), 77 (100), 51 (56). Anal. calc. for $C_{17}H_{13}ClN_2O_2$ (312.74): C 65.28, H 4.20, N 8.95; found: C 65.13, H 4.53, N 8.71.

Ethyl 3-(4-Chlorophenyl)-1-phenyl-1H-pyrazole-4-carboxylate (3j): Yield 0.28 g (86%). Pale yellow crystals. M.p. 146–148°. IR (KBr): 1685 (C=O), 1596, 1531, 1273, 1225, 1130, 1059, 1030, 953, 839, 770. 1H -NMR: 1.34 (t , $^3J = 7.1$, Me); 4.31 (q , $^3J = 7.1$, CH_2O); 7.37 (t , $^3J = 7.3$, CH); 7.42 (d , $^3J = 8.4$, 2 CH); 7.50 (t , $^3J = 7.9$, 2 CH); 7.77 (d , $^3J = 8.1$, 2 CH); 7.87 (d , $^3J = 8.4$, 2 CH); 8.50 (s , CH). ^{13}C -NMR: 14.2 (Me); 60.4 (CH_2O); 113.7 (C); 119.5 (2 CH); 127.5 (CH); 128.0 (C); 129.5 (2 CH); 130.6 (2 CH); 130.7 (2 CH); 132.3 (CH); 134.7 (C); 139.2 (C); 152.8 (C); 162.7 (COO). EI-MS: 326 (71, M^+), 281 (71), 77 (100), 51 (58). Anal. calc. for $C_{18}H_{13}ClN_2O_2$ (326.77): C 66.15, H 4.62, N 8.57; found: C 66.73, H 4.19, N 8.43.

REFERENCES

- [1] Y. K. Lee, M. J. Kim, N. J. Kim, *Tetrahedron Lett.* **2003**, 44, 6737.
- [2] S. Komeda, M. Lutz, L. A. Spek, *Inorg. Chem.* **2000**, 39, 4230.
- [3] M. Francisco, H. Seltzman, F. A. Gilliam, *J. Med. Chem.* **2002**, 45, 2708.
- [4] T. Nakamura, M. Sato, H. Kakinuma, N. Miyata, K. Taniguchi, K. Bando, A. Koda, K. Kameo, *J. Med. Chem.* **2003**, 46, 5416.
- [5] M. S. Abdel-Gawad, A. Abdel-Aziem, M. M. Ghorab, *Phosphorus, Sulfur Silicon Relat. Elem.* **2003**, 178, 1795.
- [6] I. B. Eriksson, J. D. Quinlan, *Drugs* **2006**, 66, 1411.
- [7] A. M. Farghaly, G. S. F. Soliman, A. M. M. El Semaary, F. A. Sh. Rostom, *Pharmazie* **2001**, 56, 28.
- [8] K. Seki, J. Isegawa, M. Fukuda, M. Oki, *Chem. Pharm. Bull.* **1984**, 32, 1568.
- [9] B. Cottineau, P. Toto, C. Marot, A. Pipaud, J. Chenault, *Bioorg. Med. Chem. Lett.* **2002**, 12, 2105.
- [10] C. M. Cardia, L. Corda, A. M. Fadda, A. M. Maccioni, E. Maccioni, A. Plumitallo, *Il Farmaco* **1998**, 53, 698.
- [11] D. M. Mullican, W. M. Wilson, T. D. Cannor, R. C. Kostlan, D. R. Dyer, *J. Med. Chem.* **1993**, 36, 1090.
- [12] I. Yavari, L. Moradi, *Helv. Chim. Acta* **2006**, 89, 1942.
- [13] I. Yavari, L. Moradi, A. Mokhtarporyani-Sanandaj, A. Mirzaei, *Helv. Chim. Acta* **2007**, 90, 392.
- [14] I. Yavari, A. Mirzaei, L. Moradi, N. Hosseini, *Tetrahedron Lett.* **2008**, 49, 2355.
- [15] I. Yavari, A. Moradi, L. Mokhtarporyani-Sanandaj, A. Mirzaei, *Tetrahedron* **2008**, 64, 5221.
- [16] P. Wolkoff, *Can. J. Chem.* **1975**, 53, 1333.
- [17] R. Huisgen, M. Seidel, G. Wallbillich, H. Knupfer, *Tetrahedron* **1962**, 17, 3.
- [18] R. Huisgen, R. Grashey, H. Knupfer, R. Kunz, *Chem. Ber.* **1964**, 97, 1085.
- [19] R. Huisgen, M. Morikawa, K. Herbig, E. Brunn, *Chem. Ber.* **1967**, 100, 1094.
- [20] E. Winterfeldt, D. Schumann, H. Dillinger, *Chem. Ber.* **1969**, 102, 1656.
- [21] J. H. Dillinger, G. Fengler, D. Schumann, E. Winterfeldt, *Tetrahedron* **1974**, 30, 2553.
- [22] S. J. Clovis, W. Fliege, R. Huisgen, *Chem. Ber.* **1983**, 116, 3062.

Received June 29, 2009