

Article

Model Studies to Access the [6,7,5,5]-Core of Ineleganolide Using Tandem Translactonization-Cope or Cyclopropanation-Cope Rearrangements as Key Steps

Jennifer L. Roizen, Amanda C Jones, Russell C Smith, Scott C Virgil, and Brian M. Stoltz

J. Org. Chem., **Just Accepted Manuscript** • DOI: 10.1021/acs.joc.7b02030 • Publication Date (Web): 07 Nov 2017

Downloaded from <http://pubs.acs.org> on November 8, 2017

Just Accepted

"Just Accepted" manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides "Just Accepted" as a free service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. "Just Accepted" manuscripts appear in full in PDF format accompanied by an HTML abstract. "Just Accepted" manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are accessible to all readers and citable by the Digital Object Identifier (DOI®). "Just Accepted" is an optional service offered to authors. Therefore, the "Just Accepted" Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the "Just Accepted" Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these "Just Accepted" manuscripts.




ACS Publications

Model Studies to Access the [6,7,5,5]-Core of Ineleganolide Using Tandem Translactonization-Cope or Cyclopropanation-Cope Rearrangements as Key Steps

Jennifer L. Roizen, Amanda C. Jones, Russell C. Smith, Scott C. Virgil, and Brian M. Stoltz*

Warren and Katharine Schlinger Laboratory for Chemistry and Chemical Engineering, Division of Chemistry and Chemical Engineering, California Institute of Technology, 1200 E. California Boulevard, MC 101-20, Pasadena, CA 91125 (USA).

E-mail: stoltz@caltech.edu



ABSTRACT: Recently, we reported a convergent cyclopropanation-Cope approach to the core of ineleganolide, which was the first disclosed synthesis of the core of the norditerpene natural product ineleganolide. In this complementary work, a model system for the core of ineleganolide has been prepared through a series of tandem cyclopropanation-Cope and translactonization-Cope rearrangements. Work with this model system has enriched our understanding of the cyclopropanation-Cope rearrangement sequence. Additionally, research into this model system has driven the development of tandem translactonization-Cope rearrangements.

Introduction

In 1999, Duh and co-workers isolated a norditerpene from the soft coral *Sinularia*¹ *inelegans* and named it ineleganolide (**1**, Figure 1).^{2,3} Ineleganolide demonstrates in vitro cytotoxicity against murine lymphocytic leukemia P388 cell lines ($\text{ED}_{50} = 3.82 \mu\text{g/mL}$),² yet does not demonstrate activity against human oral epidermoid (KB) and liver (Hepa59T/VGH) carcinoma cells ($\text{ED}_{50} > 20 \mu\text{g/mL}$).³ The structure of ineleganolide (**1**) was elucidated by single crystal X-ray diffraction,² revealing the relative configuration of nine stereocenters, six of which lined a central cycloheptanone core. The absolute configuration was established by Pattenden's biomimetic semi-synthesis from 5-epi-sinuleptolide (**3**), which furnished the only disclosed laboratory preparation of ineleganolide.⁴ Owing to the novel structures and bioactivity of ineleganolide (**1**), Nicolaou, Moeller,

Vanderwal, Frontier, Romo and co-workers have disclosed inventive and formidable approaches to its synthesis,⁵ none of which have provided access to its central [6,7,5,5]-core. We have accessed this core through a cyclopropanation-Cope cascade.⁶ Herein, we disclose model studies for that critical cyclopropanation-Cope sequence, as well as three complementary tandem transformations: transactonization-Cope, cyclopropanation-Cope-epoxidation and cyclopropanation-Cope-enolization cascades. Each transformation furnishes the [6,7,5,5]-core of ineleganolide.





Figure 1. A subset of metabolites (1–4) isolated from plants of the genus *Sinularia*.

Results and Discussion


From the outset, we sought to overcome the challenges of the enantioselective installation of the C(8) tertiary ether and construction of the seven-membered ring core of ineleganolide (Scheme 1). To evaluate strategies, we targeted the simpler model, des-isopropenyl-*ent*-ineleganolide (**5**). Initial retrosynthetic simplification of the C(8)–C(5) ether and other oxidation state manipulations revealed cycloheptadienone **6**. Access to model compound **6** would proceed through a strain-release Cope rearrangement of diene **7**.⁷ The cyclopropane in diene **7** would arise from vinylcyclopropane **8**. In turn, vinylcyclopropane **8** would be directly available from vinyl diazoester **9**, which itself would be the coupling product of carboxylic acid **10** and alcohol **11**. Embedded within alcohol **11** was a tertiary stereocenter at C(8), which we targeted through an enantioselective allylation strategy.⁸

Scheme 1. Initial retrosynthetic analysis of des-isopropenyl-*ent*-ineleganolide (**5**)




In the forward direction, the first critical challenge was installation of the C(8) stereocenter. To this end, we found inspiration in work within our laboratory to establish quaternary stereogenic centers through the enantioselective decarboxylative allylation of cyclic enolates.⁹ If this method could be employed with a substrate bearing α -oxygenation,^{8b} we would be able to access the C(8) center embedded within alcohol **11** (Scheme 2).^{8a} At the time, there was limited evidence that α -heteroatoms might be tolerated in the reaction.¹⁰ Coincident with early model system investigations, we developed a general enantioselective alkylation of dioxanone-derived enol ethers,^{8a} which could be used to prepare alcohol **11**^{8b} by way of chloroallyl dioxanone **15**. Treatment of triethylsilyl ether **12** with Pd(dmdba)₂ (5 mol%), tris(CF₃)-(S)-*t*-BuPHOX (**14**, 5.5 mol%),¹¹ and chloroallyl mesylate (**13**, 1.05 equiv) with an equivalent of TBAT in PhMe at 25 °C, yielded the desired tetrasubstituted ether **15** with good enantioselectivity (91% ee). Chloroolefin **15** was converted in a fickle two-step oxidative bromination/Wittig olefination sequence to volatile cyclopentenone **16**.^{8a} In turn, facile diastereoselective reduction furnished cyclopentenol **11**, which contains both C(8) and C(10) stereogenic centers.

Scheme 2. Synthesis of cyclopentenol 11




Having prepared enantioenriched cyclopentenol **11**, we sought access to its coupling partner, carboxylic acid **10** (Scheme 3). Conversion of cyclohexenone (**17**) proceeded by an efficient CeCl₃-mediated 1,2-addition of the lithium enolate of EtOAc.¹² Oxidative rearrangement with allylic transposition¹³ and saponification generated carboxylic acid **10** in 67% overall yield. Notably, TEMPO⁺BF₄⁻ could be employed as an environmentally benign alternative to affect an oxidative rearrangement, albeit in slightly lower 54% overall yield.¹⁴ To prepare for the key cyclopropanation, we coupled alcohol **11** and carboxylic acid **10** with DCC. Subsequently, diazo transfer was accomplished upon treatment with *p*-ABSA (**20**) to provide the targeted cyclopropanation precursor (**9**) in excellent yield.

Scheme 3. Toward the cyclopropanation substrate




Under standard cyclopropanation conditions, diazoester **9** did not give way to desired cyclopropane **8**. Rather, a thermal rearrangement occurred to furnish pyrazole **21** in 59% yield (Scheme 4). This type of reaction had been previously reported,¹⁵ and Padwa had proposed that pyrazole formation proceeds through a 1,5-cyclization followed by a proton shift.^{15f} Indeed, pyrazole **21** was generated cleanly when diazoester **9** was heated in benzene at reflux, in the absence of copper. The structure of pyrazole **21** was characterized by X-ray crystallographic diffraction, and this structure confirmed the *syn*-arrangement of C(8) and C(10) oxygenation.

Scheme 4. Pyrazole formation and oxidation plague initial cyclopropanation approach




At lower temperatures, neither cyclopropane **8** nor pyrazole **21** formed. One potential explanation for this lack of reactivity was that diazoester **9** did not generate the requisite metal carbenoid. To interrogate this hypothesis, we exposed diazoester **9** to $\text{Rh}_2(\text{oct})_4$ and ambient air at 0 °C, anticipating that oxygen or water could intercept a formed metal carbenoid, ultimately furnishing oxidation products. Indeed, under these conditions, oxidized **22** formed along with its acetonide-cleaved analogue (**23**) suggesting that a carbenoid was generated upon exposure of diazoester **9** to $\text{Rh}_2(\text{oct})_4$, but that this carbenoid failed to undergo cyclopropanation. Close examination of diazoester **9** and cyclopropane **8** revealed a 1,3-diaxial interaction between methyl groups at C(8) and on the acetonide that would likely disfavor cyclopropanation, as this 1,3-diaxial interaction was necessarily enhanced in cyclopropane **8**, relative to diazoester **9**.

To avoid this strain, we converted ester **19** to an alternative cyclopropanation precursor (Scheme 5). To this end, we turned to De Waard's catalytic conditions for ketal cleavage,¹⁶ which enabled cleavage of acetonide **19** without olefin isomerization.¹⁶ Silylation provided ready access to an ether, which reacted with TsN_3 to generate diazoester **24**. We were pleased to find that copper *tert*-butyl salicylaldimine ($\text{Cu}(\text{tbs})_2$) affected cyclopropanation in 74% yield.

Scheme 5. Successful cyclopropanation of silyl ether **24**


With the cyclopropane successfully installed, we sought to advance silyl ether **25a** to the requisite divinylcyclopropane Cope precursor (Scheme 6). Desilylation of cyclopropane **25a** occurred readily; however, an array of products formed under a variety of oxidation conditions. At least one problematic side reaction was documented. Upon oxidation with Dess-Martin periodinane, ether **26** was observed as the major product (Scheme 6). Presumably, this product arises through conjugate addition of the primary alcohol into the enone functionality.

Scheme 6. Byproduct formed in the desilylation–oxidation sequence




Having struggled with oxidation to provide the targeted Cope precursor (e.g., silyl ether **25a** → divinyl cyclopropane **7**, Scheme 1), we chose to mask the problematic C(8) tertiary alcohol and the C(3) ketone in cyclopropane **25a**. Initially, we targeted formation of C(8) benzyl ether and acetyl ester (e.g., **25b** and **25c**, R=Bn and Ac respectively; Scheme 7). By standard analytical measurements (e.g., ¹H- and ¹³C-NMR and IR spectroscopy, and HRMS), the formed products matched expectations for the C(8)-ester and C(8)-ether (e.g., **25b** and **25c**, R=Bn or Ac); however, careful analysis of gradient-heteronuclear multiple bond correlation (gHMBC) spectroscopy revealed that all efforts to alkylate the C(8) alcohol result in translactonization¹⁷ to furnish instead a C(8) lactone (e.g., **25a** → **27**). Specifically, benzylation of the C(10) alcohol was achieved on treatment of C(8) alcohol **25a** with BnBr, KI, and Ag₂O at room temperature forming benzylated ether **27a**, while base-mediated acetylation (Ac₂O, pyridine, DMAP) yielded the translactonized product **27b** containing a C(8) lactone and a C(10) acetyl ester.¹⁸

Scheme 7. Translactonization to form benzyl ether and acetyl ester




While these products were not originally targeted, we anticipated that acetyl-masked **27b** could still be converted to the core of ineleganolide (i.e., **6**) by way of a divinylcyclopropane (Scheme 8). Having previously recognized the problematic reactivity of the C(3) ketone (see Scheme 6), divinylcyclopropane **29** was targeted as a suitable precursor for this tandem transformation. We found that Luche reduction of ketone **27b** furnished a 1:1.5 ratio of diastereomeric alcohols **28a** and **b**. These diastereomers were separated and carried through subsequent reactions in parallel. Secondary alcohol **28b** was masked as an acetyl ester, then the silyl ether was desilylated. The resultant alcohol was oxidized to an aldehyde, and subjected to methylenation to provide divinylcyclopropane **29b** in 76% yield over four steps.

Scheme 8. Access to divinylcyclopropane **29b**

We suspected that bisacetyl-masked divinyl cyclopropane **29b** could still be converted to the central [6,7,5,5]-fused system within ineleganolide through an acetate-cleavage/translactonization/Cope rearrangement (Scheme 9). We expected that both translactonization and Cope rearrangement would be thermodynamically governed transformations (Scheme 9). Consequently, when acetyl ester cleavage revealed alcohol **29c**, it would enable the reaction to funnel to the lowest energy product, namely less-strained cycloheptadiene **32b** via initial translactonization to lactone **31b** and subsequent Cope rearrangement. A hypothetical and disfavored direct Cope rearrangement of the C(8) lactone would form highly-strained **30b**, possessing two anti-Bredt, bridgehead alkenes.¹⁹ Were hypothetical intermediate **30b** to form, it would presumably undergo translactonization to cycloheptadiene **32b**. In the event, lithium hydroxide-mediated acetate cleavage of bis-acetyl ester **29b** promoted a translactonization–Cope rear-


rangement cascade to deliver cycloheptadiene **32b** in 55% yield, producing the carbocyclic core framework of ineganolide for the first time. To our knowledge, this is also the first example of a translactonization-triggered Cope rearrangement.

Scheme 9. Translactonization-Cope approach to access cycloheptadiene **32b**



An alternative translactonization–Cope approach was pursued concurrently, and also furnished the central [6,7,5,5]-fused system within ineganolide (Scheme 10). Efforts to mask the C(3) ketone through ketalization proved fruitless under many standard conditions, as ketones **25a** and **27b** either decomposed, or formed complex product mixtures. Moreover, these ketones were unreactive toward mild conditions designed to prevent alkene isomerization (TMSOTf, (TMSOCH₂)₂, CH₂Cl₂, –78 °C). At a higher temperature (0 °C), ketalization proceeded, albeit with undesired alkene isomerization. Subsequent desilylation and oxidation of the primary alcohol to an aldehyde followed by Wittig olefination delivers divinylcyclopropane **33**, an appropriate Cope precursor.


Scheme 10. Ketalization with undesired olefin isomerization and subsequent translactonization-Cope rearrangement sequence





Although this olefin isomer was not originally designed into our synthetic approach to ineleganolide, we expected that ketal **33** would be an appropriate precursor for a tandem transactonization-Cope rearrangement. Acetyl ester **33** reacted with nucleophilic super-hydride to form desired cycloheptadiene **35**, presumably by a transactonization-Cope process (i.e., **33** \rightarrow **34** \rightarrow **35**). Importantly, this transactonization-Cope rearrangement again provided access to the [6,7,5,5]-fused core of ineleganolide, albeit with modified substitution on the cyclohexyl ring.

Having established a route to the carbon scaffold of ineleganolide (i.e., cycloheptadiene **35**), we attended to C(6) oxygenation and reduction of the C(12)–C(13) tetrasubstituted alkene (Scheme 11). To our delight, the tertiary C(8) alcohol could direct vanadium-mediated diastereoselective epoxidation of the C(6)–C(7) alkene to generate epoxide **36**. For subsequent functionalization, we anticipated that olefin reduction could prove useful. Expecting direct reduction of the C(12)–C(13) alkene to be challenging, we chose to pursue deketalization of cycloheptene **36**, which could potentially induce alkene isomerization to furnish enone **37**. We envisioned that the carbonyl in enone **37** could facilitate olefin reduction. Regrettably, selective deketalization of ketal **36** was not accomplished under a broad range of conditions.

Scheme 11. Olefin differentiation from cycloheptadiene **35**




As a result of the aforementioned challenges, we continued concurrent pursuit of a tandem cyclopropanation-Cope approach to form the [6,7,5,5]-fused core of ineleganolide. The tandem cyclopropanation-Cope^{20,21} rearrangement was developed by Davies and co-workers for application in diastereoselective intramolecular reactions.^{22,23,24,25} To investigate this reaction, we targeted diazoester **39** as the precursor to this key tandem cyclopropanation-Cope rearrangement (Scheme 12). Unfortunately, we were unable to directly convert acetone **19** to diazoester **39** due to an inability to affect the requisite olefination. We speculated that deprotonation of the acidic γ -proton of the α,β -unsaturated ketone / δ -ester system interfered with the attempted Wittig methylenation. To access diazoester **39**, we carried out the methylenation on the cyclopentene fragment prior to coupling it with the cyclohexenone carboxylic acid (Scheme 13). Thus, cyclopentenone **16** was converted to allylic benzoyl ester **40**. Reduction and benzoylation furnished benzoyl ester **40** in 98% yield over two-steps. Benzoyl ester **40** was treated with acid to cleave the ketal, oxidized with Dess-Martin periodinane and olefinated to deliver diene **41**. Alkoxide-mediated debenzoylation furnished alcohol **42**, which was coupled with carboxylic acid **10**. Subsequent diazotization provided targeted diazoester **39**.

Scheme 12. Alternative plan to access cycloheptadiene **6**Scheme 13. Alternative plan to access diazoester **39**

This diazoester was poised for a tandem cyclopropanation-Cope cascade to generate cycloheptadiene **6** directly (Scheme 14). To our surprise, treatment of diazoester **39** with Cu(tbs)₂ under N₂-atmosphere in a glovebox resulted in the formation of a compound possessing three olefins, instead of two as in desired product **6**. After careful analysis, we identified cycloheptatriene **44** as the major product of this reaction. Introduction of DBU into the reaction resulted in conversion of minor products, which were presumably olefin positional isomers of cycloheptatriene **44** to triene **44**. Cycloheptatriene **44** likely arose from air-accelerated oxidation during work-up of cycloheptadiene **43**. ¹H NMR spectroscopic analysis of the crude reaction mixture prior to exposure to air revealed the presence of double-bond isomerized enone **43**, instead of expected Cope product **6**. Presumably, the undesired oxidation and isomerization reactions are facilitated by the acidic β- and δ-protons of the C(3) ketone.


Scheme 14. Cyclopropanation-Cope-oxidation sequence with nucleophilic epoxidation



With cycloheptatriene **44** in hand, we suspected that appropriate orbital overlap could render C(5) as the reactive electrophile in a conjugated acceptor, a property we would hope to use in a total synthesis of ineleganolide. Nucleophilic epoxidation of cycloheptatriene **44** with the lithium salt of *tert*-butyl hydroperoxide yielded epoxide **45**, in a selective process that provided proof-of-concept for the anticipated reactivity of the C(4)–C(5) olefin.

Concurrently, alternative routes to the [6,7,5,5]-system were evaluated, with a focus on approaches that would avoid over-oxidation to the triene (Scheme 15). Formation of the triene could be avoided by employing lower $\text{Cu}(\text{tbs})_2$ -loadings, and intercepting the intermediate cycloheptadiene **43** prior to exposure to air. For example, diastereoselective epoxidation of the C(6)–C(7) olefin of intermediate diene **43** with $\text{VO}(\text{acac})_2$ and TBHP afforded epoxide **46** in 32% yield in a formal cyclopropanation, Cope rearrangement, epoxidation sequence. Alternatively, acetylation of the intermediate C(3) ketone furnished enol ester **47** in 44% yield in a formal cyclopropanation, Cope rearrangement, enolate trapping sequence. In turn, triene **47** could undergo selective oxidation of the C(6)–C(7) olefin either by epoxidation catalyzed by $\text{VO}(\text{acac})_2$ and TBHP to generate epoxide **48** diastereoselectively, or by oxidative rearrangement with pyridinium chlorochromate (PCC) to install C(6) ketone **49**, albeit with elimination of the critical C(8) alcohol. These additional transformations brought us tantalizingly close to a full synthesis of the des-isopropyl model system of ineleganolide (i.e., **5**). Full details of the ensuing challenges have been described in a series of manuscripts detailing a related model system.⁶

Scheme 15. Cyclopropanation-Cope-epoxidation and cyclopropanation-Cope-enolate trapping sequences




Conclusion

The construction of the ring system of ineganolide (**1**) has been demonstrated in a strategy that merges the asymmetric ketone alkylation and tandem transactonization-Cope rearrangements or formal cyclopropanation-Cope-epoxidation or cyclopropanation-Cope-enolate trapping sequences. This effort prompted the development of the previously unknown palladium-catalyzed asymmetric alkylation of dioxanones,^{8b} and a mild oxidative bromination, Wittig olefination and reduction sequence^{8a} to advance the resultant chloroalkene to enantioenriched cyclopentenol **11**. This alcohol can be coupled with cyclohexenone acid **10**, and the resultant vinylogous β -ketoesters can be advanced to rigid cyclopropanes. By necessity, we have created a rich body of chemistry exploring transactonizations in *cis*-substituted cyclopentane diols, including a transactonization-Cope cascade and a formal cyclopropanation-Cope-epoxidation sequence. This research complements a recently disclosed route to access the carbon framework of ineganolide via a formal cyclopropanation-Cope sequence.⁶ All of these cascade sequences provide access to the rigid [6,7,5]-fused scaffold of ineganolide (**1**).


EXPERIMENTAL SECTION

General Methods. Unless otherwise stated, reactions were performed in flame-dried glassware under an argon atmosphere using dry, deoxygenated solvents. Solvents were dried by passage through an activated alumina column under argon. Tetrabutylammonium triphenyldifluorosilicate (TBAT) was purchased from Sigma-Aldrich Chemical Company and azeotropically dried five times from acetonitrile prior to use. Trimethylsilyl chloride (TMSCl), pyridine, and triethylamine (NEt_3) were distilled from sodium hydride immediately prior to use. Sodium iodide was dried by heating at 90°C (2 torr) for 12 h. $\text{TEMPO}^+\text{BF}_4^-$,¹⁴ *p*-acetamidobenzenesulfonyl azide (*p*ABSA), TsN_3 ,²⁶ bis(*N*-*tert*-butylsalicylaldehydiminato) copper (II, $\text{Cu}(\text{tbs})_2$),²⁷ Dess-Martin periodinane (DMP),²⁸ diazomethane,²⁹ and Amberlyst A26 ($\text{S}_2\text{O}_3^{2-}$)³⁰ were prepared by known methods. Other reagents were used without further purification. Molecular sieves were purchased from Sigma Aldrich Chemical Company as activated 5 μm powder and stored in a 120°C drying oven until immediately prior to use unless otherwise noted. Rhodium octanoate was purchased from Strem. *N*-(3-dimethylaminopropyl)-*N'*-ethylcarbodiimide hydrochloride ($\text{EDC}\cdot\text{HCl}$) and 4-(dimethyl amino)pyridine (DMAP) and *N,N'*-dicyclohexylcarbodiimide (DCC) were purchased from Sigma Aldrich Chemical Company. Reaction temperatures were controlled by an IKA

temperature modulator. Thin-layer chromatography (TLC) was performed using E. Merck silica gel 60 F254 precoated plates (0.25 mm) and visualized by UV fluorescence quenching, anisaldehyde, or CAM staining. Florisil® (100–200 mesh) and ICN Silica gel (particle size 0.032–0.063 mm) were used for flash chromatography. Chiral HPLC analysis was performed with an Agilent 1100 Series HPLC utilizing chiralpak AD or chiralcel OD-H columns (4.6 mm x 25 cm) obtained from Daicel Chemical Industries, Ltd. with visualization at 254 nm or 220 nm. Chiral GC analysis was performed with an Agilent 6850 GC utilizing a G-TA (30 m x 0.25 cm) column (1.0 mL/min carrier gas flow). Optical rotations were measured with a Jasco P-1010 polarimeter at 589 nm. ¹H and ¹³C NMR spectra were recorded on a Varian Mercury 300, 500, or 600 NMR spectrometer (at 300, 500, or 600 MHz, and 75, or 125 MHz, respectively), and are reported relative to Me₄Si (δ 0.0). Data for ¹H NMR spectra are reported as follows: chemical shift (δ ppm) (multiplicity, coupling constant (Hz), integration). Data for ¹³C NMR spectra are reported in terms of chemical shift relative to Me₄Si (δ 0.0). If carbons were not recorded in a particular spectrum, then the missing carbons were noted by italicizing the absent carbon in a partial formula at the beginning of the spectrum. IR spectra were recorded on a Perkin Elmer Paragon 1000 spectrometer and are reported in wavenumbers (cm⁻¹). High resolution mass spectra were obtained from the Caltech Mass Spectral Facility, or acquired using an Agilent 6200 Series TOF with an Agilent G1978A Multimode source in electrospray ionization (ESI), atmospheric pressure chemical ionization (APCI), or mixed (MM) ionization mode.




2-Chloroallyl mesylate (13). A flask was charged with 2-chloroallyl alcohol (1.0 g, 10.8 mmol, 1 equiv), THF (20 mL, 0.54 M), and NEt₃ (3.0 mL, 21.5 mmol, 2 equiv) at 0 °C (ice water bath). The solution was treated dropwise with mesyl chloride (1.26 mL, 16.2 mmol, 1.5 equiv). After 2 h at 0 °C, the reaction was quenched by addition of NaHCO₃, and the mixture was extracted with Et₂O. The organic layer was washed successively with 1N HCl (20 mL), aq NaHCO₃ (20 mL), and brine (20 mL), dried over MgSO₄, filtered, and concentrated under reduced pressure. The residue was purified by column chromatography (SiO₂ (ca. 50 g) with 1:4 EtOAc:hexanes as the eluent) to provide a colorless oil (1.716 g, 93% yield). R_f 0.34 (1:2 EtOAc:hexanes); ¹H NMR (300 MHz, CDCl₃) δ 5.65 (d, 1H, J = 0.5 Hz), 5.55 (d, 1H, J = 0.5 Hz), 4.77 (s, 2H), 3.10 (s, 3H); ¹³C NMR (75 MHz, CDCl₃) δ 134.1, 117.7, 70.9, 38.4; IR (thin film/NaCl) 3033, 2943, 1639, 1360, 1175, 1010 cm⁻¹; HRMS (EI) *m/z*: [M⁺]⁺ Calcd for C₄H₇O₃SCl 169.9804; Found 169.9811.




(6*R*,7α*S*)-2,2,7α-Trimethyl-4,6,7α-tetrahydrocyclopenta[*d*][1,3]dioxin-6-ol (11). A flask was charged with a colorless solution of ketone **16** (0.110 g, 0.60 mmol, 1 equiv), MeOH (7.4 mL, 0.082 M) and CeCl₃·7H₂O (0.292 g, 0.78 mmol, 1.3 equiv). The solution was cooled to 0 °C (ice water bath), and treated with NaBH₄ (55 mg). Bubbles evolved. The solution was treated with a 1.5 M NaOH solution, extracted with EtOAc (to 150 mL), dried over Na₂SO₄, filtered, and concentrated under reduced pressure, and immediately diluted with CH₂Cl₂. Alcohol **11** was purified by silica gel chromatography (SiO₂ (ca. 16 mL)) with 1:1 pentane:Et₂O as the eluent to give pure alcohol **11** as a colorless oil (75 mg, 67% yield). The alcohol was immediately diluted in Et₂O. R_f 0.11 (50% Et₂O in hexanes; visualized with anisaldehyde); ¹H NMR (300 MHz, CDCl₃) δ 5.57 (q, J = 1.7, 1H), 4.77–4.70 (m, 1H), 4.70–4.61 (m, 2H), 2.57 (dd, J = 12.4, 6.5, 1H), 1.82 (ddd, J = 12.5, 6.8, 1.1, 1H), 1.63 (dt, J = 6.8, 3.3, 1H), 1.48 (s, 3H), 1.41 (s, 6H); ¹³C NMR (126 MHz, CDCl₃) δ 144.7, 125.7, 105.0, 100.2, 81.1, 74.0, 60.4, 53.5, 30.0, 28.3, 26.0; IR (NaCl) 3400 (b), 1197,


1156, 1107, 1088, 1061, 1037, 847; HRMS (EI) m/z : $[M-Me]^+$ Calcd for $C_9H_{13}O_3$ 169.0865; Found 169.0858; $[\alpha]_D^{21.6} -2.7^\circ$ (c 0.235, $CHCl_3$, 91% ee).



Ethyl 2-(3-oxocyclohex-1-en-1-yl)acetate (SI2).¹² To a 0.17 M acetonitrile solution of known allylic alcohol **50**¹² (1.14 g, 6.19 mmol) at ambient temperature was added $TEMPO^+BF_4^-$ (1.49 g, 6.15 mmol). After stirring for 1 hour, the orange solution was poured onto H_2O (30 mL) and Et_2O (100 mL). The organic and aqueous phases were separated and the aqueous layer further extracted with Et_2O (2 x 100 mL). The combined organic layers were dried with Na_2SO_4 , filtered and concentrated under reduced pressure. The crude material was purified by flash chromatography (1:10→1:4→1:1 EtOAc:hexanes) to yield known¹² cyclohexenone **51** (0.77 g, 69% yield) as a colorless liquid. 1H NMR of this compound was as reported in the literature.¹²




2-(3-Oxocyclohex-1-en-1-yl)acetic acid (10). To a 1:1 EtOH/ H_2O solution (0.14 M) of ester **51**¹² (0.76 g, 4.2 mmol) cooled in an ice water bath was added NaOH (aq) (1.8 mL, 3.04 M, 5.47 mmol) dropwise. The bright yellow solution was allowed to warm to room temperature and stirred for 5 hours. The solution was quenched with 10% HCl (aq) (15 mL) and extracted with EtOAc (5 x 100 mL). The combined organic layers were dried over Na_2SO_4 , filtered, and concentrated under reduced pressure. The crude oil was purified by flash chromatography (3:1→1:1→0:1 hexanes/EtOAc eluent) to yield carboxylic acid **10** (0.57 g, 89% yield) as a yellow oil. 1H NMR (500 MHz, $CDCl_3$) δ 6.02–6.00 (m, 1H), 3.28 (d, J = 0.7, 2H), 2.43–2.39 (m, 4H), 2.06–2.00 (m, 2H); ^{13}C NMR (125 MHz, $CDCl_3$) δ 200.1, 174.5, 157.1, 129.1, 43.1, 37.0, 29.8, 22.6; IR (thin film/NaCl) 3445, 1722, 1652, 1260 cm^{-1} ; HRMS (ESI-APCI) m/z : $[M+H]^+$ Calcd for $C_8H_{10}O_3$ 155.0703; Found 155.0704.




Ester 19. A concentration flask was charged with a yellow mixture of acid **10** (1.303 g, 8.4 mmol, 3 equiv), alcohol **11** (from 2.95 mmol ketone **16**, 1 equiv) and EDC·HCl (1.131g, 5.9 mmol, 2 equiv) in CH_2Cl_2 (29.5 mL, 0.1 M). The suspension was cooled to 0 °C (ice water bath) and treated with DMAP (72 mg, 0.59 mmol, 0.2 equiv). The suspension was allowed to gradually warm to room temperature (ca. 25 °C) as the ice bath melted. After 3.5 hours, the reaction was treated with additional EDC·HCl (1.131g, 5.9 mmol, 2 equiv). After 1.5 hours, the solution was cooled to 0 °C (ice water bath) and treated with 0.1 N HCl (6 mL). The layers were separated, and the organics were further rinsed with 0.1 N HCl (6 mL), 5% K_2CO_3 (aq, 6 mL x 2), and brine (6 mL) in succession. Each aqueous layer was further extracted with EtOAc (12 mL x 10). The organic extracts were dried over $MgSO_4$, filtered and concentrated under reduced pressure to give ester **19** as a yellow oil (0.923 g, 98% yield). R_f 0.72 (EtOAc; UV active; visualized with anisaldehyde stain); 1H NMR (300 MHz, $CDCl_3$) δ 5.93 (d, J = 0.7, 1H), 5.59–5.56 (m, 1H), 5.54–5.53 (m, 1H), 4.67–


4.63 (m, 2H), 3.24 (s, 2H), 2.58 (ddd, $J = 12.9, 6.7, 1.2$, 1H), 2.40–2.32 (m, 4H), 2.03 (dt, $J = 12.7, 6.5$, 2H), 1.94 (dd, $J = 12.8, 6.6$, 1H), 1.46 (s, 3H), 1.44 (s, 3H), 1.41 (s, 3H); ^{13}C NMR (125 MHz, CDCl_3) δ 199.5, 169.4, 157.1, 147.2, 129.1, 121.3, 100.4, 81.0, 77.0, 60.4, 49.3, 43.6, 37.3, 30.1, 29.8, 28.6, 25.8, 22.8; IR (thin film/ NaCl) 1732, 1667, 1169, 1127, 974 cm^{-1} ; HRMS (EI) m/z : $[\text{M}^+]^+$ Calcd for $\text{C}_{18}\text{H}_{24}\text{O}_5$ 320.1624; Found 320.1613; $[\alpha]_{\text{D}}^{24.1}$ 28.4° (c 0.11, CHCl_3 , derived from ketone **16** with 91% ee).



Diazoester 9. A vial beneath $\text{N}_2(\text{g})$ was charged with ester **19** (32 mg, 0.1 mmol, 1 equiv), CH_3CN (0.5 mL, 0.2 M) and $p\text{ABSA}$ (31 mg, 0.13 mmol, 3 equiv).³¹ The solution turned deep yellow upon dropwise addition of NEt_3 (0.04 mL, 0.3 mmol, 3 equiv). After six hours, the yellow mixture was treated with Et_2O and filtered through a plug of SiO_2 (ca. 1 mL) with copious elution. The yellow solution was concentrated under reduced pressure. The residue was diluted with EtOAc and purified by flash chromatography (SiO_2 (ca. 14 mL), 1:3 \rightarrow 1:2 EtOAc :hexanes eluent) to give diazoester **9** as a yellow oil (37 mg, >99% yield). R_f 0.38 (1:1 EtOAc :hexanes; visualized with UV); ^1H NMR (500 MHz, CDCl_3) δ 6.37 (s, 1H), 5.67 (dddd, $J = 8.5, 6.7, 3.4, 1.9$, 1H), 5.54 (dd, $J = 3.5, 1.5$, 1H), 4.69–4.65 (m, 1H), 4.65–4.61 (m, 1H), 2.58 (dd, $J = 12.8, 6.9$, 1H), 2.54–2.49 (m, 2H), 2.41–2.36 (m, 2H), 2.06 (dt, $J = 12.7, 6.3$, 2H), 1.97 (dd, $J = 12.9, 6.5$, 1H), 1.43 (s, 3H), 1.43 (s, 3H), 1.39 (s, 3H); ^{13}C NMR (125 MHz, CDCl_3 , $-\text{C}=\text{N}_2$ or $-\text{C}(\text{CH}_3)_2$) δ 197.4, 162.7, 147.5, 147.2, 121.3, 120.9, 100.5, 81.0, 77.5, 60.4, 49.4, 37.0, 30.0, 28.6, 26.8, 25.9, 22.5; IR (CH_2Cl_2) 2101, 1708, 1580, 1191, 1137, 1110, 1064, 994 cm^{-1} ; HRMS (EI) m/z : $[\text{M}^+]^+$ Calcd for $\text{C}_{18}\text{H}_{22}\text{O}_5\text{N}_2$ 346.1529; Found 346.1524; $[\alpha]_{\text{D}}^{24.4}$ 47.9° (c 0.11, CHCl_3 , derived from ketone **16** with 91% ee).




Pyrazole 21. A flask fitted with a reflux condenser was charged with diazoester **9** (20 mg, 0.054 mmol, 1 equiv) and PhMe (8 mL, 0.00675 M). The yellow solution was treated with a warm (previously refluxing) solution of copper(II) acetylacetonate (2.8 mg, 0.011 mmol, 0.2 equiv) in PhMe (2 mL, with 10 mL wash, 0.054 M in diazoester **9**). The solution was heated in an oil bath at 110 °C. After 40 minutes, the solution was removed from heat and concentrated under reduced pressure. The residue was diluted with 1:1 EtOAc :hexanes and purified by flash chromatography ($\text{SiO}_2 \sim 8$ mL; 1:1 EtOAc :hexanes eluent) to give pyrazole **21** as a white solid (10 mg, 59% yield). Colorless crystals could be obtained by slow diffusion of PhH into a solution of pyrazole **21** in CHCl_3 . Melting point analysis resulted in crystals yellowing at 149–150 °C and then converting to a red liquid as gas evolved at 170–172 °C. R_f 0.26 (1:1 EtOAc :hexanes; visualized with UV or anisaldehyde stain); ^1H NMR (500 MHz, CDCl_3) δ 11.40 (s, 1H), 5.83 (ddtd, $J = 6.8, 5.0, 3.4, 1.9$, 1H), 5.70 (dt, $J = 3.0, 1.5$, 1H), 4.73 (ddd, $J = 15.7, 3.3, 2.4$, 1H), 4.70–4.68 (m, 1H), 3.04 (t, $J = 6.1$, 2H), 2.71 (dd, $J = 12.7, 6.8$, 1H), 2.63 (dd, $J = 7.3, 5.6$, 2H), 2.23–2.13 (m, 3H), 1.66 (s, 2H), 1.49 (s, 3H), 1.48 (s, 3H), 1.40 (s, 3H); ^{13}C NMR (125 MHz, CDCl_3 , $-\text{C}=\text{O}$, $-\text{CC}(\text{O})\text{OR}$) δ 147.2, 131.8, 121.4, 110.2, 100.4, 81.0, 77.2, 60.4, 49.5, 38.7, 30.1, 29.9, 28.6, 25.9, 24.6, 21.5; IR (thin film/ NaCl) 3286, 1716, 1689, 1188, 1172, 1094, 1042, 994 cm^{-1} ; HRMS (EI) m/z : $[\text{M}+\text{H}]^+$ Calcd for $\text{C}_{18}\text{H}_{23}\text{O}_5\text{N}_2$ 347.1607; Found 347.1596; $[\alpha]_{\text{D}}^{27.4}$ +45.6° (c 0.59, acetone, derived from ketone **16** with 91% ee).



Oxidized 22a, or **22b**, and **23a**, or **23b**.^{17a} As a note of clarification, one set of data has been acquired for compound **22**, and another for compound **23**. ¹³C and IR spectra support their assignment as ketones **22a** and **23a**, respectively, while HRMS data supports their assignment as hydrates **22b** and **23b**, respectively. It is likely that these hydrates form under the conditions used to obtain HRMS data.


A vial was charged with rhodium octanoate dimer ($\text{Rh}_2(\text{oct})_4$, 0.6 mg, 0.0007 mmol, 0.01 equiv) and CH_2Cl_2 (0.2 mL, 0.375 M relative to diazoester **9**). The pale green solution was cooled to 0 °C (ice water bath). To it was added dropwise diazoester **9** (28 mg, 0.075 mmol, 1 equiv) in CH_2Cl_2 (0.22 mL, 0.341 M). After 25 minutes, the yellow solution was treated with additional $\text{Rh}_2(\text{oct})_4$ (0.4 mg, 0.0005 mmol, 0.007 equiv). After an additional 18 hours, the yellow solution was treated with a third portion of $\text{Rh}_2(\text{oct})_4$ (0.4 mg, 0.0005 mmol, 0.007 equiv). After an additional two hours, the reaction was diluted with EtOAc (10 mL), filtered through a plug of celite and concentrated under reduced pressure. The residue was diluted with EtOAc (10 mL), filtered and concentrated. This residue was purified by flash chromatography ($\text{SiO}_2 \sim 16$ mL; 1:40 \rightarrow 1:20 \rightarrow 1:5 \rightarrow 1:0 EtOAc: CH_2Cl_2 then 1:20 MeOH:EtOAc elution) to give oxidized **22a** (9.8 mg, 41% yield). R_f 0.61 (1:5 EtOAc: CH_2Cl_2 ; UV/Vis); ¹H NMR (500 MHz, CDCl_3) δ 6.62 (t, $J = 1.7$, 1H), 5.75–5.70 (m, 1H), 5.58 (dd, $J = 3.7, 1.7$, 1H), 4.70–4.62 (m, 2H), 2.64 (dd, $J = 12.9, 6.9$, 1H), 2.54 (td, $J = 6.1, 1.8$, 2H), 2.52–2.47 (m, 2H), 2.11–2.02 (m, 3H), 1.45 (d, $J = 0.5$, 3H), 1.42 (s, 3H), 1.38 (s, 3H); ¹³C NMR (125 MHz, CDCl_3) δ 200.0, 188.7, 162.7, 150.0, 148.6, 137.0, 120.2, 100.6, 81.0, 79.0, 60.3, 49.0, 38.2, 30.1, 28.7, 25.7, 23.0, 21.9; IR (thin film/NaCl) 3384, 2934, 2871, 1732, 1682, 1455, 1429, 1416, 1373, 1350, 1317, 1257, 1195, 1148, 1110, 1065, 1030, 978, 949, 902, 849, 735 cm^{-1} ; HRMS (ES) m/z : $[\text{M}+\text{H}]^+$ Calcd for **22** $\text{C}_{18}\text{H}_{24}\text{O}_7$ 353.1600; Found 353.1631; $[\alpha]_D^{26.6} +30.4^\circ$ (c 0.15, CHCl_3 , derived from ketone **16** with 91% ee).

Chromatography also furnished slightly impure acetone-cleaved **23a** or **23b**. Acetone-cleaved **23a** or **23b** was purified by flash chromatography ($\text{SiO}_2 \sim 2$ mL; 4:1 EtOAc:hexanes elution) to give acetone-cleaved **23a** (4.8 mg, 23% yield). R_f 0.40 (EtOAc; UV active, visualized with anisaldehyde stain); ¹H NMR (300 MHz, CDCl_3) δ 6.65 (t, $J = 1.7$, 1H), 5.84 (q, $J = 1.7$, 1H), 5.70–5.64 (m, 1H), 4.45–4.35 (m, 2H), 2.60 (dd, $J = 14.7, 7.2$, 2H), 2.54 (ddd, $J = 6.0, 6.0, 1.8$, 2H), 2.51–2.43 (m, 2H), 2.44 (s, 1H), 2.16 (dd, $J = 14.6, 3.3$, 1H), 2.13–2.03 (m, 3H), 1.44 (s, 3H); ¹³C NMR (125 MHz, CDCl_3) δ 200.2, 188.6, 162.6, 155.2, 150.1, 137.1, 124.4, 81.4, 78.8, 59.2, 47.9, 38.2, 26.6, 23.0, 21.9; IR (thin film/NaCl) 3382, 2968, 2935, 1732, 1682, 1193, 1110, 1092, 1065, 1030, cm^{-1} ; HRMS (ES+) m/z Calcd for **23** $\text{C}_{15}\text{H}_{18}\text{O}_6$ $[\text{M}+\text{H}]^+$: 295.1182; Found 295.1195; HRMS (ES) m/z : $[\text{M}+\text{H}]^+$ Calcd for **23** $\text{C}_{15}\text{H}_{20}\text{O}_7$ 313.1287; Found 313.1286; $[\alpha]_D^{24.9} +48.8^\circ$ (c 0.09, CHCl_3 , derived from ketone **16** with 91% ee).




Silyl ether 52. A flask was charged with acetone **19** (0.451 g, 1.41 mmol, 1 equiv) in MeOH (30 mL, 0.047 M), and the solution was cooled to 0 °C (ice water bath). The solution was treated dropwise with a solution of fumaric acid (59 mg, 0.50 mmol, 0.36 equiv) in MeOH (10 mL, 0.14 M total relative to acetone). After 3 days, the reaction was quenched by addition of a 1:1 solution of H₂O and saturated aq NaHCO₃ (40 mL each), and extracted with EtOAc (300 mL, and then 200 mL x 5). The organic extracts were dried over Na₂SO₄, filtered and concentrated under reduced pressure to give a golden/orange oil.

The resultant oil was dissolved in CH₂Cl₂ (35 mL, 0.040 M), placed beneath an Ar(g) atmosphere, and cooled to 0 °C (ice water bath). The solution was treated dropwise with imidazole (0.303 g, 4.46 mmol, 3.16 equiv) in CH₂Cl₂, followed by TIPSCl (0.91 mL, 4.25 mmol, 3.01 equiv) and DMAP (20 mg, 0.16 mmol, 0.11 equiv) in CH₂Cl₂. After 17 hours, the solution was quenched with saturated aq NH₄Cl (50 mL), and extracted with EtOAc (150 mL x 3). The organic extracts were dried over Na₂SO₄, filtered and concentrated under reduced pressure. The reaction was purified by flash chromatography (1:19 → 1:9 EtOAc:CH₂Cl₂ eluent) to give silyl ether **52** (0.371 g, 59% yield over two steps) as a yellow oil. *R*_f 0.76 (EtOAc, UV/Vis, visualized with anisaldehyde stain); ¹H NMR (500 MHz, CDCl₃) δ 5.95 (t, *J* = 1.0, 1H), 5.73 (dd, *J* = 3.5, 1.7, 1H), 5.51 (dtd, *J* = 7.2, 3.6, 1.8, 1H), 4.59–4.38 (m, 2H), 3.24 (s, 2H), 2.82 (s, 1H), 2.52 (dd, *J* = 14.4, 7.3, 1H), 2.42–2.35 (m, 4H), 2.11–1.91 (m, 3H), 1.41 (s, 3H), 1.21–1.10 (m, 3H), 1.09–1.05 (m, 18H); ¹³C NMR (125 MHz, CDCl₃) δ 199.6, 169.5, 157.3, 153.6, 129.1, 124.5, 81.1, 76.9, 60.4, 48.0, 43.7, 37.4, 29.8, 26.8, 22.8, 18.2, 12.0; IR (thin film/NaCl) 3440, 1733, 1668, 1192, 1165, 1127, 1107, 1055, cm⁻¹; HRMS (mixed EIC) *m/z*: [M+Na]⁺ Calcd for C₂₄H₄₀O₅Si 459.2537; Found 459.2539; [α]_D^{18.9} +44.8° (*c* 0.93, CHCl₃, derived from ketone **16** with 91% ee).




Diazoester 24. A flask was charged with a pale yellow solution of ester **52** (53 mg, 0.121 mmol, 1 equiv) in CH₃CN (4.7 mL, 0.026 M) with TsN₃ (0.600 g, 0.535 mmol, 4.4 equiv), and cooled to 0 °C (ice water bath). CAUTION!!! TsN₃ is SHOCK SENSITIVE AND POTENTIALLY EXPLOSIVE. The solution was treated with Et₃N (0.03 mL, 0.22 mmol, 1.8 equiv) dropwise, and immediately turned deeper yellow in color. After 24 hours, the reaction was concentrated under reduced pressure. The crude yellow mixture was purified by flash chromatography (SiO₂ ~ 16 mL; 1:5 EtOAc:PhH eluent) to give diazoester **24** (48.9 mg, 87% yield) as a yellow oil. *R*_f 0.48 (1:2 EtOAc:PhH; visualized with UV); ¹H NMR (500 MHz, CDCl₃) δ 6.37 (s, 1H), 5.71 (dd, *J* = 3.3, 1.6, 1H), 5.59 (ddd, *J* = 7.3, 3.7, 1.9, 1H), 2.82 (s, 1H), 2.58–2.49 (m, 3H), 2.41–2.35 (m, 2H), 2.10–2.00 (m, 3H), 1.40 (s, 3H), 1.13 (tt, *J* = 12.2, 6.9, 3H), 1.08–1.03 (m, 21H); ¹³C NMR (125 MHz, CDCl₃, -C=N₂) δ 197.5, 162.7, 153.7, 147.5, 124.4, 120.6, 81.0, 77.4, 60.4, 48.2, 37.0, 27.0, 26.8, 22.5, 18.2, 12.0; IR (thin film/NaCl) 3407, 2101, 1709, 1645, 1191, 1138, 1106, 1061, 993 cm⁻¹; HRMS (ES) *m/z*: [M+H]⁺ Calcd for C₂₄H₃₉N₂O₅Si 463.2628; Found 463.2640; [α]_D^{23.4} +48.5° (*c* 0.175, CHCl₃, derived from ketone **16** with 91% ee).



Cyclopropane 25a. In the glovebox, a flask was charged with Cu(tbs)₂ (92.2 mg, 0.22 mmol, 1.1 equiv) and CH₂Cl₂ (20 mL, 0.01 M). This solution was treated with diazoester **24** (93.2 mg, 0.20 mmol, 1 equiv) in CH₂Cl₂ (8 mL, 0.025 M). After 7 days, the reaction was removed from the

glovebox and concentrated under reduced pressure to yield a brown residue, which was purified by flash chromatography (SiO₂ ~ 16 mL; 1:1 EtOAc:hexanes eluent) to give cyclopropane **25a** as a yellow oil (62.8 mg, 74% yield). *R*_f 0.18 (1:1 EtOAc:hexanes; UV active; visualized with anisaldehyde stain); ¹H NMR (500 MHz, CDCl₃) δ 5.87 (s, 1H), 4.87 (td, *J* = 6.2, 2.9, 1H), 4.07 (d, *J* = 11.5, 1H), 3.49 (d, *J* = 11.5, 1H), 2.68–2.60 (m, 1H), 2.59 (d, *J* = 5.9, 1H), 2.45–2.23 (m, 2H), 2.17–1.97 (m, 3H), 1.93–1.83 (m, 1H), 1.74 (s, 1H), 1.49 (s, 3H), 1.11–0.76 (m, 21H); ¹³C NMR (125 MHz, CDCl₃) δ 199.7, 173.1, 156.4, 130.0, 93.8, 70.6, 59.9, 59.6, 53.8, 48.0, 42.8, 37.6, 29.5, 22.8, 19.1, 18.2, 12.1; IR (thin film/NaCl) 3427, 1759, 1668, 1626, 1192, 1175, 1122, 1085, 1065 cm⁻¹; HRMS (FAB) *m/z*: [M+H]⁺ Calcd for C₂₄H₃₉O₅Si 435.2567; Found 435.2586; [α]_D^{21.2} -12.6° (*c* 0.125, CHCl₃, derived from ketone **16** with 91% ee).




26
mixture of C(13)
diastereomers


Dihydropyran 26. A flask was charged with a colorless solution of silyl ether **25a** (1 equiv) in THF (0.00825 M) at 0 °C (ice water bath). The solution was treated with TBAF (1M in THF, 1.2 equiv), and the solution turned yellow in color. After 30 min, the reaction was filtered through SiO₂ (ca. 1 mL; EtOAc elution), and concentrated under reduced pressure to give a crude white paste, assigned as crude diol **53**. *R*_f 0.14 (EtOAc; UV/Vis active); ¹H NMR (500 MHz, CDCl₃) δ 5.93 (s, 1H), 4.93 (t, *J* = 6.3, 1H), 3.94 (d, *J* = 11.9, 1H), 3.66 (d, *J* = 12.0, 1H), 2.74–2.65 (m, 2H), 2.69 (d, *J* = 6.0, 1H), 2.47–2.34 (m, 3H), 2.31–2.23 (m, 2H), 2.19 (dd, *J* = 14.0, 7.0, 1H), 2.13 (d, *J* = 13.5, 1H), 2.11–2.01 (m, 2H), 1.99–1.89 (m, 2H), 1.69 (s, 3H); HRMS (EI) *m/z*: [M+]⁺ Calcd for C₁₅H₁₈O₅ 278.1154; Found 278.1143.

A flask was charged with a solution of crude diol **53**, CH₂Cl₂ (1.1 mL, 0.012 M) and pyridine (0.02 mL, 0.159 mmol, 12 equiv) at 0 °C (ice water bath), which was subsequently treated with DMP (7.9 mg, 0.018 mmol, 1.4 equiv). CAUTION! DMP is a HEAT- and SHOCK-SENSITIVE COMPOUND, showing exotherms when heated (>130 °C). ALL OPERATIONS SHOULD BE CONDUCTED BEHIND A BLAST SHIELD. After 2 days at this temperature, the mixture was diluted with EtOAc (ca. 10 mL), and concentrated under reduced pressure. The resultant semi-solid was treated with CH₂Cl₂ (ca. 1 mL) and Amberlyst A26 (S₂O₃²⁻) (19.6 mg). After 6.5 h, the mixture was treated with JandaJel (2.3 mmol/g, 17.2 mg, 0.0396 mmol, 3.0 equiv). After an additional 17.5 h, the mixture was filtered through SiO₂ (ca. 0.2 mL; EtOAc elution), and concentrated under reduced pressure. The diastereomeric mixture was purified by thin layer chromatography on a 250 μm silica gel plate (10 cm wide, 20 cm tall; Et₂O x 2 elution). The desired diastereomeric mixture was removed from the plate at *R*_f 0.10. The diastereomeric mixture of dihydropyrans (**26**) was further purified by thin layer chromatography on a 250 μm silica gel plate (10 cm wide, 20 cm tall; 1:1 EtOAc:PhH elution). The diastereomeric mixture was removed from the plate at *R*_f 0.27, as a white solid (<20% yield). Major diastereomer: ¹H NMR (500 MHz, CDCl₃) δ 4.94 (app. q, *J* = 3.7, 1H), 4.02 (d, *J* = 9.4, 1H), 3.90 (d, *J* = 9.4, 1H), 3.18 (d, *J* = 4.8, 1H), 2.77 (ddd, *J* = 14.2, 11.6, 5.1, 1H), 2.71 (d, *J* = 14.1, 1H), 2.43–2.28 (m, 2H), 2.27 (dt, *J* = 14.1, 2.3, 1H), 2.18 (dd, *J* = 14.3, 2.0, 1H), 2.08–2.03 (m, 1H), 2.04–1.86 (m, 3H), 1.76–1.70 (m, 1H), 1.52 (s, 3H); ¹³C NMR (125 MHz, CDCl₃) δ 208.5, 172.9, 84.7, 80.5, 78.0, 64.7, 55.7, 53.0, 46.2, 45.9, 43.7, 40.9, 31.5, 27.8, 20.2. Minor diastereomer: ¹H NMR (500 MHz, CDCl₃) δ 4.94 (app. q, *J* = 3.7, 1H), 4.11 (d, *J* = 9.6, 1H), 3.88 (d, *J* = 9.6, 1H), 3.61 (d, *J* = 15.1, 1H), 3.18 (d, *J* = 4.8, 1H), 2.77 (ddd, *J* = 14.2, 11.6, 5.1, 1H), 2.59 (d, *J* = 15.1, 1H), 2.43–2.28 (m, 2H), 2.18 (dd, *J* = 14.3, 2.0, 1H), 2.08–2.03 (m, 1H), 2.04–1.86 (m, 3H), 1.76–1.70 (m, 1H), 1.51 (s, 3H); ¹³C NMR (125 MHz, CDCl₃) δ 209.7, 173.0, 84.7, 80.5, 78.1, 64.7, 56.6, 52.8, 48.5, 45.6, 43.9, 40.6,

31.2, 27.8, 21.5. Diastereomeric Mixture: IR (thin film/NaCl) 3460, 1756, 1712, 1172, 1109, 1073, 1047, 1012 cm^{-1} ; HRMS (ESI-APCI) m/z : [M+Na] $^{+}$ Calcd for $\text{C}_{15}\text{H}_{18}\text{O}_5$ 301.1046; Found 301.1067.




Benzyl ether 27a.³² A flask was charged with a solution of silyl ether **25a** (1.1 mg, 0.0026 mmol, 1 equiv) in PhMe (0.2 mL, 0.013 M) beneath an Ar(g) atmosphere at room temperature (22.5 °C). The solution was treated with KI (0.9 mg, 0.0054 mmol, 2.1 equiv) and Ag_2O (0.9 mg, 0.0039 mmol, 1.5 equiv), followed by BnBr (10 μL , 0.084 mmol, 32 equiv). After 2 days, the yellow mixture was heated in an oil bath at 40 °C for 22 hours, at which point additional BnBr (10 μL , 0.084 mmol, 32 equiv) was added to the reaction. After 24 h, the suspension was treated with Ag_2O (3.0 mg, 0.013 mmol, 5 equiv) and KI (4.6 mg, 0.028 mmol, 10.6 equiv). After an additional 27 h, additional BnBr (10 μL , 0.084 mmol, 32 equiv) was added to the reaction. After an additional 27 h, the reaction was diluted with EtOAc (10 mL), filtered through SiO_2 , and concentrated under reduced pressure. The mixture was purified by thin layer chromatography on a 250 μm silica gel plate (10 cm wide, 20 cm tall; 1:10 acetone: CH_2Cl_2 elution) to give benzoyl ester **27a** (0.5 mg, 37% yield, R_f 0.69). R_f 0.22 (1:1 Et $_2\text{O}$:heptanes, UV/Vis, visualized with anisaldehyde stain); ^1H NMR (600 MHz, CDCl_3) δ 7.36–7.32 (m, 4H), 7.29–7.26 (m, 1H), 5.93 (t, J = 1.4, 1H), 4.63 (d, J = 11.7, 1H), 4.53 (dd, J = 7.5, 5.6, 1H), 4.38 (d, J = 11.6, 1H), 4.17 (d, J = 11.5, 1H), 3.57 (d, J = 11.5, 1H), 2.79–2.72 (m, 1H), 2.61 (d, J = 5.6, 1H), 2.47 (ddd, J = 16.6, 9.2, 4.7, 1H), 2.39 (ddd, J = 12.1, 7.9, 4.7, 1H), 2.29 (d, J = 13.9, 1H), 2.21–2.08 (m, 2H), 2.14 (dd, J = 13.9, 7.6, 1H), 1.95 (tdd, J = 13.8, 9.3, 4.6, 1H), 1.70 (s, 3H), 1.12–0.90 (m, 21H); ^{13}C NMR (125 MHz, C_6D_6) δ 199.2, 172.5, 156.2, 137.5, 129.7, 128.7, 128.2, 128.0, 93.1, 76.9, 71.4, 59.9, 59.4, 52.1, 48.1, 40.3, 37.6, 29.5, 22.9, 19.0, 18.2, 12.1; IR (thin film/NaCl) 1760, 1674, 1190, 1177, 1125, 1086, 1039 cm^{-1} ; HRMS-MM (ESI-APCI) m/z : [M+H] $^{+}$ Calcd for $\text{C}_{31}\text{H}_{44}\text{O}_5\text{Si}$ 525.3031; Found 525.3023; $[\alpha]_D^{22.9}$ +5.9° (c 0.20, CHCl_3 , derived from ketone **16** with 91% ee).




Secondary acetate 27b. A flask charged with CH_2Cl_2 (0.96 mL, 0.027 M), tertiary alcohol **25a** (11.2 mg, 0.0258 mmol, 1 equiv), Ac_2O (0.03 mL, 0.33 mmol, 13 equiv) and pyridine (0.03 mL, 0.38 mmol, 15 equiv) at room temperature (23 °C) was treated with DMAP (0.4 mg, 0.003 mmol, 0.13 equiv). After 1 h 10 m, the solution was extracted with EtOAc (10 mL x 4), and rinsed successively with 1N HCl (3 mL) and brine (3 mL x 2). The organic extracts were dried over MgSO_4 , filtered, and concentrated under reduced pressure to yield a secondary acetate **27b** (11.7 mg, 87.3% yield) as a pale yellow oil. R_f 0.64 (1:1 EtOAc:hexanes, UV/Vis, visualized with anisaldehyde stain); ^1H NMR (500 MHz, C_6D_6) δ 6.17 (t, J = 1.5, 1H), 5.11 (dd, J = 7.7, 5.9, 1H), 3.77 (d, J = 11.6, 1H), 3.40 (d, J = 11.6, 1H), 2.62 (d, J = 5.9, 1H), 2.46 (dddd, J = 9.4, 7.3, 5.0, 1.7, 1H), 2.11–1.96 (m, 4H), 1.80–1.72 (m, 4H), 1.58–1.44 (m, 2H), 1.42 (s, 3H), 0.93 (d, J = 1.9, 18H), 0.95–0.87 (m, 3H); ^1H NMR (500 MHz, CDCl_3) δ 5.89 (t, J = 1.5, 1H), 5.40 (td, J = 6.0, 2.1, 1H), 4.18 (d, J = 11.4, 1H), 3.52 (d, J = 11.5, 1H), 2.80 (d, J = 5.9, 1H), 2.54 (dddd, J = 18.1, 7.3, 4.9, 1.5, 1H), 2.45–2.29 (m, 2H), 2.23 (d, J = 0.9, 1H), 2.23 (d, J = 5.0, 1H), 2.21–2.13 (m, 1H), 2.09–2.01 (m, 1H), 2.00 (s, 3H), 1.99–1.89 (m, 1H), 1.69 (s, 3H), 1.10–0.92 (m, 3H), 1.01 (d, J = 3.8, 18H); ^{13}C NMR (125 MHz, C_6D_6) δ 196.5, 171.2, 170.1, 153.9, 130.1, 91.8, 73.1, 59.7, 57.9, 50.2, 47.6, 41.3,

37.1, 28.8, 22.5, 19.9, 18.5, 17.7, 11.8; IR (film) 1769, 1740, 1675, 1191, 1173, 1127, 1082, 1064 cm^{-1} ; HRMS (ESI-APCI) m/z : $[\text{M}+\text{H}]^+$ Calcd for $\text{C}_{26}\text{H}_{40}\text{O}_6\text{Si}$ 477.2667; Found 477.2667; $[\alpha]_{\text{D}}^{17.9} -11.6^\circ$ (c 0.35, CH_3CN , derived from ketone **16** with 91% ee).




Alcohols 28. Alcohol **25a** (57.2 mg, 0.131 mmol, 1 equiv) was converted to acetate **27b**, as described above. The crude **27b** as a pale yellow oil was dissolved in MeOH (12.0 mL, 0.011 M), cooled to 0 °C (ice water bath) and treated with NaBH_4 (18.6 mg, 0.491 mmol, 3.75 equiv). After 24 m, the yellow solution was treated with a saturated aq solution of NH_4Cl (0.70 mL). The mixture was diluted with EtOAc (125 mL), filtered through SiO_2 (ca. 2 mL) and concentrated under reduced pressure to an off-white solid. The solid was purified by preparative thin layer chromatography on a 250 μm analytical plate (2:1 CH_2Cl_2 :EtOAc x 2 elution) to give alcohol **28a** (21.4 mg, 34% yield) as a colorless oil and alcohol **28b** (32.2 mg, 51% yield) as a colorless oil. Alcohol **28a**: R_f 0.65 (1:1 EtOAc:hexanes, visualized by UV/Vis, or with anisaldehyde stain); ^1H NMR (500 MHz, CDCl_3) δ 5.73 (app dt, J = 3.1, 1.7, 1H), 5.41 (dt, J = 5.9, 4.1, 1H), 4.27 (d, J = 11.4, 2H), 3.45 (dd, J = 11.4, 8.7, 1H), 2.79 (d, J = 5.9, 1H), 2.26–2.21 (m, 2H), 2.19–2.11 (m, 1H), 2.00 (s, 3H), 1.91 (tdd, J = 7.8, 4.8, 3.1, 1H), 1.79–1.74 (m, 1H), 1.73 (ddd, J = 7.6, 4.3, 2.2, 1H), 1.70 (s, 3H), 1.68–1.61 (m, 1H), 1.50 (dddd, J = 12.5, 9.4, 6.5, 3.2, 1H), 1.44 (s, 1H), 1.14–0.94 (m, 21H); ^{13}C NMR (125 MHz, CDCl_3) δ 174.2, 171.3, 133.7, 133.0, 93.0, 73.8, 65.9, 60.0, 57.5, 51.0, 48.3, 41.9, 31.6, 28.1, 20.9, 19.3, 18.9, 18.2, 12.1; IR (film) 3482, 2941, 2866, 1760, 1740, 1462, 1374, 1242, 1080, 1065, 882, 683 cm^{-1} ; HRMS (ESI-APCI) m/z : $[\text{M}+\text{NH}_4]^+$ Calcd for $\text{C}_{26}\text{H}_{42}\text{O}_6\text{Si}$ 496.3089; Found 496.3071; $[\alpha]_{\text{D}}^{27.2} +8.7^\circ$ (c 1.52, CHCl_3 , derived from ketone **16** with 91% ee). Alcohol **28b**: R_f 0.61 (1:1 EtOAc:hexanes, UV/Vis, visualized with anisaldehyde stain); ^1H NMR (500 MHz, CDCl_3) δ 5.74 (dt, J = 3.3, 1.6, 1H), 5.42 (dt, J = 5.9, 4.1, 1H), 4.26 (d, J = 11.4, 1H), 4.17 (app d, 1H), 3.43 (d, J = 11.4, 1H), 2.77 (d, J = 5.9, 1H), 2.23 (d, J = 4.1, 1H), 2.19–2.14 (m, 2H), 2.02 (s, 3H), 1.82–1.72 (m, 3H), 1.69 (s, 3H), 1.66–1.51 (m, 2H), 1.11–0.95 (m, 21H); ^{13}C NMR (125 MHz, CDCl_3) δ 174.1, 171.2, 133.9, 132.6, 93.0, 73.7, 65.8, 59.9, 57.2, 51.0, 48.3, 41.7, 31.7, 28.1, 20.9, 19.3, 18.9, 18.2, 12.1; IR (thin film/ NaCl) 3447, 1763, 1740, 882 cm^{-1} ; HRMS (EI) m/z : $[\text{M}+\text{NH}_4]^+$ Calcd for $\text{C}_{26}\text{H}_{42}\text{O}_6\text{Si}$ 496.3089; Found 496.3099; $[\alpha]_{\text{D}}^{26.8} +33.1^\circ$ (c 1.01, CHCl_3 , derived from ketone **16** with 91% ee).



Bisacetates 54. For representative acetylation and reduction procedures, see acetylation of tertiary alcohol **25a** to **27c**, and reduction of **27c** to **28**. Alcohols **28** can be acetylated without attempting their purification or separation. Bisacetates **54** can be separated by preparative thin layer chromatography on a 250 μm analytical plate (eluent: 20:1 CH_2Cl_2 :EtOAc x 2) to give bisacetate **54b** as a colorless oil (6.3 mg, 51% yield over three steps) and bisacetate **54a** as a white solid (2.6 mg, 21% yield over three steps). Alternatively, bisacetates **54** can be formed by acylation in parallel. Alcohol **28b** (20.0 mg, 0.042 mmol, 1 equiv) furnished bisacetate **54b** (21.7 mg, 99% yield), while alcohol **28a** (13.9 mg, 0.029 mmol, 1 equiv) provided bisacetate **54a** (12.1 mg, 80% yield). In this case, bisacetates were carried on without further purification. Bisacetate **54a**: R_f 0.41 (1:20 EtOAc: CH_2Cl_2 x 2, visualized with anisaldehyde stain); ^1H NMR (500 MHz, CDCl_3) δ 5.68 (app dt, J = 3.5, 1.6, 1H), 5.43 (dd, J = 10.0, 4.1, 1H),

5.34–5.29 (m, 1H), 4.27 (d, $J = 11.4$, 1H), 3.45 (d, $J = 11.4$, 1H), 2.81 (d, $J = 6.0$, 1H), 2.28–2.21 (m, 3H), 2.02 (d, $J = 5.7$, 6H), 1.93–1.86 (m, 1H), 1.80–1.70 (m, 1H), 1.69 (s, 3H), 1.69–1.65 (m, 2H), 1.65–1.60 (m, 1H), 1.12–0.94 (m, 21H); ^{13}C NMR (125 MHz, CDCl_3) δ 173.8, 171.3, 170.6, 135.9, 128.3, 93.0, 73.7, 68.0, 59.8, 57.6, 51.1, 48.4, 41.5, 28.1 (2C), 21.5, 20.9, 19.1, 18.9, 18.2, 12.1; IR (thin film/ NaCl) 2944, 2867, 1770, 1738, 1732, 1667, 1463, 1433, 1373, 1321, 1297, 1241, 1200, 1172, 1129, 1100, 1080, 1065, 1045, 1025, 965, 947, 918, 901, 883, 792, 748, 730 cm^{-1} ; HRMS (ESI–APCI) m/z : $[\text{M} - \text{OAc}]^+$ Calcd for $\text{C}_{28}\text{H}_{44}\text{O}_7\text{Si}$ 461.2718; Found 461.2740; $[\alpha]_{\text{D}}^{25.8} +72.4^\circ$ (c 0.19, CHCl_3 , derived from ketone **16** with 91% ee). Bisacetate **54b**: R_f 0.56 (1:20 EtOAc: CH_2Cl_2 x 2, visualized with anisaldehyde stain); ^1H NMR (500 MHz, CDCl_3) δ 5.62 (app dt, $J = 3.5$, 1.8, 1H), 5.43 (ddd, $J = 5.9$, 4.9, 3.4, 1H), 5.23–5.18 (m, 1H), 4.26 (d, $J = 11.4$, 1H), 3.46 (d, $J = 11.4$, 1H), 2.71 (d, $J = 6.0$, 1H), 2.28–2.21 (m, 3H), 2.04 (s, 3H), 2.03 (s, 3H), 1.86–1.76 (m, 3H), 1.73–1.63 (m, 1H), 1.69 (s, 3H), 1.63–1.55 (m, 1H), 1.14–0.82 (m, 21H); ^{13}C NMR (125 MHz, CDCl_3) δ 173.8, 171.2, 171.0, 136.1, 128.1, 92.9, 73.7, 68.1, 60.1, 57.3, 51.1, 48.1, 41.6, 28.1, 27.9, 21.5, 20.9, 19.4, 18.9, 18.2, 12.1; IR (film) 1770, 1738, 883 cm^{-1} ; HRMS (ESI–APCI) m/z : $[\text{M} - \text{OAc}]^+$ Calcd for $\text{C}_{28}\text{H}_{44}\text{O}_7\text{Si}$ 461.2718; Found 461.2732; $[\alpha]_{\text{D}}^{26.0} -28.5^\circ$ (c 0.16, CHCl_3 , derived from ketone **16** with 91% ee).




Divinylcyclopropane 29b. Desilylation. A flask charged with THF (4.8 mL, 0.012 M) and bisacetate **54b** (30.0 mg, 0.0576 mmol, 1 equiv) was cooled to 0 °C (ice-water bath) and treated dropwise with TBAF (1 M in THF, 0.09 mL, 0.09 mmol, 1.56 equiv). After 30 m, the yellow solution was diluted with EtOAc (to 25 mL), filtered through SiO_2 (ca. 0.3 mL) and concentrated under reduced pressure to give desilylated compound as an oil (R_f 0.11 (1:1 EtOAc:hexanes, visualized with anisaldehyde stain)).


Oxidation. The crude oil was taken up in CH_2Cl_2 (30 mL), cooled to –20 °C (cryocool) beneath an Ar atmosphere, and treated with crushed Me-IBX (34 mg, 0.12 mmol, 2.0 equiv). After 10 m, the mixture was allowed to gradually warm to 4 °C (cold room). Two additional portions of crushed Me-IBX (34.6 and 29.5 mg, 0.12 and 0.10 mmol, 2.0 and 1.7 equiv) were added to the mixture ca. 6 and ca. 9 hours after the initial addition. On completion of the reaction (ca. 21 h), the mixture was diluted with EtOAc (to 100 mL) and filtered through SiO_2 (ca. 2 mL). The organics were concentrated under reduced pressure to give aldehyde as a cream-colored solid (R_f 0.31 (1:1 EtOAc:hexanes, visualized with anisaldehyde stain)). The solid was placed under an argon atmosphere and quickly dissolved in THF (0.5 mL).

Wittig olefination. A room temperature (ca. 24 °C) flask charged with a mixture of MePPh_3Br (0.230 g, 0.643 mmol, 11.1 equiv) and THF (30 mL, 0.0019 M) beneath an Ar atmosphere was treated dropwise with $n\text{-BuLi}$ (2.5 M in hexanes, 0.40 mL, 1 mmol, 17 equiv). The yellow solution was stirred for 30 min, and then cooled to –65 °C ($i\text{-PrOH}$ / dry ice bath). The yellow solution was treated dropwise with aldehyde in THF (0.5 mL x 2). The solution was allowed to warm to –10 °C over ca. 1 h, at which point it was quenched by addition of acetone (1 mL). The white mixture was rinsed with 1:1 H_2O : brine (ca. 5 mL) and extracted with EtOAc (to 75 mL). The extracts were dried over MgSO_4 , filtered, and concentrated under reduced pressure. The resulting solid was dissolved in EtOAc (10 mL), filtered through SiO_2 (ca. 0.3 mL), and concentrated under reduced pressure. The semi-solid was purified by preparatory thin layer chromatography on a 250 μm plate (1:15 EtOAc: CH_2Cl_2 x 2 eluent) to give divinylcyclopropane **29b** as a colorless oil (16.0 mg, 77% yield). R_f 0.17 (1:15 EtOAc: CH_2Cl_2 , visualized with anisaldehyde stain); ^1H NMR (500 MHz, CDCl_3) δ 6.09 (dd, $J = 17.0$, 10.6, 1H), 5.67–5.62 (m, 1H), 5.44 (t, $J = 6.6$, 1H), 5.30–5.23 (m, 1H), 5.17 (dd, $J = 10.5$, 1.2, 1H), 5.01 (dd, $J = 17.0$, 1.2, 1H), 3.03 (d, $J = 5.9$, 1H), 2.23 (d, $J = 14.1$, 2H), 2.17 (dd, $J = 14.4$, 7.5, 2H), 2.06 (s, 3H), 2.04 (s, 3H), 1.83–1.62 (m, 4H), 1.59 (s, 3H);


^{13}C NMR (125 MHz, CDCl_3) δ 173.4, 171.2, 171.1, 135.4, 128.7, 127.0, 118.7, 92.6, 73.2, 68.2, 58.3, 50.8, 49.8, 40.0, 28.1, 27.7, 21.6, 20.9, 20.0, 19.4; IR (film) 2937, 1760, 1738, 1733, 915 cm^{-1} ; HRMS (ESI-APCI) m/z : $[\text{M}-\text{H}]^-$ Calcd for $\text{C}_{20}\text{H}_{24}\text{O}_6$ 359.1500; Found 359.1513; $[\alpha]_{\text{D}}^{22.8} -48.0^\circ$ (c 0.13, CHCl_3 , derived from ketone **16** with 91% ee).

**32b**

Cycloheptadiene 32b. A flask charged with bisacetate **29b** (7.3 mg, 0.020 mmol, 1 equiv) in THF (5.2 mL, 0.0039 M) at 0 $^\circ\text{C}$ (ice water bath) was treated dropwise with a solution of $\text{LiOH}\cdot\text{H}_2\text{O}$ (9.1 mg, 0.22 mmol, 10.8 equiv) in H_2O (0.91 mL, 0.022 M). After 9 h, the solution was diluted with EtOAc (to 40 mL), filtered through SiO_2 (ca. 0.4 mL) and concentrated under reduced pressure. The semi-solid was purified by preparatory thin layer chromatography on a 250 μm plate (1:3 acetone: CH_2Cl_2 elution) to give cycloheptadiene **32b** (3.1 mg, 55% yield, R_f 0.14) as a white solid. R_f 0.45 (EtOAc , UV/Vis, stained blue in anisaldehyde); ^1H NMR (600 MHz, CD_2Cl_2) δ 5.71 (app td, J = 5.2, 2.7, 1H), 4.82 (dt, J = 9.2, 7.5, 1H), 4.23 (d, J = 9.0, 1H), 3.69 (ddd, J = 8.5, 7.0, 1H), 3.13–3.05 (m, 1H), 2.84–2.75 (m, 1H), 2.66 (t, J = 8.0, 1H), 2.57–2.50 (m, 1H), 2.44 (dd, J = 12.8, 7.1, 1H), 2.32 (dddd, J = 16.3, 9.2, 4.9, 2.2, 1H), 1.91 (dtd, J = 12.8, 5.4, 3.6, 1H), 1.84–1.76 (m, 1H), 1.74 (dd, J = 12.8, 7.9, 1H), 1.60–1.52 (m, 3H), 1.51–1.44 (m, 1H), 1.25 (s, 3H); ^{13}C NMR (125 MHz, CD_2Cl_2 , –lactone carbon, –disubstituted carbon of trisubstituted olefin) δ 158.4, 123.3, 118.9, 78.8, 76.1, 72.0, 49.0, 48.7, 43.7, 32.6, 28.7, 27.1, 26.1, 21.0; IR (film) 3393, 2929, 1719, 1653, 966 cm^{-1} ; HRMS (ESI-APCI) m/z : $[\text{M}+\text{H}]^+$ Calcd for $\text{C}_{16}\text{H}_{20}\text{O}_4$ 277.1434; Found 277.1427; $[\alpha]_{\text{D}}^{28.1} +33.5^\circ$ (c 0.17, CH_2Cl_2 , derived from ketone **16** with 91% ee).


**29a**

Divinylcyclopropane 29a. For a representative procedure, see silyl ether **54b** \rightarrow divinylcyclopropane **29b**. 59% yield over three steps. R_f 0.35 (1:20 acetone: CH_2Cl_2 , visualized with anisaldehyde stain); ^1H NMR (500 MHz, CDCl_3) δ 6.13 (dd, J = 17.1, 10.5, 1H), 5.67 (dt, J = 3.5, 1.7, 1H), 5.42 (dd, J = 7.1, 6.1, 1H), 5.31 (dddd, J = 7.3, 5.5, 3.7, 1.8, 1H), 5.19 (dd, J = 10.5, 1.1, 1H), 5.02 (dd, J = 17.1, 1.1, 1H), 3.06 (d, J = 5.9, 1H), 2.24 (d, J = 14.1, 1H), 2.18 (dd, J = 14.4, 7.5, 1H), 2.14–2.08 (m, 1H), 2.04 (s, 3H), 2.03 (s, 3H), 1.89 (ddd, J = 17.8, 9.6, 5.6, 1H), 1.81–1.74 (m, 1H), 1.65 (dt, J = 12.4, 6.2, 2H), 1.60 (s, 3H), 1.57–1.50 (m, 1H); ^{13}C NMR (125 MHz, CDCl_3) δ 173.4, 171.2, 170.7, 135.3, 129.3, 127.1, 118.5, 92.6, 73.3, 68.4, 58.5, 50.8, 49.7, 40.6, 28.0, 27.7, 21.5, 20.9, 20.0, 19.2; IR (film) 2937, 1759, 1738, 1733, 915 cm^{-1} ; HRMS (ESI-APCI) m/z : $[\text{M}-\text{H}]^-$ Calcd for $\text{C}_{20}\text{H}_{24}\text{O}_6$ 359.1500; Found 359.1488; $[\alpha]_{\text{D}}^{23.7} +30.4^\circ$ (c 0.18, CHCl_3 , derived from ketone **16** with 91% ee).


**32a**

Cycloheptadiene 32a (white solid). For a representative procedure see, divinylcyclopropane **29b** \rightarrow cycloheptadiene **32b**. 51% yield. R_f 0.44 (EtOAc , visualized by UV/Vis or with anisaldehyde stain); ^1H NMR (500 MHz, C_6D_6) δ 5.55 (s, 1H), 4.22 (dd, J = 16.1, 7.8, 1H), 3.50 (dd, J =

12.2, 6.7, 1H), 3.43 (d, $J = 7.7$, 1H), 3.32–3.09 (m, 1H), 2.91 (app s, 1H), 2.56 (dd, $J = 10.0$, 4.6, 1H), 2.44 (ddd, $J = 17.3$, 8.6, 4.2, 1H), 1.99 (dd, $J = 12.6$, 7.3, 1H), 1.76–1.63 (m, 1H), 1.56 (dq, $J = 19.2$, 6.4, 1H), 1.46–1.11 (m, 6H), 0.93 (s, 3H); ^{13}C NMR (125 MHz, C_6D_6) δ 168.7, 157.1, 148.5, 125.4, 119.4, 78.6, 75.7, 71.6, 48.9, 44.8, 43.3, 30.7, 28.3, 25.2, 24.4, 21.6; IR (film) 3369, 2929, 1724, 1654 cm^{-1} ; HRMS (ESI–APCI) m/z $[\text{M}+\text{H}]^+$ Calcd for $\text{C}_{16}\text{H}_{20}\text{O}_4$ 277.1434; Found 277.1445; $[\alpha]_{\text{D}}^{26.6} +100.0^\circ$ (c 0.16, acetone, derived from ketone **16** with 91% ee).



Ketal 55. A flask was charged with a colorless solution of ketone (7.0 mg, 0.015 mmol, 1 equiv), CH_2Cl_2 (2 mL, 0.0073 M), and 1,2-bis(trimethylsilyloxy)ethane (0.10 mL, 0.41 mmol, 28 equiv). The solution was cooled to 0 $^\circ\text{C}$ (ice water bath), and treated dropwise with a 1% v/v solution of TMSOTf in CH_2Cl_2 (150 μL , 0.000008 mmol, 0.0005 equiv). After two days, the solution was treated sequentially with CH_2Cl_2 (0.5 mL, 0.0294 M), 1,2-bis(trimethylsilyloxy)ethane (0.10 mL, 0.41 mmol, 28 equiv) and TMSOTf (150 μL of a solution containing TMSOTf (10 μL) in CH_2Cl_2 (1 mL), 0.000008 mmol, 0.0005 equiv). After an additional 12 h, the reaction was quenched through addition of pyridine (0.3 mL, 0.0037 mmol, 0.25 equiv), and then concentrated under reduced pressure. The resultant yellow oil was purified by thin layer chromatography on a 250 μm plate (12 cm x 20 cm; 1:1 Et_2O :hexanes, then 2:1 Et_2O :pentane elution) to furnish ketal **55** (5.1 mg, 67% yield) as a colorless oil. R_f 0.64 (1:2 EtOAc :PhH, visualized with anisaldehyde stain); ^1H NMR (600 MHz, CDCl_3) δ 5.71 (tt, $J = 3.5$, 1.6, 1H), 5.42 (ddd, $J = 5.6$, 5.6, 2.8, 1H), 4.33 (d, $J = 11.4$, 1H), 3.99–3.88 (m, 3H), 3.88–3.82 (m, 1H), 3.39 (d, $J = 11.5$, 1H), 2.81 (d, $J = 6.0$, 1H), 2.49 (d, $J = 16.8$, 1H), 2.36–2.26 (m, 1H), 2.26–2.18 (m, 3H), 2.02 (s, 3H), 1.77 (dddd, $J = 16.8$, 9.2, 4.8, 1.2, 2H), 1.70–1.60 (m, 4H), 1.56–1.51 (m, 3H), 1.10–0.98 (m, 21H); ^{13}C NMR (125 MHz, CDCl_3) δ 174.1, 171.4, 129.5, 127.9, 107.9, 92.8, 73.9, 64.7, 64.5, 59.6, 57.2, 51.2, 48.5, 42.2, 38.4, 30.6, 24.6, 21.0, 18.9, 18.2, 18.2, 12.1; IR (thin film/ NaCl) 2943, 2892, 1763, 1739, 903, 852 cm^{-1} ; HRMS (ESI–APCI) m/z : $[\text{M}+\text{H}]^+$ Calcd for $\text{C}_{28}\text{H}_{44}\text{O}_7\text{Si}$ 521.2925; Found 521.2929; $[\alpha]_{\text{D}}^{21.0} +16.8^\circ$ (c 0.37, CHCl_3 , derived from ketone **16** with 91% ee).




Divinylcyclopropane 33. A flask was charged with silyl ether **55** (15.0 mg, 0.027 mmol, 1 equiv) in THF (2.3 mL, 0.012 M) at 0 $^\circ\text{C}$ (ice water bath). The colorless solution was treated dropwise with TBAF (1M in THF, 40 μL , 0.040 mmol, 1.5 equiv), generating a yellow solution. After 1.5 h, the solution was diluted with EtOAc (to 40 mL), filtered through SiO_2 (ca. 0.1 mL) and concentrated under reduced pressure to give primary alcohol **56** as a yellow oil. This yellow oil had already been characterized analytically as a byproduct in the ketalization reaction. R_f 0.07 (1:1 EtOAc :hexanes, visualized with anisaldehyde stain); ^1H NMR (500 MHz, CDCl_3) δ 5.67 (dd, $J = 5.9$, 3.3, 1H), 5.40 (td, $J = 6.7$, 1.6, 1H), 4.02 (dd, $J = 6.1$, 2.2, 1H), 4.01 (d, $J = 6.1$, 1H), 3.94 (d, $J = 6.1$, 1H), 3.92 (dd, $J = 6.1$, 1.3, 1H), 3.85 (d, $J = 12.9$, 1H), 3.82 (d, $J = 13.0$, 1H), 2.95 (d, $J = 6.0$, 1H), 2.75–2.68 (m, 1H), 2.40–2.32 (m, 1H), 2.27–2.18 (m, 3H), 2.02 (s, 3H), 1.97 (app d, $J = 16.7$, 1H), 1.78 (dd, $J = 9.3$, 6.6, 1H), 1.83–1.73 (m, 2H), 1.69 (s, 3H); ^{13}C NMR (125 MHz, CDCl_3) δ 173.9, 171.3, 128.1, 127.1, 108.3, 92.7, 73.5, 64.8, 64.6, 58.6, 57.8, 50.9, 47.8, 41.0, 38.8,

30.0, 24.6, 20.9, 18.7; IR (thin film/NaCl) 2932, 1738 cm^{-1} ; HRMS (ESI-APCI) m/z : $[\text{M}+\text{H}]^+$ Calcd for $\text{C}_{19}\text{H}_{24}\text{O}_7$ 365.1595; Found 365.1598; $[\alpha]_{\text{D}}^{22.5} +36.7^\circ$ (c 0.13, CHCl_3 , derived from ketone **16** with 91% ee).

The crude alcohol **56** was thrice concentrated from PhH (0.15 mL each), and backfilled with Ar(g). The diol **56** was diluted with CH_2Cl_2 (16 mL, 0.0017 M), cooled to -15°C (cryocool temperature), and treated with crushed Me-IBX (15 mg, 0.051 mmol, 1.9 equiv). After 5 min, the solution was allowed to warm gradually to 4°C (cold room temperature). After 18, 24, 30, 45, and 55 h, additional portions were added of crushed Me-IBX (67.4 mg total, 0.23 mmol, 8.5 equiv). After 65 h, the reaction was diluted with EtOAc (to ca. 25 mL), filtered through SiO_2 (ca. 0.2 mL), and concentrated under reduced pressure to give crude aldehyde **57**.


A flask was charged with a mixture of Ph_3PMeBr (0.129 g, 0.36 mmol, 13 equiv) in THF (9 mL, 0.003 M) at room temperature (ca. 24°C). The mixture was treated dropwise with $n\text{-BuLi}$ (2.5 M in hexanes, 0.13 mL, 0.325 mmol, 12.0 equiv), generating an orange solution. After 55 min, the orange solution had been cooled to -78°C (acetone / dry ice) and was treated dropwise with crude aldehyde **57** in THF (0.3 mL x 2). The solution was allowed to gradually warm to -10°C (over 1.33 h), at which point it was treated with acetone (0.3 mL). The reaction was rinsed with 1:1 brine: H_2O (ca. 1 mL), and extracted with EtOAc (to ca. 50 mL). Extracts were dried over MgSO_4 , filtered and concentrated under reduced pressure. The mixture was purified by thin layer chromatography on 250 μm plates (10 cm x 20 cm; EtOAc x 2 elution) to furnish the desired divinylcyclopropane (**33**, R_f 0.79, 7.9 mg including ca. 5% other, 91% yield of slightly impure material) as an oil. ^1H NMR (500 MHz, CDCl_3) δ 6.09 (ddd, $J = 17.1, 10.5, 0.5$, 1H), 5.74 (tt, $J = 3.5, 1.7$, 1H), 5.40 (dd, $J = 6.9, 5.9$, 1H), 5.17 (dd, $J = 10.5, 1.1$, 1H), 5.03 (dd, $J = 17.1, 1.1$, 1H), 4.02–3.91 (m, 4H), 3.04 (d, $J = 5.9$, 1H), 2.38–2.30 (m, 2H), 2.24 (app dd, $J = 14.3, 0.6$, 1H), 2.26–2.19 (m, 1H), 2.17 (dd, $J = 14.4, 7.6$, 1H), 2.03 (s, 3H), 1.95–1.90 (m, 1H), 1.75 (ddd, $J = 14.0, 6.9, 3.9$, 1H), 1.64–1.59 (m, 1H), 1.59 (s, 3H); ^{13}C NMR (125 MHz, CDCl_3 , $-\text{CO}_2\text{R}$) δ 171.4, 130.1, 127.8, 127.0, 118.5, 107.9, 92.4, 73.4, 64.6, 64.5, 58.1, 50.6, 49.8, 41.0, 37.6, 30.7, 24.5, 21.0, 20.0; IR (thin film/NaCl) 2932, 1759, 1738, 958, 851 cm^{-1} ; HRMS (ESI-APCI) m/z : $[\text{M}+\text{Na}]^+$ Calcd for $\text{C}_{20}\text{H}_{24}\text{O}_6$ 383.1465; Found 383.1466; $[\alpha]_{\text{D}}^{20.4} -11.2^\circ$ (c 0.17, CHCl_3 , derived from ketone **16** with 91% ee).




Cycloheptadiene 35. A flask was charged with acetate **33** (2.0 mg, 0.0055 mmol, 1 equiv) in THF (2.0 mL, 0.0028 M) at -78°C (acetone, dry ice bath). The colorless solution was treated dropwise with LiEt_3BH (0.5 M solution in THF, 24 μL , 2.2 equiv). After 10 min, the solution was allowed to warm to -20°C (cryocool temperature). After an additional 6 h, the reaction was treated with H_2O (0.02 mL), allowed to warm to room temperature (ca. 24°C), diluted with EtOAc (to 10 mL), filtered through SiO_2 (ca. 0.2 mL), and concentrated under reduced pressure. The product was combined with product from two equivalent reactions that had been run with acetate **33** (0.4 and 1.0 mg, respectively). The reactions were purified by thin layer chromatography on a 250 μm plate (10 cm x 20 cm; EtOAc elution) to furnish the desired cycloheptadiene (**35**, R_f 0.34, slightly impure material). The material was further purified by thin layer chromatography on a 250 μm plate (10 cm x 20 cm; 1:5 acetone: CH_2Cl_2 elution) to furnish the desired cycloheptadiene (**35**, R_f 0.29, 1.8 mg, 68% yield). R_f 0.56 (EtOAc, UV/Vis, visualized with anisaldehyde stain); ^1H NMR (600 MHz, C_6D_6) δ 5.37 (ddd, $J = 2.8, 5.0, 5.0$ Hz, 1H), 4.17 (ddd, $J = 7.1, 8.0, 9.2$ Hz, 1H), 3.71 (ddd, $J = 6.2, 6.8, 6.8$ Hz, 1H), 3.68 (ddd, $J = 5.1, 6.9, 6.9$ Hz, 1H), 3.57 (br d, $J = 15$ Hz, 1H), 3.52 (ddd, $J = 4.8, 6.9, 6.9$ Hz, 1H), 3.48 (ddd, $J = 5.9, 6.9, 6.9$ Hz, 1H), 3.32 (br d, $J = 9$ Hz, 1H), 2.21 (dddd, $J = 4.5, 4.5, 8.9, 8.9$ Hz, 1H), 1.96 (dd, $J = 7.1, 12.8$ Hz, 1H), 1.93 (dddd, $J = 3.6, 3.6, 5.4, 16.7$ Hz, 1H), 1.89 (dddd, $J = 1.0, 4.3,$

6.0, 13.7 Hz, 1H), 1.73 (ddd, $J = 4.9, 10.8, 13.9$ Hz, 1H), 1.67 (dddd, $J = 2.4, 4.8, 9.5, 16.7$, 1H), 1.43 (dddd, $J = 5.4, 5.4, 5.4, 13.6$ Hz, 1H), 1.37 (dddd, $J = 4.2, 9.0, 10.6, 13.5$ Hz, 1H), 1.34 (dd, $J = 7.9, 12.8$ Hz, 1H), 0.89 (d, $J = 0.9$ Hz, 1H); ^{13}C NMR (125 MHz, CDCl_3) δ 169.6, 156.0, 149.9, 124.5, 119.1, 109.8, 78.8, 76.3, 65.0, 64.8, 48.9, 43.6, 39.6, 35.6, 34.3, 30.8, 28.5, 28.5; IR (thin film/ NaCl) 3446, 2927, 1738, 1647, 955 cm^{-1} ; HRMS (ESI-APCI) m/z : $[\text{M}+\text{H}]^+$ Calcd for $\text{C}_{18}\text{H}_{22}\text{O}_5$ 319.1540; Found 319.1530; $[\alpha]_{\text{D}}^{23.8} +23.3^\circ$ (c 0.18, CH_2Cl_2 , derived from ketone **16** with 91% ee).

Dehydration product **58**. R_f 0.68 (1:5 acetone:PhH; visualized with anisaldehyde stain); ^1H NMR (600 MHz, CD_2Cl_2) δ 5.88 (s, 1H), 5.58 (dt, $J = 2.2, 6.0$, 1H), 5.36–5.33 (m, 1H), 4.32–4.30 (m, 1H) 3.96–3.91 (m, 2H), 3.90–3.83 (m, 2H), 2.87 (ddd, $J = 16.1, 5.0, 5.0$, 1H), 2.68 (m, 1H), 2.47–2.42 (m, 1H), 2.17 (dddd, $J = 15.9, 6.5, 6.5, 1.1$, 1H), 1.85–1.73 (m, 6H), 1.63–1.71 (m, 1H); ^{13}C NMR (125 MHz, CDCl_3) δ 170.1, 154.0, 150.8, 145.7, 130.0, 123.1, 117.0, 109.3, 81.6, 64.6, 64.3, 43.6, 40.1, 36.0, 34.8, 30.8, 29.1, 12.1; IR (thin film/ NaCl) 3456 (br), 2934, 1738, 1733, 1645 cm^{-1} ; HRMS (ESI-APCI) m/z : $[\text{M}+\text{H}]^+$ Calcd for $\text{C}_{18}\text{H}_{20}\text{O}_4$ 301.1434; Found 301.1439; $[\alpha]_{\text{D}}^{25.8} -25.5^\circ$ (c 0.22, CH_2Cl_2 , derived from ketone **16** with 91% ee).




Epoxide 36. A flask was charged with cycloheptadiene **35** (0.9 mg, 0.0028 mmol, 1 equiv) and $\text{VO}(\text{acac})_2$ (0.2 mg, 0.00075 mmol, 0.27 equiv) in PhH (0.4 mL). The pale green solution was treated with TBHP, 5.5 M in decane (ca. 10 μL , 1 drop), and then turned burgundy in color. After 12 minutes, the solution was treated with saturated aq Na_2SO_3 (0.04 mL), diluted with EtOAc (to 10 mL), filtered through SiO_2 (ca. 0.2 mL), and concentrated under reduced pressure. This compound was combined with the crude product from an analogous reaction carried out with cycloheptadiene **35** (1.8 mg, 0.0056 mmol). The reactions were purified by thin layer chromatography on a 250 μm plate (10 cm x 20 cm; 1:5 acetone: CH_2Cl_2 elution) to furnish epoxide **36** (1.6 mg, 56% yield, R_f 0.34) as a colorless oil. R_f 0.40 (EtOAc, visualized by UV/Vis, or with anisaldehyde stain); ^1H NMR (500 MHz, CD_2Cl_2) δ 4.83 (ddd, $J = 8.7, 7.5, 6.6$, 1H), 4.03–3.85 (m, 5H), 3.34 (t, $J = 3.0$, 1H), 3.28 (d, $J = 15.0$, 1H), 3.07 (d, $J = 15.1$, 1H), 2.70–2.62 (m, 1H), 2.52 (dd, $J = 13.5, 7.5$, 1H), 2.27 (s, 1H), 2.07–2.03 (m, 1H), 2.05 (d, $J = 2.6$, 1H), 1.90–1.77 (m, 3H), 1.73–1.67 (m, 2H), 1.29 (s, 3H); ^{13}C NMR (125 MHz, CDCl_3) δ 169.3, 156.4, 120.0, 109.5, 75.8, 74.8, 68.7, 65.1, 64.8, 55.9, 47.6, 42.8, 39.4, 35.3, 34.0, 29.2, 29.1, 23.7; IR (thin film/ NaCl) 3473, 1739, 1662, 1268, 1120 cm^{-1} ; HRMS (ESI-APCI) m/z : $[\text{M}+\text{H}]^+$ Calcd for $\text{C}_{18}\text{H}_{22}\text{O}_6$ 335.1489; Found 335.1483; $[\alpha]_{\text{D}}^{22.1} +28.6^\circ$ (c 0.16, CH_2Cl_2 , derived from ketone **16** with 91% ee).




Benzoyl ester 40. A dilute solution of enone **16** in Et_2O was concentrated at 150 torr to 0.645 g of a 30% (w/w) solution as determined by ^1H NMR (0.19 g enone, 1.04 mmol) and dissolved in THF (8 mL, 0.13 M). [The volatility of the starting enone precluded preparation of fully concentrated samples without significant loss of material.] The solution was cooled in a dry ice/acetone bath and to it was added DIBALH (2.3 mL, 1 M toluene, 2.3 mmol). The starting material was consumed within 10 minutes according to TLC. The solution was warmed to room temperature and quenched with saturated aq sodium potassium tartrate (20 mL) and saturated aq NH_4Cl (20 mL) and stirred vigorously for 2 h. The solution was extracted with Et_2O (6 x 50 mL) and the combined organic phases dried with Na_2SO_4 , filtered and concentrated at 150 torr. The crude material was

purified by flash chromatography (1:1 pentane:Et₂O), concentrated at 150 torr and dissolved immediately in CH₂Cl₂ (17 mL, 0.06 M) for use in the next step. The CH₂Cl₂ solution of allylic alcohol was cooled in an ice water bath and to it was added DMAP (0.27 g, 2.2 mmol) and Et₃N (1.3 mL, 9.38 mmol) followed by benzoic anhydride (0.472 g, 2.1 mmol). The solution was stirred for 4 hr with warming to ambient temperature, then quenched with saturated aq NH₄Cl (20 mL) and extracted with EtOAc (3 x 75 mL). The combined organic phases were dried with Na₂SO₄, filtered and concentrated under reduced pressure. The crude material was purified by flash chromatography (2:1 hexanes:CH₂Cl₂ with 2% Et₃N) to yield benzoate ester **40** (0.294 g, 98% yield, 2 steps) as a colorless oil. *R*_f 0.66 (1:1 EtOAc:hexanes, visualized by UV/Vis, or with anisaldehyde stain); ¹H NMR (300 MHz, CDCl₃) δ 8.08–8.03 (m, 2H), 7.60–7.53 (m, 1H), 7.48–7.41 (m, 2H), 5.83–5.75 (m, 1H), 5.69–5.66 (m, 1H), 4.77–4.64 (m, 2H), 2.69 (dd, *J* = 6.9, 12.8, 1H), 2.13 (dd, *J* = 6.9, 12.9, 1H), 1.50 (s, 3H), 1.50 (s, 3H), 1.42 (s, 3H); ¹³C NMR (75 MHz, CDCl₃) δ 166.6, 146.5, 133.2, 130.4, 129.8, 128.5, 122.0, 100.3, 81.0, 76.6, 60.4, 49.4, 30.0, 28.6, 25.9; IR (thin film/NaCl) 2990, 1715, 1602, 1268, 1158 cm⁻¹; HRMS (EI) *m/z*: [M+H]⁺ Calcd for C₁₇H₂₀O₄ 288.1362; Found 288.1366; [α]_D^{24.7} +74.2° (*c* 1.0, CHCl₃, derived from ketone with 91% ee).




Allylic benzoyl ester 41. To a MeOH (3 mL, 0.037 M) solution of acetone **40** (0.031 g, 0.11 mmol), cooled in an ice water bath, was added trimethyl orthoformate (0.16 mL, 0.96 mmol) followed by a MeOH (1.5 mL) solution of fumaric acid (0.032 g, 0.28 mmol). The solution was allowed to gradually warm to ambient temperature and stirred an additional 10 hr. The solution was quenched with saturated aq NaHCO₃ (10 mL) and extracted with EtOAc (3x20 mL). The combined organic phases were dried with MgSO₄, filtered and concentrated in vacuo to yield crude diol **59** (0.028 g, >99% yield) as a colorless oil, which was used immediately.


Crude diol **59** (0.028, 0.11 mmol) was dissolved in CH₂Cl₂ (2 mL, 0.055 M) and cooled in an ice water bath. To the solution was added DMP (0.11 g, 0.27 mmol). CAUTION! DMP is a HEAT- and SHOCK-SENSITIVE COMPOUND, showing exotherms when heated (>130 °C). All operations should be carried out behind a blast shield. The solution was stirred for an additional two hr (with warming to ambient temperature) and quenched with saturated aq NaHCO₃ (3 mL) and saturated aq Na₂S₂O₃ (6 mL). The two phases were stirred vigorously for 5 min and then extracted with EtOAc (6 x 25 mL). The combined organic phases were dried with MgSO₄, filtered and concentrated in vacuo. The crude aldehyde was used immediately as is. A solution of methylenetriphenylphosphorane was prepared in THF (8 mL, 0.06 M) from triphenylphosphonium bromide (0.17 g, 0.48 mmol) and potassium *t*-butoxide (0.053 g, 0.47 mmol). The yellow solution was cooled in an ice water bath after stirring for 30 min at ambient temperature, and to it added crude aldehyde dropwise as a solution in THF (4 mL). The reaction was complete within 10 min according to TLC. The brown solution was quenched with H₂O (20 mL) and extracted with CH₂Cl₂ (3 x 30 mL). The combined organic phases were dried with Na₂SO₄, filtered and concentrated in vacuo. The crude material was purified by flash chromatography (1:20 EtOAc:hexanes elution) to yield allylic benzoyl ester **41** as a pale yellow oil (0.019 g, 69% yield, 3 steps). *R*_f 0.62 (1:2 EtOAc:hexanes, visualized with anisaldehyde stain); ¹H NMR (300 MHz, CDCl₃) δ 8.03 (dt, *J* = 8.4, 1.5, 2H), 7.56 (app t, *J* = 7.4, 1H), 7.44 (app t, *J* = 7.7, 2H), 6.36 (dd, *J* = 17.9, 11.3, 1H), 5.94 (d, *J* = 2.2, 1H), 5.84–5.77 (m, 1H), 5.75 (ddd, *J* = 7.0, 4.7, 2.0, 1H), 5.34 (dd, *J* = 11.3, 1.6, 1H), 2.75 (dd, *J* = 14.1, 7.3, 1H), 2.16 (dd, *J* = 14.1, 4.7, 1H), 1.91 (s, 1H), 1.50 (s, 3H); ¹³C NMR (75 MHz, CDCl₃, -CO₂Ph) δ 151.3, 133.2, 130.4, 129.8, 129.2, 128.6, 127.2, 119.4, 81.3, 76.1, 49.5, 27.0; IR (thin film/NaCl) 3436 (br), 3063, 2974, 2931, 2865, 1715, 1602, 924, 859, 713 cm⁻¹; HRMS (EI) *m/z*: [M+]⁺ Calcd for C₁₅H₁₆O₃ 244.1100; Found 244.1101; [α]_D^{27.3} +150.0° (*c* 0.13, CHCl₃, derived from ketone **16** with 91% ee).

**42**

Alcohol 42. To a MeOH (5 mL, 0.05 M) solution of allylic benzoyl ester **41** (0.062 g, 0.25 mmol) was added a MeOH solution of sodium hydroxide (0.85 mL, 0.64 M, 0.54 mmol) dropwise. The solution was stirred at ambient temperature for 3 hours, quenched with H₂O (10 mL) and extracted with EtOAc (6 x 25 mL). The combined organic phases were dried with Na₂SO₄, filtered and concentrated in vacuo. The crude material was purified by flash chromatography (4:1 CH₂Cl₂:EtOAc → 1:1 hexanes/EtOAc elution) to yield alcohol **42** as a white solid (0.030 g, 84% yield). *R_f* 0.13 (1:5 EtOAc:CH₂Cl₂, visualized with anisaldehyde stain); ¹H NMR (300 MHz, CDCl₃) δ 6.33 (ddt, *J* = 17.7, 11.2, 0.7, 1H), 5.82 (d, *J* = 2.1, 1H), 5.75 (dd, *J* = 17.9, 1.7, 1H), 5.29 (dd, *J* = 11.2, 1.7, 1H), 4.68 (dd, *J* = 11.5, 5.1, 1H), 2.57 (dd, *J* = 13.6, 6.8, 1H), 1.88 (app dd, *J* = 13.6, 4.8, 2H), 1.71 (d, *J* = 6.6, 1H), 1.43 (s, 3H); ¹³C NMR (75 MHz, CDCl₃) δ 149.5, 131.3, 129.5, 118.6, 81.3, 73.0, 53.1, 26.8; IR (thin film/NaCl) 3287, 3252, 2968, 2930, 2873, 926 cm⁻¹; HRMS (EI) *m/z*: [*M*+]⁺ Calcd for C₈H₁₂O₂, 140.0837; Found 140.0859; [*α*]_D^{24.2} +100.0° (*c* 0.085, MeOH, derived from ketone **16** with 91% ee).


**60**

Ester 60. A CH₂Cl₂ (15 mL, 0.026 M) solution of alcohol **42** (0.054 g, 0.39 mmol) with carboxylic acid **10** (0.16 g, 1.1 mmol) was cooled in an ice water bath. EDC•HCl (0.21 g, 1.1 mmol) was added followed by DMAP (0.011 g, 0.09 mmol). The deep yellow solution was allowed to warm to ambient temperature and stirred an additional 1 hour. The solution was quenched with aq HCl (20 mL, 0.12 M) and extracted with EtOAc (2 x 50 mL). The combined organic phases were washed with aq HCl (20 mL, 0.12 M), aq K₂CO₃ (2 x 20 mL 5 % w/v), brine (20 mL), and saturated aq NH₄Cl (20 mL). The organic phase was dried with Na₂SO₄, filtered and concentrated to provide crude ester **60** (0.125 g, 117% crude yield) as an orange oil. The crude ester was used without further purification. An analytical sample was obtained at another point, upon purification by preparatory thin layer chromatography with Et₂O x 2 as the eluent. *R_f* 0.21 (1:5 EtOAc:CH₂Cl₂, visualized with anisaldehyde stain); ¹H NMR (300 MHz, CDCl₃) δ 6.30 (dd, *J* = 17.9, 11.3, 1H), 5.94–5.91 (m, 1H), 5.80 (d, *J* = 2.2, 1H), 5.83–5.75 (m, 1H), 5.53–5.47 (m, 1H), 5.36 (dq, *J* = 1.6, 0.5, 1H), 5.32 (dq, *J* = 1.7, 0.6, 1H), 3.23 (s, 2H), 2.64 (dd, *J* = 14.1, 7.3, 1H), 2.42–2.38 (m, 4H), 2.06–1.93 (m, 3H), 1.84 (s, 1H), 1.46 (s, 3H); ¹³C NMR (75 MHz, CDCl₃) δ 199.6, 169.4, 157.2, 151.7, 129.1, 129.0, 126.5, 119.7, 81.1, 76.5, 49.2, 43.7, 37.3, 29.8, 27.0, 22.7; IR (thin film/NaCl) 3417, 2925, 2853, 1732, 1661, 959, 887 cm⁻¹; HRMS (EI) *m/z*: [*M*]⁺ Calcd for C₁₆H₂₀O₄ 276.1362; Found 276.1363; [*α*]_D^{23.2} +45.6° (*c* 0.20, CHCl₃, derived from ketone **16** with 91% ee).


**39**

Diazoester 39. To a CH₃CN (42 mL, 0.01 M) solution of ester **60** (≤0.125 g, 0.385 mmol) was added TsN₃ (0.670 g, 3.4 mmol) dropwise using a flame-dulled pipette. CAUTION!!! TsN₃ is SHOCK SENSITIVE AND POTENTIALLY EXPLOSIVE. The flask was equipped with an argon balloon and cooled in an ice water bath. Et₃N (0.8 mL, 5.8 mmol) was added dropwise causing the solution to become deep orange in color. The


solution was allowed to warm to ambient temperature and stirred an additional 11 hr. The solution was concentrated in vacuo and purified using flash chromatography (4:1 CH₂Cl₂:EtOAc) to yield diazoester **39** (0.746 g, 64% yield, 2 steps) as a bright yellow oil. *R_f* 0.18 (1:5 EtOAc:CH₂Cl₂, UV/Vis, visualized with anisaldehyde stain); ¹H NMR (500 MHz, C₆D₆) 6.75 (t, *J* = 1.3, 1H), 6.15 (dd, *J* = 17.8, 11.3, 1H), 5.72 (dd, *J* = 17.8, 1.8, 1H), 5.60 (d, *J* = 2.0, 1H), 5.46–5.24 (m, 1H), 5.11 (dd, *J* = 11.3, 1.8, 1H), 2.32 (dd, *J* = 13.7, 7.3, 1H), 2.16–2.07 (m, 2H), 1.82 (dd, *J* = 13.7, 5.2, 1H), 1.64 (m, 2H), 1.39 (dt, *J* = 12.4, 6.2, 2H), 1.37–1.20 (m, 1H), 1.17 (s, 3H); ¹³C NMR (125 MHz, C₆D₆, –CN₂) δ 195.0, 162.4, 151.7, 144.6, 129.3, 126.2, 121.6, 119.0, 80.3, 76.7, 49.2, 36.8, 26.9, 26.0, 22.1; IR (thin film/NaCl) 3407, 2930, 2103, 1707, 1644, 992 cm^{–1}; HRMS (EI) *m/z*: [M+H]⁺ Calcd for C₁₆H₁₈N₂O₄ 303.1345; Found 303.1339; [α]_D^{20.4} +67.0° (*c* 0.24, CHCl₃, derived from ketone **16** with 91% ee).




Cycloheptatriene 44. To a CH₂Cl₂ (2.4 mL) solution of Cu(tbs)₂ (0.035 g, 0.084 mmol, 3.2 eq.) in a moisture-free, oxygen-free glovebox was added diazoester **39** (0.008 g, 0.026 mmol) as a solution in CH₂Cl₂ (5.6 mL). The deep maroon solution was stirred for 5 days at 29 °C and removed from the glove box. DBN (0.01 mL, 0.08 mmol) was added and the solution stirred for 2 hr. The solution was diluted with EtOAc, filtered over a sort silica plug, concentrated in vacuo and purified by preparatory thin layer chromatography (2:1 EtOAc/benzene) to yield cycloheptatriene **44** (0.0022 g, 31% yield) as a pale yellow solid. *R_f* 0.50 (1:2 EtOAc:PhH, UV/Vis, visualized with anisaldehyde stain); ¹H NMR (500 MHz, CDCl₃) δ 7.72 (d, *J* = 6.4, 1H), 6.54 (dd, *J* = 6.4, 2.2, 1H), 5.17 (ddd, *J* = 7.9, 6.6, 4.5, 1H), 3.59 (m, 1H), 3.21 (d, *J* = 7.9, 1H), 2.94 (dt, *J* = 17.5, 5.7, 1H), 2.61 (m, 2H), 2.52 (dd, *J* = 14.5, 6.6, 1H), 2.27 (dd, *J* = 14.5, 4.4, 1H), 2.03–1.92 (m, 2H), 1.77 (s, 1H), 1.54 (s, 3H); ¹³C NMR (125 MHz, CDCl₃) δ 199.9, 168.4, 156.3, 144.8, 139.8, 139.2, 117.3, 115.3, 79.2, 78.2, 48.3, 46.4, 39.6, 28.1, 26.5, 20.9; IR (thin film/NaCl) 3416 (br), 2956, 2925, 2853, 1739, 1734, 802 cm^{–1}; HRMS (TOF) *m/z*: [M+H]⁺ Calcd for C₁₆H₁₇O₄ 273.1121; Found 273.1131; [α]_D²⁵ –86.9 (*c* 0.43, CH₂Cl₂, derived from ketone **16** with 91% ee).




Epoxide 45. In a flame-dried, argon purged RBF was added THF (4 mL) and *t*-BuOOH (0.07 mL, 5–6 M / decane, 0.35–0.42 mmol). The solution was cooled in a dry ice/acetone bath and to it added *n*-BuLi (0.1 mL, 2.3 M / hexanes, 0.25 mmol). The solution was allowed to warm to room temperature. A separate RBF was prepared with triene **44** (2.1 mg, 0.0077 mmol) in THF (0.8 mL) and cooled in a dry ice/acetone bath. The *t*-BuOOLi/THF solution (0.25 mL, 0.058 M, 0.015 mmol) was added dropwise. The reaction was allowed to warm to room temperature, then quenched with Na₂S₂O₃ (aq) and stirred vigorously for 10 min. The solution was extracted with EtOAc (3 x 5 mL), dried with Na₂SO₄ and filtered. Purification by thin-layer chromatography with 2:1 EtOAc:PhH eluent yielded the epoxide (1.4 mg, 64% yield). ¹H NMR (500 MHz, CDCl₃): δ 5.95 (dd, *J* = 4.3, 2.6 Hz, 1H), 4.99 (ddd, *J* = 8.6, 6.5, 5.7 Hz, 1H), 4.59 (dd, *J* = 8.6, 2.5 Hz, 1H), 3.90 (dd, *J* = 4.2, 0.5 Hz, 1H), 3.62 (dm, *J* = 17.2, 1H), 3.14 (dm, *J* = 17.2 Hz, 1H), 2.73 (dt, *J* = 17.8, 6.6 Hz, 1H), 2.64 (dt, *J* = 17.8, 6.6 Hz, 1H), 2.40 (dd, *J* = 13.9, 6.5 Hz, 1H), 2.06–1.97 (m, 2H), 1.99 (dd, *J* = 13.9, 5.7 Hz, 1H), 1.68 (s, 1H), 1.41 (s, 3H); ¹³C NMR (125 MHz, CDCl₃) δ 202.9, 169.0, 155.2, 146.9, 122.4, 113.5, 79.3, 78.1, 62.7, 62.5, 47.3, 43.2, 40.6, 28.5, 28.4, 20.0; IR (Neat film, NaCl) 3446 (br), 2965, 2928, 2872, 1732, 1738, 1262, 806, 735 cm^{–1}; HRMS (TOF) *m/z*: [M+H]⁺ Calcd for C₁₆H₁₇O₅ 289.1071; Found 289.1085; [α]_D²⁵ = +35.1 (*c* 0.14, CH₂Cl₂)



Epoxide 46. To a CH_2Cl_2 (2 mL) solution of $\text{Cu}(\text{tbs})_2$ (0.0062 g, 0.015 mmol, 0.4 equiv) in a moisture-free, oxygen-free glove box was added diazoester **39** (0.0108 g, 0.036 mmol) as a solution in CH_2Cl_2 (2.5 mL). The deep maroon solution was stirred for 5 days at 30–31 °C and then concentrated in vacuo (predominantly diene, R_f 0.36 in 2:1 EtOAc:benzene, visualized by UV/Vis, or with anisaldehyde stain). PhH (3 mL) was added and the solution removed from the glovebox. $\text{VO}(\text{acac})_2$ (0.010 g, 0.038 mmol) followed by *t*-BuOOH (0.05 mL, 5.5 M decane, 0.275 mmol). Upon addition of *t*-BuOOH the solution turns an even deeper maroon and over the course of 20 minutes, after which time starting material has been consumed by TLC, the solution gradually becomes a light tan color. The solution was diluted with EtOAc, filtered over a short silica plug, concentrated in vacuo and purified by preparatory TLC (2:1 EtOAc:PhH) to yield epoxide **46** (3.3 mg, 32% yield) as an off-white solid. R_f 0.23 (1:2 EtOAc:PhH, UV/Vis, visualized with KMnO_4 stain); ^1H NMR (600 MHz, CDCl_3) δ 4.84 (ddd, J = 6.0, 4.6, 1.6, 1H), 3.82 (dd, J = 18.4, 6.1, 1H), 3.52 (d, J = 6, 1H), 3.41 (dd, J = 6.3, 4.6, 1H), 3.38 (d, J = 6.4, 1H), 3.26–3.18 (m, 1H), 2.54 (ddd, J = 16.9, 7.5, 4.8, 1H), 2.52 (s, 1H), 2.46–2.36 (m, 3H), 2.24–2.14 (m, 1H), 2.12–2.04 (m, 1H), 1.97–1.90 (m, 1H), 1.37 (s, 3H); ^{13}C NMR (125 MHz, CDCl_3) δ 198.7, 172.2, 150.0, 129.9, 80.0, 75.0, 70.3, 54.3, 51.0, 46.1, 43.7, 37.9, 32.7, 26.9, 22.0, 21.8; IR (thin film/NaCl) 3472 (br), 2960, 2933, 2857, 1766, 1661, 1266, 919, 785 cm^{-1} ; HRMS (ESI–APCI) m/z : $[\text{M}+\text{H}]^+$ Calcd for $\text{C}_{16}\text{H}_{18}\text{O}_5$ 291.1227; Found 291.1228; $[\text{M}-\text{H}]^-$: Calcd for $\text{C}_{16}\text{H}_{18}\text{O}_5$ 289.1081; Found 289.1086; $[\alpha]_D^{25.4}$ +86.2 (c 0.16, CH_2Cl_2 , derived from ketone **16** with 91% ee).




Epoxide 48. To CH_2Cl_2 (0.6 mL) solution of epoxide **46** (2.9 mg, 0.01 mmol) was added DMAP (3.3 mg, 0.027 mmol) followed by Ac_2O (0.005 mL, 0.053 mmol), added in two equal batches. The solution was purified directly by thin-layer chromatography (EtOAc) to yield epoxide **48** (1.5 mg, 45% yield) as a white solid. ^1H NMR (500 MHz, CDCl_3) δ 4.83 (ddd, J = 9.4, 7.6 Hz, 1H), 3.84 (d, J = 9.4 Hz, 1H), 3.51–3.45 (m, 1H), 3.47 (d, J = 6.1 Hz, 1H), 3.29 (dd, J = 16.5, 6.3 Hz, 1H), 2.94–2.87 (m, 1H), 2.60 (dd, J = 13.0, 7.1 Hz, 1H), 2.49 (dt, J = 16.5, 2.8 Hz, 1H), 2.40–2.37 (m, 2H), 2.34 (s, OH), 2.25 (s, 3H), 1.95 (dd, J = 13.0, 7.9 Hz, 1H), 1.89–1.82 (m, 1H), 1.78–1.71 (m, 1H), 1.31 (s, 3H); ^{13}C NMR (125 MHz, CDCl_3) δ 170.4, 168.2, 156.7, 152.5, 122.6, 115.0, 74.0, 73.3, 70.9, 53.1, 48.6, 42.2, 29.3, 25.7, 23.8, 23.6, 21.3, 21.1; IR (neat film, NaCl) 3491 (br), 2972, 2935, 1757, 1732, 1274, 934, 788 cm^{-1} ; HRMS (TOF) m/z : $[\text{M}+\text{H}]^+$ Calcd for $\text{C}_{18}\text{H}_{21}\text{O}_6$ 333.1333; Found 333.1333; $[\alpha]_D^{25}$ = +265.6 (c 0.15, CH_2Cl_2).




Cycloheptadiene 47. To a CH_2Cl_2 (1 mL) solution of $\text{Cu}(\text{tbs})_2$ (0.0055 g, 0.013 mmol) in a moisture-free, oxygen-free glove box was added diazoester **39** (0.0087 g, 0.029 mmol) as a solution in CH_2Cl_2 (1.9 mL). The deep maroon solution was stirred for 2 days at 35–36 °C. The solution

was then removed from the glove box and to it added acetic anhydride (0.02 mL, 0.21 mmol) followed by DMAP (0.0056 g, 0.046 mmol) as a solution in 0.4 mL. The solution turned to a light green solution and was loaded directly onto a single analytical TLC plate and purified using 2:1 EtOAc:PhH eluent to provide enol acetate **47** (0.004 g, 0.013 mmol, 44% yield) as a yellow oil. ^1H NMR (500 MHz, CDCl_3) δ 5.82 (ddd, J = 8, 6, 2.7 Hz, 1H), 4.85 (ddd, J = 8.5, 6.7, 1H), 4.35 (d, J = 9 Hz, 1H), 3.18 (dd, J = 14.8, 8 Hz, 1H), 3.15–3.04 (m, 2H), 2.94 (ddt, J = 14.6, 3.3 Hz, 1H), 2.44 (dd, J = 13.1, 6.8 Hz, 1H), 2.40–2.30 (m, 1H), 2.24 (s, 3H), 2.11 (s, 1H), 1.97 (dd, J = 12.8, 8 Hz, 1H), 1.82–1.76 (m, 2H), 1.33 (s, 3H); ^{13}C NMR (125 MHz, CDCl_3) δ 170.4, 168.6, 155.2, 154.1, 149.9, 121.4, 119.0, 117.3, 77.4, 75.5, 49.3, 46.2, 29.1, 28.5, 26.0, 25.9, 21.5, 21.1; IR (Neat film, NaCl) 3444 (br), 2970, 2933, 2868, 1738, 1626 cm^{-1} ; HRMS (TOF) m/z : $[\text{M}+\text{H}]^+$ Calcd for $\text{C}_{18}\text{H}_{21}\text{O}_5$ 317.1384; Found 317.1388; $[\alpha]_{\text{D}}^{25.4} = +65.6$ (c 0.38, CH_2Cl_2).



Epoxide 48. Enol acetate **47** (4 mg, 0.013 mmol) was dissolved in benzene (2 mL) and to it added excess *t*-BuOOH followed by $\text{VO}(\text{acac})_2$ (3.4 mg, 0.013 mmol). The deep maroon solution was purified directly by thin-layer chromatography (EtOAc) to yield epoxide (3.3 mg, 76% yield) as a white solid.



Ketone 49. To a CH_2Cl_2 (1.8 mL) solution of allylic alcohol **47** (0.0028 g, 0.0089 mmol) was added PCC (0.031 g, 0.15 mmol). The solution was stirred for 45 minutes, diluted with ethyl acetate and filtered through a plug of silica. The crude reaction mixture was purified by preparatory thin-layer chromatography with 2:1 EtOAc / PhH eluent to provide the rearranged enone (0.001 g, 0.0032 mmol, 36% yield) as a colorless oil. ^1H NMR (500 Mhz, CDCl_3) δ 5.13 (ddd, J = 8.3, 8.3, 3.2 Hz, 1H), 4.64 (d, J = 8.3, 1H), 3.57 (dt, J = 13.7, 2.4 Hz, 1H), 3.52 (d, J = 13.7 Hz, 1H), 3.36–3.28 (m, 1H), 3.07 (dd, J = 20.1, 8.3 Hz, 1H), 3.01–2.92 (m, 1H), 2.76 (d, J = 20.1 Hz, 1H), 2.43–2.39 (m, 2H), 2.26 (s, 3H), 2.15 (app q, J \leq 1.1 Hz, 3H), 1.86–1.76 (m, 2H); ^{13}C NMR (125 MHz, CDCl_3) δ 193.8, 170.5, 168.2, 156.8, 152.6, 148.2, 133.9, 120.6, 119.6, 75.8, 51.6, 47.6, 43.6, 29.3, 25.6, 21.3, 21.1, 16.1; IR (neat film, NaCl) 2931, 1762, 1737, 1687 cm^{-1} ; HRMS (TOF) m/z : $[\text{M}+\text{H}]^+$ Calcd for $\text{C}_{18}\text{H}_{19}\text{O}_5$ 315.1227; Found 315.1233; $[\alpha]_{\text{D}}^{25} = +122.56$ (c 0.16, CH_2Cl_2).

ASSOCIATED CONTENT

Supporting Information

Spectra (^1H NMR, ^{13}C NMR, and IR). This material is available free of charge via the Internet at <http://pubs.acs.org>.

AUTHOR INFORMATION

Corresponding Author

*E-mail: stoltz@caltech.edu

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENTS

The authors wish to thank the NIH-NIGMS (R01GM080269), Amgen, the Gordon and Betty Moore Foundation, and Caltech for financial support. Additionally, the authors gratefully acknowledge Larry Henling and the late Dr. Michael Day (Caltech) for X-ray crystallographic structural determination, Dr. Mona Shahgholi and Naseem Torian (Caltech) for mass spectrometric analysis, Dr. David VanderVelde (Caltech) for NMR experimental assistance and helpful discussions, Dr. Masaki Seto, Dr. N. Sherden, and Dr. D. E. White for helpful discussions, and Dr. R. A. Craig, II for editorial suggestions. J.L.R. thanks the California Tobacco-Related Disease Research Program of the University of California, Grant Number 14DT-0004 for a predoctoral fellowship. A.C.J. thanks the NIH (F32GM082000) for a postdoctoral fellowship.

REFERENCES

-
- (1) For a review on the terpenoids that have been isolated from *Sinularia*, see (a) Kamel, H. N.; Slattery, M. *Pharmaceutical Biology*. **2005**, *43*, 253. The taxonomy of *Sinularia* has been well reviewed in (b) Coll, J. C. *Chem. Rev.* **1992**, *92*, 613.
- (2) Duh, C.-Y.; Wang, S.-K.; Chia, M.-C.; Chiang, M. Y. *Tetrahedron Lett.* **1999**, *40*, 6033.
- (3) (a) Sheu, J.-H.; Ahmed, A. F.; Shiue, R.-T.; Dai, C.-F.; Kuo, Y.-H. *J. Nat. Prod.* **2002**, *65*, 1904. (b) Ahmed, A. F.; Shiue, R.-T.; Wang, G.-H.; Dai, C.-F.; Kuo, Y.-H.; Sheu, J.-H. *Tetrahedron* **2003**, *59*, 7337. (c) Tseng, Y.-J.; Ahmed, A. F.; Hsu, C.-H.; Su, J.-H.; Dai, C.-F.; Sheu, J.-H. *J. Chinese Chem. Soc.* **2007**, *54*, 1041. (d) Cheng, S.-Y.; Chuang, C.-T.; Wen, Z.-H.; Wang, S.-K.; Chiou, S.-F.; Hsu, C.-H.; Dai, C.-F.; Duh, C.-Y. *Bioorg. Med. Chem.* **2010**, *18*, 3379. (e) Thao, N. P.; Nam, N. H.; Cuong, N. X.; Quang, T. H.; Tung, P. T.; Dat, L. D.; Chae, D.; Kim, S.; Koh, Y.-S.; Kiem, P. V.; Minh, C. V.; Kim, Y. H. *Bioorg. Med. Chem. Lett.* **2013**, *23*, 228. (f) Lillsunde, K.-E.; Festa, C.; Adel, H.; De Marino, S.; Lombardi, V.; Tilvi, S.; Nawrot, D. A.; Zampella, A.; D'Souza, L.; D'Auria, M. V.; Tammela, P. *Mar. Drugs* **2014**, *12*, 4045.
- (4) (a) Li, Y.; Pattenden, G. *Tetrahedron* **2011**, *67*, 10045. (b) Li Y.; Pattenden, G. *Nat. Prod. Rep.* **2011**, *28*, 1269. (c) Li Y.; Pattenden, G. *Nat. Prod. Rep.* **2011**, *28*, 429.
- (5) (a) C. O'Connell, Ph.D. dissertation, Queen's University of Belfast, Belfast, Northern Ireland, U.K., 2006. (b) C. E. O'Connell and A. J. Frontier, Abstracts of Papers, 32nd Northeast Regional Meeting of the American Chemical Society, Rochester, NY, Oct 31–Nov 3, 2004; American Chemical Society, 2004; GEN-088. (c) B. A. Pratt, Ph.D. dissertation, Scripps Research Institute, La Jolla, California, 2008. (d) G. Liu, Ph.D. dissertation, Texas A&M University, College Station, Texas, 2011. (e) G. Liu, and D. Romo, Abstracts of Papers, 237th American Chemical Society National Meeting, Salt Lake City, UT, Mar 22–26, 2009; American Chemical Society, 2009; ORGN-083. (f) F. Tang, Ph.D. dissertation, Washington University in St. Louis, St. Louis, Missouri, 2009. (g) F. Tang, and K. D. Moeller, *Tetrahedron* **2009**, *65*, 10863. (h) F. Tang, and K. D. Moeller, *J. Am. Chem. Soc.* **2007**, *129*, 12414. (i) E. J. Horn, J. S. Silverston and C. D. Vanderwal, *J. Org. Chem.* **2016**, *81*, 1819. (j) E. J. Horn, Ph.D. dissertation, University of California at Irvine, Irvine, CA, 2014.
- (6) Craig, II, R. A.; Roizen, J. L.; Smith, R. C.; Jones, A. C.; Virgil S. C.; Stoltz, B. M. *Chem. Sci.* **2017**, *8*, 507.

- (7)(a) Krüger S.; Gaich, T. *Beilstein J. Org. Chem.* **2014**, *10*, 183. (b) Davies, H. M. L.; Stafford, D. G.; Doan, B. D.; Houser, J. H. *J. Am. Chem. Soc.* **1998**, *120*, 3326. (c) Davies, H. M. L. *Tetrahedron* **1993**, *49*, 5203. (d) E. Vogel, *Angew. Chem., Int. Ed.* **1963**, *2*, 1.
- (8) (a) Craig, H. R. A.; Roizen, J. L.; Smith, R. C.; Jones, A. C.; Stoltz, B. M. *Org. Lett.* **2012**, *14*, 5716. (b) Seto, M.; Roizen, J. L.; Stoltz, B. M. *Angew. Chem. Int. Ed.* **2008**, *47*, 6873.
- (9) (a) D. C. Behenna, J. T. Mohr, N. H. Sherden, S. C. Marinescu, A. M. Harned, K. Tani, M. Seto, S. Ma, Z. Novák, M. R. Krout, R. M. McFadden, J. L. Roizen, J. A. Enquist, Jr., D. E. White, S. R. Levine, K. V. Petrova, A. Iwashita, S. C. Virgil, B. M. Stoltz *Chem. – Eur. J.* **2011**, *17*, 14199. (b) Mohr, J. T.; Stoltz, B. M. *Chem.–Asian J.* **2007**, *2*, 1476. (c) Mohr, J. T.; Behenna, D. C.; Harned, A. M.; Stoltz, B. M. *Angew. Chem. Int. Ed.* **2005**, *44*, 6924. (d) Behenna, D. C.; Stoltz, B. M. *J. Am. Chem. Soc.* **2004**, *126*, 15044.
- (10) Trost, B. M.; Xu, J.; Reichle, M. *J. Am. Chem. Soc.* **2007**, *129*, 282.
- (11) (a) Tani, K.; Behenna, D. C.; McFadden, R. M.; Stoltz, B. M. *Org. Lett.* **2007**, *9*, 2529. (b) Krout, M. R.; Mohr, J. T.; Stoltz, B. M. *Org. Synth.* **2009**, *86*, 181. (c) Helmchen, G.; Pfaltz, A. *Acc. Chem. Res.* **2000**, *33*, 336. (d) Williams, J. M. J. *Synlett* **1996**, 705, and references therein.
- (12) Liu, H.-J.; Zhu, B.-Y. *Can. J. Chem.* **1991**, *69*, 2008.
- (13) Herscovici, J.; Egron, M.-J.; Antonakis, K. *J. Chem. Soc., Perkin Trans. 1* **1982**, 1967.
- (14) For similar transformations, see: Shibuya, M.; Tomizawa, M.; Iwabuchi, Y. *J. Org. Chem.* **2008**, *73*, 4750.
- (15) Pyrazole formation has been previously reported in thermolysis of vinyl diazo compounds: (a) Doyle, M. P.; Yan, M.; Gao, H.-M.; Blossey, E. C. *Org. Lett.* **2003**, *5*, 561. (b) Doyle, M. P.; Yan, M. *J. Org. Chem.* **2002**, *67*, 602. (c) Hutchinson, I. S.; Matlin, S. A.; Mete, A. *Tetrahedron Lett.* **2001**, *42*, 1773. (d) Pellicciari, R.; Natalini, B.; Sadeghpour, B. M.; Marinozzi, M.; Snyder, J. P.; Williamson, B. L.; Kuethe, J. T.; Padwa, A. *J. Am. Chem. Soc.* **1993**, *115*, 9468. (e) Davies, H. M. L.; Huby, N. J. S.; Cantrell, W. R., Jr.; Olive, J. L. *J. Am. Chem. Soc.* **1993**, *115*, 9468. (f) Padwa, A.; Kulkarni, Y. S.; Zhang, Z. *J. Org. Chem.* **1990**, *55*, 4144. (g) Bailey, R. J.; Card, P. J.; Shechter, H. *J. Am. Chem. Soc.* **1983**, *105*, 6096. (h) Hurd, C. D.; Lui, S. C. *J. Am. Chem. Soc.* **1935**, *57*, 2656.
- (16) De Leeuw, J. W.; De Waard, E. R.; Beetz, T.; Huisman, H. O. *Recl. Trav. Chim. Pays-Bas* **1973**, *92*, 1047.
- (17) For related examples involving intramolecular transactonization of γ -lactones, see: (a) Raffaelli, B.; Pohjoispää, M.; Hase, T.; Cardin, C. J.; Gan, Y.; Wähälä, K. *Org. Biomol. Chem.* **2008**, *6*, 2619. (b) Mulzer, J.; Giester, G.; Gilbert, M. *Helv. Chim. Acta* **2005**, *88*, 1560. (c) Duret, P.; Figadère, B.; Hocquemiller, R.; Cavé, A. *Tetrahedron Lett.* **1997**, *38*, 8849. (d) Clardy, J.; Springer, J. P.; Buechi, G.; Matsuo, K.; Wightman, R. *J. Am. Chem. Soc.* **1975**, *97*, 663.
- (18) The immediate product of acetylation does not demonstrate the necessary gHMBC correlations; however, ketone reduction generates two diastereomers, one of which exhibits diagnostic gHMBC correlations.
- (19) Brecht, J. *Justus Liebigs Annalen der Chemie* **1924**, *437*, 1.
- (20) (a) Tei, T.; Sugimura, T.; Katagiri, T.; Tai, A.; Okuyama, T. *Tetrahedron: Asymmetry* **2001**, *12*, 2727. (b) Davies, H. M. L.; Calvo, R. L.; Townsend, R. J.; Ren, P.; Churchill, R. M. *J. Org. Chem.* **2000**, *65*, 4261. (c) Davies, H. M. L. *Tetrahedron* **1993**, *49*, 5203. (d) Davies, H. M. L.; Clark, T. J.; Smith, H. D. *J. Org. Chem.* **1991**, *56*, 3817. (e) Cantrell, W. R., Jr.; Davies, H. M. L. *J. Org. Chem.* **1991**, *56*, 723. (f) Davies, H. M. L.; Clark, D. M.; Smith, T. K. *Tetrahedron Lett.* **1985**, *26*, 5659.
- (21) Davies, H. M. L.; Smith, H. D.; Korkor, O. *Tetrahedron Lett.* **1987**, *28*, 1853.

-
- (22) (a) Davies, H. M. L.; McAfee, M. J.; Oldenburg, C. E. M. *J. Org. Chem.* **1989**, *54*, 930; (b) Davies, H. M. L.; Oldenburg, C. E. M.; McAfee, M. J.; Nordahl, J. G.; Henretta, J. P.; Romines, K. R. *Tetrahedron Lett.* **1988**, *29*, 975.
- (23) For intermolecular cascades, see: (a) Davies, H. M. L.; Stafford, D. G.; Doan, B. D.; Houser, J. H. *J. Am. Chem. Soc.* **1998**, *120*, 3326. (b) Davies, H. M. L.; Peng, Z.-Q.; Houser, J. H. *Tetrahedron Lett.* **1994**, *35*, 8939.
- (24) Davies, H. M. L.; Doan, B. D. *J. Org. Chem.* **1999**, *64*, 8501.
- (25) For asymmetric intramolecular cascades, see: (a) Davies, H. M. L.; Ahmed, G.; Churchill, M. R. *J. Am. Chem. Soc.* **1996**, *118*, 10774. For related uses of chiral auxiliaries in cyclopropanation/Cope reactions, see: (b) Davies, H. M. L.; Matasi, J. J.; Hodges, L. M.; Huby, N. J. S.; Thornley, C.; Kong, N.; Houser, J. H. *J. Org. Chem.* **1997**, *62*, 1095.
- (26) Danheiser, R. L.; Miller, R. F.; Brisbois, R. G.; Park, S. Z. *J. Org. Chem.* **1990**, *55*, 1959.
- (27) Nozaki, H.; Takaya, H.; Moriuti, S.; Noyori, R. *Tetrahedron* **1968**, *24*, 3655.
- (28) (a) Frigerio, M.; Santagostino, M.; Sputore, S. *J. Org. Chem.* **1999**, *64*, 4537; (b) Boeckman, R. K., Jr.; Shao, P.; Mullins, J. J. *Org. Syn.* **2000**, *77*, 141.
- (29) de Boer, Th. J.; Backer, H. J. *Org. Syn.* **1963**, *Coll. Vol. 4*, 250–253.
- (30) Parlow, J. J.; Case, B. L.; South, M. S. *Tetrahedron* **1999**, *55*, 6785.
- (31) Baum, J. S.; Shook, D. A.; Davies, H. M. L.; Smith, H. D. *Synth. Comm.* **1987**, *17*, 1709.
- (32) This procedure has been influenced by: Bouzide, A.; Sauv  , G. *Tetrahedron Lett.* **1997**, *38*, 5945.
-