

Borole derivatives

XXI [☆], 2,5-Diphenyl-2,5-dihydro-1*H*-boroles; structures of *tert*-butyl-2,5-diphenyl-2,5-dihydro-1*H*-borole and of bis(*tmeda*) lithium 2,5-diphenyl-2,5-dihydro-1*H*-borolediide

Gerhard E. Herberich ^{*}, Trixie Wagner, Hans-W. Marx

Institut für Anorganische Chemie der Technischen Hochschule Aachen, D-52056 Aachen, Germany

Received 11 May 1995

Abstract

The magnesium reagent $\text{Mg}(\text{C}_6\text{H}_5)_2(\text{THF})_3$ derived from (*E,E*)-1,4-diphenyl-1,3-butadiene reacts with boron dihalides $\text{BCl}_2(\text{NR}_2)$ ($\text{R} = \text{Me}, \text{Et}, ^i\text{Pr}$) and $^t\text{BuBF}_2$ to give 2,5-dihydro-2,5-diphenyl-1*H*-boroles 2,5- $\text{Ph}_2\text{C}_4\text{H}_4\text{BNR}_2$ (**1a–1c**) and 2,5- $\text{Ph}_2\text{C}_4\text{H}_4\text{B}^t\text{Bu}$ (**1d**) as mixtures of *cis–trans* isomers. *cis*-**1d** possesses a classical structure with a ring folding of 8° along the line C-2,C-5 of the C_4B ring. The dimethylamino compounds **1a** undergo metallation when treated with LDA in THF. After addition of TMEDA, the product can be crystallized from hexane–THF (2:1) as $[\text{Li}(\text{TMEDA})]_2[2,5\text{-Ph}_2\text{C}_4\text{H}_4\text{BNMe}_2]$ (**3**). The structure of **3** is that of a contact ion triple with a triple-decker-like arrangement of the borole ring and two $[\text{Li}(\text{TMEDA})]^+$ units.

Keywords: 2,5-Diphenyl-2,5-dihydro-1*H*-boroles; 2,5-dihydro-1*H*-borolediide; Borole dianion; Boron; Borole

1. Introduction

Syntheses of simple 2,5-dihydro-1*H*-boroles (3-borolenes) are known [2–4], including *C*-unsubstituted [2] and various *C*-alkyl derivatives [3,4]. Here we report some exploratory work on the 2,5-diphenyl derivatives **1a–1d** and the metallation of the 1-(dimethylamino) compound **1a** to give a new borole dianion 2^{2-} . The main aim of this paper is to communicate two structures: (i) that of the *tert*-butyl compound *cis*-**1d** which is the first structure of a *B*-alkyl-2,5-dihydro-1*H*-borole and (ii) that of $[\text{Li}(\text{tmeda})]_2(\mathbf{2}) \equiv \mathbf{3}$ which is the second structure with a borole dianion.


2. Results and discussion

2.1. Synthesis of 2,5-dihydro-2,5-diphenyl-1*H*-boroles

Simple 3-borolenes can readily be made from magnesium-butadiene and organoborondihalides [2,3]. We have applied this synthetic method to (*E,E*)-1,4-diphenyl-1,3-butadiene. This highly activated butadiene readily forms a magnesium adduct $\text{Mg}(\text{C}_6\text{H}_5)_2(\text{THF})_3$ (**4**) which is well known [5] and has been characterized structurally [6]. Treatment with dichloro(dialkylamino)boranes $\text{BCl}_2(\text{NR}_2)$ ($\text{R} = \text{Me}, \text{Et}, ^i\text{Pr}$) produces the corresponding 3-borolenes **1a–1c** in near quantitative yields (Scheme 1). The products **1a–1c** are obtained as yellowish oils which slowly solidify. Nuclear magnetic resonance (NMR) spectroscopic analysis (see below) shows that they are mixtures of *cis–trans* isomers. In the case of **1c** a single crystallization from hexane afforded crystals of the pure *cis* isomer (*cis*-**1c**).

[☆] For part XX see Ref. [1]. Dedicated to Professor Henri Brunner on the occasion of his 60th birthday.

^{*} Corresponding author.




- 1a: R = NMe₂
 1b: R = NEt₂
 1c: R = NⁱPr₂
 1d: R = ^tBu

Two difficulties were encountered during the attempted synthesis of *B*-hydrocarbyl derivatives of type 1. One was ether cleavage of the THF which, for instance, was very fast in the reaction with MeBBBr₂. The other difficulty appeared when Mg(C₄H₄Ph₂)(THF)₃ was treated with ^tBuBF₂. This reaction produced (*E,E*)-1,4-diphenyl-1,3-butadiene in mixture with *cis*-1d and *trans*-1d; fractionating crystallization afforded a small amount of pure *cis*-1d.

2.2. Distinction of *cis* and *trans* isomers 1a–1d

The new 2,5-dihydro-2,5-diphenyl-1H-boroles were essentially characterized by NMR spectroscopy. Both *cis* and *trans* isomers exhibit twofold symmetry in solution corresponding to C_s symmetry for the *cis* and C₂ symmetry for the *trans* isomers. However, the distinction of the two geometries was not straightforward. Both isomers display an AA'BB' spin system [7] for the four protons of the borole ring. This may be approximated as an AA'XX' spin system because of the relatively large shift difference (ca. 2.5 ppm) between the AA' and the BB' parts.

For 1a–1c, the XX' part (3,4-H) displays an apparent triplet for the *cis* isomers and an apparent doublet for the *trans* isomers with small satellites in both cases. The AA' part (2,5-H) is somewhat less well resolved. Keeping in mind that vicinal coupling constants always have positive signs [7] the spectra could be simulated with the coupling constants given in Fig. 1. For *cis*-1d the XX' part is seen as an apparent quartet with two



Scheme 1.





Fig. 1. Coupling constants J_{HH} for the 2,5-dihydro-1H-boroles 1a–1c.



satellites, and simulation was possible with $J_{34} = 6.1$ Hz, $J_{23} = 2.5$ Hz, $J_{24} = -2.0$ Hz, and $J_{25} = 0$ Hz.

It seemed likely that the two Ph groups would increase the torsional angle 2-H,C-2,C-3,3-H if in *cis* disposition and hence would give rise to a smaller vicinal coupling constant J_{23} for the *cis* isomer. On this basis an assignment of the two isomers was feasible. However, a safer assignment was based on a comparison with 2,5-dihydro-1,1-dimethyl-2,5-diphenylsilole (**5**). Reaction of the magnesium reagent with Me_2SiCl_2 gave *cis*-**5** as described in the literature [8] but with a small (7%) admixture of the *trans* isomer (Scheme 2). Here the assignment of the stereochemistry is unambiguous. While the dihydrosilole ring protons display the same $\text{AA}'\text{BB}'$ -type spectra as in the case of the dihydroboroles **1**, the two Si-methyl groups are chemically non-equivalent for *cis*-**5** and symmetry related for *trans*-**5**.

2.3. Structure of the *tert*-butyl compound **1d**

2,5-Dihydro-1*H*-boroles may have a classical structure **A** with a planar ring skeleton or alternatively a non-classical structure **B** with a folded ring skeleton and an interaction between the empty p_z orbital of the boron and the π orbital of the olefinic double bond. For the parent compound $\text{C}_4\text{H}_4\text{BH}$, it has been shown by means of computational chemistry that the two structures are very close in energy, the difference being smaller than 1 kcal mol^{-1} [9].

Exocyclic substituents such as *B*-amino groups stabilize the classical structure **A** in agreement with the known structures of 1-amino-2,5-dihydro-1*H*-boroles which are indeed classical [10,11]. The non-classical structure **B** has been found in bicyclic systems where the bicyclic structure strongly favours the folding of the





Fig. 2. The molecular structure of the 2,5-dihydro-1*H*-borole **1d** with the crystallographic atom numbering scheme.

dihydroborole ring [12]. The case of the phenyl derivative 3,4- $\text{Me}_2\text{C}_4\text{H}_4\text{BPh}$ is interesting in that there is very little stabilization of the boron centre through π interaction with the phenyl substituent, yet this compound also displays the classical planar ring structure [4].

The *tert*-butyl group of compound **1d** avoids π stabilization at the boron centre altogether. We also reasoned that the *cis*-2,5-diphenyl grouping would give rise to intramolecular repulsion between the two phenyl groups, and this repulsion would be avoided if the dihydroborole ring was folded. The structure determination (Fig. 2, Tables 1 and 2) reveals that this dihydroborole is nevertheless essentially planar. There is a small but significant folding of $8(1)^\circ$ along the line C1,C4 of the dihydroborole ring. Thus the non-classical interaction with the olefinic double bond of the ring cannot be important.

2.4. Metallation of 2,5- $\text{Ph}_2\text{C}_4\text{H}_4\text{BNMe}_2$ (**1a**)

1-(Dimethylamino)-2,5-dihydro-1*H*-borole $\text{C}_4\text{H}_6\text{BNMe}_2$ undergoes metallation when treated with suitable lithium dialkylamides and quaternization at boron when treated with alkyl lithium reagents [13]. The stereoisomeric 2,5-diphenyl derivatives **1a** react with LDA (LiN^iPr_2) or LiTMP (lithium 2,2,6,6-tetramethylpiperide) in THF to give a dilithio derivative which after addition of TMEDA (*N,N,N',N'*-tetramethylethylenediamine) can be crystallized as intensely yellow solid $[\text{Li}(\text{TMEDA})]_2[2,5\text{-Ph}_2\text{C}_4\text{H}_2\text{BNMe}_2]$ (**3**). Previously known compounds with borole dianions are



Scheme 2.

Table 1

Atomic coordinates of non-hydrogen atoms for **1d**

Atom	x	y	z	$B_{eq} (\times 10^4 \text{ pm}^2)^a$
C1	0.2197(5)	0.6282(2)	0.6641(1)	3.62(6)
C2	0.2878(5)	0.7122(2)	0.6977(2)	4.14(6)
C3	0.2856(4)	0.7133(2)	0.7719(2)	4.01(6)
C4	0.2177(5)	0.6301(2)	0.8075(1)	3.71(6)
C5	0.0489(5)	0.4823(2)	0.7379(2)	4.12(6)
C11	0.0517(5)	0.6337(2)	0.6002(1)	3.43(5)
C12	−0.1446(5)	0.6829(2)	0.6064(1)	4.01(6)
C13	−0.3000(5)	0.6871(2)	0.5488(2)	4.67(7)
C14	−0.2667(5)	0.6416(2)	0.4840(2)	4.85(7)
C15	−0.0747(6)	0.5932(2)	0.4761(1)	5.21(8)
C16	0.0865(6)	0.5903(2)	0.5333(2)	4.54(7)
C41	0.0516(5)	0.6394(2)	0.8719(1)	3.58(6)
C42	−0.1481(6)	0.6855(2)	0.8627(2)	4.39(7)
C43	−0.3022(6)	0.6958(2)	0.9210(2)	5.04(7)
C44	−0.2623(6)	0.6585(2)	0.9889(2)	5.44(8)
C45	−0.0647(6)	0.6111(2)	0.9992(2)	5.66(8)
C46	0.0888(5)	0.6028(2)	0.9416(2)	4.55(7)
C51	−0.2084(6)	0.4927(2)	0.7392(3)	8.9(1)
C52	0.1167(8)	0.4292(2)	0.6704(2)	7.2(1)
C53	0.1242(8)	0.4316(2)	0.8073(2)	7.0(1)
B	0.1504(5)	0.5734(2)	0.7370(2)	3.50(6)

^a The anisotropic thermal parameters are given in the form of their isotropic equivalents, defined as $B_{eq} = (4/3)[a^2\beta_{11} + b^2\beta_{22} + c^2\beta_{33}]$.

$K_2[C_4Ph_4BPh]$ [14,15], $Li_2[C_4H_4BN^iPr_2]$ [16], and $[Li_2(TMEDA)][C_4H_4BNR_2]$ ($R = Me, Et, ^iPr$) [13].

The new compound **3** displays a boron resonance at $\delta(^{11}B) = 28$ ppm in the expected region. The 7Li resonance is found at $\delta(^7Li) = -5.00$ ppm in THF (0.49 mol l^{−1}, ambient temperature). This chemical shift indicates facial bonding of the lithium to the C_4B ring system. For $LiCp$ the corresponding resonance is found at $\delta(^7Li) = -8.37$ ppm (in THF, ambient temperature) [17] while the sandwich species $[LiCp_2]^-$ exhibits a resonance at $\delta(^7Li) = -13.1$ ppm (in THF, $-107^\circ C$) [18]. The ^{13}C NMR data indicate considerable delocalization of the anion charge into the phenyl substituents.

Table 2

Selected bond distances (picometres) and bond angles (degrees) for **1d**

B–C1	161.0(4)	C4–B	158.9(4)
C1–C2	150.4(4)	C3–C4	150.9(4)
C2–C3	132.0(3)	B–C5	155.3(4)
C1–C11	150.7(4)	C4–C41	151.0(4)
C5–C51	151.7(4)	C5–C52	151.7(4)
C5–C53	153.4(4)		
C2–C1–C11	115.1(2)	C3–C4–C41	113.9(2)
C2–C1–B	102.5(2)	C3–C4–B	102.7(2)
C11–C1–B	118.3(2)	C41–C4–B	119.5(2)
C1–C2–C3	114.0(3)	C2–C3–C4	114.3(3)
C1–B–C5	126.8(3)	C4–B–C5	127.2(3)
C1–B–C4	105.9(2)		
C51–C5–B	106.3(2)	C51–C5–C52	109.4(3)
C52–C5–B	113.6(2)	C51–C5–C53	109.3(3)
C53–C5–B	112.2(2)	C52–C5–C53	106.1(2)

2.5. Structure of $[Li(TMEDA)]_2[2,5-Ph_2C_4H_2BNMe_2]$ (**3**)

The structure of **3** displays a triple-decker-like arrangement of the dianionic borole ring and two $[Li(TMEDA)]^+$ units (Fig. 3, Tables 3 and 4). The contact ion triple exhibits an approximate C_2 symmetry. The C_4BN moiety is essentially planar (maximum deviation 3.5(3) pm for C8). The various fragments of the anion deviate systematically from coplanarity to avoid strong repulsions between the *N*-methyl and the phenyl groups. Thus, the dimethylamino group is rotated by $25.9(3)^\circ$ against the C_4B plane, and the two phenyl groups are rotated by $33.5(2)^\circ$ and $33.9(2)^\circ$ respectively.

There are indications that both the dimethylamino group and the phenyl groups perturb the aromatic dihydroborole ring. The B–N distance (147.0(4) pm) is between a typical double bond (141 pm [19]) and a representative B–N single bond (156 pm, between an NR_2 group and a trigonal boron atom as in $[Li(OEt_2)]_2[Ph(Me_2N)B=B(NMe_2)Ph]$ [20]). Furthermore, the C–B bonds in the ring (C6–B, 155.0(4) pm; C9–B, 154.1(4) pm) are significantly longer than in $[Li_2(TMEDA)][C_4H_4BNEt_2]$ (152.2(3) and 152.0(3) pm [13]). A comparison with calculated bond lengths (C–B, 155.9 pm; C=B, 139.0 pm) [21] also demonstrates the reduced contribution of C–B π bonding in **3**.

The $[Li(TMEDA)]^+$ units exhibit an asymmetric slip distortion, away from the boron and towards a trihapto bonding mode. This slip distortion clearly diminishes the non-bonding repulsions with the anion. It also implies an improved overlap with three ring carbon atoms and thus will increase covalent bonding contributions. This phenomenon is known as electrostatic orbital polarization [22,23].




Fig. 3. The structure of the 2,5-dihydro-1H-borole diide **3** with the crystallographic atom numbering scheme.

3. Experimental details

Reactions were carried out under an atmosphere of dinitrogen by means of conventional Schlenk techniques. Pentane and hexane were distilled from Na–K alloy and THF from sodium benzophenone ketyl. NMR spectra were recorded on Bruker WH 270 PFT (^{13}C , 67.9 MHz), JEOL NM-PS-100 (^{11}B , 32.08 MHz), Varian VXR 300 (^1H , 300 MHz, ^{13}C , 75.4 MHz), and Varian Unity 500 (^1H , 500 MHz; ^6Li , 73.6 MHz; ^7Li , 194.3 MHz; ^{13}C , 125.7 MHz) spectrometers. Mass spectra were recorded on a Varian MAT CH-5 spectrometer.

3.1. 1-(Dimethylamino)-2,5-dihydro-2,5-diphenyl-1H-borole (**1a**)

Magnesium–butadiene reagent **4** (5.8 g, 13.0 mmol) was suspended in hexane (or pentane) at -78°C and

Table 3
Atomic coordinates of non-hydrogen atoms for **3**

Atom	x	y	z	$B_{\text{eq}} (\times 10^4 \text{ pm}^2)^a$
N2	0.091	0.19744(6)	0.123	5.15(6)
N3	−0.2274(4)	0.16445(6)	−0.0377(3)	5.13(7)
C2	−0.0494(6)	0.21179(8)	−0.0098(4)	7.1(1)
C3	−0.1674(6)	0.18771(9)	−0.1161(4)	7.6(1)
C21	0.1667(6)	0.22103(9)	0.2447(4)	7.6(1)
C22	0.2254(5)	0.1855(1)	0.0839(5)	7.4(1)
C31	−0.2967(5)	0.13632(9)	−0.1314(4)	6.7(1)
C32	−0.3551(5)	0.1779(1)	0.0036(5)	7.5(1)
Li1	−0.0191(6)	0.1547(1)	0.1764(5)	4.1(1)
B	−0.1083(4)	0.12485(8)	0.3417(3)	3.44(6)
C6	0.0416(3)	0.14863(6)	0.4211(3)	3.33(5)
C7	0.1819(3)	0.13800(6)	0.3913(3)	3.54(5)
C8	0.1324(3)	0.11203(6)	0.2915(3)	3.50(5)
C9	−0.0380(3)	0.10135(6)	0.2624(3)	3.36(5)
N1	−0.2779(3)	0.12495(6)	0.3416(3)	4.32(5)
C11	−0.4313(4)	0.11153(9)	0.2216(4)	5.84(9)
C12	−0.3113(4)	0.13829(9)	0.4612(4)	5.82(8)
C61	0.0556(3)	0.17855(6)	0.5032(3)	3.62(6)
C62	−0.0872(4)	0.19849(7)	0.4662(3)	4.40(6)
C63	−0.0758(4)	0.22719(8)	0.5378(4)	5.34(8)
C64	0.0804(5)	0.23758(7)	0.6501(4)	5.81(9)
C65	0.2230(5)	0.21898(8)	0.6877(4)	5.65(9)
C66	0.2116(4)	0.18997(7)	0.6161(3)	4.66(7)
C91	−0.1054(3)	0.07167(6)	0.1807(3)	3.53(6)
C92	−0.2122(4)	0.05162(7)	0.2177(3)	4.48(7)
C93	−0.2731(4)	0.02298(7)	0.1453(4)	5.28(8)
C94	−0.2286(5)	0.01242(8)	0.0332(4)	5.87(9)
C95	−0.1234(5)	0.03100(8)	−0.0042(4)	5.65(9)
C96	−0.0630(4)	0.06014(7)	0.0672(3)	4.52(7)
Li2	0.1472(6)	0.0953(1)	0.5076(5)	4.0(1)
N4	0.3099(3)	0.05261(6)	0.5602(3)	5.08(6)
N5	0.1520(3)	0.08552(6)	0.7207(3)	5.12(6)
C4	0.2998(6)	0.03834(8)	0.6922(4)	7.2(1)
C5	0.2913(6)	0.06248(9)	0.7999(4)	7.7(1)
C41	0.2646(5)	0.02930(9)	0.4389(5)	7.4(1)
C42	0.4818(5)	0.06474(9)	0.5978(6)	7.6(1)
C51	0.1762(5)	0.11370(9)	0.8155(4)	6.7(1)
C52	−0.0179(5)	0.0726(1)	0.6796(4)	7.7(1)

^a The anisotropic thermal parameters are given in the form of their isotropic equivalents, defined as $B_{\text{eq}} = (4/3)[a^2\beta_{11} + b^2\beta_{22} + c^2\beta_{33} + ac(\cos\beta)\beta_{13}]$.

Table 4

Selected bond distances (picometres) and bond angles (degrees) for **3**

B–C6	155.0(4)	B–C9	154.1(4)
C6–C7	144.3(4)	C8–C9	144.7(4)
C7–C8	140.2(4)	B–N1	147.0(4)
N1–C11	144.3(4)	N1–C12	143.9(4)
C6–C61	147.1(4)	C9–C91	146.0(3)
Li1–B	242.9(6)	Li2–B	243.1(5)
Li1–C6	222.6(5)	Li2–C6	243.5(5)
Li1–C7	216.9(5)	Li2–C7	222.2(6)
Li1–C8	221.6(5)	Li2–C8	217.8(6)
Li1–C9	243.0(5)	Li2–C9	223.3(5)
N2–C2	146.1(5)	N3–C3	147.4(5)
N2–C21	146.6(5)	N3–C31	145.6(4)
N2–C22	146.3(5)	N3–C32	145.1(5)
N2–Li1	221.1(6)	N3–Li1	211.4(5)
C2–C3	148.9(5)		
N4–C4	146.1(5)	N5–C5	147.4(5)
N4–C41	145.4(5)	N5–C51	146.4(5)
N4–C42	146.0(5)	N5–C52	144.9(5)
Li2–N4	220.1(5)	Li2–N5	210.7(6)
C4–C5	149.0(6)		
C7–C6–C61	120.2(2)	C8–C9–C91	120.0(2)
C7–C6–B	107.0(2)	C8–C9–B	107.3(2)
C61–C6–B	132.6(2)	C91–C9–B	132.4(2)
C6–C7–C8	111.0(2)	C7–C8–C9	110.6(2)
C6–B–N1	127.9(3)	C9–B–N1	128.3(3)
C6–B–C9	103.7(2)		
B–N1–C11	124.1(3)	B–N1–C12	124.2(2)
C11–N1–C12	111.7(2)		

$\text{BCl}_2(\text{NMe}_2)$ [24] (1.64 g, 13.0 mmol) was added. Stirring was continued while the reaction mixture was allowed to warm up to ambient temperature. After the end of the reaction (ca. 3 h, as seen by ^{11}B NMR spectroscopy and by a change of colour) the MgCl_2 was filtered off and washed with several small portions of solvent. All volatiles were carefully removed from the filtrate in a vacuum. The residue was a yellowish oil which crystallized very slowly from hexane solution to give solid **1a** (3.23 g, 95%); air and water sensitive; *cis*:*trans* = 2.2:1 (NMR).

MS: m/e ($I_{\text{rel.}}$) = 261 (100, M), 170 (51, M – C_7H_7). ^{11}B NMR (CDCl_3): δ = 49.

cis-**1a**. ^1H NMR (500 MHz, CDCl_3): δ = 7.15–7.35 (m, 2 Ph), 5.98 (m, 3-, 4-H), 3.27 (m, 2-, 5-H), 2.66 (s, NMe_2). $^{13}\text{C}\{^1\text{H}\}$ NMR (126 MHz, CDCl_3): δ = 144.59 (C_i), 137.35 (C-3, 4), 129.05 and 128.34 (C_o and C_m), 124.79 (C_p), 43.4 (C-2, 5), 40.93 (NMe_2).

trans-**1a**. ^1H NMR (500 MHz, CDCl_3): δ = 7.15–7.35 (m, 2 Ph), 5.98 (m, 3-, 4-H), 3.38 (m, 2-, 5-H), 2.53 (s, NMe_2). $^{13}\text{C}\{^1\text{H}\}$ NMR (126 MHz, CDCl_3): δ = 144.99 (C_i), 137.72 (C-3, 4), 128.64 and 128.15 (C_o and C_m), 124.65 (C_p), 43.4 (C-2, 5), 40.53 (NMe_2).

3.2. 1-(Diethylamino)-2,5-dihydro-2,5-diphenyl-1H-borole (**1b**)

Treatment of **4** (5.8 g, 13.0 mmol) with $\text{BCl}_2(\text{NEt}_2)$ [25] (1.82 g, 13.0 mmol) at 0°C , stirring at ambient temperature and workup as described for **1a** gave **1b**

(3.57 g, 95%); air and water sensitive; *cis*:*trans* = 1.1:1 (NMR).

MS: m/e (I_{rel}) = 289 (75, M), 117 (94, C_9H_8^+). ^{11}B NMR (CDCl_3): δ = 49.

cis-**1b**. ^1H NMR (500 MHz, CDCl_3): δ = 7.15–7.35 (m, 2 Ph), 5.96 (m, 3-, 4-H), 3.26 (m, 2-, 5-H). NET_2 : 2.95 and 2.93 (dq, CH_2), 0.74 (t, 2 Me), $^2J_{11}$ = 14.0, $^3J_{12}$ = 7.0 Hz. $^{13}\text{C}\{^1\text{H}\}$ NMR (126 MHz, CDCl_3): δ = 145.05 (C_i), 137.33 (C-3,4), 128.85 and 128.35 (C_o and C_m), 124.77 (C_p), 43.5 (C-2,5), 43.27 (NCH_2), 14.47 (Me).

trans-**1a**. ^1H NMR (500 MHz, CDCl_3): δ = 7.15–7.35 (m, 2 Ph), 5.99 (m, 3-, 4-H), 3.37 (m, 2-, 5-H). NET_2 : 2.92 and 2.75 (dq, CH_2), 0.48 (t, 2 Me), $^2J_{11}$ = 14.0, $^3J_{12}$ = 7.0 Hz. $^{13}\text{C}\{^1\text{H}\}$ NMR (126 MHz, CDCl_3): δ = 145.40 (C_i), 137.67 (C-3, 4), 128.49 and 128.28 (C_o and C_m), 124.64 (C_p), 43.5 (C-2, 5), 42.71 (NCH_2), 13.94 (Me).

3.3. 1-(Diisopropylamino)-2,5-dihydro-2,5-diphenyl-1H-borole (**1c**)

Treatment of **4** (5.8 g, 13.0 mmol) with BCl_2 (N^iPr_2) [26] (2.36 g, 13.0 mmol) at -78°C , stirring at ambient temperature (ca. 3 d) and workup as described for **1a** gave **1c** (3.9 g, 95%); air and water sensitive; *cis*:*trans* = 2.4:1 (NMR). A single crystallization from hexane gave 2.1 g (51%) of pure *cis*-**1c**.

MS: m/e (I_{rel}) = 317 (70, M), 77 (100, Ph^+). ^{11}B NMR (CDCl_3): δ = 49.

cis-**1c**. ^1H NMR (500 MHz, CDCl_3): δ = 7.11–7.29 (m, 2 Ph), 5.89 (m, 3-, 4-H), 3.31 (m, 2-, 5-H). N^iPr_2 : 3.64 (sept, 2 CH), 0.98 and 0.90 (d, 2 Me), 3J = 6.71 Hz. ^{13}C NMR (68 MHz, CDCl_3): δ = 145.01 (q, 7.3 Hz, C_i), 136.96 (dq, 160.4, 5.5 Hz, C-3, 4), 129.44 (dq, 156.1, 6.3 Hz, C_o), 128.30 (dd, 158.6, 7.0 Hz, C_m), 124.68 (dt, 160.2, 6.7 Hz, C_p), 44.35 (d, 115.8 Hz, C-2, 5). N^iPr_2 : 49.67 (d, 135 Hz, CH), 23.22 (qm, 125.5, 4.5 Hz, Me), 22.49 (qm, 125.4, 4.5 Hz, Me). The barrier to internal rotation about the C–N bond was determined by the coalescence method from ^{13}C spectra [27]: T_c = 243 K, $\Delta\nu$ = 612 Hz, $\Delta G^\ddagger$ = 44.2 kJ mol $^{-1}$ for NCH; T_c = 223 K, $\Delta\nu$ = 89 Hz, $\Delta G^\ddagger$ = 43.9 kJ mol $^{-1}$ for C_i ; T_c = 223 K, $\Delta\nu$ = 101 Hz, $\Delta G^\ddagger$ = 43.7 kJ mol $^{-1}$ for C-3, 4.

trans-**1c**. ^1H NMR (500 MHz, CDCl_3): δ = 7.11–7.29 (m, 2 Ph), 5.90 (m, 3-, 4-H), 3.41 (m, 2-, 5-H). N^iPr_2 : 3.45 (sept, 2 CH), 0.77 and 0.72 (d, 2 Me), 3J = 6.71 Hz. $^{13}\text{C}\{^1\text{H}\}$ NMR (126 MHz, CDCl_3): δ = 146.84 (C_i), 137.34 (C-3, 4), 129.00 (C_o), 128.25 (C_m), 124.65 (C_p), 44.3 (C-2, 5). N^iPr_2 : 49.6 (CH), 22.99 and 22.31 (Me).

3.4. 1-tert-Butyl-2,5-dihydro-2,5-diphenyl-1H-borole (**1d**)

The reaction of **4** with an equivalent of $^t\text{BuBF}_2$ [28] at -78°C , workup as usual, gave a mixture of *cis*-**1d**,

trans-**1d**, and (*E*, *E*)-1,4-diphenyl-1,3-butadiene. Fractionating crystallization from hexane first removed the diene; further cooling to -30°C afforded colourless rod-shaped crystals of *cis*-**1d** (yield not determined); very air and water sensitive.

MS: m/e (I_{rel}) = 247 (50, M), 142 (100, ?). ^{11}B NMR (CDCl_3): δ = 90 br.

cis-**1d**. ^1H NMR (500 MHz, CDCl_3): δ = 7.27 (m, 4 H_o), 7.13–7.18 (m, 4 H_m + 2 H_p), 6.03 (m, 3-, 4-H), 3.48 (m, 2-, 5-H), 0.70 (s, ^tBu). $^{13}\text{C}\{^1\text{H}\}$ NMR (126 MHz, CDCl_3): δ = 142.74 (C_i), 136.85 (C-3, 4), 129.20 and 128.46 (C_o and C_m), 125.28 (C_p), 51.2 (C-2, 5), 27.73 (Me), 27 (br, CMe_3).

trans-**1d**. ^1H NMR (300 MHz, CDCl_3): δ = 7.1–7.5 (m, 2 Ph), 6.07 (m, 3-, 4-H), 3.69 (m, 2-, 5-H), 0.87 (s, ^tBu).

3.5. 2,5-Dihydro-1,1-dimethyl-2,5-diphenylsilole (**5**)

Treatment of **4** (4.1 g, 9.2 mmol) in pentane (80 ml) with Me_2SiCl_2 (1.1 ml, 9.0 mmol) at -78°C and workup as described for **1a** gave an oil. This was chromatographed on a short column of alumina and eluted until all material with a blue fluorescence (366 nm) had been passed through. Removal of the eluent left **5** (1.5 g, 63%) a colourless oil; *cis*:*trans* = 15:1 (NMR).

cis-**5**. NMR data as published [8].

trans-**5**. ^1H NMR (500 MHz, CDCl_3): δ = 6.18 (m, 3-, 4-H), 3.29 (m, 2-, 5-H), -0.18 (s, SiMe_2), remaining signals obscured by signals of *cis* isomer.

3.6. Synthesis of $[\text{Li}(\text{TMEDA})]_2[2,5\text{-Ph}_2\text{C}_4\text{H}_2\text{BNMe}_2]$ (**3**)

1a (2.43 g, 9.3 mmol) in hexane (5 ml) was added dropwise with stirring to LiN^iPr_2 [29] (19.0 mmol, prepared in situ from NH^iPr_2 (2.7 ml) in THF (20 ml) and LiBu (11.9 ml, 1.6 m in hexane)). The colour of the mixture slowly turned red while it was allowed to warm up to ambient temperature. Stirring was continued for 24 h; monitoring the reaction by ^{11}B NMR spectroscopy is recommendable. After removal of the volatiles, the residue was washed with hexane, then dissolved with heating in a mixture of THF (20 ml), hexane (40 ml) and TMEDA (3 ml). Slow cooling to -78°C gave **3** (2.3 g, 49%) as large, vivid yellow crystals; air and water sensitive; darkens above 110°C ; melting point ca. 150°C (dec.).

^1H NMR (500 MHz, $\text{THF}-d_8$): δ = 7.44 (d br, 7.6 Hz, 4 H_o), 6.92 (t br, 7.4 Hz, 4 H_m), 6.43 (t br, 7.1 Hz, 2 H_p), 5.87 (s, 3-, 4-H), 2.87 (s, NMe_2). TMEDA: 2.16 (s, 4 NCH_2), 1.98 (s, 8 NMe). $^{13}\text{C}\{^1\text{H}\}$ NMR (126 MHz, $\text{THF}-d_8$): δ = 149.32 (C_i), 127.53 and 125.14 (C_o and C_m), 117.62 (C_p), 105.5 (br, C-2, 5), 100.07 (C-3, 4), 45.03 (NMe_2). TMEDA: 57.75 (NCH_2), 45.70 (NMe). ^7Li NMR (194.3 MHz, $\text{THF}-d_8$, 0.49 mol l $^{-1}$): δ = -5.00 . ^{11}B NMR ($\text{THF}-d_8$): δ = 28.

Table 5

Crystallographic data, data collection parameters, and refinement parameters

Compound	1d	3
Formula	C ₂₀ H ₂₃ B	C ₃₀ H ₅₀ BLi ₂ N ₅
Formula weight	274.22	505.46
Space group	P2 ₁ 2 ₁ 2 ₁ (no. 19)	Cc (no. 9)
a (pm)	586.1(2)	867.0(2)
b (pm)	1573.1(6)	4215.5(4)
c (pm)	1779.9(4)	978.4(1)
β (°)		116.288(9)
V (nm ³)	1.642(2)	3.206(2)
Z	4	4
ρ (calc.) (g cm ⁻³)	1.109	1.047
α (cm ⁻¹)	0.57	4.29
Temperature (°C)	-15	25
Radiation, λ (pm)	Mo K α, 71.07	Cu K α, 154.18
Crystal dimensions (mm ³)	0.5 × 0.9 × 0.9	0.2 × 0.5 × 0.6
Measured reflections	2596	6455
Scan range	3° ≤ θ ≤ 25°	5° ≤ θ ≤ 70°
Scan type	ω	ω-2θ
Absorption correction	None	None
Secondary extinction coefficient E	Not refined	1.46 × 10 ⁻⁵
Observed independent reflections, I > 1σ(I)	1841	2678
Parameters refined	246	342
R ^a , R _w ^b	0.055, 0.057	0.046, 0.054
Goodness of fit	1.430	1.233
Residual electron density (e pm ⁻³)	0.3 × 10 ⁻⁶	0.1 × 10 ⁻⁶

$$^a R = \sum ||F_o| - |F_c|| / \sum |F_o|, \quad ^b R_w = [\sum w(|F_o| - |F_c|)^2 / \sum w|F_o|^2]^{1/2} \text{ with } w^{-1} = \sigma^2(F_o).$$

3.7. Structure determinations of 1d and 3

For both compounds geometry and intensity data were collected on ENRAF Nonius CAD4 diffractometers equipped with graphite monochromators. The structure solutions by direct methods (SHELXS-86 [30]) were straightforward and yielded the coordinates of all non-hydrogen atoms. The structures were refined with the SDP program package [31]. All non-hydrogen atoms were refined anisotropically. Hydrogen atoms of **1d** were refined isotropically except those at the *tert*-butyl group, which were treated as riding atoms with idealized geometry; all hydrogen atoms of **3** were also treated as riding atoms (C–H, 98 pm; B(H) = 1.3B_{iso}(C)). A statistical weighting scheme $w^{-1} = \sigma^2(F_o)$ was used for all reflections. Crystal data, data collection parameters, and convergence results are compiled in Table 5.

Further details of the crystal structure investigations are available on request from the Fachinformationszentrum Karlsruhe, Gesellschaft für wissenschaftlich-technische Information m.b.H., D-76344 Eggenstein-Leopoldshafen, on quoting the depository numbers CSD-404122 (**1d**) and CSD-404121 (**3**).

Acknowledgements

We thank Dr. U. Englert for helpful discussions. This work was generously supported by the Volkswagen-Stiftung and the Fonds der Chemischen Industrie.

References

- [1] G.E. Herberich, T. Carstensen, D.P.J. Köffer, N. Klaff, R. Boese, I. Hyla-Kryspin, R. Gleiter, M. Stephan, H. Meth and U. Zenneck, *Organometallics*, **13** (1994) 619.
- [2] G.E. Herberich, W. Boveleth, B. Heßner, M. Hostalek, D.P.J. Köffer, H. Ohst and D. Söhnen, *Chem. Ber.*, **119** (1986) 420.
- [3] G. Zweifel, S.J. Backlund and T. Leung, *J. Am. Chem. Soc.*, **99** (1977) 5192. G.M. Clark, K.G. Hancock and G. Zweifel, *J. Am. Chem. Soc.*, **93** (1971) 1308. T.M. Shoup and G. Zweifel, *Organomet. Synth.*, **3** (1986) 456.
- [4] G.E. Herberich, H.-W. Marx and T. Wagner, *Chem. Ber.*, **127** (1994) 2135.
- [5] R. Baker, R.C. Cookson and A.D. Saunders, *J. Chem. Soc., Perkin Trans. I*, (1976) 1809.
- [6] Y. Kai, N. Kanehisa, K. Miki, N. Kasai, K. Mashima, H. Yasuda and A. Nakamura, *Chem. Lett.*, (1982) 1277.
- [7] H. Günther, *NMR-Spektroskopie*, Georg Thieme, Stuttgart, 2nd edn., 1983.
- [8] R.D. Rieke and H. Xiong, *J. Org. Chem.*, **56** (1991) 3109.
- [9] J.M. Schulman, R.L. Disch, P.v.R. Schleyer, M. Bühl, M. Brehmer and W. Koch, *J. Am. Chem. Soc.*, **114** (1992) 7897.
- [10] D. Bromm, D. Stalke, A. Heine, A. Meller and G.M. Sheldrick, *J. Organomet. Chem.*, **386** (1990) 1.
- [11] G.E. Herberich, T.P. Spaniol and U. Steffan, *Chem. Ber.*, **127** (1994) 1401.
- [12] P.J. Fagan, E.G. Burns and J.C. Calabrese, *J. Am. Chem. Soc.*, **110** (1988) 2979. P.J. Fagan, W.A. Nugent and J.C. Calabrese, *J. Am. Chem. Soc.*, **116** (1994) 1880.
- [13] G.E. Herberich, M. Hostalek, R. Laven, R. Boese, *Angew. Chem.*, **102** (1990) 330; *Angew. Chem., Int. Edn. Engl.*, **29** (1990) 317.
- [14] G.E. Herberich, B. Buller, B. Hessner and W. Oschmann, *J. Organomet. Chem.*, **195** (1980) 253.
- [15] J.J. Eisch, J.E. Galle and S. Kozima, *J. Am. Chem. Soc.*, **108** (1986) 379.
- [16] G.E. Herberich and H. Ohst, *Z. Naturforsch., Teil B*, **38** (1983) 1388.
- [17] R.H. Cox, H.W. Terry and L.W. Harrison, *J. Am. Chem. Soc.*, **93** (1971) 3297.
- [18] L.A. Paquette, W. Bauer, M.R. Sivik, M. Bühl, M. Feigel and P.v.R. Schleyer, *J. Am. Chem. Soc.*, **112** (1990) 8776.
- [19] P. Paetzold, *Adv. Inorg. Chem.*, **31** (1987) 123.
- [20] A. Moezzi, R.A. Bartlett and P.P. Power, *Angew. Chem.*, **104** (1992) 1075; *Angew. Chem., Int. Edn. Engl.*, **31** (1992) 1082.
- [21] P.v.R. Schleyer, P.K. Freeman, H. Jiao and B. Goldfuß, *Angew. Chem.*, **107** (1995) 332; *Angew. Chem., Int. Edn. Engl.*, **34** (1995) 337.
- [22] T. Clark, C. Rohde and P.v.R. Schleyer, *Organometallics*, **2** (1983) 1344.
- [23] P.v.R. Schleyer, A.J. Kos, D. Wilhelm, T. Clark, G. Boche, G. Decher, H. Etzrodt, H. Dietrich and W. Mahdi, *J. Chem. Soc., Chem. Commun.*, (1984) 1495.
- [24] A.J. Banister, N.N. Greenwood, B.P. Straughan and J. Walker, *J. Chem. Soc.*, (1964) 995. H. Nöth, P. Schweizer and F. Ziegelgänsberger, *Chem. Ber.*, **99** (1966) 1089.
- [25] K. Niedenzu and J.W. Dawson, *J. Am. Chem. Soc.*, **81** (1959) 3561.

- [26] W. Gerrard, H.R. Hudson and E.F. Mooney, *J. Chem. Soc.*, (1960) 5168. K. Niedenzu, H. Beyer, J.W. Dawson and H. Jenne, *Chem. Ber.*, 96 (1963) 2653.
- [27] J. Sandström, *Dynamic NMR Spectroscopy*, Academic Press, London, 1982.
- [28] R. Boese, P. Paetzold, A. Tapper and R. Ziembinski, *Chem. Ber.*, 122 (1989) 1057.
- [29] M. Fieser and L.F. Fieser, *Reagents for Organic Synthesis*, Vol. 4, Wiley Interscience, New York, 1974, p. 310.
- [30] B.A. Frenz and ENRAF Nonius, *SDP*, Version 5.0, Delft, 1988.
- [31] G.M. Sheldrick, *SHELXS-86, PROGRAM FOR CRYSTAL STRUCTURE SOLUTION*, Göttingen, 1986.