Natural Products Synthesis

Total Synthesis and Determination of the Absolute Configuration of (+)-Intricatetraol**

Yoshiki Morimoto,* Tatsuya Okita, Mamoru Takaishi, and Takeshi Tanaka

Intricatetraol (1), a member of a family of squalene-derived triterpene polyethers named oxasqualenoids,^[1] was isolated



from the red alga *Laurencia intricata* by Suzuki et al. in 1993. A crude fraction containing intricatetraol (1) as the major component exhibited cytotoxic activity against P388 "leuke-mia cells" with an IC₅₀ value of 12.5 μ gmL^{-1.[2]} The structural analysis was mainly carried out by NMR spectroscopic methods. Although it was found that the molecule has C_2 symmetry, the *cis* configuration within the tetrahydrofuran ring, the *R* configuration at C11 (C14), the relative configurations at C6 and C7 (C18 and C19), C10 and C11 (C14 and C15), and at bromine-bonded C3 (C22) remained to be determined.

There are many other types of oxasqualenoids; however, it is often difficult to determine their configurations even by highly advanced spectroscopic methods, especially for the acyclic portions that include stereogenic quaternary carbon centers, such as C6–C7 (C18–C19) and C10–C11 (C14–C15) in **1**. These difficulties have prompted attempts to determine the configurations of oxasqualenoids by chemical synthesis.^[3] Furthermore, the presence of the vicinal bromochloro functionality in **1** makes the problem of stereochemical assignment even more inaccessible. An enantioselective

 [*] Prof. Dr. Y. Morimoto, T. Okita, M. Takaishi, T. Tanaka Department of Chemistry Graduate School of Science, Osaka City University Sumiyoshi-ku, Osaka 558–8585 (Japan) Fax: (+ 81) 6-6605-2522 E-mail: morimoto@sci.osaka-cu.ac.jp

- [**] We thank Dr. M. Suzuki, University Malaysia Sabah, for kindly providing us with an authentic sample of natural intricatetraol and its spectra. This research was supported by the Novartis Foundation (Japan) for the Promotion of Science, the Saneyoshi Scholarship Foundation, and Grants-in-Aid for Scientific Research on Basic Research (C) from the JSPS and on Priority Areas 17035071 from the MEXT.
- Supporting information for this article is available on the WWW under http://www.angewandte.org or from the author.

method for the synthesis of the vicinal bromochloro functionality,^[4] which also occurs in the marine polyhalogenated monoterpene halomon (2), a promising anticancer agent,^[5] has never been developed. Herein we report the total assignment of the configuration of (+)-intricatetraol (1) as 3 through the first asymmetric total synthesis of this natural product. The synthesis features the enantioselective construction of the vicinal bromochloro functionality through a pathway in which the configuration is secured.

Suzuki et al. suggested structure **4**, which disregards the configuration at C3 (C22), as a possibility on the basis of the hypothetical biogenesis,^[2] and recently we^[6] and Ujihara^[7] independently confirmed this assignment. Our retrosynthetic



Scheme 1. Retrosynthetic analysis of the target compound **4**. AD = asymmetric dihydroxylation, AE = asymmetric epoxidation.

analysis of the target compound **4** is depicted in Scheme 1. We envisaged that it would be efficient to dimerize by olefin metathesis the functionalized fragment **5**, which represents half of the molecule, because of the C_2 symmetry of the natural product. The *R* or *S* configuration at the carbon atom attached to bromine in **5** could be introduced through epoxide chemistry. The desired trihydroxytetrahydrofuran **6** might be constructed by the stereospecific and stereoselective oxacyc-



1132

lization of the diepoxyalcohol **8** under alkaline conditions.^[8] The required stereocenters in **8** would be introduced into commercially available *trans,trans*-farnesyl acetate (9) by established methods of asymmetric oxidation.

The protection of the chiral diol 10 $(>95\% ee)^{[9]}$ as a pmethoxybenzylidene acetal (7:3 mixture) and subsequent deacetylation afforded the allylic alcohol $11^{[10]}$ (Scheme 2). Katsuki-Sharpless asymmetric epoxidation of 11 in the presence of (+)-diethyl L-tartrate (L-(+)-DET) provided the epoxyalcohol 12. Asymmetric epoxidation of the alkene 12 by the method of Shi and co-workers^[11] with ketone **13** as the chiral catalyst then gave the diepoxyalcohol 14. Treatment of the diepoxyalcohol 14 with 1M aqueous solution of lithium hydroxide and 1,4-dioxane (1:1) under reflux furnished the desired trihydroxytetrahydrofuran product 15 with high stereospecificity. It was thought from our previous studies^[3g,8] that the reaction proceeds through a Payne rearrangement, 5exo-tet formation of the ether ring, and an intermolecular attack of hydroxide ion on the epoxide in the more stable conformer C. In this reaction, the bicyclic ether 16 was also formed as a by-product. Compound 16 could be generated from the less stable conformer **D** by an intramolecular attack of the tertiary alkoxide on the epoxide ring. Selective tertbutyldimethylsilyl (TBDMS) protection of the primary hydroxy group in the triol 15, methoxymethyl (MOM) protection of the remaining hydroxy groups, deprotection of the silyl ether, Parikh-Doering oxidation of the alcohol, and Wittig methylenation of the resulting aldehyde afforded the terminal alkene 17 in good overall yield. Oxidation of the p-methoxybenzylidene acetal 17 with 2,3-dichloro-5,6-dicyano-1,4-benzoquinone (DDQ)^[12] provided a 3:7 mixture of regioisomeric benzoate esters, which upon reduction with lithium aluminum hydride gave the deprotected diol **18**.

Selective mesylation of the secondary hydroxy group in the diol **18** followed by treatment of the mesylate with potassium carbonate yielded the epoxide **19** (Scheme 3).^[13] The epoxide-opening reaction of **19** with dilithium tetrabromonickelate in THF proceeded regioselectively to produce the secondary bromide **20** with inversion of configuration at the less hindered carbon atom.^[14] It has been reported that dilithium tetrabromonickelate reacts with epoxides through an S_N2 mechanism. We confirmed that the bromohydrin **20** is reconverted into the starting epoxide **19** upon treatment with a base. Chlorination^[15] of the bromohydrin **20** and subsequent deprotection of the alcohols protected as MOM ethers afforded the fully functionalized half fragment **21** with a *cis* tetrahydrofuran ring, as confirmed by the observation of an NOE between the hydrogen atoms indicated in structure **21**.

In the chlorination of bromohydrin 20 with thionyl chloride in DMPU, the major (chlorination) and minor (dehydration) products 21 and 23 were formed in a 2:1 ratio (Scheme 4). We also prepared the regioisomeric bromohydrin 22 by treating the epoxide 19 with magnesium bromide etherate.^[16] When the bromohydrin 22 was subjected to the same chlorination conditions, products 21 and 23 were formed in almost the same ratio as in the reaction of 20. Therefore, it seems that the chlorination proceeds via a common bromonium-ion intermediate **E**, as described previously for a similar chlorination by Mioskowski and co-workers.^[4a]



Scheme 2. Reagents and conditions: a) p-MeOC₆H₄CH(OMe)₂, pyridinium p-toluenesulfonate, CH₂Cl₂, 0°C \rightarrow RT, 16 h; b) K₂CO₃, MeOH, RT, 7 h, 99% (2 steps); c) tBuOOH, Ti(OiPr)₄, L-(+)-DET, M.S. (4 Å), CH₂Cl₂, -20°C, 21 h, 95% (d.r. > 20:1); d) **13**, oxone, Bu₄NHSO₄, CH₂(OMe)₂/CH₃CN/H₂O, pH 10.5, 0°C, 2.5 h, 87% (d.r. > 6:1); e) aq. LiOH (1 M), 1,4-dioxane, 100°C, 7 h; f) TBDMSCl, Et₃N, 4-(dimethylamino)pyridine, CH₂Cl₂, RT, 26 h, 96%; g) MOMCl, iPr₂NEt, CH₂Cl₂, 0°C \rightarrow RT, 36 h, 96%; h) Bu₄NF, THF, 0°C \rightarrow RT, 15 h, 100%; j) SO₃·pyridine, Et₃N, DMSO/CH₂Cl₂, 0°C, 30 min, 100%; j) Ph₃P=CH₂, THF, 0°C, 1 h, 90%; k) DDQ, H₂O, CH₂Cl₂, RT, 2 h; l) LiAlH₄, THF, 0°C, 1.5 h, 96% (2 steps). DMSO=dimethyl sulfoxide, MP=p-methoxyphenyl.

Angew. Chem. Int. Ed. 2007, 46, 1132-1135

© 2007 Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim

Communications



(+)-intricatetraol (1)

Scheme 3. Reagents and conditions: a) methanesulfonyl chloride, pyridine, CH_2Cl_2 , $0^{\circ}C \rightarrow RT$, 14 h; then K_2CO_3 , MeOH, RT, 3 h, 90%; b) $Li_2[NiBr_4]$, THF, RT, 56 h, 76%; c) K_2CO_3 , MeOH, RT, 1 h, 92%; d) $SOCl_2$, DMPU, $0^{\circ}C$, 30 min; e) HCl, MeOH, RT, 66 h, 55% (2 steps); f) **24**, CH_2Cl_2 , $40^{\circ}C$, 7 h, 86%; g) (KOCON)_2, AcOH, MeOH, RT, 70 h, 56%. DMPU = 1,3-dimethyl-3,4,5,6-tetrahydro-2(1*H*)pyrimidinone.



Scheme 4. Possible mechanism for the chlorination of bromohydrins **20** and **22**.

The dimerization of fragment **21** was carried out in 86% yield by olefin metathesis with the Grubbs second-generation

catalyst **24** (Scheme 3; Cy = cyclohexyl).^[17] Although catalytic hydrogenation of the alkene **25** under standard conditions afforded the dehalogenated product by overreduction, diimide reduction^[18] provided the target compound **3** without dehalogenation. The respective spectra of the synthetic compound **3** ($[a]_D^{23} = +51.3$ (c = 0.41, CHCl₃)), prepared from the *R* alcohol **10**, were identical to those of the natural product ($[a]_D^{20} = +53.0$ (c = 0.625, CHCl₃)).^[2,6,7] Thus, it was found that the hitherto unknown absolute configuration of (+)-intricatetraol (**1**) is shown by the structural formula **3**.^[19]

In conclusion, we have completed the first asymmetric total synthesis of the marine bromochloro-functionalized triterpene polyether (+)-intricatetraol (1). Our synthesis features the enantioselective construction of the unique vicinal bromochloro functionality in an approach that may be applied to the synthesis of other bromochloro compounds, such as halomon (2), and an efficient olefin-metathesis strategy that takes the intrinsic molecular symmetry of the natural product into consideration. The total synthesis resulted in the assignment of the absolute configuration of intricatetraol (1), which is difficult to determine by other means. Further contributions to structure elucidation by organic synthesis are in progress.

Received: September 15, 2006 Revised: October 16, 2006 Published online: December 27, 2006

Keywords: asymmetric synthesis · configuration determination · halogen compounds · polyethers · terpenoids

- For reviews, see: a) J. D. Connolly, R. A. Hill, *Nat. Prod. Rep.* 2003, 20, 640–660; b) J. W. Blunt, B. R. Copp, M. H. G. Munro, P. T. Northcote, M. R. Prinsep, *Nat. Prod. Rep.* 2004, 21, 1–49.
- [2] M. Suzuki, Y. Matsuo, S. Takeda, T. Suzuki, *Phytochemistry* 1993, 33, 651–656.
- [3] For the assignment of the absolute configurations of oxasqualenoids by total synthesis, see: a) H. Kigoshi, M. Ojika, Y. Shizuri, H. Niwa, K. Yamada, *Tetrahedron* **1986**, *42*, 3789-3792;
 b) Y. Morimoto, T. Iwai, T. Kinoshita, J. Am. Chem. Soc. **2000**, *122*, 7124-7125; c) Z. Xiong, E. J. Corey, J. Am. Chem. Soc. **2000**, *122*, 9328-9329; d) Y. Morimoto, T. Iwai, T. Kinoshita, *Tetrahedron Lett.* **2001**, *42*, 6307-6309; e) H. Kigoshi, T. Itoh, T. Ogawa, K. Ochi, M. Okada, K. Suenaga, K. Yamada, *Tetrahedron Lett.* **2001**, *42*, 7461-7464; f) Y. Morimoto, M. Takaishi, T. Iwai, T. Kinoshita, H. Jacobs, *Tetrahedron Lett.* **2002**, *43*, 5849-5852; g) Y. Morimoto, Y. Nishikawa, M. Takaishi, J. Am. Chem. Soc. **2005**, *127*, 5806-5807.
- [4] For the total synthesis of halomon (2), see: a) T. Schlama, R. Baati, V. Gouverneur, A. Valleix, J. R. Falck, C. Mioskowski, *Angew. Chem.* 1998, 110, 2226–2228; *Angew. Chem. Int. Ed.* 1998, 37, 2085–2087; b) T. Sotokawa, T. Noda, S. Pi, M. Hirama, *Angew. Chem.* 2000, 112, 3572–3574; *Angew. Chem. Int. Ed.* 2000, 39, 3430–3432.
- [5] a) N. Ichikawa, Y. Naya, S. Enomoto, *Chem. Lett.* 1974, 1333–1336; b) B. J. Burreson, F. X. Woolard, R. E. Moore, *Chem. Lett.* 1975, 1111–1114; c) R. W. Fuller, J. H. Cardellina II, Y. Kato, L. S. Brinen, J. Clardy, K. M. Snader, M. R. Boyd, *J. Med. Chem.* 1992, *35*, 3007–3011; d) R. W. Fuller, *J. Med. Chem.* 1994, *37*, 4407–4411; see Supporting Information.
- [6] Y. Morimoto, M. Takaishi, N. Adachi, T. Okita, H. Yata, Org. Biomol. Chem. 2006, 4, 3220–3222.



- [7] K. Ujihara, PhD thesis, The University of Tokyo (Japan), 2004.
- [8] a) T. R. Hoye, S. A. Jenkins, J. Am. Chem. Soc. 1987, 109, 6196–6198; b) Y. Morimoto, T. Iwai, Y. Nishikawa, T. Kinoshita, *Tetrahedron: Asymmetry* 2002, 13, 2641–2647.
- [9] G. Vidari, A. Dapiaggi, G. Zanoni, L. Garlaschelli, *Tetrahedron Lett.* 1993, 34, 6485-6488.
- [10] All new compounds were characterized by ¹H NMR, ¹³C NMR, and IR spectroscopy, and by low- and high-resolution mass spectrometry (see Supporting Information).
- [11] Z.-X. Wang, Y. Tu, M. Frohn, J.-R. Zhang, Y. Shi, J. Am. Chem. Soc. 1997, 119, 11224–11235.
- [12] Y. Oikawa, T. Yoshioka, O. Yonemitsu, *Tetrahedron Lett.* 1982, 23, 889–892.

- [13] E. J. Corey, M. C. Noe, W.-C. Shieh, *Tetrahedron Lett.* 1993, 34, 5995-5998.
- [14] R. D. Dawe, T. F. Molinski, J. V. Turner, *Tetrahedron Lett.* 1984, 25, 2061–2064.
- [15] B. Chen, R. Y. Y. Ko, M. S. M. Yuen, K.-F. Cheng, P. Chiu, J. Org. Chem. 2003, 68, 4195–4205.
- [16] P. J. Mohr, R. L. Halcomb, J. Am. Chem. Soc. 2003, 125, 1712– 1713.
- [17] T. M. Trnka, R. H. Grubbs, Acc. Chem. Res. 2001, 34, 18-29.
- [18] D. J. Pasto, R. T. Taylor in *Organic Reactions, Vol. 40* (Eds.: L. A. Paquette et al.), Wiley, New York, **1991**, pp. 91–155; see Supporting Information.
- [19] For further discussion on the possibility of other diastereomers, see Supporting Information.