

Catalytic Dealkylative Synthesis of Cyclic Carbamates and Ureas via Hydrogen Atom Transfer and Radical-Polar Crossover

Takuya Nagai, Nao Mimata, Yoshihiro Terada, Chikayoshi Sebe, and Hiroki Shigehisa*



Cite This: <https://dx.doi.org/10.1021/acs.orglett.0c01872>



Read Online

ACCESS |



Metrics & More




Article Recommendations



Supporting Information

ABSTRACT: Guided by the transition-metal hydrogen atom transfer and radical-polar crossover concepts, we developed a functional-group-tolerant and scalable method for the synthesis of cyclic carbamates and ureas, which are found in the structures of bioactive compounds. This method provides not only a common five-membered ring but also six-to-eight-membered ring products. The reaction proceeds through the intramolecular displacement of an alkylcobalt(IV) intermediate and dealkylation by 2,4,6-collidine; the activation energies of these steps were calculated by DFT.



Much effort has been devoted in modern synthetic organic chemistry to the development of diverse methods for hydrofunctionalizing alkenes through transition-metal hydrogen-atom-transfer (TM-HAT) processes.¹ A transition metal hydride generated *in situ* from a catalyst and a hydrogen source reacts with the alkene unit to chemoselectively generate a carbon-centered radical that then becomes involved in diverse transformations (Scheme 1A).² Recently, this mechanism has been shown to operate in conjunction with other transition-metal catalysis mechanisms, thereby expanding the scope of the transformation.³

Our group has independently shown that the addition of an *N*-fluorocollidinium salt to a commonly used cobalt–Schiff base catalyst and a silane facilitates radical-polar crossover (RPC) to generate a cationic intermediate. To date, we have disclosed that oxygen,⁴ nitrogen,⁵ carbon,⁶ and sulfur⁷ nucleophiles react with these cationic species in TM-HAT chemistry. Among them, the formation of a lactone from an alkenyl ester is a profound example for us (Scheme 1B).^{4b} Importantly, Shenvi investigated the mechanism of their bimetallic coupling reaction in the presence of an *N*-fluorocollidinium salt and suggested the involvement of an alkylcobalt(IV) intermediate.^{3b} In light of Shenvi and related reports^{2u,w,7,8} for the lactone formation, it is plausible that the sp² oxygen atom of the carbonyl group attacks the reactive cationic carbon of the alkylcobalt(IV) intermediate to release a cobalt(II) complex, after which removal of the methyl group by 2,4,6-collidine led to the lactone. Indeed, we computed the activation energies for the steps involving TS-a and TS-b to be only 11.0 and 14.1 kcal/mol, which suggests that these two steps are possible. With the aim of expanding the substrate scope, the promising reactivity of the TM-

HAT/RPC mechanism prompted us to design further cyclization reactions that involve poorly nucleophilic species.

Herein, we report the dealkylative cyclizations of alkenyl carbamates and alkenyl isoureas to afford cyclic carbamates and cyclic ureas, respectively (Scheme 1C). Cyclic carbamates and ureas are important structural motifs found in pharmaceuticals and bioactive agents, such as linezolid (antibiotic active against VRE and MRSA),⁹ efavirenz (anti-HIV drug),¹⁰ biotin (cofactor),¹¹ and aquileidine (natural product).¹² This observation encouraged synthetic chemists to develop various preparative methods for cyclic carbamates¹³ and cyclic ureas.¹⁴ However, to the best of our knowledge, the substrate scope of most reactions is limited to only common five- and six-membered rings, and examples of medium ring formation are rather rare.^{13g,14g} On the other hand, the method described here provides five- to eight-membered ring compounds.

We commenced by determining an appropriate carbamate structure using previously developed reaction conditions: cobalt catalyst C1, *N*-fluoro-2,4,6-collidinium trifluoromethanesulfonate (3), and 1,1,3,3-tetramethyldisiloxane in benzotrifluoride at room temperature (Scheme 2). Although the desired oxazolidinone 2 was obtained from all substrates 1a–1c, we found that *tert*-butyl carbamate 1c gave the best results. It should be noted that the methyl ester gave a better


Received: June 4, 2020

Scheme 1. TM-HAT and RPC Mechanisms



(A) TM-HAT catalyzed reaction





(B) Cyclization of esters to afford lactones via TM-HAT/RPC and DFT calculations for transition states of displacement and dealkylation^a



(C) Cyclization to afford cyclic carbamates & ureas via TM-HAT/RPC and biologically active compounds and natural product (This Work)



Scheme 2. Initial Attempt of Oxazolidine Synthesis^a




^aThe reactions of **1a–1c** (0.20 mmol) were performed in the presence of **C1** (3 mol %), **3** (2.0 equiv), and 1,1,3,3-tetramethyldisiloxane (2.0 equiv) in benzotrifluoride (2.0 mL) at room temperature for 18 h under argon. Isolated yields are shown. DFT calculations were performed for the synthesis of **5** to reduce calculational costs.

yield than the corresponding *tert*-butyl ester in the previously reported dealkylative cyclization that affords lactones.^{4b} We also observed coproducts **4** in 98% yield (based on **3**). Therefore, the deprotonation of the *tert*-butyl group by 2,4,6-collidine occurs after cyclization of the alkylcobalt(IV) intermediate. The activation energies for the steps involving **TS-c** and **TS-d** were calculated to be only 12.6 and 7.2 kcal/mol (without the OMe group to reduce calculational costs), which suggests that the mechanism shown in [Scheme 2](#) is feasible.

Encouraged by this result, we next examined the alkenyl carbamate scope for the formation of oxazolidinones (Scheme 3). Starting materials bearing hydrogen (i.e., **5**) or a methyl group (i.e., **6**) in the *p*-position of the aniline unit gave good yields of the desired products. On the other hand, the introduction of electron-withdrawing chloro, fluoro, and

trifluoromethyl groups (7–9) led to significantly lower yields and the recovery of the alkenyl *tert*-butyl carbamate. Fortunately, we solved this problem by replacing the carbamate unit with a methyl group. Clearly, these results contrast with those from the reaction that produces **2** with the electron-donating methoxy group. These aryl substituents would affect the rate of the intramolecular nucleophilic displacement and subsequent dealkylation, although their exact roles are not immediately clear. We also examined the functional group tolerance of this reaction using substrate **10** bearing a methylthio group, acid-sensitive acetal **11**, and fluoride-sensitive silyl ether **12**. The alkenyl *tert*-butylcarbamate bearing a disubstituted alkene, *p*-methoxybenzylamine, or *N*-tosyl-tryptamine cyclized to afford the desired products **13–15**. Moreover, amantadine derivative **16** was synthesized from the corresponding methyl carbamate on the 5 mmol


Scheme 3. Scope of Alkenyl Carbamates Affording Five-Membered Ring Products^c

^aMethyl carbamate was used instead. ^b1.25 g (5 mmol) of starting material was used. ^cConditions: alkenyl carbamate (0.2 mmol), $C1$ (3 mol %), 3 (2 equiv.), 1,1,3,3-tetramethyldisiloxane (2 equiv.), benzotrifluoride (2.0 mL), room temperature, 18 h.

scale. The corresponding *tert*-butyl carbamate was not as efficiently synthesized as the methyl carbamate (see SI).

We found that this method was able to form rings other than five-membered. Various cyclic carbamates **17–19** were prepared using the same reaction conditions from *tert*-butyl carbamates bearing *p*-methoxyaniline units (Scheme 4). To our great delight, these yields were improved by replacing the catalyst with $C2$, which was previously developed for the formation of medium rings by our group.^{4b} Unfortunately, the yield of **19** was unsatisfactory. The *tert*-butyl carbamates bearing *o*-methoxyaniline were cyclized to afford the seven-membered ring product **20** with $C2$. The methyl carbamate bearing the *p*-chloroaniline unit and the *tert*-butyl carbamate bearing the *p*-methoxybenzylamine unit afforded the six-membered ring products **21** and **23** in good yields. However, we found that **22** and **24** were synthesized inefficiently using this method, even though $C2$ was used.

We next applied the same concept to the preparation of cyclic ureas from alkenyl isoureas. Alkenyl *N*-tosyl-*O*-methylisoureas were synthesized from secondary amines in three steps (see SI). We found that the *p*-toluenesulfonyl group was required to produce cyclic ureas. Irrespective of the electronic character of the substituent on the aniline unit, the yields of imidazolidinones **25–30** were generally good to excellent (Scheme 5). Alkenyl isoureas bearing a disubstituted alkene or a *p*-methoxybenzylamine unit were also cyclized efficiently to give **31** and **32**. Six- and seven-membered cyclic ureas **33** and **34** were also prepared by this method in good

Scheme 4. Scope of Alkenyl Carbamates Affording Products of Six-, Seven-, and Eight-Membered Rings^c


^a*tert*-Butyl carbamate was used instead. ^bMethyl carbamate was used. ^cConditions: alkenyl carbamate (0.2 mmol), $C1$ or $C2$ (3 mol %), 3 (2 equiv.), 1,1,3,3-tetramethyldisiloxane (6 equiv.), benzotrifluoride (2.0 mL), room temperature, 18 h.

yields. A complex product mixture was formed, and starting material was recovered in experiments aimed at forming eight-membered rings. We also investigated the cyclization of an alkenyl urea with the aim of potentially producing a cyclic urea.^{14b} The cyclization of unpurified urea **36**, synthesized from *N*-allylaniline (**35**) and *p*-toluenesulfonyl isocyanate in ethanol, afforded cyclic isourea **37** (without **26**) in excellent yield from **35**. Therefore, it is clear that the dealkylative cyclization approach is valuable from the perspective of O/N selectivity.


With the aim of further extending synthetic utility, the *p*-toluenesulfonyl group in the product needed to be removable under mild conditions; however, we were unable to remove this group from **39** under common reaction conditions (Scheme 6). This observation prompted us to replace the *p*-toluenesulfonyl group with other groups, such as *o*-nitrobenzenesulfonyl and trifluoroacetyl. The yields of the cyclized products from **40** and **42** were acceptable, and these transformations were scalable (1.00 g). Both protecting groups were removed to afford the same product **44** under mild conditions. The seven-membered cyclic urea **47** was also prepared by the same method. In a preliminary attempt, the same reaction conditions were applied to the cyclization of alkenyl phosphoramidate **48**. Although there is much room for improving both yield and selectivity, we obtained the cyclic phosphoramidate **49**, together with a complex product mixture.

Scheme 5. Scope of Alkenyl Isooureas Affording Cyclic Ureas^b

(A) Cyclization of alkenyl isooureas



(B) Cyclization of alkenyl urea



^aC2 was used instead. ^b(A) Conditions: alkenyl carbamate (0.2 mmol), C1 (3 mol%), 3 (2 equiv.), 1,1,3,3-tetramethyldisiloxane (2 equiv.), benzotrifluoride (2.0 mL), room temperature, 18 hours.

In summary, we developed a catalytic, Markovnikov-selective, functional-group-tolerant, scalable method for the synthesis of cyclic carbamates using a TM-HAT/RPC approach. This reaction proceeds through the cyclization of an alkylcobalt(IV) intermediate and dealkylation by 2,4,6-collidine, and the mechanism was determined to be reasonable by DFT calculations. Various cyclic carbamates were efficiently synthesized under mild conditions. Cyclic ureas and a cyclic phosphoramidate were also synthesized in the same way. The HAT-initiated reaction developed here enables the construction of medium ring carbamates and ureas. We are currently investigating enantioselective versions of these dealkylative cyclizations using a chiral cobalt catalyst.

■ ASSOCIATED CONTENT

■ Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/acs.orglett.0c01872>.

Experimental procedures and analytical data (¹H and ¹³C NMR) for all new compounds (PDF)


■ AUTHOR INFORMATION

Corresponding Author


Hiroki Shigehisa – Faculty of Pharmacy, Musashino University, Nishitokyo-shi, Tokyo 202-8585, Japan;

Scheme 6. Additional Experiment for Cyclic Urea Synthesis

(A) Synthesis of protective group free cyclic urea



(B) Cyclization of alkenyl phosphoramidate



orcid.org/0000-0003-3034-1159; Email: cgehis@musashino-u.ac.jp

Authors

Takuya Nagai – Faculty of Pharmacy, Musashino University, Nishitokyo-shi, Tokyo 202-8585, Japan

Nao Mimata – Faculty of Pharmacy, Musashino University, Nishitokyo-shi, Tokyo 202-8585, Japan

Yoshihiro Terada – Faculty of Pharmacy, Musashino University, Nishitokyo-shi, Tokyo 202-8585, Japan

Chikayoshi Sebe – Faculty of Pharmacy, Musashino University, Nishitokyo-shi, Tokyo 202-8585, Japan

Complete contact information is available at: <https://pubs.acs.org/doi/10.1021/acs.orglett.0c01872>

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

We thank Prof. Kou Hiroya (Musashino University) for our liberal research environment. We thank Dr. Yasunori Toda (Shinshu University) for helpful advice on how to prepare cyclic phosphoramidate. We thank Dr. Ryo Takita (University of Tokyo) for helpful advice on how to find TS-d. This work was supported by JSPS KAKENHI Grant number 17K15426, the Takeda Science Foundation, and the Research

Foundation for Pharmaceutical Sciences. Computational chemistry was carried out using the computer facilities at the Research Institute for Information Technology, Kyushu University.

REFERENCES

- (1) (a) Green, S. A.; Crossley, S. W. M.; Matos, J. L. M.; Vásquez-Céspedes, S.; Shevick, S. L.; Shenvi, R. A. The High Chemofidelity of Metal-Catalyzed Hydrogen Atom Transfer. *Acc. Chem. Res.* **2018**, *51*, 2628–2640. (b) Crossley, S. W.; Obradors, C.; Martinez, R. M.; Shenvi, R. A. Mn-, Fe-, and Co-Catalyzed Radical Hydrofunctionalizations of Olefins. *Chem. Rev.* **2016**, *116*, 8912–9000.
- (2) (a) Waser, J.; Carreira, E. M. Convenient synthesis of alkylhydrazides by the cobalt-catalyzed hydrohydrazination reaction of olefins and azodicarboxylates. *J. Am. Chem. Soc.* **2004**, *126*, 5676–5677. (b) Waser, J.; Nambu, H.; Carreira, E. M. Cobalt-Catalyzed Hydroazidation of Olefins: Convenient Access to Alkyl Azides. *J. Am. Chem. Soc.* **2005**, *127*, 8294–8295. (c) Waser, J.; Gaspar, B.; Nambu, H.; Carreira, E. M. Hydrazines and Azides via the Metal-catalyzed Hydrohydrazination and Hydroazidation of Olefins. *J. Am. Chem. Soc.* **2006**, *128*, 11693–11712. (d) Gaspar, B.; Carreira, E. M. Catalytic hydrochlorination of unactivated olefins with paratoluenesulfonyl chloride. *Angew. Chem., Int. Ed.* **2008**, *47*, 5758–5760. (e) Gaspar, B.; Carreira, E. M. Cobalt Catalyzed Functionalization of Unactivated Alkenes: Regioselective Reductive C-C Bond Forming Reactions. *J. Am. Chem. Soc.* **2009**, *131*, 13214–13215. (f) Ishikawa, H.; Colby, D. A.; Seto, S.; Va, P.; Tam, A.; Kakei, H.; Rayl, T. J.; Hwang, I.; Boger, D. L. Total Synthesis of Vinblastine, Vincristine, Related Natural Products, and Key Structural Analogues. *J. Am. Chem. Soc.* **2009**, *131*, 4904–4916. (g) Barker, T. J.; Boger, D. L. Fe(III)/NaBH₄-mediated free radical hydrofluorination of unactivated alkenes. *J. Am. Chem. Soc.* **2012**, *134*, 13588–13591. (h) Leggans, E. K.; Barker, T. J.; Duncan, K. K.; Boger, D. L. Iron(III)/NaBH₄-mediated Additions to Unactivated Alkenes: Synthesis of Novel 20'-vinblastine Analogues. *Org. Lett.* **2012**, *14*, 1428–1431. (i) Lo, J. C.; Gui, J.; Yabe, Y.; Pan, C. M.; Baran, P. S. Functionalized Olefin Cross-coupling to Construct Carbon-carbon Bonds. *Nature* **2014**, *516*, 343–348. (j) Lo, J. C.; Yabe, Y.; Baran, P. S. A Practical and Catalytic Reductive Olefin Coupling. *J. Am. Chem. Soc.* **2014**, *136*, 1304–1307. (k) King, S. M.; Ma, X.; Herzon, S. B. A Method for the Selective Hydrogenation of Alkenyl Halides to Alkyl Halides. *J. Am. Chem. Soc.* **2014**, *136*, 6884–6887. (l) Iwasaki, K.; Wan, K. K.; Oppedisano, A.; Crossley, S. W. M.; Shenvi, R. A. Simple, Chemoselective Hydrogenation with Thermodynamic Stereocontrol. *J. Am. Chem. Soc.* **2014**, *136*, 1300–1303. (m) Crossley, S. W. M.; Barabé, F.; Shenvi, R. A. Simple, Chemoselective, Catalytic Olefin Isomerization. *J. Am. Chem. Soc.* **2014**, *136*, 16788–16791. (n) Dao, H. T.; Li, C.; Michaudel, Q.; Maxwell, B. D.; Baran, P. S. Hydromethylation of Unactivated Olefins. *J. Am. Chem. Soc.* **2015**, *137*, 8046–8049. (o) Gui, J.; Pan, C. M.; Jin, Y.; Qin, T.; Lo, J. C.; Lee, B. J.; Spergel, S. H.; Mertzman, M. E.; Pitts, W. J.; La Cruz, T. E.; Schmidt, M. A.; Darvatkar, N.; Natarajan, S. R.; Baran, P. S. Practical Olefin Hydroamination with Nitroarenes. *Science* **2015**, *348*, 886–891. (p) Ma, X.; Herzon, S. B. Non-classical Selectivities in the Reduction of Alkenes by Cobalt-mediated Hydrogen Atom Transfer. *Chem. Sci.* **2015**, *6*, 6250–6255. (q) Ma, X.; Herzon, S. B. Intermolecular Hydropyridylation of Unactivated Alkenes. *J. Am. Chem. Soc.* **2016**, *138*, 8718–8721. (r) Obradors, C.; Martinez, R. M.; Shenvi, R. A. Ph(*i*-PrO)SiH₂: An Exceptional Reductant for Metal-Catalyzed Hydrogen Atom Transfers. *J. Am. Chem. Soc.* **2016**, *138*, 4962–4971. (s) Shen, Y.; Qi, J.; Mao, Z.; Cui, S. Fe-Catalyzed Hydroalkylation of Olefins with para-Quinone Methides. *Org. Lett.* **2016**, *18*, 2722. (t) Lo, J. C.; Kim, D.; Pan, C. M.; Edwards, J. T.; Yabe, Y.; Gui, J.; Qin, T.; Gutierrez, S.; Giacoboni, J.; Smith, M. W.; Holland, P. L.; Baran, P. S. Fe-Catalyzed C-C Bond Construction from Olefins via Radicals. *J. Am. Chem. Soc.* **2017**, *139*, 2484–2503. (u) Touney, E. E.; Foy, N. J.; Pronin, S. V. Catalytic Radical-Polar Crossover Reactions of Allylic Alcohols. *J. Am. Chem. Soc.* **2018**, *140*, 16982–16987. (v) Matos, J. L. M.; Vasquez-Céspedes, S.; Gu, J.; Oguma, T.; Shenvi, R. A. Branch-Selective Addition of Unactivated Olefins into Imines and Aldehydes. *J. Am. Chem. Soc.* **2018**, *140*, 16976–16981. (w) Discolo, C. A.; Touney, E. E.; Pronin, S. V. Catalytic Asymmetric Radical-Polar Crossover Hydroalkoxylation. *J. Am. Chem. Soc.* **2019**, *141*, 17527–17532. (x) Green, S. A.; Huffman, T. R.; McCourt, R. O.; van der Puyl, V.; Shenvi, R. A. Hydroalkylation of Olefins To Form Quaternary Carbons. *J. Am. Chem. Soc.* **2019**, *141*, 7709–7714. (y) Sun, H.-L.; Yang, F.; Ye, W.-T.; Wang, J.-J.; Zhu, R. Dual Cobalt and Photoredox Catalysis Enabled Intermolecular Oxidative Hydrofunctionalization. *ACS Catal.* **2020**, *10*, 4983–4989. (z) Wu, B.; Zhu, R. Radical Philicity Inversion in Co- and Fe-Catalyzed Hydrogen-Atom-Transfer-Initiated Cyclizations of Unsaturated Acylsilanes. *ACS Catal.* **2020**, *10*, 510–515. (aa) Zhou, X. L.; Yang, F.; Sun, H. L.; Yin, Y. N.; Ye, W. T.; Zhu, R. Cobalt-Catalyzed Intermolecular Hydrofunctionalization of Alkenes: Evidence for a Bimetallic Pathway. *J. Am. Chem. Soc.* **2019**, *141*, 7250–7255. (ab) Shen, X.; Chen, X.; Chen, J.; Sun, Y.; Cheng, Z.; Lu, Z. Ligand-promoted cobalt-catalyzed radical hydroamination of alkenes. *Nat. Commun.* **2020**, *11*, 783. (ac) Wang, Y. Y.; Bode, J. W. Olefin Amine (OLA) Reagents for the Synthesis of Bridged Bicyclic and Spirocyclic Saturated N-Heterocycles by Catalytic Hydrogen Atom Transfer (HAT) Reactions. *J. Am. Chem. Soc.* **2019**, *141*, 9739–9745. (ad) Jiang, H.; Lai, W.; Chen, H. Generation of Carbon Radical from Iron-Hydride/Alkene: Exchange-Enhanced Reactivity Selects the Reactive Spin State. *ACS Catal.* **2019**, *9*, 6080–6086.
- (3) (a) Green, S. A.; Matos, J. L. M.; Yagi, A.; Shenvi, R. A. Branch-Selective Hydroarylation: Iodoarene-Olefin Cross-Coupling. *J. Am. Chem. Soc.* **2016**, *138*, 12779–12782. (b) Shevick, S. L.; Obradors, C.; Shenvi, R. A. Mechanistic Interrogation of Co/Ni-Dual Catalyzed Hydroarylation. *J. Am. Chem. Soc.* **2018**, *140*, 12056–12068.
- (4) (a) Shigehisa, H.; Aoki, T.; Yamaguchi, S.; Shimizu, N.; Hiroya, K. Hydroalkoxylation of Unactivated Olefins with Carbon Radicals and Carbocation Species as Key Intermediates. *J. Am. Chem. Soc.* **2013**, *135*, 10306–10309. (b) Shigehisa, H.; Hayashi, M.; Ohkawa, H.; Suzuki, T.; Okayasu, H.; Mukai, M.; Yamazaki, A.; Kawai, R.; Kikuchi, H.; Satoh, Y.; Fukuyama, A.; Hiroya, K. Catalytic Synthesis of Saturated Oxygen Heterocycles by Hydrofunctionalization of Unactivated Olefins: Unprotected and Protected Strategies. *J. Am. Chem. Soc.* **2016**, *138*, 10597–10604.
- (5) Shigehisa, H.; Koseki, N.; Shimizu, N.; Fujisawa, M.; Niitsu, M.; Hiroya, K. Catalytic Hydroamination of Unactivated Olefins Using a Co Catalyst for Complex Molecule Synthesis. *J. Am. Chem. Soc.* **2014**, *136*, 13534–13537.
- (6) Shigehisa, H.; Ano, T.; Honma, H.; Ebisawa, K.; Hiroya, K. Co-Catalyzed Hydroarylation of Unactivated Olefins. *Org. Lett.* **2016**, *18*, 3622–3625.
- (7) Date, S.; Hamasaki, K.; Sunagawa, K.; Koyama, H.; Sebe, C.; Hiroya, K.; Shigehisa, H. Catalytic Direct Cyclization of Alkenyl Thioester. *ACS Catal.* **2020**, *10*, 2039–2045.
- (8) (a) Anderson, S. N.; Ballard, D. H.; Chrastowski, J. Z.; Dodd, D.; Johnson, M. D. Inversion of Configuration in the Nucleophilic Displacement of Cobalt from Alkylcobalt(IV) Complexes and Its Relevance to the Halogenation of the Corresponding Alkylcobalt-(III) Complexes. *J. Chem. Soc., Chem. Commun.* **1972**, 685–686. (b) Magnuson, R. H.; Halpern, J.; Levitin, I. Y.; Vol'pin, M. E. Stereochemistry of the Nucleophilic Cleavage of Cobalt-carbon Bonds in Organocobalt(IV) Compounds. *J. Chem. Soc., Chem. Commun.* **1978**, 44–46. (c) Vol'pin, M. E.; Levitin, I. Y.; Sigan, A. L.; Halpern, J.; Tom, G. M. Reactivity of Organocobalt(IV) Chelate Complexes Toward Nucleophiles: Diversity of Mechanisms. *Inorg. Chim. Acta* **1980**, *41*, 271–277.
- (9) Mukhtar, T. A.; Wright, G. D. Streptogramins, Oxazolidinones, and Other Inhibitors of Bacterial Protein Synthesis. *Chem. Rev.* **2005**, *105*, 529–542.

- (10) Young, S. D.; Britcher, S. F.; Tran, L. O.; Payne, L. S.; Lumma, W. C.; Lyle, T. A.; Huff, J. R.; Anderson, P. S.; Olsen, D. B.; Carroll, S. S. L-743, 726 (DMP-266): a novel, highly potent nonnucleoside inhibitor of the human immunodeficiency virus type 1 reverse transcriptase. *Antimicrob. Agents Chemother.* **1995**, *39*, 2602–2605.
- (11) Cronan, J. E. Advances in synthesis of biotin and assembly of lipoic acid. *Curr. Opin. Chem. Biol.* **2018**, *47*, 60–66.
- (12) Chen, S.-B.; Gao, G.-Y.; Leung, H.-W.; Yeung, H.-W.; Yang, J.-S.; Xiao, P.-G. Aquileidine and Isoaquileidine, Novel Flavonoid Alkaloids from *Aquilegia ecalcarata*. *J. Nat. Prod.* **2001**, *64*, 85–87.
- (13) (a) Robles-Machín, R.; Adrio, J.; Carretero, J. C. Gold-Catalyzed Synthesis of Alkylidene 2-Oxazolidinones and 1,3-Oxazin-2-ones. *J. Org. Chem.* **2006**, *71*, 5023–5026. (b) Jagtap, S. R.; Patil, Y. P.; Fujita, S.-I.; Arai, M.; Bhanage, B. M. Heterogeneous base catalyzed synthesis of 2-oxazolidinones/2-imidiazolidinones via transesterification of ethylene carbonate with β -aminoalcohols/1,2-diamines. *Appl. Catal., A* **2008**, *341*, 133–138. (c) Ishida, T.; Kobayashi, R.; Yamada, T. Novel Method of Tetramic Acid Synthesis: Silver-Catalyzed Carbon Dioxide Incorporation into Propargylic Amine and Intramolecular Rearrangement. *Org. Lett.* **2014**, *16*, 2430–2433. (d) Paddock, R. L.; Adhikari, D.; Lord, R. L.; Baik, M.-H.; Nguyen, S. T. [(Salcen)Cr(III) + Lewis base]-catalyzed synthesis of N-aryl-substituted oxazolidinones from epoxides and aryl isocyanates. *Chem. Commun.* **2014**, *50*, 15187–15190. (e) Zhang, Y.; Zhang, Y.; Ren, Y.; Ramström, O. Synthesis of chiral oxazolidinone derivatives through lipase-catalyzed kinetic resolution. *J. Mol. Catal. B: Enzym.* **2015**, *122*, 29–34. (f) Wang, P.; Qin, J.; Yuan, D.; Wang, Y.; Yao, Y. Synthesis of Oxazolidinones from Epoxides and Isocyanates Catalyzed by Rare-Earth-Metal Complexes. *ChemCatChem* **2015**, *7*, 1145–1151. (g) Niemi, T.; Perea-Buceta, J. E.; Fernández, I.; Hiltunen, O.-M.; Salo, V.; Rautiainen, S.; Räisänen, M. T.; Repo, T. A One-Pot Synthesis of N-Aryl-2-Oxazolidinones and Cyclic Urethanes by the Lewis Base Catalyzed Fixation of Carbon Dioxide into Anilines and Bromoalkanes. *Chem. - Eur. J.* **2016**, *22*, 10355–10359. (h) Zhang, Y.; Zhang, Y.; Xie, S.; Yan, M.; Ramström, O. Lipase-catalyzed kinetic resolution of 3-phenyloxazolidin-2-one derivatives: Cascade O- and N-alkoxycarbonylations. *Catal. Commun.* **2016**, *82*, 11–15. (i) Toda, Y.; Gomyou, S.; Tanaka, S.; Komiyama, Y.; Kikuchi, A.; Suga, H. Tetraarylphosphonium Salt-Catalyzed Synthesis of Oxazolidinones from Isocyanates and Epoxides. *Org. Lett.* **2017**, *19*, 5786–5789. (j) Niemi, T.; Fernández, I.; Steadman, B.; Mannisto, J. K.; Repo, T. Carbon dioxide-based facile synthesis of cyclic carbamates from amino alcohols. *Chem. Commun.* **2018**, *54*, 3166–3169. (k) Yousefi, R.; Struble, T. J.; Payne, J. L.; Vishe, M.; Schley, N. D.; Johnston, J. N. Catalytic, Enantioselective Synthesis of Cyclic Carbamates from Dialkyl Amines by CO₂-Capture: Discovery, Development, and Mechanism. *J. Am. Chem. Soc.* **2019**, *141*, 618–625.
- (14) (a) Streuff, J.; Hövelmann, C. H.; Nieger, M.; Muñoz, K. Palladium(II)-Catalyzed Intramolecular Diamination of Unfunctionalized Alkenes. *J. Am. Chem. Soc.* **2005**, *127*, 14586–14587. (b) Rao, W.-H.; Yin, X.-S.; Shi, B.-F. Catalyst-Controlled Amino- versus Oxy-Acetoxylation of Urea-Tethered Alkenes: Efficient Synthesis of Cyclic Ureas and Isoureas. *Org. Lett.* **2015**, *17*, 3758–3761. (c) Kondoh, A.; Kamata, Y.; Terada, M. Synthesis of Enantioenriched γ -Amino- α,β -unsaturated Esters Utilizing Palladium-Catalyzed Rearrangement of Allylic Carbamates for Direct Application to Formal [3 + 2] Cycloaddition. *Org. Lett.* **2017**, *19*, 1682–1685. (d) Hinds, E. M.; Wolfe, J. P. A Cross-Metathesis/Aza-Michael Reaction Strategy for the Synthesis of Cyclic and Bicyclic Ureas. *J. Org. Chem.* **2018**, *83*, 10668–10676. (e) Wu, M.-S.; Fan, T.; Chen, S.-S.; Han, Z.-Y.; Gong, L.-Z. Pd(II)-Catalyzed Asymmetric Oxidative 1,2-Diamination of Conjugated Dienes with Ureas. *Org. Lett.* **2018**, *20*, 2485–2489. (f) Struble, T. J.; Lankswert, H. M.; Pink, M.; Johnston, J. N. Enantioselective Organocatalytic Amine-Isocyanate Capture-Cyclization: Regioselective Alkene Iodoamination for the Synthesis of Chiral Cyclic Ureas. *ACS Catal.* **2018**, *8*, 11926–11931. (g) Zhang, H.; Tian, P.; Ma, L.; Zhou, Y.; Jiang, C.; Lin, X.; Xiao, X. Remote Directed Isocyanation of Unactivated C(sp³)-H Bonds: Forging Seven-Membered Cyclic Ureas Enabled by Copper Catalysis. *Org. Lett.* **2020**, *22*, 997–1002.