

Synthesis of *cis*-2-Fluorocyclopropylamine by Stereoselective Cyclopropanation Under Phase-transfer Conditions

Jun-ichi Matsuo,* Yu-ichirou Tani,[†] and Yu-ichirou Hayakawa[‡]


Center for Basic Research, The Kitasato Institute, 6-15-5, Toshima, Kita-ku, Tokyo 114-0003

[†]Chemical Technology Research Laboratories, Daiichi Pharmaceutical Co., Ltd.,
1-16-13, Kita-Kasai, Edogawa-ku, Tokyo 134-8630

(Received February 2, 2004; CL-040125)


cis-2-Fluorocyclopropylamine is stereoselectively synthesized by cyclopropanation of 3-aryl-2-vinyl-3-(methoxy)isoindol-1-one by treating dibromofluoromethane with saturated aqueous KOH solution in the presence of 18-crown-6 in dichloromethane, followed by removal of a bromine atom of the formed bromofluorocyclopropane derivative with Raney Ni, and successive three steps-deprotection procedures for generating an amino group on the cyclopropane ring.

Quinolonecarboxylic acids are widely used for therapy of various infections,¹ and sitafloxacin (**1**) was recently found as a new generation of quinolone antibacterial agent having excellent antibacterial activity.² In order to synthesize **1** efficiently, stereoselective synthesis of (1*R*,2*S*)-2-fluorocyclopropylamine (**2**) is an important key step. Therefore, several synthetic trials have been reported:^{3,4} e.g., Terashima et al. reported that cyclopropanation of an *N*-vinylcarbamate with zinc-monofluorocarbene generated from fluorodiiodomethane and diethylzinc gave *N*-(2-fluorocyclopropyl)carbamates stereoselectively (*cis/trans* = 93:7).^{3a,c,e} However, pyrophoric nature of diethylzinc is a major drawback of the above method, especially in large-scale synthesis. Then, stereoselective cyclopropanation under phase-transfer conditions was planned to establish a useful and safe method. Only one non-stereoselective cyclopropanation of enamide under phase-transfer conditions affording an equal amount of *cis*- and *trans*-fluorocyclopropylamines was reported to date.⁴ We would like to describe here *cis*-selective cyclopropanation of cyclic enamides with bromofluorocarbene under phase-transfer conditions and transformation of thus-obtained bromofluorocyclopropane derivative to *cis*-2-fluorocyclopropylamine salt.



Cyclopropanation of commercially available *N*-vinylphthalimide was tried first under phase-transfer conditions by using Br₂CHF, saturated aq KOH, and a catalytic amount of BnBu₃N⁺Cl⁻: however, only a trace amount of the desired bromofluorocyclopropane derivative was obtained⁴ probably because electron density of the vinyl group of *N*-vinylphthalimide was insufficient for the reaction with the electrophilic bromofluorocarbene. Then, one of the two carbonyl groups of *N*-vinylphthalimide was reduced in order to increase the electron density

of the vinyl group. 2-Vinyl-3-(methoxy)isoindol-1-one (**4**) was afforded in 95% yield by the following procedure; *N*-vinylphthalimide was reduced by using NaBH₄ in EtOH to afford 2-vinyl-3-hydroxyisoindol-1-one (**3**),⁵ followed by methylation with NaH and MeI (Scheme 1).



Scheme 1. a) NaH, MeI, DMF (95%); b) CHBrF₂, sat. KOH, BnN⁺Bu₃Cl, CHCl₂ (92%); c) Raney Ni, H₂, EtOH (90%).

Cyclopropanation of thus-obtained **4** under the above-mentioned phase-transfer conditions proceeded smoothly to give the desired compound **5** in 92% yield as a mixture of four diastereomers (60:27:8:5). Since it was difficult to determine *cis/trans* selectivity of **5** directly, it was determined after a bromine atom of **5** was reductively removed by using Raney Ni in EtOH under a hydrogen atmosphere: a fluorocyclopropane derivative **6** was then obtained as a mixture of diastereomers (60:25:9:6), and its *cis/trans* selectivity was found to be *cis/trans* = 31:69 by ¹H NMR analysis (Scheme 1). In order to improve the *cis/trans* selectivity, the reaction conditions concerning phase-transfer catalysts (Aliquatoo[®] 336, 18-crown-6), solvents (1,2-dichloroethane, benzene, toluene, benzotrifluoride, petroleum ether, ether, CCl₄), and bases (NaOH, CsOH) were further investigated. However, not only the *cis/trans* selectivity but the diastereomer ratio did not change drastically. On the other hand, protecting groups of the hydroxy group of **3** gave slight influence on the diastereomer ratio: that is, only two diastereomers were formed when the 3-hydroxy group was protected with trityl group (dr = 70:30:0:0, *cis/trans* = 30:70).⁶ It was then suggested that when the 3-hydroxy group was protected with the bulky group, bromofluorocarbene would approach to the *N*-vinyl group from the π -face opposite to the trityl group, while the direction of a fluorine group was still not controlled on that π -face. Then, a bulky substituent was introduced at the 3-carbon of **5** in order to control the orientation of the fluorine group *cis* to the 2-nitrogen group.

3-Aryl(alkyl)-2-vinyl-3-methoxyisoindol-1-ones **7** were prepared by 1,2-addition of Grignard reagent to vinylphthalimide, followed by protection of a hydroxy group by using NaH and MeI in DMF. Cyclopropanation of thus-prepared enamides **7** with bromofluorocarbene under phase-transfer conditions was shown in Table 1. By introducing phenyl group at


Table 1. Cyclopropanation of **7**

Entry	R	8		9	
		Yield/%	Yield/%	D.r. ^a	<i>Cis/trans</i> ^a
1	Ph	60	90	46:26:14:14	72/28
2	<i>o</i> -MeC ₆ H ₄	42	90	44:33:13:10	77/23
3	<i>m</i> -MeC ₆ H ₄	78	80	50:28:12:10	78/22
4	<i>p</i> -MeC ₆ H ₄	44	87	49:29:13:9	78/22
5	<i>o</i> -MeOC ₆ H ₄	52	86	47:26:15:12	73/27
6 ^b	<i>o</i> -MeOC ₆ H ₄	88	86	47:26:15:12	73/27
7	<i>p</i> -MeOC ₆ H ₄	64	71	46:32:12:10	78/22
8	1-Naph	44	73	50:34:9:7	84/16
9	ⁱ Pr	86	79	40:16:30:14	56/44
10	^t Bu	67	82	33:33:18:16	66/34

^aRatios were determined by ¹H NMR. ^bCyclopropanation was performed at 0 °C.

the 3-carbon, the *cis/trans* ratio was inversed compared to the case using **5**, and a *cis*-fluorocyclopropylamine derivative was preferentially formed (*cis/trans* = 72/28, Entry 1). After screening aryl or alkyl groups to be introduced at the 3-carbon, 1-naphthyl group gave the best result (*cis/trans* = 84/16). By introducing alkyl groups such as *tert*-butyl or isopropyl group at the 3-carbon, the corresponding cyclopropane derivative was formed in a less selective manner. It was further found that the use of other protecting groups such as ethyl, benzyl, and isopentyl groups instead of the methyl group of **7** did not improve the *cis/trans* selectivity.

The obtained *cis*-product **9a** was converted to a *cis*-fluorocyclopropylamine salt as a model (Scheme 2). Namely, methyl



Scheme 2. a) 1 N HCl, THF, reflux (79%); b) NaBH₄, *i*-PrOH-H₂O (60%); c) *p*-TsOH, Et₂O (71%).

group was deprotected in refluxing aqueous 1 N HCl-THF to give a hemiaminal **10a** which was in turn reduced to amido alcohol **11a** with NaBH₄ in *i*-PrOH-H₂O. Finally, *cis*-fluorocyclopropylamine *p*-toluenesulfonic acid salt was obtained by treating **11a** with *p*-TsOH in Et₂O.

We are grateful to Professor Teruaki Mukaiyama for valuable discussion and advices.

References and Notes

- 1 L. A. Mitscher, R. M. Zavod, P. V. Devasthale, D. T. W. Chu, L. L. Shen, P. N. Sharma, and A. G. Pernet, *CHEM-TECH*, **50**, 29 (1991).
- 2 Y. Kimura, S. Atarashi, K. Kawakami, K. Sato, and I. Hayakawa, *J. Med. Chem.*, **37**, 3344 (1994), and references are cited therein.
- 3 a) O. Tamura, M. Hashimoto, Y. Kobayashi, T. Katoh, K. Nakatani, M. Kamada, I. Hayakawa, T. Akiba, and S. Terashima, *Tetrahedron Lett.*, **33**, 3483, 3487 (1992). b) O. Tamura, M. Hashimoto, Y. Kobayashi, T. Katoh, K. Nakatani, M. Kamada, I. Hayakawa, T. Akiba, and S. Terashima, *Tetrahedron*, **50**, 3889 (1994). c) T. Akiba, O. Tamura, M. Hashimoto, Y. Kobayashi, T. Katoh, K. Nakatani, M. Kamada, I. Hayakawa, and S. Terashima, *Tetrahedron*, **50**, 3905 (1994). d) S. Kaneko, A. Toyota, and I. Hayakawa, Jpn. Kokai Tokkyo Koho, JP 10072402 (1998); *Chem. Abstr.*, **128**, 217121 (1998). e) S. Terashima, O. Tamura, Y. Kobayashi, M. Hashimoto, T. Katoh, K. Nakatani, J. Yukimoto, T. Ehara, T. Akiba, and H. Kawanishi, Jpn. Kokai Tokkyo Koho, JP 06092911 (1994); *Chem. Abstr.*, **121**, 108042 (1994). f) Y. Kimura, H. Takahashi, M. Takemura, I. Hayakawa, K. Kawakami, T. Yabune, T. Tojo, and T. Iketani, Jpn. Kokai Tokkyo Koho, JP 05194323 (1993); *Chem. Abstr.*, **120**, 76935 (1994). g) Y. Yamamoto, K. Shimokawa, Y. Kimura, and I. Hayakawa, Jpn. Kokai Tokkyo Koho, JP 05301827 (1993); *Chem. Abstr.*, **121**, 8749 (1994).
- 4 N. Lui, A. Marhold, and S. Boehm, Eur. Pat. Appl., EP 711750 (1996); *Chem. Abstr.*, **125**, 86229 (1996).
- 5 Cyclopropanation of **3** and Br₂CHF under phase-transfer conditions did not proceed.
- 6 Diastereomer and *cis/trans* ratios are as follows concerning other protecting groups: Bn (dr = 65:26:5:4, *cis/trans* = 30/70), 1-NaphCH₂ (dr = 63:26:6:5, *cis/trans* = 31/69), 2-NaphCH₂ (dr = 65:25:6:4, *cis/trans* = 29/71).