



Accepted Article

Title: Decarboxylative Giese-type reaction of carboxylic acids promoted by visible-light: a sustainable and photoredox neutral protocol.

Authors: Jose Carlos Gonzalez-Gomez and Nieves P. Ramirez

This manuscript has been accepted after peer review and appears as an Accepted Article online prior to editing, proofing, and formal publication of the final Version of Record (VoR). This work is currently citable by using the Digital Object Identifier (DOI) given below. The VoR will be published online in Early View as soon as possible and may be different to this Accepted Article as a result of editing. Readers should obtain the VoR from the journal website shown below when it is published to ensure accuracy of information. The authors are responsible for the content of this Accepted Article.

To be cited as: *Eur. J. Org. Chem.* 10.1002/ejoc.201601478

Link to VoR: <http://dx.doi.org/10.1002/ejoc.201601478>

FULL PAPER

Decarboxylative Giese-type reaction of carboxylic acids promoted by visible-light: a sustainable and photoredox neutral protocol.

Nieves P. Ramirez,^[a] and Jose C. Gonzalez-Gomez^{*[a]}

Dedication ((optional))

Abstract: This work describes a transition-metal free method for decarboxylative generation of radicals from carboxylic acids and their 1,4-addition to Michael acceptors. The Fukuzumi catalyst ([Acr⁺-Mes]) enabled this transformation under visible light irradiation, at room temperature and with CO₂ as the only byproduct. Scope and limitations of this protocol were examined using a range of Michael acceptors (15 examples) and a diverse array of carboxylic acids (18 examples). The use of 3-hydroxypivalic acid in this protocol allowed the straightforward formation of a diastereomerically pure δ -lactone. Moreover, when a homoallylic acid was used, a radical cascade took place with the formation of three C-C bonds.

Introduction

The addition of radicals to α,β -unsaturated carbonyl compounds takes place selectively at the β position without any 1,2-addition. This selectivity in the formation of a C-C bond was first discovered by Giese and has found several applications in organic synthesis.¹ The reaction is very efficient when alkyl radicals (high-lying SOMO) are added to electron-deficient olefins (low-lying LUMO), becoming a good alternative to the Michael reaction.² This strategy benefits from some issues associated to radical processes, such as: (a) carbonyl, hydroxyl and free amino groups are generally tolerated; (b) since free-radicals are neutral species, they are poorly solvated and tertiary radicals can be used to generate quaternary centers.

Carboxylic acids have gathered much attention as radical sources since they are abundant, non-toxic and renewable feedstocks, which make them ideal starting materials in organic synthesis.³ Moreover, radicals are produced upon decarboxylation and the elimination of CO₂, as a traceless by-product, does not impact on their reactivity. Decarboxylative radical generation dated back to the classical works of Kolbe, Hunsdiecker and Barton, among others.⁴ In the last two decades, the use of transition-metal catalysts has expanded the scope of this strategy, but still high temperatures are required and only scarce examples of C_{sp3}-CO₂H decarboxylation were reported for cross-coupling reactions.⁵

Very recently, visible-light-promoted photoredox catalysis has emerged as a powerful and sustainable tool for the generation of radicals from carboxylic acids at room temperature.⁶ Interestingly, if a photoexcited catalyst is oxidant enough to remove an electron from a carboxylate anion [E(RCOO[•]/RCOO⁻) \approx +1.1 to +1.5 V vs SCE],⁷ after rapid decarboxylation of aliphatic acyloxy radicals,⁸ this radical can be added to electron-poor alkenes and the new α -acyl radical could be reduced [E(R[•]/R⁻) \approx -0.7 to -0.6 V vs SCE]⁹ and protonated. In this last step, the 1,4-adduct is delivered, while regenerating the photocatalyst and also the base (Scheme 1a). Since the process is redox-neutral, stoichiometric amounts of chemicals and wastes (except CO₂) are avoided, being a valuable strategy to increase molecular complexity within the principles of "Green-Chemistry".¹⁰ This strategy was first put into practice by Yoshimi's group, using 10 mol % of 1,4-dicyanonaphthalene and phenanthrene as a catalytic system, but still using UV light and very dilute conditions.¹¹ Almost at the same time, Nishibayashi and coworkers found an iridium photocatalyst [Ir(ppy)₂bpy]⁺ competent for the visible-light-promoted decarboxylative addition of arylacetic acids to Michael acceptors.¹² Soon after, fine tuning of the iridium photocatalyst structure and reaction conditions allowed to MacMillan's group to find a protocol for this transformation amenable to a wide range of carboxylic acids and electron-deficient olefins.¹³ Very recently, iridium-based photocatalysts have also been successful in the decarboxylative Giese-type reaction of α -aminoacids.¹⁴ Given the low abundance of iridium in earth, iridium catalysts are expensive. Consequently, cheaper and more sustainable catalysts are desired for this transformation. In this context, the group of Koike and Akita has reported that organic [Acr-Mes]ClO₄ (Fukuzumi catalyst)¹⁵ is also a competent photocatalyst in this neutral photoredox process. However, only four acids were examined with the same Michael acceptor and the efficiency of the reaction was rather low for secondary acids (Scheme 1b).¹⁶ Encouraged by these results, we decided to reexamine the reaction conditions in order to improve the applicability of this organic photoredox catalyst in the decarboxylative alkylation of carboxylic acids.¹⁷

Results and Discussion

Giving the strong oxidant capacity of the Fukuzumi catalyst,¹⁵ it is not surprising that carboxylic acids, or more efficiently the corresponding carboxylates, can act as reductive quenchers of this photocatalyst. Following the general mechanism for decarboxylative alkylation depicted in Scheme 1a, the turnover of [Acr⁺-Mes] is proposed to occur by ET from the [Acr⁺-Mes] to the


[a] Departamento de Química Orgánica, Facultad de Ciencias and Instituto de Síntesis Orgánica (ISO), Universidad de Alicante, Apdo. 99, 03080 Alicante, Spain.

E-mail: josecarlos.gonzalez@ua.es.


Supporting information for this article is given via a link at the end of the document.

FULL PAPER

α -acyl radical. Since the acridinyl radical is a fairly weak reductant [$E_{1/2}(\text{PC}^+/\text{PC}^\bullet) = -0.57 \text{ V vs SCE}$],¹⁵ it was not clear at the outset of this work that a range of α -substituted alkyl radicals could efficiently regenerate the photocatalyst. For example, according to the reported reduction potential of $\text{CH}_3\text{CH}^\bullet\text{-CO}_2\text{Me}$ ($E_{1/2} = -0.66 \text{ V vs SCE}$),⁹ the reaction with the acridinyl radical would operate against a moderate potential gradient (about 100 mV). We reasoned that a final fast and thermodynamically favor protonation would compensate this issue for some acceptors.









(b) Previous work from Akita's group (Reference 16)



Scheme 1. (a) Plausible mechanism for a photoredox neutral decarboxylative Giese-type reaction catalyzed by $[\text{Acr}^+-\text{Mes}]$ and base. (b) Previous work.

With the precedents above mentioned, we decided to screen different Michael acceptors with adamantane-1-carboxylic acid (**1a**). The results of this study are shown in Scheme 2. It is worth mentioning that due to the steric bulkiness of adamantane, radical chemistry is very convenient for its derivatization and this moiety has been quintessential for the development of new drugs and catalysts.¹⁸ With apparently minor modifications from Akita's conditions (Scheme 1b), we [0.2 equiv of Na_2CO_3 , under air atmosphere and using one bulb of 12 x 1 W blue leds and 0.3 M of **2**] were able to isolate compound **3aa** in similar excellent yields but in a significant shorter time (19 h vs 60 h). Importantly, careful deoxygenation and the use of inert atmosphere were not necessary to achieve this goal, simplifying this procedure even more. Under these conditions, other double activated Michael acceptors gave the corresponding products (**3ab-3ae**) in very good yields after only 15 to 21 h of irradiation. Notably, 4-chromenone (**2f**) and butyl methacrylate (**2g**) afforded compounds **3af** and **3ag** in good yields, although longer reaction times were necessary. Apparently, the push-pull effect in 4-chromenone and the increased stability of the α -Me- α -acyl radical (or its higher nucleophilicity) with methacrylate acceptor, had a positive impact on the reactivity. We were pleased to find that acyclic (**2h-2k**) and cyclic (**2l, 2m**) α,β -unsaturated ketones

furnished the expected products (**3ah-3am**) in very good isolated yields after reasonable reaction times (12 – 36 h). Vinylsulfone **2n** was also a suitable substrate under the reaction conditions to obtain **3an** in good yield and, more importantly, compound **3ao** was obtained in very good yield after only 14 h when crotonaldehyde (**2o**) was used as acceptor. On the contrary, sterically demanding acceptor **2p** failed to add the bulky adamantane moiety. In addition, substrates substituted at α -position by a single ester group (**2q, 2r** and **2s**) also failed to react, probably because the α -acyl radical is unable to reoxidize $[\text{Acr}^+-\text{Mes}]$. Moreover, (*E*)-chalcone (**2t**) failed to give any product under the same reaction conditions, likely due to the competitive generation of a stable benzylic radical that is not capable to recycle the catalyst.



Scheme 2 Scope of electron-deficient alkenes. ^aIsolated pure product. ^b1.7:1 MeOH/EtOAc (0.185 M) was used as solvent mixture to improve the solubility of substrates. ^c3 equiv of volatile MVK were used.

Regarding the carboxylic acid partners, it was reported that the reaction of secondary substrates examined (**1e** and **1f**) under the conditions disclosed by Koike and Akita never went to completion (Scheme 1b).¹⁶ Consequently, we decided to examine this issue in more detail using cyclohexane carboxylic acid (**1e**) and diethyl

FULL PAPER

2-ethylidene malonate (**2a**) as acceptor in a model reaction. The results of this study are summarized in Table 1.



Table 1. Optimization of reaction conditions.

1e (0.30 M)	2a (0.20 M)	3ea
Entry	Modification from standard conditions	Yield (%) ^a
1	none	92
2	in MeOH (0.3 M)	77
3	in CF ₃ CH ₂ OH (0.3 M)	0
4	in <i>t</i> -BuOH (0.3 M)	12
5	in [CH ₂ Cl] ₂ /MeOH (2:1)	72
6	in Me ₂ CO/H ₂ O (2:1)	32
7	in MeCN/H ₂ O (1:1)	20
8	in MeCN/H ₂ O (3:1)	50
9	under argon atmosphere	59
10	under oxygen atmosphere (1 atm)	71
11	1e (1 equiv) : 2a (1 equiv)	57
12	1e (1 equiv) : 2a (1.5 equiv)	58
13	blue LED 2 bulbs (12 W)	89
14	white LED bulb (13.5 W)	77
15	Sunlight (10 h)	70 ^b
16	CFL light bulb (18 W)	0
17	in the dark	0
18	5 mol % of photocatalyst (10 h)	100
19	1 mol % photocatalyst	25
20	without photocatalyst	0
21	with 20 mol % of Cs ₂ CO ₃	90
22	with 20 mol % of DBU	55
23	without base	23

[a] ^aYield determined by GC using adamantane as internal standard. [b] See SI for details. DBU = 1,8-diazabicyclo[5.4.0]undec-7-ene.

After some experimentation, we have found that the solvent system has a significant impact on the reaction (entries 1-9), obtaining the best results in 2:1 MeCN/H₂O and observing some competitive Michael addition of MeOH when it is used as solvent. To our delight, when the reaction was run under air atmosphere, it was completed after only 17 h (entry 1). Surprisingly, an argon atmosphere (entry 9) had an even more deleterious effect on the yield than an oxygen atmosphere (entry 10).¹⁹ Regarding the stoichiometry, a slight excess of acid (1.5 equiv.) was found beneficial (entries 1, 11 and 12) and it was also helpful to work under concentrated conditions (0.2 – 0.3 M). It was clearly demonstrated that the reaction progress depends highly on the source of light (entries 1, 13-17), obtaining optimal results with intense blue LEDs (12 x 1W). This could also explain the difference in the reaction rate observed under Akita's conditions (3W blue LEDs, Scheme 1b) and ours. Since it has been recently reported that the reductive quenching of [Acr⁺-Mes]⁺ by carboxylates has low efficiency, we examined the use of two blue

LED bulbs, but the yield of the reaction was similar (entry 13 vs 1).²⁰ Control experiments revealed that the reaction absolutely requires the photocatalyst (entry 20). In addition, while full conversion is achieved after only 10 h using 5 mol % of catalyst (entry 18), similar results are obtained with 2.5 mol % after 17 h and the reaction is significantly slower when the load of photocatalyst is reduced even further (entry 19). It was also found that the reaction progress poorly without base (entry 23). The use of 20 mol % of Cs₂CO₃ leads to similar results than with Na₂CO₃, while lower yields were achieved with DBU (entries 21 and 22).




Scheme 3. Scope of acids. ^aIsolated pure product. ^b2 equivalents of acid **1** were used. ^cAnother portion of photocatalyst (2.5 mol %) was added after 48 h. ^dAnother portion of acid **1** (1.5 equiv.) was added after 48 h. ^eSee SI for details.

Having found these new optimized conditions, we examined the scope of acids with this protocol, using acceptor **2a** (Scheme 3). Tertiary carboxylic acids, including natural clofibric acid (**1d**), participate efficiently in the reaction (products **3ba-3da**) to generate quaternary centers. Secondary cyclic and acyclic acids were also suitable substrates with this protocol (**1e-1k**). Notably,

FULL PAPER


secondary α -oxyacids **1h** and **1i** furnished the corresponding conjugate adducts (**3ha** and **3ia**) in good to excellent yields. Since ethers are formed, this protocol represents a good alternative to the classic Williamson ether synthesis. In addition, natural *N*-Boc protected secondary α -aminoacids provide the corresponding α -alkylamines (**3ja** and **3ka**) in synthetically useful yields. Primary α -amino (from glycine derivative **1l**) and α -oxyacids (from phenoxyacetic derivatives **1m** and **1n**) are also suitable substrates. This procedure allowed the formation of aminomethyl derivative **3la** and aryloxymethyl derivative **3ma** and **3na** in good to excellent yields. Importantly, for phenoxyacetic derivatives, electron-donating groups in the aromatic ring accelerate the reaction (**1m** vs **1n**). Remarkably, α -ketoacids were well tolerated, not only aromatic 2-oxophenylacetic acid (**1o**), but also the aliphatic 2-oxoethyl acetic acid (**1p**), gave the corresponding products (**3oa** and **3pa**) in reasonable good yields. This method to build 1,4-dicarbonyl compounds is complementary to the classic Stetter reaction.²¹ Very recently, the visible-light mediated decarboxylative 1,4-addition of a range of aromatic α -ketoacids to different Michael acceptors was successfully accomplished using an iridium photocatalyst.²² Although our protocol furnished compound **3pa** in a moderate yield, we have not found in the literature other examples for the decarboxylative coupling of aliphatic α -ketoacids with Michael acceptors. It is worth mentioning that (*S*)-lactic acid (**1q**) failed in this protocol, most likely due to formation of acetaldehyde after oxidative decarboxylation. Electron-poor phenoxyacetic derivatives (such as **1r**), as well as common primary aliphatic acids (**1s**) and arylacetic acids (e.g. **1t**) failed to provide the conjugate adduct in synthetically useful yields.²³



Scheme 4. Mechanistic studies

To gain insight into the reaction pathway, we confirmed that the reaction of **1e** and **2a** was completely inhibited in the presence of TEMPO (Scheme 4a) and cyclohexyl radical was trapped to form compound **4** (detected by LC-MS, see SI). Moreover, when the reaction was conducted in $\text{CD}_3\text{CN}/\text{H}_2\text{O}$, incorporation of deuterium was not observed in product **3ea**, excluding the possibility of hydrogen abstraction from the solvent (Scheme 4b).

In contrast, high content of deuterium was observed for the product when D_2O was used as co-solvent. This result is in accordance with a single electron reduction of the α -acyl radical and final deuteration with D_2O and, in a minor extend, protonation with the *in-situ* formed small amount of H_2O to regenerate the base (Scheme 1).



Scheme 5. Synthetic applications.

Some synthetic applications accomplished with the developed protocol are depicted in Scheme 5. When hydroxypivalic acid (**1r**) was used under standard conditions, the free hydroxyl group was well tolerated and after conjugated addition, the lactonization took place smoothly to afford compound **3ra** as a single diastereoisomer (according to ^1H - and ^{13}C -NMR). Since poor diastereoselection was observed for other studied examples, we think that this should be the result of a thermodynamic equilibration by deprotonation-protonation at C_3 . In addition, when acid **1q** was used in our decarboxylative alkylation protocol with acceptor **2a**, compound **5** was isolated in very good yield, as well

FULL PAPER

as minor amounts of disulfide **6** (Scheme 5b). Based on the radical nature of this process, we presume that after decarboxylation, β -fragmentation takes place efficiently to release isobutene and the aromatic thiyl radical, which is quickly trapped by acceptor **2a**.²⁴ Finally, we took advantage of our protocol to develop a radical cascade where two molecules of acceptor **2a** are incorporated in the product and three new σ C-C bonds are formed (Scheme 5c). For this transformation, homoallylic acid **1s** was put into react with acceptor **2a** (2.5 equiv.) under standard conditions, after decarboxylation and radical addition, fast 5-exo-trig cyclization took place to generate a primary nucleophilic radical that was trapped by another molecule of Michael acceptor. Upon formation of the new electrophilic radical, electron transfer to [Acr⁺-Mes] could occur, regenerating the photocatalyst, and final protonation allowed formation of compound **7** in good yield. Remarkably, only two diastereoisomers out of four are formed (GC-MS, ¹H, ¹³C-NMR). It seems reasonable that the diastereocontrol is achieved in the cyclization step, following a chairlike transition state akin to the Beckwith stereoelectronic model.²⁵

Conclusions

In conclusion, we have demonstrated the applicability of visible-light induced decarboxylative Giese-type reaction, catalysed by [Acr⁺-Mes], for a range of substrates. The transformation is redox neutral and the only residue is CO₂. Regarding the acceptor, not only alkylidene malonates/malonitriles were suitable substrates, but also α -substituted acrylates, conjugated ketones/aldehydes and conjugated sulfones. The scope of carboxylic acids includes mainly tertiary and secondary substrates diversely substituted, as well as α -ketoacids and primary α -amino and α -oxy acids. Compare to other existing methods, some salient features of this protocol are: (a) is free of noble-metal catalysts; (b) only 20 mol % of cheap Na₂CO₃ is used; (c) an aqueous solvent mixture is used and neither deoxygenation nor inert atmosphere is required. We hope that this work contributes to the transformation of carboxylic acids –renewable feedstocks– into pharmaceuticals or fine chemicals using Green Chemistry.

Experimental Section

General Remarks: TLCs were performed on silica gel 60 F₂₅₄, using aluminium plates and visualized by exposure to ultraviolet light or using different stains (PMA, KMnO₄ or Ninhydrin). Flash chromatographies (FC) were carried out on handpacked columns of silica gel 60 (230 – 400 mesh). Infrared (IR) analysis was performed with a spectrophotometer equipped with an ATR component; wavenumbers are given in cm⁻¹. LRMS were performed in a mass spectrometer coupled with a gas chromatographer (GC); the mobile phase was helium (2 mL/min); HP-1 column of 12 m was used; temperature program starts at 80 °C for 3 min, then up to 270 °C with a rate of 20 °C/min, and 17.5 min at 270 °C. HRMS analyses were carried out using the Electron Impact (EI) mode at 70 eV by Q-TOF. ¹H NMR spectra were recorded at 300 or 400 MHz for ¹H-NMR and 75 or 100 MHz for ¹³C-NMR, using CDCl₃ as solvent and TMS as an internal standard (0.00 ppm). ¹³C-NMR spectra were recorded with ¹H-decoupling at 100 MHz and referenced to CDCl₃ at 77.16 ppm.

General procedure A (GPA) for the preparation of adamantane derivatives (Scheme 3): In a microwave tube, equipped with a stirring bar, were introduced adamantane-1-carboxylic acid (**1a**, 99 mg, 0.55 mmol, 1.10 equiv.), Na₂CO₃ (4.06 mg, 0.10 mmol, 20 mol %) and [Acr-Mes]ClO₄ (5.2 mg, 2.5 mol %), followed by a solution of the desired Michael acceptor (0.50 mmol) in MeOH (1.7 mL). The yellow solution was irradiated using blue LED's and stirred at room temperature, without any inert atmosphere, until complete conversion was observed (monitored by TLC and/or GC). The reaction mixture was concentrated under reduced pressure and the residue was purified by column chromatography.

Diethyl 2-((3*r*, 5*r*, 7*r*)-adamantan-1-yl)ethyl)malonate (3aa**, Scheme 2):¹³ Prepared according to GPA, after 19 h. It was purified by FC (Hexane/EtOAc 97:3 to 90:10) and obtained as a colorless oil (128 mg, 0.40 mmol, 80%); TLC *R*_f 0.12 (Hexane/EtOAc 97:3); IR ν 2908, 2848, 1756, 1727, 1447, 1147, 1032, 733 cm⁻¹; ¹H-NMR (300 MHz, CDCl₃) δ 4.19 (q, *J* = 7.1 Hz, 2H), 4.18 (q, *J* = 7.1 Hz, 2H), 3.57 (d, *J* = 5.2 Hz, 1H), 2.08 (dd, *J* = 7.4, 5.5 Hz, 1H), 1.97 (br s, 3H), 1.71 - 1.59 (m, 7H), 1.52 (br s, 5H), 1.28 (t, *J* = 7.1 Hz, 3H), 1.27 (t, *J* = 7.1 Hz, 3H), 0.99 (d, *J* = 7.3 Hz, 3H); ¹³C-NMR (75 MHz, CDCl₃) δ 170.5 (C), 169.9 (C), 61.4 (CH₂), 61.0 (CH₂), 52.0 (CH), 43.2 (CH), 39.5 (3 \times CH₂), 37.1 (3 \times CH₂), 35.3 (C), 28.7 (3 \times CH), 14.19 (CH₃), 14.15 (CH₃), 10.5 (CH₃); GC *R*_T 19.896 min; LRMS (EI) *m/z* (%) = 277 (M⁺-C₂H₅O, 4), 276 (8), 136 (11), 135 (100), 93 (10).**

(Diethyl 2-((3*r*, 5*r*, 7*r*)-adamantan-1-yl(phenyl)methyl)malonate (3ab**, Scheme 2):** Prepared according to GPA, after 21 h. It was purified by FC (Hexane/EtOAc 99:1 to 85:15) and obtained as a pale yellow oil (166 mg, 0.43 mmol, 86%); TLC *R*_f 0.19 (Hexane/EtOAc 97:3); IR ν 2905, 2847, 1752, 1724, 1257, 729, 684 cm⁻¹; ¹H-NMR (300 MHz, CDCl₃) δ 7.26 - 7.07 (m, 5H), 4.32 - 4.14 (m, 2H), 4.00 (d, *J* = 10.8 Hz, 1H), 3.77 - 3.61 (m, 2H), 3.34 (d, *J* = 10.8 Hz, 1H), 1.91 (br s, 3H), 1.66 - 1.46 (m, 12H), 1.30 (t, *J* = 7.1 Hz, 3H), 0.78 (t, *J* = 7.1 Hz, 3H); ¹³C-NMR (75 MHz, CDCl₃) δ 169.7 (C), 168.5 (C), 138.9 (C), 127.3 (2 \times CH), 126.6 (3 \times CH), 61.8 (CH₂), 61.3 (CH₂), 56.2 (CH), 53.7 (CH), 40.0 (3 \times CH₂), 36.8 (3 \times CH₂), 36.2 (C), 28.7 (3 \times CH), 14.1 (CH₃), 13.5 (CH₃); GC *R*_T 25.638 min; LRMS (EI) *m/z* (%) = 384 (M⁺, 2), 136 (11), 135 (100), 93 (6); HRMS (EI) *Calcd.* for C₂₄H₃₂O₄ 384.2301, found 384.2302.

Diethyl 2-((3*r*, 5*r*, 7*r*)-adamantan-1-yl(4-methoxyphenyl)methyl)malonate (3ac**, Scheme 2):** Prepared according to GPA, after 15 h. It was purified by FC (Hexane/EtOAc 95:5 to 85:15) and obtained as a white solid (141 mg, 0.34 mmol, 68%); TLC *R*_f 0.15 (Hexane/EtOAc 97:3); IR ν 2971, 2888, 2300, 1767, 1716, 1457, 1309, 1194, 1047, 1033, 761 cm⁻¹; ¹H-NMR (300 MHz, CDCl₃) δ 7.04 (d, *J* = 8.1 Hz, 2H), 6.76 (d, *J* = 8.8 Hz, 2H), 4.31 - 4.13 (m, 2H), 3.96 (d, *J* = 10.7 Hz, 1H), 3.77 (s, 3H), 3.76 - 3.68 (m, 2H), 3.29 (d, *J* = 10.7 Hz, 1H), 1.91 (s, 3H), 1.70 - 1.42 (m, 12H), 1.29 (t, *J* = 7.1 Hz, 3H), 0.84 (t, *J* = 7.1 Hz, 3H); ¹³C-NMR (75 MHz, CDCl₃) δ 169.8 (C), 168.6 (C), 158.2 (C), 130.9 (C), 112.7 (4 \times CH), 61.8 (CH₃), 61.2 (CH₂), 55.4 (CH₂), 55.2 (CH), 53.7 (CH), 40.0 (3 \times CH₂), 36.9 (3 \times CH₂), 36.3 (C), 28.7 (3 \times CH₃), 14.1 (CH₃), 13.7 (CH₃); GC *R*_T 16.558 min; LRMS (EI) *m/z* (%) = 414 (M⁺, 18), 279 (16), 278 (18), 255 (13), 136 (11), 135 (100); HRMS (EI) *Calcd.* for C₂₅H₃₄O₅ 414.2406, found 414.2414.

2-((3*r*, 5*r*, 7*r*)-adamantan-1-yl(phenyl)methyl)malononitrile (3ad**, Scheme 2):** Prepared according to GPA, after 21 h, but in this case MeOH/EtOAc (1.7 mL/1 mL) was the solvent mixture to improve the solubility of benzyldenemalonitrile. It was purified by FC (Hexane/EtOAc 97:3 to 85:15) and obtained as a white solid (93 mg, 0.32 mmol, 64%); TLC *R*_f 0.08 (Hexane/EtOAc 97:3); IR ν 2937, 2900, 1713, 1048, 703 cm⁻¹; ¹H-NMR (300 MHz, CDCl₃) δ 7.38 (s, 5H), 4.25 (d, *J* = 5.4 Hz, 1H), 2.80 (d, *J* = 5.4 Hz, 1H), 2.02 (s, 3H), 1.78 - 1.48 (m, 12H); ¹³C-NMR (75 MHz, CDCl₃) δ 135.4 (CH), 129.8 (C), 128.68 (2 \times CH), 128.6 (2 \times CH), 113.6 (C), 113.4 (C), 58.1 (CH), 40.5 (3 \times CH₂), 36.6 (C), 36.4 (3 \times CH₂), 28.5 (3 \times CH₃), 23.9 (CH); GC *R*_T 22.913 min; LRMS (EI) *m/z* (%) = 290 (M⁺, 1),

FULL PAPER

136 (11), 135 (100), 93 (10); HRMS (EI) *Calcd.* for $C_{20}H_{22}N_2$ 290.1783, found 290.1782.

Ethyl 4-((3*r*, 5*r*, 7*r*)-adamantan-1-yl)-2-oxochroman-3-carboxylate (3ae, Scheme 2): Prepared according to GPA, after 16 h, but in this case the solvent mixture was MeOH/EtOAc (1.7 mL/1 mL) to improve the solubility of ethyl 2-oxo-2*H*-chromene-3-carboxylate. It was purified by FC (Hexane/EtOAc 95:5 to 90:10) and obtained as a white solid (124 mg, 0.35 mmol, 70%, >98:2 *dr* according to $^1\text{H-NMR}$): TLC *R_f* 0.06 (Hexane/EtOAc 97:3); IR ν 2971, 2888, 2300, 1767, 1716, 1457, 1309, 1194, 1033, 761 cm^{-1} ; $^1\text{H-NMR}$ (300 MHz, CDCl_3) δ 7.34–7.25 (m, 1H), 7.15–7.05 (m, 3H), 4.12–4.04 (m, 1H), 4.02 (d, J = 1.3 Hz, 1H), 4.00–3.91 (m, 1H), 2.96 (d, J = 0.7 Hz, 1H), 1.98 (br s, 3H), 1.70–1.52 (m, 12H), 0.98 (t, J = 7.1 Hz, 3H); $^{13}\text{C-NMR}$ (75 MHz, CDCl_3) δ 168.3 (C), 166.2 (C), 151.8 (C), 131.4 (CH), 129.0 (CH), 124.0 (CH), 120.6 (C), 117.0 (CH), 62.3 (CH₂), 51.7 (CH), 47.2 (CH), 39.5 (3 \times CH₂), 36.6 (3 \times CH₂), 36.2 (C), 28.4 (3 \times CH), 13.9 (CH₃); GC *R_T* 23.503 min; LRMS (EI) m/z (%) = 282 (M^+ + $\text{C}_3\text{H}_5\text{O}_2$, 4), 281 (M^+ - $\text{C}_3\text{H}_5\text{O}_2$, 1), 136 (12), 135 (100), 93 (12), 91 (10), 79 (10); HRMS (EI) *Calcd.* for $\text{C}_{22}\text{H}_{26}\text{O}_4$ 354.1831, found 354.1812.

2-((3*r*, 5*r*, 7*r*)-adamantan-1-yl)chroman-4-one (3af, Scheme 2):²⁶ Prepared according to GPA, after 67 h. It was purified by FC (Hexane/EtOAc 97:3 to 95:5) and obtained as a colorless oil (105 mg, 0.37 mmol, 74%); TLC *R_f* 0.22 (Hexane/EtOAc 97:3); IR ν 2908, 2859, 1678, 1601, 1461, 1313, 1229, 1038, 752 cm^{-1} ; $^1\text{H-NMR}$ (300 MHz, CDCl_3) δ 7.86 (dd, J = 8.1, 1.8 Hz, 1H), 7.46 (ddd, J = 8.5, 7.2, 1.8 Hz, 1H), 7.02–6.94 (m, 2H), 3.91 (dd, J = 13.6, 3.1 Hz, 1H), 2.67 (qd, J = 16.5, 8.3 Hz, 2H), 2.06 (br s, 3H), 1.94–1.40 (m, 12H); $^{13}\text{C-NMR}$ (75 MHz, CDCl_3) δ 193.9 (C), 162.4 (C), 135.9 (CH), 127.0 (CH), 121.0 (C + CH), 118.0 (CH), 85.6 (CH), 37.9 (3 \times CH₂), 37.24 (CH₂), 37.17 (3 \times CH₂), 36.0 (C), 28.3 (3 \times CH); GC *R_T* 23.746 min; LRMS (EI) m/z (%) = 283 (M^+ + 1, 19), 282 (M^+ , 94), 281 (M^+ - 1, 15), 147 (50), 136 (11), 135 (100), 121 (12), 120 (10), 93 (14), 92 (10), 91 (12), 79 (12).

Butyl 3-((3*r*, 5*r*, 7*r*)-adamantan-1-yl)-2-methylpropanoate (3ag, Scheme 2): Prepared according to GPA, after 70 h. It was purified by FC (Hexane/EtOAc 97:3 to 95:5) and obtained as a colorless oil (68 mg, 0.25 mmol, 50%); TLC *R_f* 0.43 (Hexane/EtOAc 97:3); IR ν 2957, 2898, 2846, 1727, 1451, 1181, 1121, 1095 cm^{-1} ; $^1\text{H-NMR}$ (300 MHz, CDCl_3) δ 4.12–3.98 (m, 2H), 2.54 (dq, J = 14.1, 7.1, 3.1 Hz, 1H), 1.92 (brs, 3H), 1.79–1.55 (m, 9H), 1.54–1.33 (m, 8H), 1.13 (d, J = 7.1 Hz, 3H), 1.01 (dd, J = 14.2, 3.1 Hz, 1H), 0.94 (t, J = 7.3 Hz, 3H); $^{13}\text{C-NMR}$ (75 MHz, CDCl_3) δ 178.3 (C), 64.2 (CH₂), 48.7 (CH₂), 42.4 (3 \times CH₂), 37.1 (3 \times CH₂), 34.5 (CH), 32.8 (C), 30.8 (CH₂), 28.8 (3 \times CH), 20.7 (CH₂), 19.3 (CH₃), 13.9 (CH₃); GC *R_T* 17.869 min; LRMS (EI) m/z (%) = 278 (M^+ , 5), 143 (12), 136 (12), 135 (100); HRMS (EI) *Calcd.* for $\text{C}_{18}\text{H}_{30}\text{O}_2$ - $\text{C}_4\text{H}_9\text{O}$ 204.1514, found 204.1517.

4-((3*r*, 5*r*, 7*r*)-adamantan-1-yl)pentan-2-one (3ah, Scheme 2): Prepared according to GPA, after 12 h, but in this case the reaction was performed using 1-adamantanecarboxylic acid (90 mg, 0.50 mmol) and 3-penten-2-one (103 μL , 0.75 mmol, 1.5 equiv.). It was purified by FC (Hexane/EtOAc 95:5) and obtained as a colorless oil (88 mg, 0.40 mmol, 79%); TLC *R_f* 0.38 (Hexane/EtOAc 97:3); IR ν 1707, 1255, 723, 703 cm^{-1} ; $^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 2.59 (dd, J = 15.7, 2.6 Hz, 1H), 2.14 (s, 3H), 2.06 (dd, J = 15.7, 10.3 Hz, 3H), 1.97 (br s, 3H), 1.74–1.57 (m, 6H), 1.48 (s, 5H), 0.79 (d, J = 6.9 Hz, 3H); $^{13}\text{C-NMR}$ (101 MHz, CDCl_3) δ 210.2 (C), 45.5 (CH₂), 39.4 (3 \times CH₂), 39.0 (CH₃), 37.4 (3 \times CH₂), 34.4 (C), 30.5 (CH), 28.8 (3 \times CH), 13.7 (CH₃); GC *R_T* 16.080 min; LRMS (EI) m/z (%) = 220 (M^+ , 1), 202 (15), 136 (11), 135 (100), 93 (14), 79 (14); HRMS (EI) *Calcd.* for $\text{C}_{15}\text{H}_{24}\text{O}$ 220.1827, found 220.1836.

4-((3*r*, 5*r*, 7*r*)-adamantan-1-yl)-3-methylpentan-2-one (3ai, Scheme 2): Prepared according to GPA, after 23 h. It was purified by FC

(Hexane/EtOAc 97:3 to 90:10) and obtained as a pale yellow oil (70.2 mg, 0.30 mmol, 60%, >98:2 *dr* according to NMR and GC experiments): TLC *R_f* 0.22 (Hexane/EtOAc 97:3); IR ν 2908, 2849, 1702, 1442, 1342, 1087, 1048, 871 cm^{-1} ; $^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 2.18 (qd, J = 7.1, 3.4 Hz, 1H), 2.17 (s, 3 H), 1.97 (br s, 3H), 1.73–1.47 (m, 12H), 1.32–1.23 (m, 1H), 1.13 (d, J = 7.1 Hz, 3H), 0.95 (d, J = 7.4 Hz, 3H); $^{13}\text{C-NMR}$ (101 MHz, CDCl_3) δ 213.9 (C), 48.7 (CH), 46.3 (CH), 40.3 (3 \times CH₂), 37.3 (3 \times CH₂), 36.1 (C), 30.1 (CH₃), 28.8 (3 \times CH), 18.3 (CH₃), 10.4 (CH₃); GC *R_T* 16.223 min; LRMS (EI) m/z (%) = 216 (M^+ - H_2O , 11), 136 (11), 135 (100), 93 (10), 79 (10); HRMS (EI) *Calcd.* for $\text{C}_{16}\text{H}_{26}\text{O}$ 234.1984, found 234.1983.

4-((3*r*, 5*r*, 7*r*)-adamantan-1-yl)butan-2-one (3aj, Scheme 2):²⁷ Prepared according to GPA, after 36 h, but in this case the reaction was performed using 1-adamantanecarboxylic acid (90 mg, 0.50 mmol) and methyl vinyl ketone (125 μL , 1.5 mmol, 3 equiv.). It was purified by FC (Hexane/EtOAc 97:3 to 90:10) and obtained as a colorless oil (83 mg, 0.40 mmol, 80%); TLC *R_f* 0.24 (Hexane/EtOAc 97:3); IR ν 2902, 2844, 1713, 1269, 737, 700 cm^{-1} ; $^1\text{H-NMR}$ (300 MHz, CDCl_3) δ 2.44–2.31 (m, 2H), 2.15 (s, 3H), 1.95 (br s, 3H), 1.76–1.54 (m, 6H), 1.46–1.45 (m, 6H), 1.40–1.30 (m, 2H); $^{13}\text{C-NMR}$ (101 MHz, CDCl_3) δ 210.2 (C), 42.3 (3 \times CH₂), 37.9 (CH₂), 37.6 (CH₂), 37.2 (3 \times CH₂), 31.9 (C), 30.0 (CH₃), 28.8 (3 \times CH); GC *R_T* 15.479 min; LRMS (EI) m/z (%) = 206 (M^+ , 1), 189 (5), 188 (33), 136 (11), 135 (100), 93 (13), 79 (13).

3-((3*r*, 5*r*, 7*r*)-adamantan-1-yl)-1-phenylbutan-1-one (3ak, Scheme 2): Prepared according to GPA, after 17 h. It was purified by FC (Hexane/EtOAc 97:3 to 85:15) and obtained as a pale yellow oil (90 mg, 0.32 mmol, 64%); TLC *R_f* 0.44 (Hexane/EtOAc 97:3); IR ν 2898, 2849, 1678, 1442, 1205, 999, 743, 674 cm^{-1} ; $^1\text{H-NMR}$ (300 MHz, CDCl_3) δ 8.05–7.88 (m, 2H), 1H), 7.58–7.51 (m, 1H), 7.50–7.41 (m, 2H), 3.16 (dd, J = 15.4, 2.9 Hz, 1H), 2.57 (dd, J = 15.4, 10.4 Hz, 1H), 2.00 (br s, 3H), 1.89–1.78 (m, 1H), 1.76–1.59 (m, 6H), 1.57–1.56 (m, 6H), 0.83 (d, J = 6.8 Hz, 3H); $^{13}\text{C-NMR}$ (101 MHz, CDCl_3) δ 201.7 (C), 137.6 (C), 132.9 (CH), 128.6 (2 \times CH), 128.3 (2 \times CH), 40.2 (CH₂), 39.6 (3 \times CH₂), 37.4 (3 \times CH₂), 34.8 (C + CH), 28.8 (3 \times CH), 13.7 (CH₃); GC *R_T* 22.617 min; LRMS (EI) m/z (%) = 282 (M^+ , 10), 162 (18), 147 (22), 146 (13), 136 (11), 135 (100), 105 (19), 93 (11), 79 (11), 77 (17); HRMS (EI) *Calcd.* for $\text{C}_{20}\text{H}_{26}\text{O}$ 282.1984, found 282.1999.

3-((3*r*, 5*r*, 7*r*)-adamantan-1-yl)cyclohexanone (3al, Scheme 2):²⁸ Prepared according to GPA, after 28 h. It was purified by FC (Hexane/EtOAc 97:3 to 90:10) and obtained as a pale yellow oil (108 mg, 0.46 mmol, 92%); TLC *R_f* 0.31 (Hexane/EtOAc 97:3); IR ν 2898, 2848, 1701, 1451, 1231, 1058 cm^{-1} ; $^1\text{H-NMR}$ (300 MHz, CDCl_3) δ 2.50–2.03 (m, 5H), 1.98 (br s, 4H), 1.74–1.59 (m, 6H), 1.52–1.51 (m, 6H), 1.41–1.19 (m, 3H); $^{13}\text{C-NMR}$ (75 MHz, CDCl_3) δ 213.5 (C), 49.8 (CH), 42.2 (CH₂), 41.6 (CH₂), 39.5 (3 \times CH₂), 37.3 (3 \times CH₂), 34.5 (C), 28.7 (3 \times CH₃), 25.8 (CH₂), 24.7 (CH₂); GC *R_T* 19.014 min; LRMS (EI) m/z (%) = 232 (M^+ , 4), 136 (11), 135 (100), 93 (10), 79 (10).

3-((3*r*, 5*r*, 7*r*)-adamantan-1-yl)-2-pentylcyclopentan-1-one (3am, Scheme 2): Prepared according to GPA, after 23 h. It was purified by FC (Hexane/EtOAc 98:2 to 95:5) and obtained as a pale-yellow oil (108 mg, 0.375 mmol, 75%, 95:5 *dr* according GC); TLC *R_f* 0.32 (Hexane/EtOAc 97:3); IR ν 2900, 2846, 1734, 1451, 1168, 753 cm^{-1} ; $^1\text{H-NMR}$ (300 MHz, CDCl_3) *mixture of diastereoisomers* δ 2.48–2.04 (m, 3H), 1.98 (brs, 3H), 1.94–1.85 (m, 1H), 1.84–1.57 (m, 9H), 1.54–1.53 (m, 7H), 1.38–1.20 (m, 7H), 0.88 (t, J = 6.9 Hz, 3H); $^{13}\text{C-NMR}$ (75 MHz, CDCl_3) *mixture of diastereoisomers* δ 223.0 (C=O), 220.6 (C=O), 52.8, 51.4, 51.2, 48.6, 41.6, 40.2, 38.2, 37.3, 36.8, 34.9, 34.6, 32.2, 28.8, 28.6, 27.0, 26.3, 26.2, 22.7, 22.6, 20.6, 19.8, 14.2; GC *R_T* 24.300 min and 24.515 min (*two diastereoisomers*); LRMS (EI) m/z (%) (*Diastereoisomer A*) = 218 (M^+ - C_5H_{11} , 31), 136 (11), 135 (100), 93 (16), 83 (28), 82 (98), 79 (16); m/z (%) (*Diastereoisomer B*) = 218 (M^+ - C_5H_{11} , 18), 136 (11), 135 (100), 93 (12),

FULL PAPER

83 (16), 82 (63), 79 (13); HRMS (EI) *Calcd.* for $C_{20}H_{32}O$ - C_8H_{10} 218.1671, found 218.1678.

(3*r*, 5*r*, 7*r*)-1-(2-(phenylsulfonyl)ethyl)adamantane (3an, Scheme 2):²⁹ Prepared according to *GPA*, after 36 h. It was purified by FC (Hexane/EtOAc 95:5 to 80:20) and obtained as a white solid (102 mg, 0.335 mmol, 67%): TLC *R_f* 0.10 (Hexane/EtOAc 97:3); IR ν 1734, 1373, 1237, 1046, 914, 730 cm^{-1} ; 1H -NMR (300 MHz, $CDCl_3$) δ 7.97 - 7.96 (m, 2H), 7.70 - 7.63 (m, 1H), 7.61 - 7.54 (m, 2H), 3.06 (d, J = 17.3 Hz, 1H), 3.06 (dd, J = 5.1, 3.8 Hz, 1H), 1.94 (br s, 3H), 1.75 - 1.53 (m, 6H), 1.52 - 1.43 (m, 2H), 1.41 - 1.40 (m, 6H); ^{13}C -NMR (75 MHz, $CDCl_3$) δ 139.3 (C), 133.7 (CH), 129.3 (2 \times CH), 128.1 (2 \times CH), 51.4 (CH₂), 42.0 (3 \times CH₂), 36.9 (3 \times CH₂), 36.0 (CH₂), 31.9 (C), 28.5 (3 \times CH); GC *R_T* 23.559 min; LRMS (EI) *m/z* (%) = 136 (M^+ + 1 - $C_8H_8O_2S$, 11), 135 (100).

3-((3*r*, 5*r*, 7*r*)-adamantan-1-yl)butanal (3ao, Scheme 2): Prepared according to *GPA*, after 14 h. It was purified by FC (Hexane/EtOAc 97:3 to 90:10) and obtained as a colorless oil (80.340 mg, 0.39 mmol, 78%): TLC *R_f* 0.32 (Hexane/EtOAc 97:3); IR ν 2924, 2848, 1723, 1707, 1309, 1208, 967, 733 cm^{-1} ; 1H -NMR (300 MHz, $CDCl_3$) δ 9.71 (dd, J = 3.4, 1.3 Hz, 1H), 2.56 (dd, J = 16.2, 3.3 Hz, 1H), 2.09 - 1.99 (m, 1H), 1.99 - 1.90 (m, 4H), 1.72 - 1.75 (m, 6H), 1.46 (s, 6H), 0.83 (dd, J = 6.9, 0.5 Hz, 3H); ^{13}C -NMR (75 MHz, $CDCl_3$) δ 203.9 (C=O), 45.6 (CH₂), 39.4 (3 \times CH₂), 37.8 (CH), 37.2 (3 \times CH₂), 34.4 (C), 28.7 (3 \times CH), 14.0 (CH₃); GC *R_T* 15.605 min; LRMS (EI) *m/z* (%) = 206 (M^+ , 1), 136 (11), 135 (100), 93 (11), 79 (11); HRMS (EI) *Calcd.* for $C_{14}H_{22}O$ 206.1671, found 206.1674.

General procedure B (GPB) for the decarboxylative addition of different carboxylic acids to Michael acceptor 2a (Scheme 3): The method is similar to *GPA*, but in this case a solution of the desired carboxylic acid (0.75 mmol, 1.5 equiv.) and diethyl 2-ethylidene malonate (2a, 92 μ L, 0.50 mmol) in MeCN (1.7 mL) were charged in the microwave tube, followed by H₂O (0.85 mL).

Diethyl 2-((3, 3-dimethylbutan-2-yl)malonate (3ba, Scheme 3):¹⁶ Prepared according to *GPB*, after 22 h, but in this case 2 equiv. of pivalic acid were used (1b, 103 mg, 1 mmol). It was purified by FC (Hexane/EtOAc 97:3 to 85:15) and obtained as a pale yellow oil (103 mg, 0.42 mmol, 84%): TLC *R_f* 0.27 (Hexane/EtOAc 97:3); IR ν 2962, 1757, 1731, 1466, 1370, 1289, 1147, 1024 cm^{-1} ; 1H -NMR (300 MHz, $CDCl_3$) δ 4.19 (q, J = 7.1 Hz, 2H), 4.18 (q, J = 7.1 Hz, 2H), 3.52 (d, J = 5.4 Hz, 1H), 2.30 - 2.19 (m, 1H), 1.28 (t, J = 7.1 Hz, 3H), 1.26 (t, J = 7.1 Hz, 3H), 1.01 (d, J = 7.2 Hz, 3H), 0.90 (s, 9H); ^{13}C -NMR (75 MHz, $CDCl_3$) δ 170.3 (C), 169.7 (C), 61.4 (CH₂), 61.0 (CH₂), 53.5 (CH), 42.7 (CH), 33.7 (C), 27.6 (3 \times CH₃), 14.14 (CH₃), 14.12 (CH₃); GC *R_T* 13.178 min; LRMS (EI) *m/z* (%) = 229 (M^+ - CH₃, 16), 199 (41), 189 (21), 188 (39), 173 (10), 160 (15), 155 (44), 143 (24), 142 (89), 147 (10), 127 (10), 116 (10), 115 (100), 114 (15), 109 (10), 99 (18), 87 (29), 86 (14), 85 (16), 69 (23), 57 (20), 55 (10).

Diethyl 2-(3, 3-dimethyl-4-phenylbutan-2-yl)malonate (3ca, Scheme 4): Prepared according to *GPB*, after 24 h. It was purified by FC (Hexane/EtOAc 100% Hexane to 97: 3 Hexane) and obtained as a colorless oil (120 mg, 0.375 mmol, 75%): TLC *R_f* 0.22 (Hexane/EtOAc 97:3); IR ν 2977, 1754, 1725, 1463, 1289, 1148, 697 cm^{-1} ; 1H -NMR (300 MHz, $CDCl_3$) δ 7.30 - 7.17 (m, 3H), 7.13 - 7.10 (m, 2H), 4.20 (q, J = 7.1 Hz, 2H), 4.18 (q, J = 7.1 Hz, 2H), 3.65 (d, J = 4.9 Hz, 1H), 2.56 (d, J = 1.6 Hz, 2H), 2.38 (qd, J = 7.2 Hz, 4.9 Hz, 1H), 1.29 (t, J = 7.1 Hz, 3H), 1.25 (t, J = 7.1 Hz, 3H), 1.13 (d, J = 7.2 Hz, 3H), 0.83 (s, 3H), 0.82 (s, 3H); ^{13}C -NMR (75 MHz, $CDCl_3$) δ 170.2 (C), 169.7 (C), 138.8 (C), 130.9 (2 \times CH), 127.8 (2 \times CH), 126.1 (CH), 61.6 (CH₂), 61.2 (CH₂), 53.1 (CH), 45.7 (CH₂), 42.0 (CH), 37.5 (C), 24.5 (CH₃), 24.1 (CH₃), 14.20 (CH₃), 14.15 (CH₃), 12.04 (CH₃); GC *R_T* 19.440 min; LRMS (EI) *m/z* (%) = 275 (M^+ - C₂H₅O, 9), 230 (19), 229 (100), 201 (12), 187 (73), 156 (15), 155 (100), 145 (11), 137 (39),

127 (25), 117 (11), 115 (17), 109 (30), 105 (11), 91 (68), 69 (12); HRMS (EI) *Calcd.* for $C_{19}H_{28}O_4$ 319.1908, found 319.1909.

Diethyl 2-(3-(4-chlorophenoxy)-3-methylbutan-2-yl)malonate (3da, Scheme 3): Prepared according to *GPB*, after 43 h. It was purified by FC (Hexane/EtOAc 97:3 to 95:5) and obtained as a colorless oil (107 mg, 0.3 mmol, 60%): TLC *R_f* 0.29 (9:1 Hexane/EtOAc); IR ν 2986, 1724, 1482, 1375, 1234, 1145, 1094, 1036, 850 cm^{-1} ; 1H -NMR (400 MHz, $CDCl_3$) δ 7.26 - 7.14 (m, 2H), 6.98 - 6.84 (m, 2H), 4.24 - 4.16 (m, 2H), 4.12 (q, J = 7.1 Hz, 2H), 3.78 (d, J = 6.8 Hz, 1H), 2.77 (p, J = 7.1 Hz, 1H), 1.28 (t, J = 7.2 Hz, 3H), 1.25 (s, 3H), 1.19 (t, J = 7.1 Hz, 3H), 1.19 (s, 3H), 1.12 (d, J = 7.1 Hz, 3H); ^{13}C -NMR (101 MHz, $CDCl_3$) δ 169.6 (C), 169.3 (C), 153.3 (C), 129.0 (C + 2 \times CH), 125.7 (2 \times CH), 83.0 (C), 61.4 (CH₂), 61.2 (CH₂), 53.6 (CH), 43.7 (CH), 25.7 (CH₃), 21.9 (CH₃), 14.19 (CH₃), 14.12 (CH₃), 13.2 (CH₃); GC *R_T* 20.674 min; LRMS (EI) *m/z* (%) = 229 (M^+ - C₆H₄ClO, 27), 156 (12), 155 (100), 137 (29), 136 (24), 130 (24), 128 (81), 127 (31), 111 (17), 109 (39), 108 (17), 81 (14), 69 (24), 65 (25); HRMS (EI) *Calcd.* for $C_{18}H_{25}ClO_5$ - C₆H₄ClO 229.1440, found 229.1447.

Diethyl 2-(1-cyclohexylethyl)malonate (3ea, Scheme 3):¹⁶ Prepared according to *GPB*, after 22 h. It was purified by FC (97:3 to 85:15 Hexane/EtOAc) and obtained as a pale yellow oil (97 mg, 0.38 mmol, 76%): TLC *R_f* 0.16 (Hexane/EtOAc 97:3), stained with KMnO₄; IR ν 2977, 2926, 2848, 1730, 1451, 1147, 1032 cm^{-1} ; 1H -NMR (300 MHz, $CDCl_3$) δ 4.25 - 4.14 (m, 4H), 3.39 (d, J = 9.2 Hz, 1H), 2.18 (dq, J = 13.8, 6.9, 4.4 Hz, 1H), 1.82 - 1.49 (m, 6H), 1.27 (t, J = 7.1 Hz, 6H), 1.20 - 0.93 (m, 5H), 0.90 (d, J = 7.0 Hz, 3H); ^{13}C -NMR (75 MHz, $CDCl_3$) δ 169.4 (C), 169.2 (C), 61.3 (CH₂), 61.2 (CH₂), 55.9 (CH), 40.4 (CH), 38.7 (CH), 31.6 (CH₂), 27.5 (CH₂), 26.9 (CH₂), 26.7 (CH₂), 26.6 (CH₂), 14.2 (2 \times CH₃), 13.0 (CH₃); GC *R_T* 15.943 min; LRMS (EI) *m/z* (%) = 225 (M^+ - C₂H₅O, 13), 187 (48), 161 (91), 160 (100), 141 (26), 133 (39), 132 (11), 115 (39), 114 (10), 110 (10), 109 (19), 90 (10), 87 (18), 81 (16), 69 (24), 67 (11), 55 (16).

Diethyl 2-(3-methylpentan-2-yl)malonate (3fa, Scheme 3):¹⁶ Prepared according to *GPB*, after 28 h, but using 2 equiv. of volatile (\pm)-2-methylbutyric acid (111 μ L, 1 mmol). It was purified by FC (Hexane/EtOAc 95:5 to 85:15) and obtained as a pale yellow oil (95 mg, 0.39 mmol, 78%, 51:49 *dr* according to 1H -NMR): TLC *R_f* 0.35 (Hexane/EtOAc 97:3), stained with KMnO₄; IR ν 2972, 2939, 2880, 1747, 1725, 1463, 1035, 912, 735 cm^{-1} ; 1H -NMR (300 MHz, $CDCl_3$) δ 4.19 (q, J = 7.1 Hz, 5H), 3.39 (d, J = 9.4 Hz, 0.48H), 3.31 (d, J = 10.4 Hz, 0.46H), 2.44 - 2.31 (m, 0.47H), 2.25 (dq, J = 13.8, 7.0, 4.0 Hz, 0.52H), 1.53 - 1.31 (m, 2H), 1.27 (t, J = 7.1 Hz, 3H), 1.26 (t, J = 7.1 Hz, 3H), 0.95 - 0.86 (m, 6H), 0.82 (d, J = 6.9 Hz, 1.46H), 0.79 (d, J = 6.7 Hz, 1.44H); ^{13}C -NMR (101 MHz, $CDCl_3$) δ 169.3 (C), 169.11 (C), 169.09 (C), 169.0 (C), 61.2 (CH₂), 61.1 (CH₂), 56.9 (CH), 56.1 (CH), 38.8 (CH), 36.9 (CH), 36.5 (CH), 36.2 (CH), 28.1 (CH₂), 23.8 (CH₃), 17.5 (CH₃), 14.2 (3 \times CH₃), 13.4 (CH₃), 12.5 (CH₃), 12.1 (CH₃), 13.0 (CH₃), 11.2 (CH₃); GC *R_T* 13.429 min; LRMS (EI) *m/z* (%) = 199 (M^+ - C₂H₅O, 19), 187 (11), 161 (30), 160 (100), 141 (18), 133 (37), 132 (11), 115 (39), 114 (13), 88 (12), 87 (20), 86 (12), 69 (23).

Diethyl 2-(1-(1-*tert*-butyl)piperidin-4-yl)ethyl)malonate (3ga, Scheme 3):¹⁶ Prepared according to *GPB*, after 72 h, but in this case another portion of photocatalyst (5.2 mg, 2.5 mol %) was added after 48 h of reaction. It was purified by FC (Hexane/EtOAc 97:3 to 85:15) and obtained as a colorless oil (113 mg, 0.305 mmol, 61%): TLC *R_f* 0.10 (Hexane/EtOAc 97:3); IR ν 1746, 1720, 1681, 1426, 1364, 1263, 732, 701 cm^{-1} ; 1H -NMR (300 MHz, $CDCl_3$) *mixture of rotamers* δ 4.21 (q, J = 7.1 Hz, 2H), 4.20 (q, J = 7.1 Hz, 2H), 3.39 (d, J = 8.6 Hz, 1H), 2.65 - 2.52 (m, 2H), 2.30 - 2.14 (m, 1H), 1.67 - 1.46 (m, 2.42H), 1.45 (s, 9.82H), 1.37 - 1.30 (m, 1H), 1.27 (t, J = 7.1 Hz, 7H), 1.22 - 1.01 (m, 2H), 0.93 (d, J = 7.0 Hz, 3H); ^{13}C -NMR (75 MHz, $CDCl_3$) *mixture of rotamers* δ 169.2 (C), 168.8 (C), 154.9 (C), 79.5 (C), 61.4 (CH₂), 61.3 (CH₂), 55.5 (CH), 44.3 (CH₂), 44.0 (CH₂), 38.9 (CH), 37.9 (CH), 30.4 (CH₂), 28.6 (3 \times CH₃), 27.1 (CH₂), 14.2 (2 \times CH₃),

FULL PAPER

13.1 (CH₃); LRMS (DIP) m/z (%) = 314 (M⁺-C₄H₉, 11), 298 (12), 270 (31), 226 (22), 198 (39), 188 (10), 187 (100), 180 (11), 160 (15), 156 (28), 153 (11), 141 (23), 112 (18), 85 (13), 84 (10), 82 (13), 57 (72), 41 (12).

Diethyl 2-(1-(2,3-dihydrobenzo[b][1,4]dioxin-2-yl)ethyl)malonate (3ha, Scheme 3): Prepared according to *GPB* after 15 h. It was purified by FC (Hexane/EtOAc 95:5 to 85:15) and obtained as a pale yellow oil (139 mg, 0.43 mmol, 86%, 54:46 *dr* according to ¹H-NMR): TLC *R_f* 0.17 (Hexane/EtOAc); IR ν 2986, 1737, 1594, 1491, 1262, 1175, 743 cm⁻¹; ¹H-NMR (300 MHz, CDCl₃) *mixture of diastereoisomers* δ 6.89 - 6.79 (m, 4H), 4.31 (dd, *J* = 11.3, 2.1 Hz, 0.55H), 4.27 - 4.13 (m, 6H), 4.08 - 3.93 (m, 1H), 3.84 (d, *J* = 5.3 Hz, 0.53H), 3.67 (d, *J* = 8.9 Hz, 0.46H), 2.74 - 2.50 (m, 1H), 1.35 - 1.19 (m, 6H), 1.15 (d, *J* = 7.0 Hz, 1.65H), 1.15 (d, *J* = 7.0 Hz, 1.49H); ¹³C-NMR (75 MHz, CDCl₃) *mixture of diastereoisomers* δ 169.0 (C), 168.5 (C), 168.4 (C), 168.3 (C), 143.7 (C), 143.4 (C), 143.3 (C), 143.0 (C), 121.6 (CH), 121.5 (2 × CH), 121.4 (CH), 117.3 (CH), 117.14 (2 × CH), 117.07 (CH), 74.6 (CH), 73.7 (CH), 66.4 (CH₂), 66.2 (CH₂), 61.7 (CH₂), 61.6 (CH₂), 61.5 (CH₂), 61.3 (CH₂), 54.5 (CH), 52.2 (CH), 34.6 (CH), 34.4 (CH), 14.21 (CH₃), 14.18 (CH₃), 14.16 (CH₃), 14.12 (CH₃), 13.1 (CH₃), 11.9 (CH₃); GC *R_T* 19.393 and 19.460 min (*two diastereoisomers*); LRMS (EI) m/z (*Diastereoisomer A*) (%) = 322 (M⁺, 34), 277 (17), 231 (15), 163 (15), 162 (100), 161 (12), 149 (18), 139 (12), 135 (43), 121 (23); m/z (*Diastereoisomer B*) (%) = 322 (M⁺, 32), 277 (18), 231 (10), 163 (15), 162 (100), 161 (15), 149 (18), 139 (15), 135 (45), 121 (26); HRMS (EI): *Calcd.* for C₁₇H₂₂O₆ - C₇H₁₁O₄ 163.0759, found 163.0761.

Diethyl 2-(3-phenoxybutan-2-yl)malonate (3ia, Scheme 3): Prepared according to *GPB*, after 44 h. It was purified by FC (Hexane/EtOAc 95:5) and obtained as a colorless oil (108 mg, 0.35 mmol, 70%, 62:38 *dr* according to ¹H-NMR): TLC *R_f* 0.14 (Hexane/EtOAc 97:3); IR ν 1717, 1599, 1494, 1265, 1232, 1027, 732, 727, 692 cm⁻¹; ¹H-NMR (300 MHz, CDCl₃) *mixture of diastereoisomers* δ 7.33 - 7.15 (m, 2H), 6.98 - 6.80 (m, 3H), 4.51 (qd, *J* = 6.3, 3.0 Hz, 0.35H), 4.40 (dq, *J* = 12.3, 6.1 Hz, 0.59H), 4.33 - 3.99 (m, 4H), 3.69 (d, *J* = 6.5 Hz, 0.51H), 3.54 (d, *J* = 9.2 Hz, 0.31H), 2.69 - 2.58 (m, 0.56H), 2.5 - 2.39 (m, 0.40H), 1.33 - 1.05 (m, 12H); ¹³C-NMR (75 MHz, CDCl₃) *mixture of diastereoisomers* δ 169.3 (C), 169.00 (C), 168.96 (C), 168.7 (C), 158.1 (C), 157.7 (C), 129.6 (2 × CH), 121.0 (CH), 120.8 (CH), 116.2 (CH), 115.9 (CH), 75.0 (CH), 73.8 (CH), 61.4 (CH₂), 61.2 (CH₂), 55.2 (CH), 53.7 (CH), 38.82 (CH), 38.78 (CH), 17.0 (CH₃), 16.9 (CH₃), 14.2 (2 × CH₃), 14.1 (CH₃), 13.9 (CH₃), 12.9 (CH₃), 11.6 (CH₃); GC *R_T* 17.695 and 17.796 min (*two diastereoisomers*); LRMS (EI) m/z (*Diastereoisomer A*) (%) = 216 (M⁺ + 1 - C₆H₅O, 11), 215 (88), 187 (11), 169 (24), 141 (100), 123 (24), 121 (16), 113 (49), 97 (22), 95 (16), 94 (31), 77 (20), 69 (10); m/z (*Diastereoisomer B*) (%) = 216 (M⁺ + 1 - C₆H₅O, 11), 215 (90), 187 (6), 169 (21), 141 (100), 123 (23), 121 (13), 113 (49), 97 (17), 95 (14), 94 (27), 77 (18), 69 (10); HRMS (EI) *Calcd.* for C₁₇H₂₄O₅ 308.1624, found 308.1624.

Diethyl 2-(3-(*tert*-butylamino)-4-phenylbutan-2-yl)malonate (3ja, Scheme 3):¹³ Prepared according to *GPB*, after 72 h, but in this case another portion of photocatalyst (5.2 mg, 2.5 mol %) was added after 48 hours of reaction. It was purified by FC (Hexane/EtOAc 95:5 to 70:30) and obtained as a pale yellow oil (122.3 mg, 0.30 mmol, 60%, 57:43 *dr* according to ¹H-NMR): TLC *R_f* 0.10 (Hexane/EtOAc 97:3); IR ν 2975, 1708, 1502, 1239, 1167, 1028, 702 cm⁻¹; ¹H-NMR (300 MHz, CDCl₃) *mixture of diastereoisomers and rotamers* δ 7.36 - 7.12 (m, 5H), 4.52 (d, *J* = 9.7 Hz, 0.52H), 4.40 (d, *J* = 10.1 Hz, 0.55H), 4.25 - 4.13 (m, 4H), 3.93 - 3.72 (m, 0.44H), 3.51 (d, *J* = 6.8 Hz, 0.43H), 3.35 (d, *J* = 9.9 Hz, 0.41H), 2.97 (dd, *J* = 13.9, 4.7 Hz, 0.44H), 2.77 (d, *J* = 6.7 Hz, 0.71H), 2.69 - 2.57 (m, 0.49H), 2.52 - 2.37 (m, 0.96H), 1.55 - 1.19 (m, 16H), 1.14 (d, *J* = 6.9 Hz, 1.87H), 0.95 (d, *J* = 7.0 Hz, 1.41H); ¹³C-NMR (75 MHz, CDCl₃) *mixture of diastereoisomers and rotamers* δ 169.6 (C), 169.0 (C), 168.4 (C), 155.6 (C), 155.4 (C), 138.04 (C), 137.99 (C), 129.4 (CH), 129.2 (CH), 128.49 (CH), 128.47 (CH), 126.5 (CH), 79.3 (C), 79.1 (C), 61.6 (CH₂), 61.4 (CH₂), 55.7 (CH), 55.3 (CH), 54.3 (CH), 52.8 (CH), 39.9 (CH₂), 39.1 (CH₂), 37.2

(CH), 35.8 (CH), 28.3 (CH₃), 28.1 (CH₃), 15.1 (CH₃), 14.2 (CH₃), 14.1 (CH₃), 11.2 (CH₃); LRMS (DIP) m/z (%) = 316 (M⁺-C₇H₇, 23), 262 (15), 260 (17), 217 (12), 216 (100), 170 (18), 142 (21), 131 (12), 124 (25), 120 (17), 91 (21), 57 (45).

Diethyl 2-(1-(1-*tert*-butoxycarbonyl)pyrrolidin-2-yl)ethyl)malonate (3ka, Scheme 3):¹³ Prepared according to *GPB*, after 72 h. It was purified by FC (Hexane/EtOAc 97:3 to 85:15) and obtained as a pale yellow oil (107 mg, 0.3 mmol, 60%, 82:18 *dr* according to ¹H-NMR): TLC *R_f* 0.10 (Hexane/EtOAc); IR ν 1726, 1678, 1382, 1264, 1166, 723, 704 cm⁻¹; ¹H-NMR (300 MHz, CDCl₃) *mixture of diastereoisomers and rotamers* δ 4.26 - 4.10 (m, 4H), 4.01 - 3.23 (m, 3H), 3.25 - 2.99 (m, 1H), 2.81 - 2.66 (m, 0.76H), 2.62 - 2.50 (m, 0.17H), 2.09 - 1.61 (m, 4H), 1.47 (s, 9H), 1.31 - 1.22 (m, 6H), 0.94 (d, *J* = 6.7 Hz, 3H); ¹³C-NMR (101 MHz, CDCl₃) *mixture of diastereoisomers and rotamers* δ 169.0 (C), 168.9 (C), 168.8 (C), 155.6 (C), 155.4 (C), 155.3 (C), 80.0 (C), 79.9 (C), 61.4 (CH₂), 61.3 (CH₂), 60.7 (CH), 60.4 (CH), 55.7 (CH), 54.9 (CH), 54.2 (CH), 48.1 (CH₂), 47.7 (CH₂), 47.3 (CH₂), 47.0 (CH₂), 37.2 (CH), 37.0 (CH), 28.6 (CH₃), 28.58 (CH₃), 28.3 (CH₃), 24.3 (CH₂), 23.9 (CH₂), 23.8 (CH₂), 14.22 (CH₃), 14.16 (CH₃), 13.6 (CH₃); LRMS (DIP) m/z (%) = 256 (M⁺ - C₄H₉, 10), 212 (26), 197 (16), 170 (18), 142 (11), 141 (19), 115 (17), 114 (100), 70 (89), 57 (47), 41 (12).

Diethyl 2-(1-((*tert*-butoxycarbonyl)amino)propan-2-yl)malonate (3la, Scheme 3):¹³ Prepared according to *GPB*, after 96 h, but in this case another portion of photocatalyst (2.5 mol %, 5.2 mg) was added after 48 h of reaction. It was purified by FC (Hexane/EtOAc 95:5 to 75:25) and obtained as a pale yellow oil (117 mg, 0.37 mmol, 74%): TLC *R_f* 0.10 (Hexane/EtOAc); IR ν 2981, 1718, 1518, 1243, 1167, 1039, 784 cm⁻¹; ¹H-NMR (300 MHz, CDCl₃) *mixture of rotamers* δ 4.74 (br s, 0.59H), 4.39 - 4.27 (m, 0.68H), 4.21 (q, *J* = 7.1 Hz, 2H), 4.20 (q, *J* = 7.1 Hz, 2H), 3.58 - 3.38 (m, 0.36H), 3.31 (d, *J* = 7.4 Hz, 1H), 3.19 - 3.17 (m, 1.78H), 2.45 (p, *J* = 6.8 Hz, 1H), 1.43 (s, 9H), 1.27 (t, *J* = 7.1 Hz, 6H), 1.02 (d, *J* = 6.9 Hz, 3H); ¹³C-NMR (75 MHz, CDCl₃) *mixture of rotamers* δ 168.9 (C), 168.7 (C), 156.1 (C), 79.4 (C), 61.54 (CH₂), 61.48 (CH₂), 55.2 (CH), 44.4 (CH), 34.2 (CH₂), 28.54 (CH₃), 28.45 (CH₃), 28.4 (CH₃), 28.1 (CH₃), 15.7 (CH₃), 14.22 (CH₃), 14.20 (CH₃); LRMS (DIP) m/z (%) = 244 (M⁺ - C₂H₅O, 14), 216 (40), 198 (15), 172 (20), 161 (23), 160 (32), 142 (29), 126 (19), 115 (29), 101 (26), 87 (15), 69 (13), 57 (100), 41 (14).

Diethyl 2-(1-phenoxypropan-2-yl)malonate (3ma, Scheme 3): Prepared according to *GPB*, after 85 h, but in this case, after 48 h, another portion of photocatalyst (5.2 mg, 2.5 mol %) and more of phenoxyacetic acid (230 mg, 1.5 mmol) were added. It was purified by FC (Hexane/EtOAc 97:3 to 90:10) and obtained as a colorless oil (96 mg, 0.325 mmol, 65%): TLC *R_f* 0.10 (Hexane/EtOAc 97:3); IR ν 1741, 1369, 1234, 1035, 752 cm⁻¹; ¹H-NMR (300 MHz, CDCl₃) δ 7.31 - 7.23 (m, 2H), 6.98 - 6.84 (m, 3H), 4.28 - 4.13 (m, 4H), 4.04 - 3.90 (m, 2H), 3.59 (d, *J* = 7.7 Hz, 1H), 2.85 - 2.69 (m, 1H), 1.26 (t, *J* = 7.1 Hz, 3H), 1.22 (t, *J* = 7.1 Hz, 3H), 1.16 (d, *J* = 7.0 Hz, 3H); ¹³C-NMR (75 MHz, CDCl₃) δ 168.8 (C), 168.7 (C), 158.9 (C), 129.5 (2 × CH), 120.9 (CH), 114.7 (2 × CH), 70.2 (CH₂), 61.5 (CH₂), 61.4 (CH₂), 54.2 (CH), 33.6 (CH), 15.0 (CH₃), 14.2 (CH₃), 14.1 (CH₃); GC *R_T* 17.778 min; LRMS (EI) m/z (%) = 249 (M⁺ - C₂H₅O, 203 (38), 202 (11), 201 (100), 173 (50), 145 (51), 138 (22), 127 (89), 109 (11), 99 (10), 94 (30), 83 (38), 57 (32), 69 (11), 55 (22); HRMS (EI) *Calcd.* for C₁₆H₂₂O₅ 294.1467, found 294.1476.

Diethyl 2-(1-(*p*-tolylloxy)propan-2-yl)malonate (3na, Scheme 3): Prepared according to *GPB*, after 50 h. It was purified by FC (Hexane/EtOAc 97:3 to 90:10) and obtained as a colorless oil (142 mg, 0.46 mmol, 92%): TLC *R_f* 0.10 (Hexane/EtOAc 97:3), visualized by exposure to UV light; IR ν 1738, 1507, 1236, 1174, 1034, 812 cm⁻¹; ¹H-NMR (300 MHz, CDCl₃) δ 7.07 - 7.04 (m, 2H), 6.79 - 6.76 (m, 2H), 4.24 - 4.13 (m, 4H), 3.98 - 3.87 (m, 2H), 3.58 (d, *J* = 7.7 Hz, 1H), 2.81 - 2.66 (m, 1H), 2.27 (s, 3H), 1.26 (t, *J* = 7.1 Hz, 3H), 1.22 (t, *J* = 7.1 Hz, 3H), 1.14 (d,

FULL PAPER

$J = 6.9$ Hz, 3H); ^{13}C -NMR (75 MHz, CDCl_3) δ 168.8 (C), 168.7 (C), 156.8 (C), 130.1 (C), 130.0 (2 \times CH), 114.6 (2 \times CH), 70.4 (CH_2), 61.5 (CH_2), 64.4 (CH_2), 54.2 (CH), 33.7 (CH), 20.6 (CH_3), 14.9 (CH_3), 14.2 (CH_3), 14.17 (CH_3); GC R_T 18.585 min; LRMS (EI) m/z (%) = 308 (M^+ , 4), 263 (10), 217 (28), 202 (11), 201 (100), 173 (56), 148 (11), 145 (51), 127 (82), 109 (11), 108 (35), 107 (18), 99 (11), 91 (18), 83 (30), 55 (14); HRMS (EI) *Calcd.* for $\text{C}_{17}\text{H}_{24}\text{O}_5$ 308.1624, found 308.1634.

Diethyl 2-(1-oxo-1-phenylpropan-2-yl)malonate (30a, Scheme 3): Prepared according to *GPB*, after 38 h, but in this case 1 equiv. of ketoacid was added and after 8 h, another 1 equiv. was added. It was purified by FC (Hexane/EtOAc 95:5) and obtained as a pale yellow oil (98 mg, 0.335 mmol, 67%); TLC R_f 0.15 (Hexane/EtOAc 97:3); IR ν 2977, 1744, 1727, 1681, 1457, 1288, 1179, 1024, 981, 714 cm^{-1} ; ^1H -NMR (300 MHz, CDCl_3) δ 8.04 - 7.99 (m, 2H), 7.63 - 7.55 (m, 1H), 7.53 - 7.45 (m, 2H), 4.36 - 4.05 (m, 5H), 3.99 (d, $J = 10.8$ Hz, 1H), 1.32 (t, $J = 7.1$ Hz, 3H), 1.20 (d, $J = 7.3$ Hz, 3H), 1.17 (t, $J = 7.1$ Hz, 3H); ^{13}C -NMR (75 MHz, CDCl_3) δ 201.7 (C), 168.9 (C), 168.5 (C), 135.7 (C), 133.4 (CH), 128.8 (2 \times CH), 128.7 (2 \times CH), 61.8 (2 \times CH_2), 56.4 (CH), 40.6 (CH), 16.0 (CH_3), 14.3 (CH_3), 14.0 (CH_3); GC R_T 17.797 min; LRMS (EI) m/z (%) = 247 ($\text{M}^+ - \text{C}_2\text{H}_5\text{O}$, 5), 201 (7), 106 (8), 105 (100), 77 (18).

Diethyl 2-(3-oxopentan-2-yl)malonate (3pa, Scheme 3): Prepared according to *GPB*, after 69 h, but in this case another portions of photocatalyst (1 mol %, 2.08 mg) and Na_2CO_3 (1 equiv., 0.5 mmol, 53 mg) were added after 48 h of reaction. Moreover, in this case 4 \times 1 equiv. (207 mg, 2 mmol) of 2-ketobutyric acid were added every 12 h. It was purified by FC (Hexane/EtOAc 97:3 to 85:15) and obtained as a pale yellow oil (66 mg, 0.27 mmol, 54%); TLC R_f 0.10 (Hexane/EtOAc 97:3); IR ν 1746, 1720, 1264, 1090, 1046, 916, 732 cm^{-1} ; ^1H -NMR (300 MHz, CDCl_3) δ 4.27 - 4.08 (m, 4H), 3.77 (d, $J = 10.6$ Hz, 1H), 3.28 (dq, $J = 10.6$ Hz, 7.2 Hz, 1H), 2.63 (qd, $J = 7.3$, 3.2 Hz, 2H), 1.29 (t, $J = 7.1$ Hz, 3H), 1.23 (t, $J = 7.1$ Hz, 3H), 1.11 (d, $J = 7.1$ Hz, 3H), 1.08 (t, $J = 7.1$ Hz, 3H); ^{13}C -NMR (75 MHz, CDCl_3) δ 212.3 (C), 168.8 (C), 168.7 (C), 61.8 (CH_2), 61.7 (CH_2), 54.7 (CH), 44.8 (CH), 34.8 (CH_2), 15.0 (CH_3), 14.3 (CH_3), 14.1 (CH_3), 7.8 (CH_3); GC R_T 13.745 min; LRMS (EI) m/z (%) = 215 ($\text{M}^+ - \text{C}_2\text{H}_5$, 51), 200 (76), 189 (89), 188 (17), 187 (93), 171 (11), 169 (10), 159 (11), 154 (94), 143 (25), 142 (47), 122 (21), 115 (53), 114 (11), 113 (23), 99 (14), 87 (29), 86 (14), 68 (43), 57 (89); HRMS (EI) *Calcd.* for $\text{C}_{12}\text{H}_{20}\text{O}_5 - \text{C}_2\text{H}_5$ 215.0919, found 215.0921.

Tandem decarboxylation-Giese reaction-lactonization (Scheme 5a).

Ethyl (3S*, 4R*)-4, 5, 5-trimethyl-2-oxotetrahydro-2H-pyran-3-carboxylate ((±)-trans-3ra): Prepared according to *GPB*, after 14 h. It was purified by FC (Hexane/EtOAc 95:5 to 80:20) and obtained as a pale yellow oil (86 mg, 0.40 mmol, 80%, >98:2 *dr* according ^1H and ^{13}C -NMR); TLC R_f 0.10 (Hexane/EtOAc 95:5); IR ν 2977, 1724, 1481, 1379, 1249, 1174, 1154, 1090, 1046 cm^{-1} ; ^1H -NMR (300 MHz, CDCl_3) δ 4.24 (q, $J = 7.1$ Hz, 2H), 4.09 (d, $J = 11.1$ Hz, 1H), 3.97 (d, $J = 11.1$ Hz, 1H), 3.18 (d, $J = 11.4$ Hz, 1H), 2.17 (dq, $J = 11.4$, 6.8 Hz, 1H), 1.31 (t, $J = 7.1$ Hz, 3H), 0.98 (s, 3H), 0.97 (d, $J = 6.7$ Hz, 3H), 0.96 (s, 3H); ^{13}C -NMR (101 MHz, CDCl_3) δ 169.5 (C), 167.3 (C), 79.8 (CH_2), 62.0 (CH_2), 53.8 (CH), 40.0 (CH), 32.5 (C), 23.5 (CH_3), 17.4 (CH_3), 14.2 (CH_3), 14.1 (CH_3); GC R_T 15.882 min; LRMS (EI) m/z (%) = 214 (M^+ , 6), 199 (13), 169 (32), 159 (10), 155 (18), 142 (13), 141 (91), 127 (15), 125 (16), 123 (12), 115 (55), 113 (18), 109 (23), 97 (37), 96 (21), 95 (26), 87 (45), 83 (27), 82 (23), 81 (18), 70 (100), 69 (73), 67 (16), 56 (30), 55 (69); HRMS (DIP) *Calcd.* for $\text{C}_{11}\text{H}_{18}\text{O}_4 - \text{CH}_3$ 199.0970, found 199.0977.

β -Fragmentation after addition of 1q to 2a (Scheme 5b).

Diethyl 2-(1-((3,4-dichlorophenyl)thio)ethyl)malonate (5): Prepared according to the *GPB*, after 24 h. It was purified by FC (Hexane/EtOAc 95:5 to 90:10) and obtained as a colorless oil (135 mg, 0.37 mmol, 74%); TLC R_f 0.28 (95:5 Hexane/EtOAc); IR ν 2989, 1730, 1456, 1258, 1219, 1133, 1050, 1025, 809 cm^{-1} ; ^1H -NMR (300 MHz, CDCl_3) δ 7.58 (d, $J = 2.0$ Hz, 1H), 7.39 (d, $J = 8.3$ Hz, 1H), 7.32 (dd, $J = 8.3$, 2.1 Hz, 1H), 4.23 (q, $J = 7.2$ Hz, 2H), 4.22 (q, $J = 7.2$ Hz, 2H), 3.75 (dq, $J = 8.6$, 6.9 Hz, 1H), 3.48 (d, $J = 8.6$ Hz, 1H), 1.39 (d, $J = 6.9$ Hz, 3H), 1.29 (t, $J = 7.1$ Hz, 3H), 1.28 (t, $J = 7.1$ Hz, 3H); ^{13}C -NMR (75 MHz, CDCl_3) δ 167.5 (C), 167.2 (C), 134.9 (CH), 133.7 (C), 133.0 (C), 132.7 (CH), 132.5 (C), 130.8 (CH), 62.0 (CH_2), 61.9 (CH_2), 57.7 (CH), 43.5 (CH), 19.5 (CH_3), 14.2 (2 \times CH_3); LRMS (DIP) m/z (%) = 368 (M^{+37}Cl , 12), 367 (10), 366 (63), 365 (16), 364 (M^{+35}Cl , 87), 247 (32), 245 (42), 217 (31), 207 (49), 205 (68), 179 (48), 178 (39), 177 (64), 142 (77), 141 (100), 135 (39), 113 (46), 97 (31), 95 (32), 87 (51), 85 (67), 73 (58), 69 (97), 57 (40), 55 (38), 43 (34), 41 (46); HRMS (DIP) *Calcd.* for $\text{C}_{15}\text{H}_{18}\text{Cl}_2\text{O}_4\text{S}$ 364.0303, found 364.0301.

1, 2-bis(3,4-dichlorophenyl)disulfane (6, Scheme 5b):³⁰ Compound 6 was obtained as side-product of 5, following the procedure described above. It was purified by FC (Hexane/EtOAc 97:3) and obtained as a white solid (45 mg, 0.13 mmol, 17%); TLC R_f 0.65 (Hexane/EtOAc 97:3); IR ν 2913, 1563, 1454, 1362, 1143, 1117, 1084, 1030, 858, 799, 675 cm^{-1} ; ^1H -NMR (300 MHz, CDCl_3) δ 7.56 (d, $J = 2.1$ Hz, 2H), 7.39 (d, $J = 8.4$ Hz, 2H), 7.29 (dd, $J = 8.4$, 2.2 Hz, 2H); ^{13}C -NMR (75 MHz, CDCl_3) δ 136.3 (2 \times C), 133.6 (2 \times C), 132.1 (2 \times C), 131.1 (2 \times CH), 129.3 (2 \times CH), 127.0 (2 \times CH); GC R_T 25.335 min; LRMS (EI) m/z (%) = 360 (M^{+37}Cl , 11), 358 (40), 357 (11), 356 (M^{+35}Cl , 72), 354 (54), 179 (69), 177 (100), 144 (30), 142 (76).

Radical Cascade of Homoallylic Acid 1s with 2a (Scheme 5c)

Diethyl (2S,5R)-5-(4-ethoxy-3-(ethoxycarbonyl)-2-methyl-4-oxobutyl)-2,3,3-trimethylcyclopentane-1,1-dicarboxylate (7): Prepared according to *GPB*, after 24 h, but in this case the reaction was performed employing 1 equiv. of homoallylic acid 1s (65 mg, 0.5 mmol) and 2.5 equiv. of 2a (235 mg, 1.25 mmol). It was purified by FC (97:3 to 90:10 Hexane/EtOAc) and obtained as a colorless oil (148 mg, 0.33 mmol, 65%, 1:1 *dr* according NMR and GC experiments); TLC R_f 0.2 (Hexane/EtOAc 95:5); IR ν 2978, 1727, 1366, 1243, 1179, 1036, 856 cm^{-1} ; ^1H -NMR (300 MHz, CDCl_3) *mixture of diastereoisomers* δ 4.30 - 4.07 (m, 8H), 3.34 (d, $J = 5.9$ Hz, 0.40H), 3.16 (d, $J = 8.4$ Hz, 0.40H), 3.09 - 2.94 (m, 1H), 2.66 (q, $J = 7.4$ Hz, 1H), 2.36 - 2.12 (m, 1H), 1.78 (p, $J = 6.2$ Hz, 1H), 1.71 - 1.53 (m, 1H), 1.30 - 1.21 (m, 13H), 1.07 - 0.99 m, 7H), 0.88 (d, $J = 3.2$ Hz, 1.50H), 0.86 (d, $J = 3.2$ Hz, 1.50H), 0.98 (s, 3H); ^{13}C -NMR (75 MHz, CDCl_3) *mixture of diastereoisomers* δ 172.0 (C), 171.8 (2 \times C), 171.7 (C), 169.3 (C), 168.8 (C), 168.7 (C), 168.6 (C), 67.7 (C), 67.6 (C), 61.32 (CH_2), 61.27 (2 \times CH_2), 61.13 (CH_2), 61.07 (CH_2), 61.0 (CH_2), 60.94 (CH_2), 60.87 (CH_2), 58.8 (CH), 56.1 (CH), 49.5 (CH), 49.2 (CH), 47.2 (CH_2), 45.7 (CH_2), 41.3 (CH), 40.6 (CH), 40.2 (C), 40.1 (C), 37.2 (CH), 36.5 (CH), 32.3 (CH_3), 31.8 (CH_3), 29.21 (CH_3), 29.17 (CH_3), 22.8 (CH_3), 22.6 (CH_3), 18.1 (CH_3), 16.4 (CH_3), 14.21 (5 \times CH_3), 14.15 (CH_3), 11.62 (CH_3), 11, 57 (CH_3); GC R_T 22.008 min and 22.224 min (*two diastereoisomers*); LRMS (EI) m/z (*diastereoisomer A*) (%) = 411 ($\text{M}^+ - \text{C}_2\text{H}_5$, 9), 337 (25), 297 (43), 222 (19), 196 (21), 187 (100), 149 (44), 141 (50), 107 (19), 69 (19); m/z (*diastereoisomer B*) (%) = 411 ($\text{M}^+ - \text{C}_2\text{H}_5\text{O}$, 15), 337 (38), 336 (20), 297 (45), 205 (24), 196 (19), 187 (100), 149 (45), 141 (50), 107 (19), 69 (19); HRMS (EI) *Calcd.* for $\text{C}_{24}\text{H}_{40}\text{O}_8 - \text{C}_2\text{H}_5\text{O}$ 411.2383 found 411.2392.

Acknowledgements

We thank the Ministerio de Economía y Competitividad (CTQ2015-66624-P) and to the University of Alicante (VIGROB-

FULL PAPER


173) for financial support. N. P. R. thanks the ISO for a grant. We also thank to Irene Bosque for preliminary investigations.

Keywords: carboxylic acids • synthetic methods • radical reactions • photocatalysis • sustainable chemistry

- [1] For selected reviews on radical conjugated additions, see: a) B. Giese in *Radicals in Organic Synthesis: Formation of Carbon–Carbon Bonds*, Pergamon: Oxford, 1986; b) J. Fossey, D. Lefort, J. Sorba in *Free Radicals in Organic Chemistry*, Wiley: Chichester, UK, 1995; c) P. Renaud, M. Gerster *Angew. Chem., Int. Ed.* **1998**, *37*, 2562; d) P. Renaud in M. Sibi, *Radicals in Organic Synthesis*, Wiley-VCH: Weinheim, 2001; e) S. Z. Zard, *Radical Reactions in Organic Synthesis*, Oxford University Press: New York NY, 2003; f) G. S. C. Srikanth, S. L. Castle *Tetrahedron* **2005**, *61*, 10377.
- [2] a) A. Michael, *J. Prakt. Chem.* **1887**, *35*, 349; b) A. Michael, *J. Prakt. Chem.* **1894**, *49*, 20.
- [3] a) N. Rodriguez, L. J. Goossen, *Chem. Soc. Rev.* **2011**, *40*, 5030; b) P. Gallezot, *Chem. Soc. Rev.* **2012**, *41*, 1538; c) A. J. J. Straathof, *Chem. Rev.* **2014**, *114*, 1871.
- [4] a) H. Kolbe, *Liebigs Ann. Chem.* **1849**, *69*, 257; b) H. Kolbe, *Liebigs Ann. Chem.* **1848**, *64*, 339; c) H. Hunsdiecker, C. Hunsdiecker *Chem. Ber.* **1942**, *75*, 291; d) D. H. R. Barton, D. Bridon, I. Fernandez-Picot, S. Z. Zard, *Tetrahedron* **1987**, *43*, 2733; e) D. H. R. Barton, D. Crich, W. B. Motherwell, *J. Chem. Soc., Chem. Commun.* **1983**, 939.
- [5] a) A. G. Myers, D. Tanaka, M. R. Mannion, *J. Am. Chem. Soc.* **2002**, *124*, 11250; b) D. Tanaka, S. P. Romeril, A. G. Myers, *J. Am. Chem. Soc.* **2005**, *127*, 10323; c) L. J. Goossen, G. Deng, L. M. Levy, *Science* **2006**, *313*, 662; d) N. Rodriguez, L. J. Goossen, *Chem. Soc. Rev.* **2011**, *40*, 5030; e) R. Shang, L. Liu, *Sci. China Chem.* **2011**, *54*, 1670.
- [6] For a recent review, see: J. Xuan, Z.-G. Zhang, W.-J. Xiao, *Angew. Chem., Int. Ed.* **2015**, *54*, 15632.
- [7] H. G. Roth, N. A. Romero, D. A. Nicewicz, *Synlett* **2016**, 27, 714.
- [8] This fragmentation is reported to be exceedingly fast for aliphatic carboxyl radicals ($k \approx 10^9 \text{ s}^{-1}$), see: J. K. Kochi, T. M. Bockman, S. M. Hubig, *J. Org. Chem.* **1997**, *62*, 2210.
- [9] N. Bortolamei, A. A. Isse, A. Gennaro, *Electrochimica Acta* **2010**, *55*, 8312.
- [10] P. T. Anastas, J. C. Warner in *Green Chemistry: Theory and Practice*, Oxford University Press, New York, 1998.
- [11] Y. Yoshimi, S. Washida, Y. Okita, K. Nishikawa, K. Maeda, S. Hayashi, T. Morita, *Tetrahedron Lett.* **2013**, *54*, 4324.
- [12] Y. Miyake, K. Nakajima, Y. Nishibayashi, *Chem. Comm.* **2013**, 49, 7854.
- [13] L. Chu, C. Ohta, Z. Zuo, D. W. C. MacMillan, *J. Am. Chem. Soc.* **2014**, *136*, 10886.
- [14] A. Millet, Q. Lefebvre, M. Rueping, *Chem. Eur. J.* **2016**, *22*, 13464.
- [15] a) S. Fukuzumi, H. Kotani, K. Ohkubo, S. Ogo, N. V. Tkachenko, H. Lemmetyinen, *J. Am. Chem. Soc.* **2004**, *126*, 1600; b) S. Fukuzumi, K. Ohkubo, T. Suenobu, *Acc. Chem. Res.* **2014**, *47*, 1455; c) S. Fukuzumi, K. Ohkubo, *Org. Biomol. Chem.* **2014**, *12*, 6059; d) D. A. Nicewicz, D. S. Hamilton, *Synlett* **2014**, 25, 1191; e) For a recent application from our group on photocatalytic oxidation of benzoic acids, see: N. P. Ramirez, I. Bosque, J. C. Gonzalez-Gomez, *Org. Lett.* **2015**, *17*, 4550; f) For a recent comprehensive review on organic photoredox catalysis, see: N. A. Romero, D. A. Nicewicz, *Chem. Rev.* **2016**, *116*, 10075.
- [16] T. Chinzei, K. Miyazawa, Y. Yasu, T. Koike, M. Akita *RSC Adv.* **2015**, *5*, 21297. In this work, the authors demonstrate the ability of the photocatalyst [Acr-Mes]⁺ to induce the decarboxylative Giese-type reaction, but it is concluded that the scope of the reaction is rather limited.
- [17] For selected examples of photocatalytic decarboxylative functionalization of carboxylic acids, using [Acr-Mes]⁺ as photocatalyst, see: a) C. Cassani, G. Bergonzini, C.-J. Wallentin, *Org. Lett.* **2014**, *16*, 4228; b) X. Wu, C. Meng, X. Yuan, X. Jia, X. Qian, J. Ye, *Chem. Commun.*, **2015**, 51, 11864; c) S. B. Lang, K. C. Cartwright, R. S. Welter, T. M. Locascio, J. A. Tunge, *Eur. J. Org. Chem.* **2016**, 3331; d) For a recent synthesis and photophysical characterization of acridinium-based photocatalysts, see: A. Joshi-Pangu, F. Levesque, H. G. Roth, S. F. Oliver, L.-C. Campeau, D. Nicewicz, D. A. DiRocco, *J. Org. Chem.* **2016**, *81*, 7244.
- [18] a) L. Wanka, K. Iqbal, P. L. Shreiner, *Chem. Rev.* **2013**, *113*, 3516; b) K. A. Agnew-Francis, C. M. Williams, *Adv. Synth. Catal.* **2016**, *358*, 675.
- [19] Very recently has been reported that this photocatalytic system enables the decarboxylative hydroxylation of carboxylic acids using molecular oxygen. Under our reaction conditions, in the presence of the Michael acceptor, this possible competitive reaction was not observed. See: H.-T. Song, W. Ding, Q.-Q. Zhou, J. Liu, L.-Q. Lu, W.-J. Xiao, *J. Org. Chem.* **2016**, *81*, 7250.
- [20] J. D. Griffin, M. A. Zeller, D. A. Nicewicz, *J. Am. Chem. Soc.* **2015**, *137*, 11340.
- [21] a) H. Stetter, M. Schreckenberger *Angew. Chem., Int. Ed. Engl.* **1973**, *12*, 81; b) H. Stetter, *Angew. Chem., Int. Ed. Engl.* **1976**, *15*, 639; c) For a recent review, see: X. Bugaut, F. Glorius, *Chem. Soc. Rev.* **2012**, *41*, 3511.
- [22] G.-Z. Wang, R. Shang, W.-M. Cheng, Y. Fu, *Org. Lett.* **2015**, *17*, 4830.
- [23] For a recent successful photocatalyzed benzylation of Michael acceptors via decarboxylation of arylacetic acids, see: L. Capaldo, L. Buzzetti, D. Merli, M. Fagnoni, D. Ravelli, *J. Org. Chem.* **2016**, *81*, 7102.
- [24] For a comprehensive review on thyl radicals in organic synthesis, see: F. Denes, M. Pichowicz, G. Povie, P. Renaud, *Chem. Rev.* **2014**, *114*, 2587.
- [25] a) A. L. J. Beckwith, C. J. Easton, A. K. Serelis, *J. Chem. Soc. Chem. Commun.*, **1980**, 482; b) A. L. J. Beckwith, C. H. Schiesser, *Tetrahedron* **1985**, *41*, 3925.
- [26] J. F. J. Engbersen, A. Koudijs, H. C. Der Van Plas, *J. Heterocyclic Chem.* **1982**, *19*, 1281.
- [27] D. Pappo, S., Shimony, Y. Kashman, *J. Org. Chem.* **2005**, *70*, 199.
- [28] D. Dondi, A. M. Cardarelli, M. Fagnoni, A. Albini, *Tetrahedron* **2006**, *62*, 5527.
- [29] H. Togo, S. Matsubayashi, O. Yamazaki, M. Yokoyama, *J. Org. Chem.* **2000**, *65*, 2816.
- [30] K. M. Khan, M. Taha, F. Rahim, M. Ali, W. Jamil, S. Perveen, M. I. Choudhary, *Lett. Org. Chem.* **2010**, *7*, 415.

FULL PAPER

Entry for the Table of Contents



The decarboxylative alkylation of carboxylic acids can be efficiently conducted under air atmosphere, at room temperature, using [Acr-Mes]⁺ as photocatalyst and promoted by visible-light. A range of readily available starting materials has been well tolerated with this transition-metal-free protocol.