

# One-Pot Synthesis of Benzo[*b*]furan and Indole Inhibitors of Tubulin Polymerization

Bernard L. Flynn,<sup>\*,†</sup> Ernest Hamel,<sup>‡</sup> and M. Katherine Jung<sup>§</sup>

Department of Chemistry, The Faculties, Australian National University, Canberra, ACT, 0200, Australia, Screening Technologies Branch, Developmental Therapeutics Program, Division of Cancer Treatment and Diagnosis, National Cancer Institute at Frederick, National Institutes of Health, Frederick, Maryland 21702, and Science Applications International Corporation—Frederick, National Cancer Institute at Frederick, National Institutes of Health, Frederick, Maryland 21702


Received February 14, 2002

Benzo[*b*]furan and indole analogues of some recently identified benzo[*b*]thiophene inhibitors of tubulin polymerization have been prepared, and their biological activity has been assessed. Several very potent analogues were identified.

Compounds that interfere with the microtubule–tubulin equilibria in cells are useful in the treatment of human disease.<sup>1</sup> Examples are the antiinflammatory activity of colchicine,<sup>1a</sup> the anthelmintic properties of the benzimidazole carbamates,<sup>1b</sup> and the antineoplastic use of the *Catharanthus* (formerly *Vinca*) alkaloids<sup>1c,d</sup> and the taxoids.<sup>1e,f</sup> More recently, it has been established that some tubulin binders selectively target the vascular system of tumors.<sup>2</sup> These agents induce morphological changes in the endothelial cells of the tumor's blood vessels so as to occlude flow.<sup>2i</sup> As a result, virtually complete vascular shutdown is achieved within the tumor in a matter of minutes. Combretastatin A-4 (CA4, Figure 1) is a powerful inhibitor of tubulin polymerization and a potent cytotoxin (Table 1).<sup>3</sup> The prodrug form of CA4, combretastatin A-4 disodium phosphate (CA4P), is undergoing clinical trials as a tumor vascular targeting agent.<sup>2</sup> The prodrug, CA4P, is itself inactive with purified tubulin and is therefore believed to require *in vivo* dephosphorylation to CA4 before it can act.<sup>2</sup> This process may contribute to its selectivity, with greater rates of dephosphorylation occurring in the tumor vascular systems than in normal vasculature.<sup>2c,j</sup>

The success of tubulin polymerization inhibitors as anticancer agents has stimulated significant interest in the identification of new compounds that may be more potent or more selective in targeted tissues or tumors.


In 1999, Pinney and co-workers<sup>4</sup> reported a benzo[*b*]thiophene, compound **1** (Figure 1), that exhibited some activity as a tubulin polymerization inhibitor and appeared to bind weakly to the colchicine binding domain of tubulin. However, **1** only inhibited the rate and not the extent of tubulin polymerization. The limited activity of **1** was hypothesized to come from poor solubility.<sup>4a</sup> Using our novel method for the flexible and concise synthesis of benzo[*b*]thiophenes, we prepared analogues of **1**.<sup>5</sup> The most potent of these was **2** (Figure 1, Table 1).<sup>5a</sup> More recently, we have developed a one-pot, multicomponent coupling approach to benzo[*b*]furans and indoles.<sup>6</sup> Here, we report on the application of this



**Figure 1.**

method to the preparation of various benzo[*b*]furan and indole analogues of **1** and **2** and their effects on tubulin polymerization.

Benzo[*b*]furans **8** and **9** were prepared by multicomponent coupling of 2-iodo-5-methoxyphenol **3**, 4-methoxyphenol **4**, 5-methoxyphenol **5**, and 2-iodo-4-methoxyphenol **7** (Scheme 1<sup>a</sup>).




<sup>a</sup> Reagents and conditions: (i) MeMgCl 2 equiv, Pd(PPh<sub>3</sub>)<sub>2</sub>Cl<sub>2</sub> 3 mol %, THF, 65 °C, 1.5 h under N<sub>2</sub>(g); (ii) cool to room temp, add **7** and DMSO, then heat to 80 °C, 16–18 h; (iii) same as (ii) except exchange N<sub>2</sub>(g) atmosphere for CO(g) atmosphere (balloon); (iv) AlCl<sub>3</sub> 3 equiv, CH<sub>2</sub>Cl<sub>2</sub>.

\* To whom correspondence should be addressed. Phone: 61 2 6125 2017. Fax: ++61 2 6125 0760. E-mail: flynn@rsc.anu.edu.au.

<sup>†</sup> Australian National University.

<sup>‡</sup> Division of Cancer Treatment and Diagnosis, National Cancer Institute at Frederick, National Institutes of Health.

<sup>§</sup> Science Applications International Corporation—Frederick, National Cancer Institute at Frederick, National Institutes of Health.

**Table 1.** Effects of Indole and Benzo[*b*]furan Compounds on Tubulin Polymerization, Colchicine Binding, and Growth of MCF-7 Human Breast Carcinoma Cells<sup>8</sup>



compound	X	Y	R	inhibition of tubulin polymerization, <sup>a</sup> IC <sub>50</sub> ± SD (μM)	inhibition of colchicine binding <sup>b</sup> (% inhibition ± SD)	inhibition of cell growth, <sup>c</sup> IC <sub>50</sub> ± SD (nM)
CA4				2.1 ± 0.1 <sup>d</sup>	91 ± 2	11 ± 4 <sup>d</sup>
<b>1</b>	S	C=O	H	>40 <sup>e,f</sup>		640 ± 10 <sup>d</sup>
<b>2</b>	S	C=O	OH	3.4 ± 0.2 <sup>d</sup>	21 ± 10 <sup>d</sup>	520 ± 400 <sup>d</sup>
<b>8</b>	O	nothing	H	>40		>1000
<b>9</b>	O	C=O	H	0.41 ± 0.1	77 ± 3	34 ± 10
<b>11</b>	O	C=O	OH	1.3 ± 0.2	80 ± 2	42 ± 10
<b>13</b>	NH	nothing	H	>40 <sup>e</sup>		460 ± 5
<b>17</b>	NH	nothing	OH	4.1 ± 0.6	28 ± 8	370 ± 2
<b>19</b>	NH	C=O	OH	1.6 <sup>g</sup>	54 ± 0.7	45 ± 5
<b>20</b>	O	CH(OH)	H	>40 <sup>e</sup>		>1000
<b>21</b>	O	CH(OH)	OH	8.8 ± 1.4		560 ± 60
<b>22</b>	O	CH <sub>2</sub>	H	2.5 ± 0.3	17 ± 10	400 ± 100

<sup>a</sup> The tubulin concentration was 10 μM. Inhibition of extent of assembly, after a 20 min incubation at 30 °C, was the parameter measured.

<sup>b</sup> The tubulin concentration was 1.0 μM, and the [<sup>3</sup>H]colchicine and inhibitor concentrations were both 5.0 μM. Incubation was for 10 min at 37 °C. At least two independent experiments were performed with each of the compounds evaluated. <sup>c</sup> Cells were grown for 48 h at 37 °C in a humidified 5% CO<sub>2</sub> atmosphere. Cell protein was the parameter measured. At least two independent experiments were performed with each compound. <sup>d</sup> Data from ref 5. <sup>e</sup> The rate, but not the extent of assembly, was reduced by compound concentrations as high as 40 μM. <sup>f</sup> Data from ref 4a. <sup>g</sup> Same value in two experiments.

oxyphenylethyne **4**, and 3,4,5-trimethoxyiodobenzene **7** (Scheme 1).<sup>6</sup> This process involved initial deprotonation of **3** and **4** with 2 equiv of methylmagnesium chloride to give the corresponding magnesium phenolate and acetylide, respectively (not shown). Coupling occurred upon addition of a catalytic amount of Pd(PPh<sub>3</sub>)<sub>2</sub>Cl<sub>2</sub> and heating to 65 °C for 1.5 h to give the intermediate *o*-alkynylphenolate **6**. The solution was diluted with dimethyl sulfoxide (DMSO), aryl iodide **7** was added, and the reaction was heated to 80 °C for 16 h under N<sub>2</sub>(g). Under these conditions, the palladium(0) present undergoes oxidative addition to **7**, followed by palladium(II)-induced cyclization of **6** and reductive elimination to afford **8** in an excellent yield (88%). When the N<sub>2</sub>(g) atmosphere was exchanged for one of CO(g) at the point of introduction of **7**, a carbonyl is inserted between palladium(II) and the trimethoxyphenyl group, leading to the carbonylated product **9** (64%).<sup>6</sup> Using alkyne **5** in place of **4** under these carbonylative conditions gave **10** (63%). The isopropyl ether in **10** was cleaved using AlCl<sub>3</sub> to give the desired phenol **11** (97%).<sup>7</sup>


Indoles could be prepared by using *o*-iodoacetanilides in place of the *o*-iodophenols in the multicomponent coupling reaction.<sup>6</sup> Accordingly, reaction of **12** with **4** and **7**, under the multicomponent coupling conditions described above, gave **13** (85%) (eq 1).<sup>6</sup> The acyl group



was lost during the course of the reaction.<sup>6</sup>

Alternatively, indoles could be obtained at room temperature in a one-pot reaction using a variant of the stepwise sequence described by Cacchi.<sup>9</sup> Accordingly,


### Scheme 2<sup>a</sup>



<sup>a</sup> Reagents and conditions: (i) Pd(PPh<sub>3</sub>)<sub>2</sub>Cl<sub>2</sub> 3 mol %, Et<sub>3</sub>N 2 equiv, CuI 6 mol %, CH<sub>3</sub>CN, 18 °C, 1 h under N<sub>2</sub>(g); (ii) K<sub>2</sub>CO<sub>3</sub> 5 equiv, **7**, 18 °C, 18 h; (iii) same as (ii) except exchange N<sub>2</sub>(g) atmosphere for CO(g) atmosphere (balloon); (iv) AlCl<sub>3</sub> 4 equiv, CH<sub>2</sub>Cl<sub>2</sub>.

the *o*-iodotrifluoroacetanilide **14** was coupled to alkyne **5** under Sonogashira conditions in acetonitrile to give **15** (Scheme 2). When this coupling was complete (approximately 2 h), the aryl iodide **7** and 5 equiv of K<sub>2</sub>CO<sub>3</sub> were added and the solution was left stirring for a further 18 h to give **16** (77%). As before, when the N<sub>2</sub>(g) atmosphere was exchanged for one of CO(g) at the point of introduction of **7**, the product obtained was the carbonylated product **18** (73%). Both **16** and **18** were deprotected to the corresponding phenols **17** and **19**, respectively, using AlCl<sub>3</sub>.<sup>7,10,11</sup> These alternative multicomponent coupling conditions are not applicable to the synthesis of 2,3-disubstituted benzo[*b*]furans because, under the Sonogashira conditions of coupling, the initially formed *o*-alkynylphenols rapidly cyclize to the simple C3-unsubstituted benzo[*b*]furans.<sup>6</sup>

Additional analogues were generated by reduction of ketones **9** and **11** to give alcohols **20** and **21**, respec-

Scheme 3<sup>a</sup>

<sup>a</sup> Reagents and conditions: (i) NaBH<sub>4</sub> 2 equiv, EtOH; (ii) CF<sub>3</sub>CO<sub>2</sub>H 10 equiv, Et<sub>3</sub>SiH 5 equiv, CH<sub>2</sub>Cl<sub>2</sub>.

tively, using sodium borohydride (Scheme 3). Alcohol **20** was further reduced to the simple methylene system **22** by reaction with trifluoroacetic acid and triethylsilane in excellent yield.

Compounds **8**, **9**, **11**, **13**, **17**, **19**, and **20–22** were evaluated for inhibition of tubulin assembly (Table 1). Those that displayed a significant inhibitory effect (defined as IC<sub>50</sub> < 5.0 μM) were also examined for an inhibitory effect on the binding of [<sup>3</sup>H]colchicine to tubulin (the tubulin used in these studies was purified to electrophoretic homogeneity from bovine brain). These compounds were also evaluated for cytotoxicity against MCF-7 human breast carcinoma cells (Table 1).

Compound **8** did not inhibit tubulin assembly at concentrations as high as 40 μM or cell growth at 1 μM and was not further examined, while compounds **13**, **17**, **20**, and **21** showed moderate to poor activity both as tubulin polymerization inhibitors and as cytotoxins. Compounds **9**, **11**, **19**, and **22** were all similar to or more potent than CA4 as inhibitors of tubulin polymerization. Compound **9** was the most active, exhibiting 5 times the level of potency of CA4. However, all of these compounds (**9**, **11**, **19**, and **22**) were less potent than CA4 as inhibitors of [<sup>3</sup>H]colchicine binding to tubulin and as cytotoxins (against MCF-7 human breast carcinoma cells). It is nonetheless noteworthy that the cytotoxicity of **9**, **11**, and **19** was only slightly less than that of CA4.

In terms of a structure–activity relationship (SAR) for benzofused heterocyclic compounds, those containing either a nitrogen or an oxygen in the heterocyclic ring are much more effective than their sulfur analogues in all three biological assays (compare **1** and **2** with **9**, **11**, and **19**). As had been previously demonstrated for the sulfur series, the absence of a one-carbon linker between C3 of the benzofused heterocycle and the trimethoxyphenyl ring leads to a complete or near-complete loss in activity (compare **8**, **13**, and **17** with **9**, **11**, and **19**).<sup>5</sup> This carbon linker is much more effective as a carbonyl group than as a carbinol or simple methylene group (compare **9** and **11** with **20–22**).

## Conclusion

Multicomponent coupling of *o*-iodophenols (or *o*-iodoacetanilides) with terminal alkynes and aryl iodides provides rapid access to potent benzo[*b*]furan- and indole-based tubulin polymerization inhibitors. These systems represent valuable new leads in the pursuit of anticancer chemotherapies.

**Acknowledgment.** The authors thank the Australian Research Council for financial support, including an Australian Research Fellowship to B.L.F. This work

was supported in part by NCI, National Institutes of Health Contract N01-CO-1240.

**Supporting Information Available:** Procedures for the synthesis of all new compounds in Table 1, including spectroscopic data. This material is available free of charge via the Internet at <http://pubs.acs.org>.

## References

- (1) (a) Ben-Chetrit, E.; Levy, M. Colchicine: 1998 update. *Semin. Arthritis Rheum.* **1998**, *28*, 48–59. (b) McKellar, Q. A.; Scott, E. W. The benzimidazole anthelmintic agents—a review. *J. Vet. Pharmacol. Ther.* **1990**, *13*, 223–247. (c) Rowinsky, E. K.; Donehower, R. C. The clinical pharmacology and use of antimicrotubule agents in cancer chemotherapeutics. *Pharmacol. Ther.* **1992**, *52*, 35–84. (d) Gregory, R. K.; Smith, I. E. Vinorelbine—a clinical review. *Br. J. Cancer* **2000**, *82*, 1907–1913. (e) Rowinsky, E. K.; Donehower, R. C. Paclitaxel (taxol). *N. Engl. J. Med.* **1995**, *13*, 1004–1014. (f) Cortes, J. E.; Pazdur, R. Docetaxel. *J. Clin. Oncol.* **1995**, *13*, 2643–2655.
- (2) (a) Chaplin, D. J.; Pettit, G. R.; Parkins, C. S.; Hill, S. A. Antivascular approaches to solid tumour therapy: evaluation of tubulin binding agents. *Br. J. Cancer* **1996**, *74*, S86–S88. (b) Dark, G. G.; Hill, S. A.; Prise, V. E.; Tozer, G. M.; Pettit, G. R.; Chaplin, D. J. Combretastatin A-4, an agent that displays potent and selective toxicity toward tumor vasculature. *Cancer Res.* **1997**, *57*, 1829–1834. (c) Tozer, G. M.; Prise, V. E.; Wilson, J.; Locke, R. J.; Vojnovic, B.; Stratford, M. R. L.; Dennis, M. F.; Chaplin, D. J. Combretastatin A-4 phosphate as a tumor vascular-targeting agent: early effects in tumors and normal tissues. *Cancer Res.* **1999**, *59*, 1626–1634. (d) Iyer, S.; Chaplin, D. J.; Rosenthal, D. S.; Boulares, A. H.; Li, L.-Y.; Smulson, M. E. Induction of apoptosis in proliferating human endothelial cells by the tumor-specific antiangiogenesis agent combretastatin A-4. *Cancer Res.* **1998**, *58*, 4510–4514. (e) Grosios, K.; Holwell, S. E.; McGown, A. T.; Pettit, G. R.; Bibby, M. C. In vivo and in vitro evaluation of combretastatin A-4 and its sodium phosphate prodrug. *Br. J. Cancer* **1999**, *81*, 1318–1327. (f) Chaplin, D. J.; Dougherty, G. J. Tumor vasculature as a target for cancer therapy. *Br. J. Cancer* **1999**, *80*, 57–64. (g) Horsman, M. R.; Murata, R.; Breidahl, T.; Nielsen, F. U.; Maxwell, R. J.; Stodkilde-Jorgensen, H.; Overgaard, J. Combretastatins: novel vascular targeting drugs for improving anticancer therapy. Combretastatins and conventional therapy. *Adv. Exp. Med. Biol.* **2000**, *476*, 311–323. (h) Chaplin, D. J.; Pettit, G. R.; Hill, S. A. Anti-vascular approaches to solid tumor therapy: evaluation of combretastatin A4 phosphate. *Anticancer Res.* **1999**, *19*, 189–196. (i) Galbraith, S. M.; Chaplin, D. J.; Lee, F.; Stratford, M. R. L.; Locke, R. J.; Vojnovic, B.; Tozer, G. M. Effects of combretastatin A4 phosphate on endothelial cell morphology in vitro and relationship to tumor vascular targeting activity in vivo. *Anticancer Res.* **2001**, *21*, 93–102. (j) Pero, R. W.; Sherris, D. PCT Int. Appl. WO 20000484006, 2000; *Chem. Abstr.* **2000**, *133*, 198575.
- (3) (a) Pettit, G. R.; Singh, S. B.; Hamel, E.; Lin, C. M.; Alberts, D. S.; Garcia-Kendall, D. Isolation and structure of the strong cell growth and tubulin inhibitor combretastatin A-4. *Experientia* **1989**, *45*, 209–211. (b) Lin, C. M.; Ho, H. H.; Pettit, G. R.; Hamel, E. Antimitotic natural products combretastatin A-4 and combretastatin A-2: studies on the mechanism of their inhibition of the binding of colchicine to tubulin. *Biochemistry* **1989**, *28*, 6984–6991. (c) Pettit, G. R.; Singh, S. B.; Boyd, M. R.; Hamel, E.; Pettit, R. K.; Schmidt, J. M.; Hogan, F. Antineoplastic agents. 291. Isolation and synthesis of combretastatins A-4, A-5, and A-6(1a). *J. Med. Chem.* **1995**, *38*, 1666–1672.
- (4) (a) Pinney, K. G.; Bounds, A. D.; Dingeman, K. M.; Mocharla, V. P.; Pettit, G. R.; Bai, R.; Hamel, E. A new anti-tubulin agent containing the benzo[*b*]thiophene ring system. *Bioorg. Med. Chem. Lett.* **1999**, *9*, 1081–1086. (b) Pinney, K. G.; Pettit, G. R.; Mocharla, V. P.; Del Pilar Majia, M.; Shirali, A. PCT Int. Appl. WO 9839323, 1998; *Chem. Abstr.* **1998**, *129*, 245037. (c) Mullica, D. F.; Pinney, K. G.; Mocharla, V. P.; Dingeman, K. M.; Bounds, A. D.; Sappenfield, E. L. Characterization and structural analyses of trimethoxy and triethoxybenzo[*b*]thiophene. *J. Chem. Crystallogr.* **1998**, *28*, 289–295. (d) Pinney, K. G.; Chen, Z.; Mocharla, V. P.; Choony, N.; Strong, T. Synthesis of a benzo[*b*]thiophene-based vascular targeting prodrug and related anti-tubulin ligands. Presented at the 220th American Chemical Society, National Meeting, Division of Organic Chemistry, Washington, DC, August 20–24, 2000; Abstract 196.
- (5) (a) Flynn, B. L.; Verdier-Pinard, P.; Hamel, E. A novel palladium-mediated coupling approach to 2,3-disubstituted benzo[*b*]thiophenes and its application to the synthesis of tubulin binding agents. *Org. Lett.* **2001**, *3*, 651–654. (b) Flynn, B. L.; Flynn, G. P.; Hamel, E.; Jung, M. K. The synthesis and tubulin binding activity of thiophene-based analogues of combretastatin A-4. *Bioorg. Med. Chem. Lett.* **2001**, *11*, 2341–2343.

- (6) Chaplin, J. H.; Flynn, B. L. A multi-component coupling approach to benzo[b]furans and indoles. *Chem. Commun.* **2001**, 1594–1595.
- (7) Banwell, M. G.; Flynn, B. L.; Stewart, S. G. Selective cleavage of isopropyl aryl ethers by aluminum trichloride. *J. Org. Chem.* **1998**, *63*, 9139–9144.
- (8) For experimental procedures, see: Verdier-Pinard, P.; Lai, J.-Y.; Yoo, H.-D.; Yu, J.; Marquez, B.; Nagle, D. G.; Nambu, M.; White, J. D.; Falck, J. R.; Gerwick, W. H.; Day, B. W.; Hamel, E. Structure–activity analysis of the interaction of curacin A, the potent colchicine site antimitotic agent, with tubulin and effects of analogs on the growth of MCF-7 breast cancer cells. *Mol. Pharmacol.* **1998**, *53*, 62–76.
- (9) (a) Arcadi, A.; Cacchi, S.; Marinelli, F. A versatile approach to 2,3-disubstituted indoles through the palladium-catalyzed cyclization of *o*-alkynyltrifluoroacetanilides with vinyl triflates and aryl halides. *Tetrahedron Lett.* **1992**, *33*, 3915–3918. (b) Arcadi, A.; Cacchi, S.; Carnicelli, V.; Marinelli, F. 2-Substituted-3-acylindoles through the palladium-catalyzed carbonylative cyclization of 2-alkynyltrifluoroacetanilides with aryl halides and vinyl triflates. *Tetrahedron* **1994**, *50*, 437–452.
- (10) During the course of these studies, indole **19** was also prepared and tested by K. G. Pinney and co-workers using conventional synthetic methods: Pinney, K. G.; Wang, F.; Del Pilar Mejia, M. PCT Int. Appl. WO 2001019794, 2001; *Chem. Abstr.* **2001**, *134*, 237388.
- (11) Von Angerer and co-workers have previously prepared and tested a series of 3-formyl-2-phenylindole inhibitors of tubulin polymerization: Gastpar, R.; Goldbrunner, M.; Marko, D.; von Angerer, E. Methoxy-substituted 3-formyl-2-phenylindoles inhibit tubulin polymerization. *J. Med. Chem.* **1998**, *41*, 4965–4972.

JM020077T