

Nucleosides, Nucleotides and Nucleic Acids

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/Incn20>

Microwave-assisted Ribosylation of modified heterocyclic bases by Vorbrüggen method

Nadja V. Nikolaus^a, Jelena Božilović^a & Joachim W. Engels^a

^a Institute for Organic Chemistry and Chemical Biology, Johann Wolfgang Goethe University, Frankfurt am Main, Germany

Published online: 10 Dec 2007.

To cite this article: Nadja V. Nikolaus, Jelena Božilović & Joachim W. Engels (2007) Microwave-assisted Ribosylation of modified heterocyclic bases by Vorbrüggen method, *Nucleosides, Nucleotides and Nucleic Acids*, 26:8-9, 889-892, DOI: [10.1080/15257770701505485](https://doi.org/10.1080/15257770701505485)

To link to this article: <http://dx.doi.org/10.1080/15257770701505485>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

MICROWAVE-ASSISTED RIBOSYLATION OF MODIFIED HETEROCYCLIC BASES BY VORBRÜGGEN METHOD

Nadja V. Nikolaus, Jelena Božilović, and Joachim W. Engels □ *Institute for Organic Chemistry and Chemical Biology, Johann Wolfgang Goethe University, Frankfurt am Main, Germany*

□ *During the last decades the nucleoside synthesis has proven to be important. The modified silyl-Hilbert-Johnson nucleoside synthesis modified by Vorbrüggen is one of the most often used methods. We have studied N-glycosilation of modified heterocyclic bases by Vorbrüggen method with microwave irradiation and we were able to shorten the reaction time and obtain higher yields. The method was demonstrated by fluoroquinolone and purine.*

Keywords Ribosylation; modified heterocyclic bases


INTRODUCTION

Nucleosides and nucleotides have provided a productive area of chemical and biological research. The monomeric units of DNA and RNA are involved in the regulation of a myriad of cellular metabolic pathways and have been the subject of intense areas of research seeking to identify therapeutic agents for a variety of diseases including viral infection, cancer, cardiovascular diseases, central nervous system diseases, etc. Among the various synthetic methods the reaction of silylated heterocyclic bases with peracylated sugars in the presence of Lewis acid catalysts has become the standard procedure which affords nucleosides routinely in high yields.^[1] In short, nucleoside synthesis includes the silylation of a heterocyclic base, the generative sugar cation from peracylated ribose and the nucleophilic attack of a heterocyclic base on C1.^[2]

Here we present some examples of ribosylation of modified heterocyclic bases (1–4) by the conventional Vorbrüggen method and with the use of

We would like to thank the EU project (EU Leuren LSHB-CT-2003-503480/TrioH) for financial support.

Address correspondence to Joachim W. Engels, Institute for Organic Chemistry and Chemical Biology, Johann Wolfgang Goethe University, Max von Laue Str. 7, 60438 Frankfurt am Main, Germany. E-mail: joachim.engels@chemie.uni-frankfurt.de



SCHEME 1 Reagents and conditions: a) 1,2,3,5-*O*-tetraacetyl-D-ribofuranose, MeCN, bis(trimethylsilyl)acetamide, trimethylsilyl triflate, Δ, 3 hours, 82–92% or 1,2,3,5-*O*-tetraacetyl-D-ribofuranose, MeCN, bis(trimethylsilyl)acetamide, trimethylsilyl triflate, MW, 50 minutes, 80–96%.

microwave irradiation (MWI). The last approach permits us to get higher yields of target nucleosides in a shorter time compared to the standard method (Scheme 1 and Table 1).

RESULTS AND DISCUSSION

In the synthesis of the heterocycles high temperatures are needed, and we decided to use microwave applicator. Therefore, we modified the procedure of Kidwai et al.^[3] on the first step. 2-, 3-, and 4-Fluoroanilines were *N*-substituted with diethyl (ethoxymethylidene)malonate to give diethyl[(fluoroanilino)methylidene]malonates. The electrophilic ring closure was performed in diphenyl ether according to Koga's procedure.^[4] The synthesis of fluoroquinolones was reduced from a two-step synthesis by hydrolysis and decarboxylation to a single-step, one-pot reaction.^[5] Using this method we were able to shorten the synthesis by one step and to obtain very good yields (87–96%).

Several attempts to synthesize the nucleosides failed due to the lack of solubility of the fluoroquinolones during Vorbrüggen reaction. A slightly modified Vorbrüggen method of Moore et al.^[6] gave better results than general procedure, but we improved efficiency of the method by using microwave irradiation on steps of the silylation of heterocyclic bases and ribosylation of them. We do explain these results by a combination of optimal solubility and heat distribution due to the microwave irradiation.

The microwave-assisted glycosylation is the best method for the preparation of fluoroquinolone nucleosides **7** and **8** because the yields are higher and the reaction time is much shorter compared to the conventional method. But in the case of ethyl-6-fluoro-4-oxo-1,4-dihydroquinoline-3-carboxylate, the yield of nucleoside **6** microwave irradiated synthesis is

TABLE 1

Compound			Compound			Compound			Compound		
6			7			8			9 and 10		
Conventional	Microwave-assisted		Conventional	Microwave-assisted		Conventional	Microwave-assisted		Conventional	Microwave-assisted	
Silylation			Silylation			Silylation			Silylation		
80 °C reflux	80 °C		80 °C reflux	80 °C		80 °C reflux	80 °C		80 °C reflux	80 °C	
30 min	15 min		30 min	15 min		30 min	15 min		15 min	15 min	
—	150 W		—	150 W		—	150 W		—	150 W	
Nucleoside synthesis			Nucleoside synthesis			Nucleoside synthesis			Nucleoside synthesis		
80 °C reflux	80 °C		80 °C reflux	80 °C		80 °C reflux	80 °C		80 °C reflux	80 °C	
3 h	50 min		3 h	50 min		3 h	50 min		2.5 h	50 min	
—	150 W		—	150 W		—	150 W		—	150 W	
Yield			Yield			Yield			Yield		
92%	63%		39%	80%		64%	96%		66% 9	64.2% 9	
									11.6% 10	17.1% 10	

lower than the one of the conventional method. It is still a good method to prepare a nucleoside due to the shorter reaction time and so we can save time in the synthesis. The synthesis of compound **10** with making use of MWI also is preferential, because it seems to favor a kinetic controlled reaction and the ratio of desired nucleoside **10** in the total yield of isomeric N-7 and N-9 nucleosides is changed from 1:5.7 to 1:3.7 (compound **10** to compound **9**).

REFERENCES

1. Vorbrüggen, H.; Höfle, G. On the mechanism of nucleoside synthesis. *Chem. Ber.* **1981**, 114, 1256–1268.
2. Zivkovic, A. RNA recognition by fluoro aromatic substituted nucleic acid analogues. *Dissertation* **2005**, 49–50.
3. Kidwai, M.; Misra, P.; Kumar, R.; Safena, R. K.; Gupta, R.; Bradoo, S. Microwave assisted synthesis and antibacterial activity of new quinolone derivatives. *Monatsh. Chem.* **1998**, 129, 961–965.
4. Koga H.; Itoh, A.; Murayama, S.; Suzue, S.; Irikura, T. Structure-activity relationships of antibacterial 6,7- and 7,8-disubstituted 1-alkyl-1,4-dihydro-4-oxoquinoline-3-carboxylic acids. *J. Med. Chem.* **1980**, 23, 1358–1363.
5. Adams, M.M.; Bats, J.W.; Nikolaus, N.V.; Witvrouw, M.; Debeyser, Z.; Engels, J. W. Microwave-assisted synthesis of fluoroquinolone nucleosides as inhibitors of HIV integrase. *Collection of Czechoslovak Chemical Communications* **2006**, 71 (7), 978–990.
6. Moore, C.L.; Zivkovic, A.; Engels, J.W.; Kuchta, R.D.; Human DNA primase uses Watson-Crick hydrogen bonds to distinguish between correct and incorrect nucleoside triphosphates. *Biochemistry* **2004**, 43, 12367–12374.