A Highly Active Catalyst for the Hydrogenation of Amides to Alcohols and Amines**

Jeremy M. John and Steven H. Bergens*

Alcohols and amines are ubiquitous in the synthesis of agrochemicals, pharmaceuticals (e.g. protection, deprotection), flavorings, fragrances, and advanced materials.^[1] One approach to accessing these compounds is through the reduction of amides. Amides are, however the most stable carboxylic acid derivative.^[2] Consequently, the reduction of amides typically requires stoichiometric amounts of active Al–H,^[3] B–H,^[3] or Si–H^[4] reducing agents that often cause reductive cleavage of the C=O bond.

Numerous heterogeneous catalysts have been developed to hydrogenate amides. These include copper/chromite systems that give mixtures of amine products under 350 atm of H₂ at temperatures of 250–400 °C.^[5] Co-catalysts of Rh or Ru with Re, W, or Mo hydrogenate amides either by reductive cleavage of the C=O bond (100 atm H₂, 160–180 °C),^[6] or by selectively hydrogenating primary amides to the corresponding primary amines (20–100 atm H₂, 130–160 °C).^[7]

There are a handful of homogeneous systems that catalyze the hydrogenation of amides or amide derivatives. The first is a Ru/triphos system (triphos = 1,1,1-tris(diphenylphosphinomethyl)ethane) that hydrogenates primary amides with a preference for the reductive cleavage of the C=O bond in the presence of NH₃ (40 atm H₂, 140-164 °C, 14 h).^[8] Beginning in 2006, Ikariya et al. reported dihydrogenations of cyclic imides,^[9a,c] N-acylcarbamates, N-sulfonyllactams, N-acylsulfonamides,^[9b] N-phenyllactams, and benzamides^[9d,e] with reductive cleavage of the C-N bond catalyzed by [Cp*RuCl(PN)] (Cp* = η^5 -C₅(CH₃)₅, e.g. PN = Ph₂P- $(CH_2)_2NH_2$ or [Cp*RuCl(LN)] (e.g. $LN = 2-C_5H_4NCH_2$ -NH₂) under the reported reaction conditions (tBuOH or 2-PrOH, 80-100°C, 30-50 atm, KOtBu 1-2.5 equiv, 2-72 h). Our group recently reported the enantioselective monohydrogenation of meso-cyclic imides to give hydroxy lactams using trans-[Ru(H)₂(binap)(dpen)] (binap = 2,2'-bis(diphenylphosphino)-1,1'-binaphthyl, dpen = 1,2-diphenylethylenediamine) and related complexes in THF at low temperatures

[*] J. M. John, Prof. Dr. S. H. Bergens Department of Chemistry, University of Alberta Edmonton, Alberta, T6G 2G2 (Canada) E-mail: steve.bergens@ualberta.ca

[**] This work was supported by the Natural Sciences and Engineering Research Council of Canada (NSERC), the Government of the Republic of Trinidad and Tobago (GoRTT), and the University of Alberta. We gratefully appreciate the assistance of Mark Miskolzie, Nupur Dabral, and Mickey Richards at the University of Alberta High Field NMR laboratory and we would like to express our gratitude to Dr. Satoshi Takebayashi for his advice and numerous discussions.

Supporting information for this article is available on the WWW under http://dx.doi.org/10.1002/anie.201103137.

(0.1 mol % [Ru], 0°C, 50 atm H₂, 9 mol % *t*BuOK, 17– 57 h).^[10] The most active system to date, reported by Milstein and co-workers, is the dearomatized, bipyridyl-based PNN/ Ru complex (PNN = (2-(di-*tert*-butylphosphinomethyl)-6-(diethylaminomethyl)pyridine) that hydrogenates a variety of secondary amides, and tertiary amides having ether groups to give the alcohol and amine products (1 mol % [Ru] in THF, base free, 110 °C, 10 atm H₂, 48 h).^[11]

We recently reported the low-temperature preparation and study of the Noyori ketone hydrogenation catalyst *trans*- $[Ru((R)-binap)(H)_2((R,R)-dpen)]$ (1).^[12] Compound 1 is remarkably active towards carbonyl reduction. For example, 1 adds acetophenone upon mixing and adds γ -butyrolactone within minutes at -80 °C to form the alkoxide *trans*-[Ru((R)binap)(H)(OCH(CH₃)(Ph))((R,R)-dpen)],^[12b] and the corresponding Ru/hemiacetaloxide of γ -butyrolactone. The complex 1 also catalyzes the hydrogenation of ethyl hexanoate under 4 atm H₂ below a temperature of 0 °C,^[13] and the monohydrogenation of *meso*-cyclic imides at 0 °C.^[10] We report herein the results of our study of 1 and related compounds as catalysts for the hydrogenation of amides.

We found that the activity of **1** towards the activated amides *N*-methylsulfonylpyrrolidin-2-one (**2a**) and *N*-acetylpyrrolidin-2-one (**2b**) was low to moderate. Compound **2a** was hydrogenated with a turnover number (TON) of approximately 27 to give the ring-opened *N*-methanesulfonyl amino alcohol product when using 2 mol% [Ru] in THF (100 °C, 50 atm, 20 mol% KOtBu, 39 h) [Eq. (1); Ms = methanesulfonyl]. Substrate **2b** formed mixtures of pyrrolidine-2-one (major) and the ring-opened N-acetyl amino alcohol with a TON of approximately 45 using 2 mol% [Ru] (80 °C, 50 atm H₂, 20 mol% KN[Si(CH₃)₃]₂, 16 h). *N*-phenylpyrrolidin-2-one (**2c**) was inactive under our reaction conditions.



These results are in contrast to the high activity of **1** towards the reduction of ketones, esters, and imides in THF.^[10,12,13] We reasoned that catalysts such as **1** are intrinsically active towards amide hydrogenation, but they decompose^[12] at the higher temperaturerequired for this transformation. We hypothesized that tethering the amine and

Communications

phosphine groups would maintain activity, and prevent dissociative loss of the diamine at high temperatures. We found that reaction between 2 equivalents of $Ph_2P(CH_2)_2NH_2$ (3) and the Ru precursor *cis*-[Ru(CH₃CN)₂(η^3 -C₃H₅)-(cod)]BF₄ (4; cod = 1,5-cyclooctadiene) in THF at 60 °C forms isomers of the π -allyl complex **5** in near-quantitative yield (in solution) by displacement of the cod and MeCN ligands [Eq. (2)].^[14,15,17]



51% combined yield of isolated products

Table 1 and Table 2 summarize the results of our amide hydrogenations using **5** and KN[Si(CH₃)₃]₂ as the added base in THF. All hydrogenations were carried out with 0.1 mol% [Ru] , 4–5 mol% KN[Si(CH₃)₃]₂, and 50 atm H₂ at 100 °C, for 24 hours. To our pleasant surprise, *N*-phenylpyrrolidin-2-one (**2c**) was hydrogenated to give *N*-phenyl-4-aminobutan-1-ol in 100% yield, or with a TON of 1000 under these reaction conditions (entry 1, Table 1). The N–Me (entry 2) and N–H (entry 3) derivatives were much less active than **2c**, whereas the six-membered N–Ph derivative **6** reacted in 100% yield (entry 4). The seven-membered unsubstituted lactam **7** (entry 5) was more reactive than the five-membered lactam **2e** (entry 3), as expected from the greater stability of fiveover seven-membered rings.

Table 1: Hydrogenation of lactams using 5 and KN[Si(CH₃)₃]₂.^[a,b]

	R + 2 H ₂ 7 50 atm	0.1 mol% 5 5 mol% [(C⊦ THF, 100°C,	H ₃)₃Si]₂NK 24h	но	¦ `R
2c 2d	:: R = Ph, <i>n</i> = 1 I: R = Me, <i>n</i> = 1	2e : R = I	H, <i>n</i> = 1	6 : R = Ph, <i>n</i> = 2 7 : R = H, <i>n</i> = 3	
Entry	Subs	trate	Yield	[%] ^[c]	TON
1	2c		100		1000
2	2 d		5		50
3	2e		0		0
4	6		100		1000
5	7		23		230

[a] Performed using in situ prepared catalyst precursor **5**. [b] Reaction conditions: H₂ at 50 atm, 100 °C, **5**/KN[Si(CH₃)₃]₂=1:50, [Substrate] = 0.626 $\,M$ in THF. [c] Determined by ¹H NMR spectroscopy.

The order of reactivity among the acyclic benzamides was $-N(Ph)_2 \approx -N(Ph)Me > -N(Me)_2$ (entries 1–3, Table 2). This

Table 2: Hydrogenation of acyclic amides using **5** and $KN[Si(CH_3)_3]_2^{[a,b]}$

J	$R^{2} + 2H_{2}$	0.1 m	ol% 5	-		\mathbb{R}^2
R ¹ 50 atm 8		4 mol% [(CH ₃) ₃ Si] ₂ NK THF, 100℃, 24h				
Entry	Substrate	R	R ¹	R ²	Yield [%] ^[c]	TON
1	8a	Ph	Ph	Ph	100 ^[d]	1000
2	8 b	Ph	Ph	Me	96	960
3	8c	Ph	Me	Me	50	500
4	8 d	Ph	-(CH ₂) ₅ -	-	82	820
5	8e	Ph	Ph	Н	50	500
6	8 f	Ph	Me	н	27	270
7	8 g	Me	Ph	Ph	100	1000
8	8 h	Me	Ph	Me	100	1000
9	8i	Me	Me	Me	50 ^[e]	500
10	8j	Me	Ph	Н	70	700

[a] Performed using in situ prepared catalyst precursor **5**. [b] Reaction conditions: H_2 at 50 atm, 100°C, **5**/KN[Si(CH₃)₃]₂=1:40, [Substrate] = 0.626 M in THF. [c] Determined by ¹H NMR spectroscopy. [d] 72% benzyl alcohol, 14% benzyl benzoate. [e] Anthracene used as an internal standard.

order is consistent with the differences in the extent of donation from the lone pair of electrons on the nitrogen atom to the carbonyl carbon center of these substrates. 1-Benzoyl-piperidine (8d; entry 4) was more active than 8c (entry 3), whereas secondary amides were somewhat less reactive than tertiary amides (entry 5 versus entry 1 and entry 6 versus entry 3). Similar results were obtained with acyclic acetamides. Specifically, the reactivity was on the order of $-N(Ph)_2 \approx -N(Ph)Me > -N(Me)_2$ (entries 7–9). The secondary acetamide -N(Ph)H 8j (entry 10) was less reactive than the corresponding tertiary amide (entry 8). The lower reactivity of secondary versus tertiary amides may arise from reaction of the secondary amide with the added base.

In preliminary experiments, we found that **5** reacts with H_2 (ca. 1 atm) and KN[Si(CH₃)₃]₂ (ca. 3 equiv) in [D₈]THF starting at approximately 0 °C to form propylene and three ruthenium monohydrides.^[16] The known^[17] dichloride [Ru(Cl)₂(Ph₂P(CH₂)₂NH₂)₂] (**11**) gives a similar mixture of monohydride species under these reaction conditions. This mixture reacts further (ca. 4 atm H₂, ca. 10 equiv KN[Si-(CH₃)₃]₂) at room temperature to generate a symmetrical dihydride as the major product, which we tentatively assign to be an isomer of *trans*-[Ru(H)₂(Ph₂P(CH₂)₂NH₂)₂] (**9**).^[18,19] Saudan et al. reported that **11** forms an active ester hydrogenation catalyst with NaOMe as the base in THF.^[17] Indeed, we found that **5** and **11** (0.01 mol%) both hydrogenate **2c** with remarkable TONs of 7120 and 6760, respectively, in the presence of 5 mol% NaOMe (Table 3).

On the basis of prior studies with ketone and ester substrates and $\mathbf{1}$,^[12,13] we propose the following mechanism for the hydrogenation of amides with **9** (Scheme 1). The first step is a bifunctional-type addition of the amide to **9**, thus forming the ruthenium hemiaminaloxide **12** as the net product. Base-assisted elimination from **12** forms the ruthenium/amide **10**, and the free hemiaminaloxide **13** which regenerates the base and eliminates aldehyde and H-NR¹₂. Addition of H₂ to **10**



Table 3: Hydrogenation of N-phenylpyrrolidin-2-one using 5 or 11 and NaOMe. $^{\rm [a,b]}$

1 2	5 11		71.2 67.6	7120 6760
Entry	Catalyst precursor		Yield [%] ^[c]	TON
2c	1 + 2 H ₂ 50 atm	5 mol% NaOMe THF, 100°C, 24h	→ HO ~	∽ Ph
		0.01 mol% 5 or 11		H N. DL

[a] Performed using the isolated catalyst precursor **5**. [b] Reaction conditions: H_2 at 50 atm, 100 °C, **5** or **11**/NaOMe = 1:500, [Substrate] = 2.08 μ in THF. [c] Determined by ¹H NMR spectroscopy.



Scheme 1. Proposed mechanism for the hydrogenation of amides.

regenerates 9 and hydrogenation of the aldehyde forms the alcohol. Consistent with this mechanism is the formation of benzyl benzoate by the Tishchenko reaction of benzaldehyde during the hydrogenation of 8a (entry 1, Table 2).We have shown that 5 and 11 are remarkably active towards the hydrogenation of a series of amides without strongly activating functional groups. Significantly, the commonly inert, unfunctionalized amide dimethyl acetamide was hydrogenated with TONs of 500, and *N*-phenylpyrrolin-2-one was hydrogenated with TONs of up to 7120. Studies on catalyst variations, the mechanism, and the synthetic scope of these hydrogenations are underway in our laboratories.

Received: May 6, 2011 Revised: June 23, 2011 Published online: August 26, 2011

Keywords: amides \cdot green chemistry \cdot homogeneous catalysis \cdot hydrogenation \cdot N,P ligands

 a) A. Ricci, Modern Amination Methods, Wiley, New York, 2000; b) Modern reduction methods (Eds.: P. G. Andersson, I. J. Munslow), Wiley, New York, **2008**; c) J. S. Carey, D. Laffan, C. Thomson, M. T. Williams, *Org. Biomol. Chem.* **2006**, *4*, 2337.

- [2] a) C. M. Breneman, M. Martinov in *The Amide Linkage: Structural Significance in Chemistry, Biochemistry and Materials Science* (Eds.: A. Greenberg, C. M. Breneman, J. F. Liebman), Wiley-VCH, Hoboken, **2003**, pp. 1–33; b) M. B. Robin, F. A. Bovey, H. Basch in *The Chemistry of Amides* (Ed.: J. Zabicky), Interscience, New York, **1970**, pp. 1–72.
- [3] a) J. Seyden-Penne, Reductions by the Alumino and Borohydrides in Organic Synthesis, 2nd ed., Wiley-VCH, Weinheim, 1997; G. W. Gribble, Chem. Soc. Rev. 1998, 27, 395.
- [4] For recent examples, see: a) G. Pelletier, W. S. Bechara, A. B. Charette, J. Am. Chem. Soc. 2010, 132, 12817; b) S. Das, D. Addis, S. Zhou, K. Junge, M. Beller, J. Am. Chem. Soc. 2010, 132, 1770; c) Y. Sunada, H. Kawakami, T. Imaoka, Y. Matoyama, H. Nagashima, Angew. Chem. 2009, 121, 9675; Angew. Chem. Int. Ed. 2009, 48, 9511.
- [5] a) B. Wojcik, H. Adkins, J. Am. Chem. Soc. 1934, 56, 2419;
 b) R. M. King, US Pat. 4,448,998, May 15th, 1984.
- [6] C. Hirosawa, N. Wakasa, T. Fuchikami, *Tetrahedron Lett.* 1996, 37, 6749.
- [7] a) G. Beamson, A. J. Papworth, C. Philipps, A. M. Smith, R. J. Whyman, *J. Catal.* 2011, 278, 228; G. Beamson, A. J. Papworth, C. Philipps, A. M. Smith, R. J. Whyman, *Adv. Synth. Catal.* 2010, 352, 869; G. Beamson, A. J. Papworth, C. Philipps, A. M. Smith, R. J. Whyman, *J. Catal.* 2010, 296, 93.
- [8] a) M. Kilner, D. V. Tyers, S. P. Crabtree, M. A. Wood, PCT Int. Pat. Appl. WO 03/093208A1, Nov 13th, 2003; b) A. A. N. Magro, G. R. Eastham, D. Cole-Hamilton, *Chem. Commun.* 2007, 3154; A. A. N. Magro, G. R. Eastham, D. Cole-Hamilton, US Pat. 2010/0010261A1, Jan 14th, 2010.
- [9] a) M. Ito, A. Sakaguchi, C. Kobayashi, T. Ikariya, J. Am. Chem. Soc. 2007, 129, 290; b) M. Ito, L. W. Koo, A. Himizu, C. Kobayashi, A. Sakaguchi, T. Ikariya, Angew. Chem. 2009, 121, 1350; Angew. Chem. Int. Ed. 2009, 48, 1324; c) M. Ito, C. Kobayashi, A. Himizu, T. Ikariya, J. Am. Chem. Soc. 2010, 132, 11414; d) T. Ikariya, M. Ito, T. Ootsuka, PCT Int. Pat. Appl. WO 2010/073974 A1, July 1st, 2010; e) M. Ito, T. Ootsuka, R. Watari, A. Shiibashi, A. Himizu, T. Ikariya, J. Am. Chem. Soc. 2011, 133, 4240.
- [10] a) S. Takebayashi, J. M. John, S. H. Bergens, *J. Am. Chem. Soc.* **2010**, *132*, 12832; b) S. Takebayashi, S. H. Bergens, PCT Int. Pat. Appl. WO 2010/145024 A1, June 17th, **2010**.
- [11] E. Balaraman, B. Gnanaprakasam, L. J. W. Shimon, D. Milstein, J. Am. Chem. Soc. 2010, 132, 16756.
- [12] a) R. J. Hamilton, C. G. Leong, G. Bigam, M. Miskolzie, S. H. Bergens, J. Am. Chem. Soc. 2005, 127, 4152; b) R. J. Hamilton, S. H. Bergens, J. Am. Chem. Soc. 2006, 128, 13700; R. J. Hamilton, S. H. Bergens, J. Am. Chem. Soc. 2008, 130, 11979.
- [13] S. Takebayashi, S. H. Bergens, Organometallics 2009, 28, 2349.
- [14] J. A. Wiles, C. J. A. Daley, R. J. Hamilton, C. G. Leong, S. H. Bergens, Organometallics 2004, 23, 4564.
- [15] Compound 5 was identified by ¹H, ³¹P, gCOSY, ¹H{³¹P} gCOSY, ¹H-³¹P gHSQC, TROESY, and gTOCSY NMR experiments, mass spectrometry, and elemental analysis.
- [16] Overlap among the signals corresponding to the arene, N–H, and aliphatic protons in the ¹H NMR spectra made a conclusive identification impossible. Their reactivity with H₂, and that they are formed from 5 and 11 suggests that they are isomers of the ruthenium/amide [Ru(H)(Ph₂P(CH₂)₂NH₂)(Ph₂P(CH₂)₂NH)] (10).
- [17] a) L. Saudan, C. M. Saudan, C. Debieux, P. Wyss, Angew. Chem.
 2007, 119, 7617; Angew. Chem. Int. Ed. 2007, 46, 7473; b) L. Saudan, P. Dupau, J. Riedhauser, P. Wyss, PCT Int. Pat. Appl. WO 2006/106484 A1, Oct 12th, 2006; c) L. Saudan, C. Saudan, PCT Int. Pat. Appl. WO 2008/065588 A1, June 5th, 2008; L.

Angew. Chem. Int. Ed. 2011, 50, 10377–10380

Communications

Saudan, C. Saudan, PCT Int. Pat. Appl. WO 2010/038209A1, April 8th, 2010; d) L. Saudan, Acc. Chem. Res. 2007, 40, 1309.

[18] This preliminary assignment is based upon the similarities between the ³¹P{¹H} and ¹H (hydride) NMR spectra between **9** and **1** (see Ref. [12]). The peaks for compound **9** were assigned using ¹H, ³¹P, gCOSY, ¹H-³¹P gHSQC, and gTOCSY NMR experiments. ³¹P{¹H} NMR (161.903 MHZ, [D₈]THF, 27°C): 56.2 ppm (s). ¹H NMR (399.951 MHZ, [D₈]THF, 27°C): $\delta =$

8.26 ppm (Ru-*H*, t, J = 14.8 Hz), $\delta = 1.25$ ppm, $\delta = 2.43$ ppm, $\delta = 2.73$ ppm.

[19] Use of 4 as a catalyst precursor, or 5 without added base (both at 10 mol%), did not result in hydrogenation. Use of Ru black (10 mol%) resulted only in hydrogenation of the arene ring in 2c. It is therefore unlikely that Ru nanoparticles are the active catalyst in these hydrogenations.