Ŕ

Up to 81% yield, 9/1 d.r

Ee: >99/99%

# Palladium(II)-Catalyzed Formal [3 + 2] Cycloaddition of Aziridines with 3-Substituted Indoles: Synthesis of Enantioenriched Pyrroloindolines

You-ming Huang, Chang-wu Zheng, Lu Pan, Qiao-wen Jin, and Gang Zhao\*

Key Laboratory of Synthetic Chemistry of Natural Substances, Shanghai Institute of Organic Chemistry, Chinese Academy of Sciences, 345 Lingling Road, Shanghai, 200032, China

**Supporting Information** 

**ABSTRACT:** A Pd-catalyzed enantiospecific formal [3 + 2] cycloaddition between chiral aziridines and indoles has been developed. With this method, chiral pyrroloindolines in enantiomerically pure forms were constructed in high yields and diastereoselectivities under mild conditions.

# INTRODUCTION

Aziridines have been extensively utilized as one of the most useful three membered ring functionalities in the reactions with a variety of carbon and heteroatom nucleophiles for the construction of various amino acids, chiral ligands, nitrogencontaining heterocycles, and alkaloids.<sup>1,2</sup> During the past few years, with the easily accessible enantiopure aziridines, a couple of stereoselective ring-opening reactions have been reported to provide nonracemic products directly. Ghorai and co-workers<sup>3</sup> have reported extensive work on the Lewis acid catalyzed S<sub>N</sub>2type ring-opening of enantiopure N-activated aziridines to provide the corresponding products with partial loss of enantiopurity in almost all cases. Recently, Takeda and coworkers developed a Pd-catalyzed enantiospecific crosscoupling of chiral 2-arylaziridines with arylboronic acids for the construction of 2-arylphenethylamines in an enantiopure form using N-heterocyclic carbene (NHC) ligands to promote the coupling.<sup>4</sup> In this context, an efficient method for regioselective and stereospecific ring-opening of chiral aziridines is still in great demand.

Pyrroloindoline backbones are present in an array of fascinating natural products with interesting biological activities such as (–)-physostigmine, <sup>5a</sup> (–)-debromoflustramine E, <sup>5b</sup> (–)-gliocladin C, <sup>5c</sup> (+)-folicanthine, <sup>5d</sup> (–)-bionectin A, <sup>5e</sup> and (+)-nocardioazine B<sup>5f</sup> (Figure 1). Considerable efforts have been devoted to developing elegant strategies for the construction of hexahydropyrrolo[2,3-*b*]indoles, which, to the best of our knowledge, can be accessed through four general strategies concerning the enantioselective synthesis.

The first strategy is started with tryptophan or tryptamine derivatives by using biomimetic methods<sup>6</sup> (Scheme 1, route a). The second strategy by using 2-oxoindoles as the starting materials undergoes alkylation and a series of functional group interconversions and cyclization<sup>7</sup> (Scheme 1, route b). The third pathway is developed by Michael-type Friedel–Crafts alkylations of indoles with *N*-acylated  $\alpha,\beta$ -dehydroamino esters<sup>8</sup> (Scheme 1, route c). Recently, Davies and co-workers reported

the Rh-catalyzed [3 + 2]-annulation of C3-substituted indoles with 4-aryl-1-sulfonyl-1,2,3-triazoles to provided chiral didehydropyrroloindolines<sup>9</sup> (Scheme 1, route d). In 2000, Nakagawa's group reported the [3 + 2] cycloaddition of the unsubstituted aziridine with N-methyl-3-methylindoles promoted by stoichimetric amount of Sc(OTf)<sub>3</sub> with TMSCl as the additive.<sup>10</sup> Wang and co-workers studied the cycloaddition between mesoaziridines and C3-alkylindoles mediated by magnesium catalyst.<sup>11</sup> Using their methods, pyrroloindolines with no substitution or with alkyl-substituted groups in the pyrrole rings were constructed smoothly (Scheme 1, route e). However, the syntheses of pyrroloindolines with arylsubstituted groups are not sufficiently investigated; herein, we would like to present our contributions to the construction of these backbones via a [3 + 2] cycloaddition reaction between C3-substituted indoles and achiral or chiral 2-arylaziridines catalyzed by the simple Pd(II) salts.<sup>12</sup>

Pd(PhCN)<sub>2</sub>Cl<sub>2</sub>

regioselective

mild conditions

no ligands no epimerization

# RESULTS AND DISCUSSION

Our initial investigation was commenced with the reaction between C3-methylindole 1a and aziridine 2a (as a racemic mixture) catalyzed by various Lewis acids in toluene (Table 1). When the commonly used  $Zn(OTf)_2$ ,  $Mg(OTf)_2$ , and  $Cu-(OAc)_2$  were employed as the catalysts, no products were observed (Table 1, entries 1–3). Fortunately, the alternative  $Cu(OTf)_2$  and  $Sc(OTf)_3$  did provide the desired products, although the yields and diastereoselectivities were low (Table 1, entries 4, 5). The Pd(II) complexes, which are more popular in transition metal catalysis, have also been applied as Lewis acids to promote a variety of reactions.<sup>13</sup> Using PdCl<sub>2</sub> as the catalyst, the reaction indeed happened with the yield and diastereoselectivity largely improved. Thus, various Pd(II) catalysts were then examined and Pd(PhCN)<sub>2</sub>Cl<sub>2</sub> was found to give the best results (69% yield and 6:1 dr, Table 1, entry 7). In the

Received: August 19, 2015



Figure 1. Representative natural products containing the pyrroloindolines.

Scheme 1. Retrosynthetic Approaches



following studies, solvents such as MeCN, DCM, THF, or DMF were screened; however, the yields and diastereoselectivities were not improved. The study of the effect of temperature on the diastereoselectivity was also made; the lower reaction temperature gave the better diastereoselectivity (see Supporting Information, page S2). Finally, when the ratio of **2a:1a** was increased to 2:1, the corresponding product could be obtained in 81% yield and 10:1 dr (Table 1, entries 12). In addition to the Pd(II) catalyst, another transition metal PtCl<sub>2</sub> was also evaluated, and the reaction underwent smoothly; however, only 5:1 dr was observed (Table 1, entry 13).

Under the optimized conditions, the reaction scope was examined (Table 2). The reaction worked well with a series of indoles. Using the methyl group instead of benzyl group to protect the nitrogen of indole, the yield and diastereoselectivity were preserved (**3b**). Indoles with various types of substituents on the phenyl rings provided the products with good to excellent yield, and moderate to good dr (3c-3g), regardless of their electronic nature or positions. In addition, the reactions of

cyclopenta- and cyclohexa-fused indoles also proceeded smoothly, giving tetracyclic indolines **3h** and **3i** containing two quaternary bridging carbons in high yields with excellent diastereoselectivities. What is more, indole C3 substituents other than methyl group are also compatible in the reaction, albeit with moderate yields and stereoselectivities (3j-3l). As for the aziridines, the electronic nature of substituents on the 2aryl group did not affect the reactivity significantly (3m-3q). However, the *o*-Br-substituted 2-aryl aziridine gave the corresponding product (3r) with decreased diastereoselectivity compared to the *p*-Br-substituted 2-aryl aziridine, probably because of the steric effect of bromine at the ortho position of the phenyl ring. In addition, reactions with aziridines protected by different sulfonyl groups also proceed smoothly (3s-3u).

Further exploration was also conducted to construct chiral hexahydropyrrolo[2,3-*b*]indole backbones. For this purpose, enantiomerically pure aziridines were employed as the substrates. To our great delight, the ring-opening reaction between chiral aziridines and indoles with various substituents

Table 1. Optimization of the Reaction Conditions<sup>a</sup>

N Bn 1a	+ N Cat. (10 Ph Toluene, 2a	mol%) 10h, r.t.	Ph N Ts <b>3a</b>
entry	catalyst	yield [%] <sup>b</sup>	dr <sup>c</sup>
1	$Zn(OTf)_2$	NR	d
2	$Mg(OTf)_2$	NR	d
3	$Cu(OAc)_2$	NR	d
4	$Cu(OTf)_2$	40	1:1
5	Sc(OTf) <sub>2</sub>	24	1:1
6	PdCl <sub>2</sub>	65	6:1
7	$Pd(PhCN)_2Cl_2$	69	6:1
8	$Pd(OAc)_2$	43	9:1
9	$Pd(O_2CCF_3)_2$	24	5:1
10	$Pd(NO_3)_2$	53	5:1
11	$Pd(PPh_3)_2Cl_2$	NR	d
$12^e$	$Pd(PhCN)_2Cl_2$	81	10:1
13 <sup>e</sup>	PtCl <sub>2</sub>	82	5:1

<sup>*a*</sup>Reaction conditions: **1a** (0.1 mol), **2a** (0.1 mol), catalyst (10 mol %), toluene (1.0 mL). <sup>*b*</sup>Yield of the isolated product after column chromatography. <sup>*c*</sup>Determined by <sup>1</sup>H NMR analysis. <sup>*d*</sup>Not determined. <sup>*e*</sup>**1a** (0.1 mol), **2a** (0.2 mol).

on the phenyl rings afforded the corresponding pyrroloindolines in enantiomerically pure form with good yields and diastereoselectivities (Table 3). The absolute configuration of (S,S,S)-3c was unambiguously determined by single-crystal Xray analysis (see Supporting Information), confirming the complete stereoinversion at the C-2 of aziridine. Other chiral products can therefore be assigned by analogy.

In the following studies, stereochemical outcomes of the reactions were probed with electron-rich 1,3,5-trimethoxybenzene 4 and (R)-2a or (S)-2a (Schemes 2a and 2b). As predicted, both of the products could be attained as a single enantiomer. This represents great progress compared to the same initial study performed by Ghorai (Scheme 2c).<sup>3</sup> Therefore, we tried to explore the mechanism difference between Pd(II) and Sc(OTf)<sub>3</sub>.

A variety of control experiments were first performed to verify the Pd(II) catalyst acting as the Lewis acid. Under strictly oxygen-free conditions, the reaction could go smoothly; while the reaction of C3-methylindole 1a and benzyl-substituted aziridine was examined and no product was observed under the same conditions<sup>14</sup> (Scheme 3a). Using Pd(0) catalysts such as Pd(PPh<sub>3</sub>)<sub>4</sub> or Pd<sub>2</sub>(dba)<sub>3</sub>, no desired product was observed and the two substrates were recovered quantitatively (Scheme 3b). Furthermore, when the Pd(PPh<sub>3</sub>)<sub>2</sub>Cl<sub>2</sub> or PdCl<sub>2</sub> coupled with phosphines or amines ligands were used as the catalysts, the reaction did not proceed probably due to the addition of ligands decreasing the Lewis acidity of palladium by resonance of unpaired electrons to the empty orbit of metal palladium (Scheme 3c).

Three more reactions of enantiopure aziridine in toluene with  $Sc(OTf)_3$  or  $Pd(PhCN)_2Cl_2$  were investigated, and results are shown in Scheme 4. It can be seen that the ee of (*R*)-2a was dropped to 39% with  $Sc(OTf)_3$  after 10 min (Scheme 4a), while the racemization was not observed with  $Pd(PhCN)_2Cl_2$  as the catalyst (Scheme 4b). Furthermore, in agreement with the above results,  $Sc(OTf)_3$ -catalyzed [3 + 2] cycloaddition of indole 1a with chiral aziridine (S)-2a provided low stereo-

selectivity (Scheme 4c). These results were probably because of the difference of Lewis acidity between Sc(III) and Pd(II).

Kinetic experiments conducted showed that the reaction with electron-rich substrates, with electron-rich substituents on either the sulfonamides or the 2-aromatic rings, has a large initial reaction rate, as the electron-rich substrate is more liable to coordinate with palladium; thus the reaction is faster (Figure 2).

Based on the above results, a plausible mechanism is proposed. The Pd catalyst coordinated to the aziridine,<sup>14</sup> generating a highly reactive species, which undergoes nucleophilic attack by indoles in an  $S_N 2$  fashion to provide stereoinverted iminium ion, followed by cyclization to obtain the desired product (Scheme 5).

### CONCLUSIONS

In summary, we have developed a Pd-catalyzed formal [3 + 2] cycloaddition of achiral and chiral aziridines with indoles. The reaction allows for preparation of chiral hexahydropyrrolo[2,3-b] indoles in enantiomerically pure forms. Further studies will concentrate on a detailed mechanism as well as extending the synthetic utility of this strategy.

### EXPERIMENTAL SECTION

The melting points recorded are uncorrected. Nuclear magnetic resonance spectra were recorded at 400 MHz. All chemical shifts ( $\delta$ ) are given in ppm. Data are reported as follows: chemical shift, integration, multiplicity (s = single, d = doublet, t = triplet, q = quartet, br = broad, m = multiplet), and coupling constants (Hz). Ee values were determined by chiral HPLC analysis using Chiral ID-3, Chiral OD-H, or Chiral AD-H connected with Chiral PC-2. The mobile phase was a binary mixture, *n*-hexane/*i*-PrOH. Optical rotations ( $\alpha$ ) were measured on a polarimeter with a sodium lamp in the given solvent at the indicated concentration (*c*, *g*/100 mL) and temperature (°C). High resolution mass spectra were recorded under TOF conditions. All solvents were used after redistillation according to standard procedures. All reactions were carried out employing ovendried glassware. 3-Substituted indoles 1 were prepared according to procedures reported previously.<sup>15</sup> Aziridines 2 were synthesized using reported procedures.<sup>16</sup> Compound 4 is commercially available.

**2-(4-Chlorophenyl)-1-tosylaziridine (2b):** <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  = 2.34 (d, *J* = 4.0 Hz, 1H), 2.44 (s, 3H), 2.98 (d, *J* = 7.2 Hz, 1H), 3.74 (dd, *J* = 4.0 Hz, *J* = 7.2 Hz, 1H), 7.15 (d, *J* = 8.4 Hz, 2H), 7.25–7.27 (m, 2H), 7.33 (d, *J* = 8.0 Hz, 2H), 7.85 (d, *J* = 8.4 Hz, 2H).

**2-(4-Bromophenyl)-1-tosylaziridine (2c):** <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  = 2.34 (d, *J* = 4.4 Hz, 1H), 2.44 (s, 3H), 2.98 (d, *J* = 7.2 Hz, 1H), 3.72 (dd, *J* = 4.4 Hz, *J* = 7.2 Hz, 1H), 7.09 (d, *J* = 8.0 Hz, 2H), 7.34 (d, *J* = 8.0 Hz, 2H), 7.42 (d, *J* = 8.4 Hz, 2H), 7.86 (d, *J* = 8.4 Hz, 2H).

**2-(2-Bromophenyl)-1-tosylaziridine (2g):** <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  = 2.27 (d, *J* = 4.4 Hz, 1H), 2.45 (s, 3H), 3.03 (d, *J* = 7.2 Hz, 1H), 3.99 (dd, *J* = 4.4 Hz, *J* = 7.2 Hz, 1H), 7.12–7.22 (m, 3H), 7.36 (d, *J* = 7.6 Hz, 2H), 7.51 (d, *J* = 7.6 Hz, 1H), 7.90 (d, *J* = 8.0 Hz, 2H).

**1-((4-Methoxyphenyl)sulfonyl)-2-phenylaziridine (2h):** <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  = 2.38 (d, *J* = 4.8 Hz, 1H), 2.96 (d, *J* = 7.6 Hz, 1H), 3.75 (dd, *J* = 4.8 Hz, *J* = 7.6 Hz, 1H), 3.87 (s, 3H), 6.98 (d, *J* = 8.8 Hz, 2H), 7.21-7.23 (m, 2H), 7.26-7.30 (m, 3H), 7.92 (d, *J* = 8.8 Hz, 2H).

**1-((4-Bromophenyl)sulfonyl)-2-phenylaziridine (2i):** <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  = 2.43 (d, *J* = 4.4 Hz, 1H), 3.02 (d, *J* = 7.2 Hz, 1H), 3.81 (dd, *J* = 4.4 Hz, *J* = 7.2 Hz, 1H), 7.21–7.23 (m, 2H), 7.30–7.31 (m, 3H), 7.68 (d, *J* = 8.0 Hz, 2H), 7.85 (d, *J* = 8.4 Hz, 2H).

**1-((4-Nitrophenyl)sulfonyl)-2-phenylaziridine (2j):** <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  = 2.49 (d, *J* = 4.8 Hz, 1H), 3.10 (d, *J* = 7.2 Hz, 1H), 3.88 (dd, *J* = 4.8 Hz, *J* = 7.2 Hz, 1H), 7.19–7.21 (m, 2H), 7.29–7.31 (m, 3H), 8.17 (d, *J* = 8.4 Hz, 2H), 8.36 (d, *J* = 8.4 Hz, 2H).

## Table 2. Reaction Scope<sup>a</sup>



<sup>*a*</sup>Reaction conditions: 1a (0.1 mol), 2a (0.2 mol), Pd(PhCN)<sub>2</sub>Cl<sub>2</sub> (10 mol %), toluene (1.0 mL). <sup>*b*</sup>Yield of the isolated product after column chromatography. <sup>*c*</sup>Determined by <sup>1</sup>H NMR analysis.

# Table 3. Construction of Chiral Hexahydropyrrolo[2,3-b]indoles<sup>a</sup>

		+ N Ph <sup>ww</sup> Pd(PhCl Tolu <b>2</b>	$\frac{N_{2}Cl_{2} (10 \text{ mol}\%)}{\text{Jene, rt, 10h}} R \frac{\Pi}{U}$	Ph + R 3 BnH Pg	N BnH Pg	"Ph
entry	R	2	products	yield [%] <sup>b</sup>	dr <sup>c</sup>	ee [%] <sup>d</sup>
1	Н	(S)-2a	( <i>S,S,S</i> )- <b>3a</b>	81	9/1	>99/99
2	Н	(R)-2a	(R,R,R)- <b>3a</b>	83	9/1	>-99/-99
3	H, <i>N</i> -Me	(S)-2a	( <i>S,S,S</i> )- <b>3b</b>	67	5/1	>99/99
4	4-Br	(S)-2a	( <i>S</i> , <i>S</i> , <i>S</i> )- <b>3</b> c	75	8/1	>99/99
5	6-Br	(S)-2a	( <i>S</i> , <i>S</i> , <i>S</i> )- <b>3e</b>	63	9/1	>99/99
6	5-F	(S)-2a	( <i>S</i> , <i>S</i> , <i>S</i> )- <b>3f</b>	65	8/1	>99/99
7	5-OMe	(S)-2a	( <i>S</i> , <i>S</i> , <i>S</i> )- <b>3</b> g	62	6/1	>99/99
8	Н	(S)- <b>2h</b>	( <i>S,S,S</i> )- <b>3s</b>	75	7/1	>99/99
9	Н	(S)-2i	( <i>S,S,S</i> )- <b>3</b> t	73	7/1	>99/99
10	Н	(S)- <b>2</b> j	( <i>S</i> , <i>S</i> , <i>S</i> )- <b>3</b> u	70	6/1	>99/99

"Reaction conditions: 1a (0.1 mol), 2a (0.2 mol), Pd(PhCN)<sub>2</sub>Cl<sub>2</sub> (10 mol %), toluene (1.0 mL). <sup>b</sup>Yield of the isolated product after column chromatography. <sup>c</sup>Determined by <sup>1</sup>H NMR analysis. <sup>d</sup>Ee % of 3/3' determined by chiral HPLC analysis.

General Procedure for the [3 + 2] Cycloaddition. To a mixture of 3-substituted indole 1 (0.1 mmol, 1.0 equiv) and Pd(PhCN)<sub>2</sub>Cl<sub>2</sub>

(0.01 mmol, 0.1 equiv) in toluene (1.0 mL) was added aziridine 2 (0.2 mmol, 2.0 equiv) at room temperature. The reaction was monitored

Scheme 2. Stereochemical Study with the Reaction of 1,3,5-Trimethoxybenzene



Scheme 3. Control Experiments



Scheme 4. Racemization Studies and Sc(OTf)<sub>3</sub>-Catalyzed Reaction of Chiral Aziridine



by TLC analysis. At the end of the reaction, the mixture was quenched with  $H_2O$  (5 mL) and extracted with ethyl acetate (3 × 5 mL). The combined organic layer was dried over  $Na_2SO_4$ , filtered, and

concentrated to afford the corresponding product 3 after flash column chromatography (hexanes/ethyl acetate = 10/1-5/1).

(3S, 3aS, 8aS)-8-Benzyl-3a-methyl-3-phenyl-1-tosyl-1,2,3,3a,8,8a-hexahydropyrrolo[2,3-b]indole (3a): 40 mg, yield 81%; white solid; mp = 98–100 °C;  $[\alpha]_D^{26} = -52.9$  (*c* = 0.9 in CHCl<sub>3</sub>); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  = 1.24 (s, 3H), 2.44 (s, 3H), 2.76 (dd, J = 6.4 Hz, J = 12.8 Hz, 1H), 3.50 (t, J = 12.8 Hz, 1H), 3.87–3.92 (m, 1H), 4.70 (AB, J = 16.0 Hz, 2H), 5.48 (s, 1H), 5.56 (d, J = 7.2 Hz, 1H), 6.27 (t, J = 7.2 Hz, 1H), 6.32 (d, J = 8.0 Hz, 1H), 6.84 (d, J = 7.2 Hz, 2H), 6.94 (t, J = 7.6 Hz, 1H), 7.21-7.34 (m, 10H), 7.73 (d, J = 8.0 Hz, 2H); <sup>13</sup>C NMR (100 MHz, CDCl<sub>2</sub>)  $\delta$  = 21.6, 26.4, 48.1, 51.2, 55.4, 56.9, 89.9, 105.4, 116.5, 125.7, 127.0, 127.3, 127.4, 127.6, 128.0, 128.4, 128.5, 128.91, 128.93, 135.8, 137.2, 138.7, 143.8, 150.2 ppm; IR (neat)  $\nu$  1603, 1494, 1455, 1349, 1162, 1088, 1036, 960 cm<sup>-1</sup>; ESI-MS (m/z) 495 (M + H<sup>+</sup>); MALDI/DHB calcd for C<sub>21</sub>H<sub>21</sub>N<sub>2</sub>O<sub>2</sub>S<sup>+</sup> 495.2101, found 495.2113; dr = 10/1. HPLC-separation conditions: Chiralcel AD-H/PC-2, 25 °C, 254 nm, 90/10 hexane/i-PrOH, 1.0 mL/min; trans diastereoisomer, t = 23.8 min, >99% ee; cis diastereoisomer, t = 22.4 min, >99% ee.

(3*R*,3a*R*,8a*R*)-8-Benzyl-3a-methyl-3-phenyl-1-tosyl-1,2,3,3a,8,8a-hexahydropyrrolo[2,3-b]indole (3a): 41 mg, yield 83%;  $[\alpha]_D^{25} = +29.3 (c = 0.7 \text{ in CHCl}_3);$  <sup>1</sup>H NMR (400 MHz, CDCl}\_3)  $\delta = 1.24$  (s, 3H), 2.44 (s, 3H), 2.76 (dd, J = 6.4 Hz, J = 12.8 Hz, 1H), 3.50 (t, J = 12.8 Hz, 1H), 3.87–3.92 (m, 1H), 4.70 (AB, J = 16.0 Hz, 2H), 5.48 (s, 1H), 5.56 (d, J = 7.2 Hz, 1H), 6.27 (t, J = 7.2 Hz, 1H), 6.32 (d, J = 8.0 Hz, 1H), 6.84 (d, J = 7.2 Hz, 2H), 6.94 (t, J = 7.6 Hz, 1H), 7.21–7.34 (m, 10H), 7.73 (d, J = 8.0 Hz, 2H); ESI-MS (m/z) 495 (M + H<sup>+</sup>); trans diastereoisomer, t = 20.7 min, >99% ee; cis diastereoisomer, t = 25.7 min, >99% ee.

(3S,3aS,8aS)-3a,8-Dimethyl-3-phenyl-1-tosyl-1,2,3,3a,8,8ahexahydropyrrolo[2,3-b]indole (3b): 28 mg, yield 67%; white solid; mp = 110–112 °C;  $[\alpha]_{D}^{26} = -10.9$  (c = 0.35 in CHCl<sub>3</sub>); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  = 1.27 (s, 3H), 2.47 (s, 3H), 2.73 (dd, J = 6.8 Hz, J = 12.8 Hz, 1H), 3.03 (t, J = 12.8 Hz, 3H), 3.39 (t, J = 12.8Hz, 1H), 3.72-3.82 (m, 1H), 5.36 (s, 1H), 5.57 (d, J = 7.2 Hz, 1H), 6.28 (t, J = 7.6 Hz, 1H), 6.36 (d, J = 7.6 Hz, 1H), 6.82 (d, J = 8.0 Hz, 2H), 7.01 (t, J = 7.6 Hz, 1H), 7.20-7.26 (m, 3H), 7.34-7.40 (m, 2H), 7.82 (d, J = 8.0 Hz, 2H); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>)  $\delta = 21.6, 26.0,$ 31.3, 51.2, 55.0, 56.7, 91.1, 105.0, 116.4, 125.5, 127.4, 127.5, 128.0, 128.4, 128.6, 128.9, 130.0, 135.8, 137.2, 143.8, 150.9 ppm; IR (neat) v 1605, 1493, 1453, 1347, 1303, 1162, 1092, 1024, 1001, 957 cm<sup>-1</sup>; ESI-MS (m/z) 441 (M + Na<sup>+</sup>); MALDI/DHB calcd for C<sub>25</sub>H<sub>26</sub>N<sub>2</sub>O<sub>2</sub>S<sup>+</sup> 418.1710, found 418.1712; dr = 5/1. HPLC-separation conditions: Chiralcel AD-H/PC-2, 25 °C, 254 nm, 90/10 hexane/i-PrOH, 1.0 mL/min; trans diastereoisomer, t = 49.2 min, >99% ee; cis diastereoisomer, t = 54.2 min, >99% ee.

(3S,3aS,8aS)-8-Benzyl-4-bromo-3a-methyl-3-phenyl-1-tosyl-1,2,3,3a,8,8a-hexahydropyrrolo[2,3-b]indole (3c): 43 mg, yield 75%; white solid; mp = 129–131 °C;  $[\alpha]_D^{25}$  = +25.2 (c = 0.78 in CHCl<sub>3</sub>); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  = 1.42 (s, 3H), 2.43 (s, 3H), 3.19 (t, I = 8.0 Hz, 1H), 3.61 (dd, I = 8.0 Hz, I = 12.0 Hz, 1H), 3.96(dd, J = 8.0 Hz, J = 12.0 Hz, 1H), 4.63 (AB, J = 16.8 Hz, 2H), 5.49 (s, 1H), 6.19 (d, J = 8.0 Hz, 1H), 6.48 (d, J = 8.0 Hz, 1H), 6.73 (t, J = 8.0 Hz, 1H), 6.97 (d, J = 7.6 Hz, 2H), 7.05 (t, J = 7.6 Hz, 2H), 7.12 (t, J = 7.2 Hz, 2H), 7.25–7.35 (m, 7H), 7.70 (d, J = 8.0 Hz, 2H); <sup>13</sup>C NMR  $(100 \text{ MHz}, \text{CDCl}_3) \delta = 21.6, 25.1, 49.6, 54.7, 57.1, 59.1, 92.3, 105.5,$ 120.9, 122.3, 127.0, 127.2, 127.4, 127.6, 127.7, 128.6, 129.2, 129.7, 129.9, 130.0, 136.2, 138.1, 138.2, 144.1, 152.2 ppm; IR (neat) v 1596, 1569, 1495, 1454, 1351, 1304, 1164, 1088, 1024, 963 cm<sup>-1</sup>; ESI-MS (m/z) 573 (M + H<sup>+</sup>); MALDI/DHB calcd for C<sub>31</sub>H<sub>30</sub>N<sub>2</sub>O<sub>2</sub>SBr<sup>+</sup> 573.1206, found 573.1202; dr = 8/1. HPLC-separation conditions: Chiralcel ID-3, 25 °C, 214 nm, 80/20 hexane/i-PrOH, 0.7 mL/min; trans diastereoisomer, t = 8.8 min, >99% ee; cis diastereoisomer, t =10.3 min, >99% ee.

(35,3a5,8a5)-8-Benzyl-5-bromo-3a-methyl-3-phenyl-1-tosyl-1,2,3,3a,8,8a-hexahydropyrrolo[2,3-*b*]indole (3d): 52 mg, yield 91%; white solid; mp = 143–145 °C; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  = 1.21 (s, 3H), 2.45 (s, 3H), 2.72 (dd, *J* = 7.0 Hz, *J* = 12.4 Hz, 1H), 3.47 (t, *J* = 12.4 Hz, 1H), 3.84 (dd, *J* = 7.0 Hz, *J* = 12.4 Hz, 1H), 4.66 (AB, *J* = 16.4 Hz, 2H), 5.47 (s, 1H), 5.53 (s, 1H), 6.17 (d, *J* = 8.4 Hz,



Figure 2. Impacts of sulfonamide and phenyl substituents on ring opening rate.

Scheme 5. A Plausible Mechanism



1H), 6.83 (d, *J* = 6.4 Hz, 2H), 7.02 (dd, *J* = 2.0 Hz, *J* = 8.4 Hz, 1H), 7.25–7.34 (m, 10H), 7.73 (d, *J* = 8.0 Hz, 2H); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>)  $\delta$  = 21.6, 26.2, 48.1, 51.0, 55.3, 56.9, 89.8, 106.8, 107.9, 127.2, 127.4, 127.9, 128.1, 128.7, 128.8, 130.0, 130.7, 130.9, 135.2, 137.0, 138.1, 144.0, 149.0 ppm; IR (neat)  $\nu$  1703, 1599, 1490, 1453, 1345, 1304, 1257, 1160, 1093, 1050, 963 cm<sup>-1</sup>; ESI-MS (*m*/*z*) 573 (M + H<sup>+</sup>); MALDI/DHB calcd for C<sub>31</sub>H<sub>30</sub>N<sub>2</sub>O<sub>2</sub>SBr<sup>+</sup> 573.1206, found 573.1218; dr = 9/1.

(35,3a5,8a5)-8-Benzyl-6-bromo-3a-methyl-3-phenyl-1-tosyl-1,2,3,3a,8,8a-hexahydropyrrolo[2,3-b]indole (3e): 36 mg, yield 63%; white solid; mp = 152–154 °C;  $[α]_D^{26} = -6.67$  (*c* = 0.22 in CHCl<sub>3</sub>); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  = 1.21 (s, 3H), 2.45 (s, 3H), 2.75 (dd, *J* = 2.8 Hz, *J* = 12.8 Hz, 1H), 3.47 (t, *J* = 12.8 Hz, 1H), 3.83 (dd, *J* = 6.4 Hz, *J* = 12.8 Hz, 1H), 4.66 (AB, *J* = 16.4 Hz, 2H), 5.35 (d, *J* = 7.6 Hz, 1H), 5.47 (s, 1H), 6.38 (d, *J* = 8.4 Hz, 1H), 6.45 (d, *J* = 0.8 Hz, 1H), 6.83 (d, *J* = 6.8 Hz, 2H), 7.22–7.35 (m, 10H), 7.71 (d, *J* = 8.0 Hz, 2H); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>)  $\delta$  = 21.6, 26.3, 47.8, 51.0, 55.2, 56.5, 89.8, 108.3, 119.3, 122.6, 126.9, 127.18, 127.2, 127.4, 127.7, 128.1, 128.7, 128.8, 130.0, 135.4, 137.0, 137.9, 144.0, 151.4 ppm; IR (neat)  $\nu$  1598, 1494, 1453, 1349, 1259, 1162, 1089, 1053, 1035, 961, 899 cm<sup>-1</sup>; ESI-MS (*m*/*z*) 573 (M + H<sup>+</sup>); MALDI/DHB calcd for  $C_{31}H_{30}N_2O_2SBr^+$  573.1206, found 573.1224; dr = 9/1. HPLCseparation conditions: Chiralcel ID-3, 25 °C, 214 nm, 80/20 hexane/*i*-PrOH, 0.7 mL/min; trans diastereoisomer,  $t_{major}$  = 8.1 min,  $t_{minor}$  = 7.8 min, >99% ee; cis diastereoisomer,  $t_{major}$  = 8.5 min,  $t_{minor}$  = 9.0 min, >99% ee.

(3S,3aS,8aS)-8-Benzyl-5-fluoro-3a-methyl-3-phenyl-1-tosyl-1,2,3,3a,8,8a-hexahydropyrrolo[2,3-b]indole (3f): 34 mg, yield 65%; white solid; mp = 127–129 °C;  $[\alpha]_{\rm D}^{26} = -22.2$  (c = 0.42 in CHCl<sub>3</sub>); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  = 1.21 (s, 3H), 2.45 (s, 3H), 2.74 (dd, J = 6.4 Hz, J = 12.4 Hz, 1H), 3.48 (t, J = 12.4 Hz, 1H), 3.84 (dd, J = 6.4 Hz, J = 12.4 Hz, 1H), 4.67 (AB, J = 16.4 Hz, 2H), 5.47 (s, 1H), 5.54 (s, 1H), 6.17 (d, J = 8.4 Hz, 1H), 6.83 (d, J = 6.8 Hz, 2H), 7.02 (d, J = 6.8 Hz, 1H), 7.25-7.37 (m, 10H), 7.73 (d, J = 8.0 Hz, 2H); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>)  $\delta$  = 21.6, 26.2, 48.1, 51.0, 55.3, 56.9, 89.8, 106.8, 108.0, 127.1, 127.2, 127.4, 127.9, 128.1, 128.6, 128.7  $(J_{C-F} = 207 \text{ Hz})$ ,128.8 130.0, 130.7, 130.9, 135.2, 137.0, 138.1, 144.0, 149.0 ppm; IR (neat) v 1598, 1495, 1454, 1348, 1305, 1259, 1212, 1186, 1092, 1035, 960, 911 cm<sup>-1</sup>; IR (neat) v 1598, 1492, 1454, 1416, 1349, 1290, 1162, 1089, 1052, 960 cm<sup>-1</sup>; IR (neat)  $\nu$  1699, 1598, 1490, 1454, 1416, 1349, 1262, 1205, 1170, 1090, 1051, 961, 913 cm<sup>-1</sup>; MALDI/DHB (m/z) 513  $(M + H^+)$ ; MALDI/DHB calcd for  $C_{31}H_{29}N_2O_2SFNa^+$  535.1825, found 535.1824; dr = 8/1. HPLCseparation conditions: Chiralcel ID-3, 25 °C, 214 nm, 80/20 hexane/i-PrOH, 0.7 mL/min; trans diastereoisomer, *t* = 26.4 min, >99% ee; cis diastereoisomer, *t* = 23.0 min, >99% ee.

(3S,3aS,8aS)-8-Benzyl-5-methoxy-3a-methyl-3-phenyl-1tosyl-1,2,3,3a,8,8a-hexahydropyrrolo[2,3-b]indole (3g): 32 mg, yield 62%; white solid; mp = 61–63 °C;  $[\alpha]_{D}^{26} = -22.7$  (c = 0.67 in CHCl<sub>3</sub>); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  = 1.26 (s, 3H), 2.44 (s, 3H), 2.81 (dd, J = 6.4 Hz, J = 12.4 Hz, 1H), 3.34 (s, 3H), 3.51 (t, J = 12.4 Hz 1H), 3.83 (dd, J = 6.4 Hz, J = 12.4 Hz, 1H), 4.65 (dd, J = 16.4 Hz, 2H), 5.19 (dd, J = 2.0 Hz, 1H), 5.48 (s, 1H), 6.20 (d, J = 8.8 Hz, 1H), 6.51 (dd, J = 2.4 Hz, J = 8.8 Hz, 1H), 6.88 (d, J = 6.0 Hz, 2H), 7.24-7.31 (m, 10H), 7.74 (d, J = 8.0 Hz, 2H); <sup>13</sup>C NMR (100 MHz,  $CDCl_3$ )  $\delta = 21.6, 26.2, 49.2, 51.0, 55.3, 55.8, 57.0, 90.5, 106.1, 112.4,$ 114.2, 126.9, 127.3, 127.4, 127.6, 128.1, 128.9, 129.8, 129.9, 135.9, 137.2, 138.9, 143.7, 144.6, 151.6 ppm; IR (neat) v 1597, 1495, 1454, 1429, 1347, 1262, 1219, 1160, 1093, 1038, 956, 909 cm<sup>-1</sup>; ESI-MS (m/z) 525 (M + H<sup>+</sup>); MALDI/DHB calcd for C<sub>32</sub>H<sub>32</sub>N<sub>2</sub>O<sub>3</sub>SNa<sup>+</sup> 547.2026, found 547.2026; dr = 6/1. HPLC-separation conditions: Chiralcel PC-2/AD-H, 25 °C, 254 nm, 90/10 hexane/i-PrOH, 1.0 mL/min; trans diastereoisomer, t = 38.6 min, >99% ee; cis diastereoisomer, t = 22.5 min, >99% ee.

4-Benzyl-9-phenyl-11-tosyl-1,2,3,4-tetrahydro-3a,8b-(epiminoethano)cyclopenta[b]indole (3h): 37 mg, yield 71%; white solid; mp = 99–100 °C; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  = 1.52-1.55 (m, 2H), 1.71-1.75 (m, 1H), 1.91-1.94 (m, 1H), 2.36 (s, 3H), 2.45-2.49 (m, 1H), 2.75-2.79 (m, 1H), 3.45-3.54 (m, 2H), 3.87 (t, J = 8.8 Hz, 1H), 4.48 (d, J = 16.8 Hz, 1H), 5.40 (d, J = 16.8 Hz, 1H), 6.15 (d, J = 8.0 Hz, 1H), 6.64 (t, J = 7.2 Hz, 1H), 6.83 (d, J = 7.2 Hz, 1H), 7.12 (t, J = 7.2 Hz, 1H), 7.15-7.27 (m, 4H), 7.32-7.40 (m, 4H), 7.32 (t, J = 7.2 Hz, 2H), 7.39 (d, J = 7.2 Hz, 2H), 7.52 (d, J = 8.4 Hz, 2H); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>)  $\delta$  = 21.5, 26.7, 34.0, 38.0, 49.4, 51.5, 53.5, 70.5, 104.7, 107.0, 117.7, 122.5, 126.5, 126.8, 127.4, 127.7, 128.4, 128.5, 128.7, 129.4, 133.7, 137.1, 137.7, 139.7, 143.5, 150.5 ppm; IR (neat) v 1703, 1603, 1486, 1464, 1453, 1384, 1324, 1217, 1160, 1088, 1032, 996, 932, 918 cm<sup>-1</sup>; ESI-MS (m/z) 521 (M + H<sup>+</sup>); MALDI/DHB calcd for  $C_{33}H_{33}N_2O_2S^+$  521.2257, found 521.2263; dr > 20/1.

**9-Benzyl-12-phenyl-10-tosyl-6,7,8,9-tetrahydro-5***H***-8a,4b-(epiminoethano)carbazole (3i): 33 mg, yield 62%; white solid; mp = 186–188 °C; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) \delta = 1.00–1.05 (m, 1H), 1.16–1.21 (m, 1H), 1.41–1.48 (m, 3H), 1.61–1.68 (m, 1H), 2.34 (s, 3H), 3.23 (d,** *J* **= 14.8 Hz, 1H), 3.71–3.82 (m, 2H), 4.06 (t,** *J* **= 9.2 Hz, 1H), 4.14 (d,** *J* **= 16.8 Hz, 1H), 5.28 (d,** *J* **= 4.0 Hz, 1H), 6.19 (d,** *J* **= 7.6 Hz, 1H), 6.65–6.71 (m, 2H), 7.00–7.04 (m, 3H), 7.10 (d,** *J* **= 8.0 Hz, 2H), 7.22–7.35 (m, 7H), 7.67 (d,** *J* **= 8.0 Hz, 2H); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>) \delta = 20.7, 21.0, 21.4, 25.2, 28.9, 48.2, 50.4, 51.8, 58.4, 98.7, 108.4, 117.6, 122.2, 126.7, 127.2, 127.9, 128.3, 128.4, 128.8, 129.4, 131.1, 139.6, 143.3, 148.7 ppm; IR (neat) \nu 1607, 1495, 1481, 1454, 1338, 1170, 1096, 1065, 1025, 991 cm<sup>-1</sup>; MALDI/DHB (***m***/***z***) 535 (M + H<sup>+</sup>); MALDI/DHB calcd for C<sub>34</sub>H<sub>35</sub>N<sub>2</sub>O<sub>2</sub>S<sup>+</sup> 535.2414, found 535.2419; dr > 20/1.** 

8-Benzyl-3a-cyclohexyl-3-phenyl-1-tosyl-1,2,3,3a,8,8ahexahydropyrrolo[2,3-b]indole (3j): 20 mg, yield 36%; white solid; mp = 74-75 °C; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  = 0.66-1.06 (m, 6H), 1.35–1.73 (m, 5H), 2.42 (s, 3H), 3.44–3.58 (m, 2H), 3.77–3.81 (m, 1H), 4.74 (AB, J = 16.8 Hz, 2H), 5.54 (d, J = 7.2 Hz, 1H), 5.60 (s, 1H), 6.23–6.31 (m, 2H), 6.84 (d, J = 6.8 Hz, 1H), 6.94 (t, J = 7.2 Hz, 1H), 7.02-7.33 (m, 1H), 7.19-7.33 (m, 10H), 7.71 (d, J = 8.0 Hz, 2H); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>)  $\delta$  = 21.4, 21.5, 25.6, 25.8, 26.1 26.2, 26.3, 26.5, 26.7, 27.7, 28.1, 28.2, 28.6, 29.7, 41.3, 43.5, 48.4, 48.6, 50.0, 53.8, 54.8, 64.7, 65.0, 68.0, 84.5, 87.3, 105.1, 107.2, 116.1, 117.3, 124.3, 126.3, 126.6, 126.8, 126.9, 127.1, 127.2, 127.3, 127.4, 127.6, 127.8, 128.1, 128.2, 128.3, 128.4, 128.9, 129.2, 129.5, 129.7, 130.9, 136.5, 137.2, 138.2, 138.5, 138.9, 143.0 149.7, 151.1 ppm; IR (neat) v 1703, 1601, 1488, 1466, 1452, 1402, 1345, 1260, 1203, 1165, 1092, 1051, 928 cm<sup>-1</sup>; ESI-MS (m/z) 563 (M + H<sup>+</sup>); MALDI/DHB calcd for  $C_{36}H_{39}N_2O_2S^+$  563.2727, found 563.2738; dr = 2.5/1.

8-Benzyl-3a-(2-(benzyloxy)ethyl)-3-phenyl-1-tosyl-1,2,3,3a,8,8a-hexahydropyrrolo[2,3-b]indole (3k): 31 mg, yield 50%; white solid; mp = 58–60 °C; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  = 1.65–1.70 (m, 1H), 2.02–2.07 (m, 1H), 2.43 (s, 3H), 2.77 (dd, *J* = 6.4 Hz, *J* = 12.8 Hz, 1H), 3.01–3.03 (m, 2H), 3.46 (t, *J* = 8.8 Hz, 1H), 3.80 (dd, *J* = 6.4 Hz, *J* = 12.8 Hz, 1H), 4.16 (s, 2H), 4.60 (AB, *J* = 16.4 Hz, 2H), 5.48 (d, *J* = 7.2 Hz, 1H), 5.86 (s, 1H), 6.25 (t, *J* = 7.2 Hz, 1H), 6.33 (d, *J* = 8.0 Hz, 1H), 6.78 (d, *J* = 7.2 Hz, 2H), 6.96 (t, *J* = 7.6 Hz, 1H), 7.16–7.39 (m, 15H), 7.72 (d, *J* = 8.0 Hz, 2H); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>)  $\delta$  = 21.6, 39.0, 48.4, 50.2, 55.7, 59.4, 67.2, 87.4, 105.3, 116.3, 125.6, 126.1, 127.0, 127.4, 127.5, 127.6, 127.9, 128.4, 128.5, 128.6, 129.2, 129.9, 135.5, 137.2, 138.3, 143.8, 138.8, 151.0 pm; IR (neat)  $\nu$  1602, 1494, 1454, 1349, 1270, 1162, 1095, 1030, 969, 912 cm<sup>-1</sup>; MALDI/DHB (*m*/*z*) 637 (M + Na<sup>+</sup>); MALDI/DHB calcd for C<sub>39</sub>H<sub>38</sub>N<sub>2</sub>O<sub>3</sub>SNa<sup>+</sup> 637.2495, found 637.2508; dr = 7/1.

**8-Benzyl-3a-isopropyl-3-phenyl-1-tosyl-1,2,3,3a,8,8a-hexahydropyrrolo**[**2**,3-**b**]indole (**3**]): 18 mg, yield 34%; white solid; mp = 65–66 °C; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  = 0.52 (d, *J* = 6.8 Hz, 3H), 0.87 (d, *J* = 6.8 Hz, 3H), 1.96–2.00 (m, 1H), 2.42 (s, 3H), 3.37–3.58 (m, 2H), 3.77–3.81 (m, 1H), 4.70 (AB, *J* = 16.4 Hz, 2H), 5.56 (d, *J* = 7.2 Hz, 1H), 5.60 (s, 1H), 6.26–6.31 (m, 2H), 6.87 (d, *J* = 6.8 Hz, 1H), 7.07 (t, *J* = 6.8 Hz, 1H), 7.19–7.63 (m, 10H), 7.71 (d, *J* = 8.4 Hz, 2H); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>)  $\delta$  = 15.0, 15.6, 15.8, 16.2, 18.87, 18.94, 28.2, 30.7, 45.8, 47.2, 48.5, 51.4, 52.3, 62.2, 81.3, 84.9, 102.5, 104.6, 113.6, 114.9, 121.9, 123.7, 123.9, 124.2, 124.5, 124.71, 124.75, 124.8, 125.0, 125.2, 125.7, 125.8, 126.3, 126.6, 126.9, 127.2, 133.8, 134.6, 136.0, 136.8, 141.6, 148.5 ppm; IR (neat)  $\nu$  1702, 1602, 1487, 1467, 1454, 1390, 1342, 1261, 1204, 1168, 1092, 1031, 919 cm<sup>-1</sup>; MALDI/DHB (*m*/*z*) 508 (M + H<sup>+</sup>); MALDI/DHB calcd for C<sub>32</sub>H<sub>32</sub>N<sub>2</sub>O<sub>2</sub>S<sup>+</sup> 508.2166, found 508.2172; dr = 2.5/1.

**8**-Benzyl-3-(4-chlorophenyl)-3a-methyl-1-tosyl-1,2,3,3a,8,8a-hexahydropyrrolo[2,3-*b*]indole (3m): 37 mg, yield 70%; white solid; mp = 146–148 °C; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$ = 1.22 (s, 3H), 2.44 (s, 3H), 2.78 (dd, *J* = 6.4 Hz, *J* = 12.4 Hz, 1H), 3.42 (t, *J* = 12.4 Hz, 1H), 3.81 (dd, *J* = 6.4 Hz, *J* = 12.4 Hz, 1H), 4.68 (AB, *J* = 16.4 Hz, 2H), 5.49 (s, 1H), 5.64 (d, *J* = 7.2 Hz, 1H), 6.32 (t, *J* = 8.4 Hz, 2H), 6.76 (d, *J* = 8.4 Hz, 2H), 6.96 (t, *J* = 7.6 Hz, 1H), 7.19–7.34 (m, 9H), 7.72 (d, *J* = 8.0 Hz, 2H); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>)  $\delta$  = 21.6, 26.4, 48.2, 55.0, 56.9, 89.9, 105.6, 116.7, 125.6, 127.0, 127.3, 127.4, 130.0, 128.2, 128.6, 128.8, 129.9, 130.0, 133.4, 134.4, 137.1, 138.6, 143.9, 150.2 ppm; IR (neat)  $\nu$  1603, 1485, 1453, 1349, 1216, 1170, 1091, 1036, 1015, 960, 899 cm<sup>-1</sup>; ESI-MS (*m*/*z*) 529 (M + H<sup>+</sup>); MALDI/DHB calcd for C<sub>31</sub>H<sub>30</sub>N<sub>2</sub>O<sub>2</sub>SCl<sup>+</sup> 529.1711, found 529.1710; dr = 15/1.

**8-Benzyl-3-(4-bromophenyl)-3a-methyl-1-tosyl-1,2,3,3a,8,8a-hexahydropyrrolo**[**2,3-b**]indole (3n): 40 mg, yield 70%; white solid; mp = 149–150 °C; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$ = 1.22 (s, 3H), 2.44 (s, 3H), 2.76 (dd, J = 6.4 Hz, J = 12.4 Hz, 1H), 3.42 (t, J = 12.4 Hz, 1H), 3.81 (dd, J = 6.4 Hz, J = 12.4 Hz, 1H), 4.67 (AB, J = 16.4 Hz, 2H), 5.49 (s, 1H), 5.65 (d, J = 7.2 Hz, 1H), 6.32 (t, J= 8.4 Hz, 2H), 6.71 (d, J = 8.0 Hz, 2H), 6.85 (d, J = 8.0 Hz, 2H), 6.96 (t, J = 8.0 Hz, 1H), 7.24–7.76 (m, 9H), 7.85 (d, J = 8.0 Hz, 2H); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>)  $\delta$  = 21.6, 26.4, 48.2, 51.2, 55.1, 56.8, 89.9, 105.6, 116.7, 121.5, 125.6, 127.0, 127.3, 127.4, 128.1, 128.57, 128.6, 130.0, 130.7, 131.1, 134.9, 137.1, 138.5, 143.9, 150.2 ppm; IR (neat)  $\nu$ 1703, 1603, 1489, 1453, 1409, 1349, 1261, 1162, 1090, 1010, 961, 899 cm<sup>-1</sup>; ESI-MS (m/z) 573 (M + H<sup>+</sup>); MALDI/DHB calcd for C<sub>31</sub>H<sub>30</sub>N<sub>2</sub>O<sub>2</sub>SBr<sup>+</sup> 573.1206, found 573.1193; dr = 11/1.

**8** - Ben zyl-3-(3-br om ophenyl)-3a-methyl-1-tosyl-1,2,3,3a,8,8a-hexahydropyrrolo[2,3-b]indole (30): 51 mg, yield 90%; white solid; mp = 151–153 °C; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$ = 1.24 (s, 3H), 2.45 (s, 3H), 2.77 (dd, J = 6.4 Hz, J = 12.4 Hz, 1H), 3.43 (t, J = 12.4 Hz, 1H), 3.82 (dd, J = 6.4 Hz, J = 12.4 Hz, 1H), 4.68 (AB, J = 16.4 Hz, 2H), 5.49 (s, 1H), 5.64 (d, J = 7.2 Hz, 1H), 6.38 (t, J= 7.6 Hz, 2H), 6.74 (d, J = 8.0 Hz, 1H), 6.91–6.96 (m, 2H), 7.04– 7.16 (m, 1H), 7.24–7.41 (m, 8H), 7.72 (d, J = 8.0 Hz, 2H); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>)  $\delta$  = 21.6, 26.4, 48.3, 51.2, 55.3, 57.0, 89.8, 105.7, 116.7, 122.1, 125.6, 127.0, 127.3, 127.4, 127.5, 128.0, 128.6, 128.7, 129.5, 130.0, 130.6, 132.0, 137.1, 138.4, 138.6, 144.0, 150.1 ppm; IR (neat)  $\nu$  1702, 1602, 1567, 1493, 1453, 1402, 1348, 1260, 1203, 1170, 1089, 961, 904 cm<sup>-1</sup>; ESI-MS (m/z) 573 (M + H<sup>+</sup>); MALDI/DHB calcd for C<sub>31</sub>H<sub>30</sub>N<sub>2</sub>O<sub>2</sub>SBr<sup>+</sup> 573.1206, found 573.1202; dr = 7/1. **8-Benzyl-3-(4-fluorophenyl)-3a-methyl-1-tosyl-1,2,3,3a,8,8a-hexahydropyrrolo**[**2,3-b**]indole (**3p**): 40 mg, yield 78%; white solid; mp = 131–133 °C; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  = 1.22 (s, 3H), 1.45 (s, 3H), 2.79 (dd, J = 6.4 Hz, J = 12.4 Hz, 1H), 3.42 (t, J = 12.4 Hz, 1H), 3.82 (dd, J = 6.4 Hz, J = 12.4 Hz, 1H), 4.68 (AB, J = 16.4 Hz, 2H), 5.49 (s, 1H), 5.60 (d, J = 7.6 Hz, 1H), 6.28–6.34 (m, 2H), 6.78–6.81 (m, 2H), 6.90–6.97 (m, 3H), 6.24–6.32 (m, 7H), 7.72 (d, J = 8.0 Hz, 2H); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>)  $\delta$  = 21.6, 26.4, 48.2, 51.4, 54.9, 56.8, 89.8, 105.6, 114.9 ( $J_{C-F}$  = 21.4 Hz), 116.6, 125.6, 127.0, 127.3, 127.4, 128.3, 128.5, 128.6, 129.9, 130.4 ( $J_{C-F}$  = 7.9 Hz), 131.6, 137.1, 138.6, 143.9, 150.2, 162.3 ( $J_{C-F}$  = 245 Hz) ppm; IR (neat)  $\nu$  1604, 1512, 1494, 1453, 1349, 1260, 1225, 1163, 1106, 1088, 1059, 1036, 960, 900 cm<sup>-1</sup>; MALDI/DHB (m/z) 513 (M + H<sup>+</sup>); MALDI/DHB calcd for C<sub>31</sub>H<sub>30</sub>N<sub>2</sub>O<sub>2</sub>SF<sup>+</sup> 513.2007, found 513.2006; dr = 9/1.

**8-Benzyl-3a-methyl-3-(p-tolyl)-1-tosyl-1,2,3,3a,8,8a-hexahydropyrrolo[2,3-b]indole (3q):** 47 mg, yield 92%; white solid; mp = 57–59 °C; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  = 1.22 (s, 3H), 2.32 (s, 3H), 2.44 (s, 3H), 2.70 (dd, *J* = 6.8 Hz, *J* = 12.8 Hz, 1H), 3.46 (t, *J* = 12.8 Hz, 1H), 3.79 (dd, *J* = 6.8 Hz, *J* = 12.8 Hz, 1H), 4.71 (AB, *J* = 16.4 Hz, 2H), 5.46 (s, 1H), 5.63 (d, *J* = 7.2 Hz, 1H), 6.27–6.32 (m, 2H), 6.73 (d, *J* = 7.6 Hz, 2H), 6.94 (d, *J* = 7.6 Hz, 1H), 7.03–7.10 (m, 3H), 7.74 (d, *J* = 8.0 Hz, 2H), 7.23–7.34 (m, 6H), 7.72 (d, *J* = 8.0 Hz, 2H); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>)  $\delta$  = 21.1, 21.6, 26.4, 48.1, 51.3, 55.0, 56.8, 89.9, 105.4, 116.5, 125.8, 126.9, 127.3, 127.4, 128.3, 128.5, 128.7, 129.9, 132.6, 137.2, 138.7, 143.8, 150.2 ppm; IR (neat)  $\nu$  1603, 1516, 1494, 1454, 1402, 1348, 1260, 1217, 1185, 1163, 1091, 1036, 960, 900 cm<sup>-1</sup>; ESI-MS (*m*/*z*) 509 (M + H<sup>+</sup>); MALDI/DHB calcd for C<sub>32</sub>H<sub>33</sub>N<sub>2</sub>O<sub>2</sub>S<sup>+</sup> 509.2257, found 509.2262; dr = 12/1.

8-Benzyl-3-(2-bromophenyl)-3a-methyl-1-tosyl-1,2,3,3a,8,8a-hexahydropyrrolo[2,3-b]indole (3r): 38 mg, yield 67%; white solid; mp = 63–65 °C; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  = 1.22 (s, 3H), 2.45 (s, 3H), 3.40-3.45 (m, 1.4H), 3.57-3.75 (m, 0.3H), 3.75-3.84 (m, 1H), 4.28-4.31 (m, 0.3H), 4.66-4.71 (m, 2H), 5.40 (s, 0.3H), 5.46 (s, 0.7H), 5.85 (d, J = 7.6 Hz, 0.7H), 6.30-6.35 (m, 1.4H), 6.43 (d, J = 8.0 Hz, 0.3H), 6.55 (dd, J = 2.0 Hz, J = 7.2 Hz, 0.7H), 6.72 (t, J = 7.2 Hz, 0.3H), 6.94-7.11 (m, 3H), 7.18-7.27 (m, 3H), 7.32-7.36 (m, 4H), 7.46-7.55 (m, 1.5H), 7.71 (d, J = 8.0 Hz, 0.6H), 7.78 (d, J = 8.0 Hz, 1.4H), 7.78 (d, J = 8.0 Hz, 0.2H); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>)  $\delta$  = 20.6, 21.6, 27.0, 48.2, 50.7, 52.6, 53.8, 57.8, 58.1, 89.5, 90.7, 105.6, 108.3, 116.4, 118.8, 122.9, 125.8, 125.9, 126.4, 127.0, 127.1, 127.2, 127.4, 127.7, 128.0, 128.4, 126.5, 128.6, 128.7, 128.8, 129.6, 130.2, 131.0, 133.0, 133.1, 135.4, 136.9, 138.5, 138.6, 144.0, 150.1 ppm; IR (neat) v 1651, 1603, 1538, 1494, 1488, 1471, 1455, 1349, 1163, 1091, 1026, 958 cm<sup>-1</sup>; ESI-MS (m/z) 573 (M + H<sup>+</sup>); DART Positive calcd for  $C_{21}H_{30}N_2O_2BrS^+$  573.1211, found 573.1196; dr = 2.5/1.

(3S, 3aS, 8aS)-8-Benzyl-1-((4-methoxyphenyl)sulfonyl)-3amethyl-3-phenyl-1,2,3,3a,8,8a-hexahydropyrrolo[2,3-b]indole (3s): 38 mg, yield 75%; white solid; mp = 69–70 °C;  $[\alpha]_{\rm D}^{26}$  = -39.5 (*c* = 1.0 in CHCl<sub>3</sub>); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  = 1.26 (s, 3H), 2.88 (dd, *J* = 6.4 Hz, *J* = 12.4 Hz, 1H), 3.49 (t, *J* = 12.4 Hz, 1H), 3.79–3.87 (m, 4H), 4.70 (AB, J = 16.4 Hz, 2H), 5.47 (s, 1H), 5.57 (d, J = 7.6 Hz, 1H), 6.27 (t, J = 7.6 Hz, 1H), 6.32 (d, J = 8.0 Hz, 1H), 6.85 (d, J = 7.2 Hz, 2H), 6.91–6.96 (m, 3H), 7.21–7.32 (m, 7H), 7.77 (d, J = 8.8 Hz, 2H); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>)  $\delta$  = 26.5, 48.2, 51.2, 55.5, 55.7, 56.9, 89.9, 105.4, 114.5, 116.5, 125.7, 127.0, 127.3, 127.6, 128.0, 128.4, 128.5, 128.9, 129.5, 131.8, 135.9, 138.7, 150.2, 163.2 ppm; IR (neat) v 1597, 1495, 1455, 1348, 1260, 1159, 1095, 1027, 960 cm<sup>-1</sup>; ESI-MS (m/z) 511 (M + H<sup>+</sup>); MALDI/DHB calcd for C<sub>31</sub>H<sub>31</sub>N<sub>2</sub>O<sub>3</sub>S<sup>+</sup> 511.2050, found 511.2056; dr = 7/1. HPLC-separation conditions: Chiralcel AD-H/PC-2, 25 °C, 254 nm, 90/10 hexane/i-PrOH, 1.0 mL/min; trans diastereoisomer, t = 43.9 min, >99% ee; cis diastereoisomer, t = 34.1 min, >99% ee.

(35,3a5,8a5)-8-Benzyl-1-((4-bromophenyl)sulfonyl)-3amethyl-3-phenyl-1,2,3,3a,8,8a-hexahydropyrrolo[2,3-b]indole (3t): 41 mg, yield 73%; white solid; mp = 68–70 °C;  $[\alpha]_D^{26} = -31.8$  (c = 1.0 in CHCl<sub>3</sub>); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  = 1.28 (s, 3H), 2.88 (dd, J = 6.4 Hz, J = 12.4 Hz, 1H), 3.49 (t, J = 12.4 Hz, 1H), 3.78–3.83 (m, 1H), 4.68 (AB, J = 16.4 Hz, 2H), 5.49 (s, 1H), 5.59 (d, J = 7.2 Hz, 1H), 6.28–6.35 (m, 2H), 6.87 (d, J = 6.8 Hz, 2H), 6.96 (t, J = 7.2 Hz, 1H), 7.24–7.29 (m, 7H), 7.60–7.68 (m, 4H); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>)  $\delta = 26.5$ , 48.5, 51.1, 55.6, 57.0, 90.0, 105.6, 116.8, 125.8, 127.2, 127.7, 127.9, 128.1, 128.4, 128.6, 128.8, 128.9, 132.5, 135.5, 138.5, 139.3, 150.1 ppm; IR (neat)  $\nu$  1603, 1574, 1494, 1453, 1389, 1353, 1164, 1087, 1067, 1009, 960 cm<sup>-1</sup>; ESI-MS (m/z) 559 (M + H<sup>+</sup>); MALDI/DHB calcd for C<sub>30</sub>H<sub>28</sub>N<sub>2</sub>O<sub>2</sub>SBr<sup>+</sup> 559.1049, found 559.1062; dr = 7/1. HPLC-separation conditions: Chiralcel AD-H/PC-2, 25 °C, 254 nm, 90/10 hexane/*i*-PrOH, 1.0 mL/min; trans diastereoisomer, t = 19.0 min, >99% ee; cis diastereoisomer, t = 17.2 min, >99% ee.

(3S,3aS,8aS)-8-benzyl-3a-methyl-1-((4-nitrophenyl)sulfonyl)-3-phenyl-1,2,3,3a,8,8a-hexahydropyrrolo[2,3-b]indole (3u): 37 mg, yield 70%; white solid; mp = 78-79 °C;  $[\alpha]_{D}^{26}$  = -41.2 (c = 1.0 in CHCl<sub>3</sub>); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  = 1.31 (s, 3H), 2.99 (dd, J = 6.4 Hz, J = 12.4 Hz, 1H), 3.52 (t, J = 12.4 Hz, 1H), 3.85-3.91 (m, 1H), 4.65 (s, 2H), 5.56 (s, 1H), 5.61 (d, J = 7.6 Hz, 1H), 6.30–6.35 (m, 2H), 6.88 (d, J = 6.8 Hz, 2H), 6.97 (t, J = 7.6 Hz, 2H), 7.23–7.29 (m, 7H), 7.94 (d, J = 8.4 Hz, 2H), 8.26 (d, J = 8.8 Hz, 2H); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>)  $\delta$  = 26.6, 49.2, 51.0, 56.0, 57.2, 90.3, 106.0, 117.3, 124.4, 125.8, 127.0, 127.2, 127.8, 128.1, 128.3, 128.6, 128.64, 128.9, 135.2, 138.2, 146.1 150.05, 105.08 ppm; IR (neat) v 1604, 1531, 1494, 1454, 1401, 1350, 1311, 1261, 1166, 1087, 1055, 1012, 960, 900 cm<sup>-1</sup>; ESI-MS (m/z) 526 (M + H<sup>+</sup>); MALDI/ DHB calcd for  $C_{30}H_{28}N_3O_4S^+$  526.1795, found 526.1793; dr = 6/1. HPLC-separation conditions: Chiralcel AD-H/PC-2, 25 °C, 254 nm, 90/10 hexane/*i*-PrOH, 1.0 mL/min; trans diastereoisomer, t = 99.9min, >99% ee; cis diastereoisomer, t = 53.3 min, >99% ee.

**4-Methyl-***N*-(**2-phenyl-2-(2,4,6-trimethoxyphenyl)ethyl)**benzenesulfonamide (5): <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  = 2.41 (s, 3H), 3.60 (s, 6H), 3.66–3.73 (m, 2H), 3.78 (s, 3H), 4.41 (t, J = 6.4 Hz, 1H), 4.66 (dd, J = 8.8, 7.2 Hz, 1H), 6.05 (s, 2H), 7.09–7.29 (m, 5H), 7.24 (d, J = 8.0 Hz, 2H), 7.64 (d, J = 8.4 Hz, 2H). HPLCseparation conditions: Chiralcel OD-H, 25 °C, 254 nm, 90/10 hexane/*i*-PrOH, 1.0 mL/min; For (*S*)-**5**: *t* = 27.9 min, >99% ee, (*R*)-**5**: *t* = 31.9 min, >99% ee.

## ASSOCIATED CONTENT

## **S** Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/acs.joc.5b01931.

**3c** (CIF) <sup>1</sup>H and <sup>13</sup>C NMR spectra and HPLC traces (PDF)

#### AUTHOR INFORMATION

#### Corresponding Author

\*Fax: 0086-21-64166128. Tel: 0086-21-54925182. E-mail: zhaog@sioc.ac.cn.

#### Notes

The authors declare no competing financial interest.

#### ACKNOWLEDGMENTS

Financial support from National Basic Research Program of China (973 Program, 2010CB833200) and the National Natural Science Foundation of China (Nos. 21272247, 21290184) is gratefully acknowledged.

## REFERENCES

 (1) (a) Kametani, T.; Honda, T. Adv. Heterocycl. Chem. 1986, 39, 181. (b) Rogers, G. A.; Parsons, S. M.; Anderson, D. C.; Nilsson, L. M.; Bahr, B. A.; Kornreich, W. D.; Kaufman, R.; Jacobs, R. S.; Kirtman, B. J. Med. Chem. 1989, 32, 1217. (c) Tanner, D. Angew. Chem., Int. Ed. Engl. 1994, 33, 599. (d) Chng, B. L.; Ganesan, A. Bioorg. Med. Chem. Lett. 1997, 7, 1511. (e) Jeong, J. U.; Tao, B.; Sagasser, I.; Henniges, H.; Sharpless, K. B. J. Am. Chem. Soc. 1998, 120, 6844. (f) Stamm, H. J. Prakt. Chem. 1999, 341, 319. (g) McCoull, W.; Davis, F. A. Synthesis **2000**, 2000, 1347. (h) Zwanenburg, B.; Holte, P. T. *Top. Curr. Chem.* **2001**, 216, 93. (i) Lee, W. K.; Ha, H.-J. *Aldrichimica Acta* **2003**, 36, 57. (j) Cardillo, G.; Gentilucci, L.; Tolomelli, A. *Aldrichimica Acta* **2003**, 36, 39.

(2) For recent reviews see: (a) Hu, X.-E. Tetrahedron 2004, 60, 2701.
(b) Lu, P.-F. Tetrahedron 2010, 66, 2549. (c) Wang, P.-A. Beilstein J. Org. Chem. 2013, 9, 1677. (d) Huang, C.-Y. D.; Doyle, A. G. Chem. Rev. 2014, 114, 8153. (e) Callebaut, G.; Meiresonne, T.; De Kimpe, N.; Mangelinckx, S. Chem. Rev. 2014, 114, 7954.

(3) (a) Ghorai, M. K.; Ghosh, K. Tetrahedron Lett. 2007, 48, 3191.
(b) Ghorai, M. K.; Das, K.; Kumar, A. Tetrahedron Lett. 2007, 48, 4373.
(c) Ghorai, M. K.; Das, K.; Shukla, D. J. Org. Chem. 2007, 72, 5859.
(d) Ghorai, M. K.; Shukla, D.; Das, K. J. Org. Chem. 2009, 74, 7013.
(e) Ghorai, M. K.; Kumar, A.; Tiwari, D. P. J. Org. Chem. 2010, 75, 137.
(f) Ghorai, M. K.; Shukla, D.; Bhattacharyya, A. J. Org. Chem. 2012, 77, 3740.
(g) Ghorai, M. K.; Tiwari, D. P.; Jain, N. J. Org. Chem. 2013, 78, 7121.

(4) Takeda, Y.; Ikeda, Y.; Kuroda, A.; Tanaka, S.; Minakata, S. J. Am. Chem. Soc. 2014, 136, 8544.

(5) (a) Jobst, J.; Hesse, O. Justus Liebigs Ann. Chem. 1864, 129, 115.
(b) Zheng, C.-J.; Kim, C.-J.; Bae, K. S.; Kim, Y.-H.; Kim, W.-G. J. Nat. Prod. 2006, 69, 1816. (c) Overman, L. E.; Shin, Y. Org. Lett. 2007, 9, 339. (d) Raju, R.; Piggott, A. M.; Huang, X.-C.; Capon, R. J. Org. Lett. 2011, 13, 2770. (e) Ozawa, T.; Kanematsu, M.; Yokoe, H.; Yoshida, M.; Shishido, K. J. Org. Chem. 2012, 77, 9240. (f) Tayu, M.; Higuchi, K.; Ishizaki, T.; Kawasaki, T. Org. Lett. 2014, 16, 3613.

(6) (a) Austin, J. F.; Kim, S.-G.; Sinz, C. J.; Xiao, W.-J.; MacMillan, D. W. C. Proc. Natl. Acad. Sci. U. S. A. 2004, 101, 5482. (b) Newhouse, T.; Baran, P. S. J. Am. Chem. Soc. 2008, 130, 10886. (c) He, B.; Song, H.; Du, Y.; Qin, Y. J. Org. Chem. 2009, 74, 298. (d) Kim, J.; Ashenhurst, J. A.; Movassaghi, M. Science 2009, 324, 238.

(7) (a) Matsuura, T.; Overman, L. E.; Poon, D. J. Am. Chem. Soc.
1998, 120, 6500. (b) Overman, L. E.; Paone, D. V.; Stearns, B. A. J. Am. Chem. Soc. 1999, 121, 7702. (c) Huang, A.; Kodanko, J. J.; Overman, L. E. J. Am. Chem. Soc. 2004, 126, 14043. (d) Fuchs, J. R.; Funk, R. L. Org. Lett. 2005, 7, 677. (e) Trost, B. M.; Zhang, Y. J. Am. Chem. Soc. 2006, 128, 4590. (f) Pinto, A.; Jia, Y.; Neuville, L.; Zhu, J. Chem. - Eur. J. 2007, 13, 961.

(8) (a) Repka, L. M.; Ni, J.; Reisman, S. E. J. Am. Chem. Soc. 2010, 132, 14418. (b) Kieffer, M. E.; Repka, L. M.; Reisman, S. E. J. Am. Chem. Soc. 2012, 134, 5131. (c) Ni, J.; Wang, H.; Reisman, S. E. Tetrahedron 2013, 69, 5622.

(9) Spangler, J. E.; Davies, H. M. L. J. Am. Chem. Soc. 2013, 135, 6802.

(10) Nakagawa, M.; Kawahara, M. Org. Lett. 2000, 2, 953.

(11) (a) Wang, L.-Q.; Yang, D.-X.; Han, F.-X.; Li, D.; Zhao, D.-P.; Wang, R. Org. Lett. **2015**, *17*, 176. (b) Yang, D.-X.; Wang, L.-Q.; Han, F.-X.; Li, D.; Zhao, D.-P.; Cao, Y.-M.; Ma, Y.-X.; Kong, W.-D.; Sun, Q.-T.; Wang, R. Chem. - Eur. J. **2014**, *20*, 16478.

(12) During the submission of this manuscript, a similar work was reported by Chai and co-workers. See: Chai, Z.; Zhu, Y.-M.; Yang, P.-J.; Wang, S.-Y.; Wang, S.-W.; Liu, Z.; Yang, G.-S. J. Am. Chem. Soc. **2015**, 137, 10088. In our work, the palladium(II) was utilized to catalyze the enantiospecific reactions between indoles and chiral aziridines. No racemization of products was found. The ee values of the reactions between indoles and chiral aziridines were decreased in Chai's work.

(13) Mikami, K.; Hatano, M.; Akiyama, K. Top. Organomet. Chem. 2005, 14, 279.

(14) The reaction between C3-methyl indole 1a and benzylsubstituted aziridine was examined, and no product was observed under the standard condition. The result could be a proof of the proposed mechanism.

(15) (a) Shao, C.-D.; Shi, G.-F.; Zhang, Y.-H.; Pan, S.-L.; Guan, X.-H. Org. Lett. **2015**, *17*, 2652. (b) Xiong, H.; Xu, H.; Liao, S.-H.; Xie, Z.-W.; Tang, Y. J. Am. Chem. Soc. **2013**, *135*, 7851. (c) Han, X.-P; Wu, J. Angew. Chem., Int. Ed. **2013**, *52*, 4637. (d) Yagoubi, M.; Cruz, A. C. F.; Nichols, P. L.; Elliott, R. L.; Willis, M. C. Angew. Chem., Int. Ed. **2010**, *49*, 7958.

161.