

Enantioselective Total Synthesis of Hyperforin

Brian A. Sparling, David C. Moebius, and Matthew D. Shair*

Department of Chemistry and Chemical Biology, Harvard University, Cambridge, Massachusetts 02138, United States


S Supporting Information

ABSTRACT: A modular, 18-step total synthesis of hyperforin is described. The natural product was quickly accessed using latent symmetry elements, whereby a group-selective, Lewis acid-catalyzed epoxide-opening cascade cyclization was used to furnish the bicyclo[3.3.1]nonane core and set two key quaternary stereocenters.

Hyperforin (**1**)^{1,2} was first isolated in 1971 from St. John's wort (*Hypericum perforatum* L.) and is now considered to be the constituent of the medicinal herb responsible for its antidepressant activity.^{3,4} Hyperforin blocks the re-uptake of a variety of neurotransmitters through a unique mechanism of action, possibly by selectively activating TRPC6 (classical transient receptor potential protein).⁵ TRPC6 activation leads to a cellular influx of Na⁺ and Ca²⁺, diminishing membrane electrochemical gradient (thus indirectly inhibiting neuronal neurotransmitter reuptake) and triggering cell differentiation. As a selective activator of TRPC6, hyperforin is a powerful probe of TRPC6 biology, a critical lead for the treatment of depression and possibly other human diseases. However, the therapeutic potential of hyperforin is severely handicapped by its poor water solubility, facile oxidative degradation upon exposure to light and air,⁶ and potent activation of pregnane X receptor,⁷ leading to increased expression of many genes involved in xenobiotic metabolism. Access to a wide variety of hyperforin analogues is critical for mitigating these shortcomings while maintaining TRPC6 activation. While limited semisynthetic manipulation of isolated hyperforin is feasible,⁸ total synthesis is the only possible means of obtaining diverse hyperforin analogues.

Structurally, hyperforin is one of over 200 polycyclic polyprenylated acylphloroglucinol (PPAP) natural products.⁹ Many PPAPs are characterized by a highly oxidized bicyclo[3.3.1]nonane core densely substituted with terpenoid side chains. Widespread interest in their bioactivity and structural complexity has culminated in the total syntheses of several PPAPs.¹⁰ While these PPAPs contain a geminal dimethyl group at the C8 position (hyperforin numbering), differential substitution at the hyperforin C8 position renders these otherwise effective strategies inapplicable toward hyperforin total synthesis. Numerous approaches toward hyperforin¹¹ have only resulted in one total synthesis of *ent*-hyperforin, accomplished by the Shibasaki group in 2010.¹² Given the considerable length of this route (51 steps from propargyl bromide), we set out to devise a new enantioselective approach that would not only incorporate elements of modularity but also exploit latent symmetry. A practical total synthesis would enable the first full SAR study of hyperforin.

Our strategy for the construction of hyperforin is shown in Scheme 1. The synthesis can be deconstructed into the

Scheme 1. (A) Synthesis Strategy for Hyperforin (**1**) and (B) Comparison of This Work to Hyperforin Biosynthesis

stepwise fusion of six easily obtainable chemicals (Scheme 1A). We postulated that **1** would be accessible from alcohol **2** through bridgehead acylation at C1 and prenylations at C3 and C7. We would access key intermediate **2** from cyclohexadiene **3** via a group-selective, Lewis acid-mediated epoxide-opening cascade cyclization. Cyclohexadiene **3** would be synthesized in two steps through the regioselective coupling of 1,5-dimethoxy-1,4-cyclohexadiene **4** with prenyl chloride **5** and known epoxygeranyl bromide **6**.¹³

In developing our hyperforin synthesis, we drew inspiration from its proposed biosynthesis¹⁴ (Scheme 1B). In nature, hyperforin is formed from alkylation of a nucleophilic,


Received: December 12, 2012

polyketide-derived acylphoroglucinol with geranyl pyrophosphate. Stereochemical fidelity over the course of this coupling in the biosynthesis is governed by enzyme catalysis.

We envisioned utilizing a similar bond disconnection in the key cyclization of our synthesis, specifically one involving a geraniol-derived fragment. While unable to exploit enzyme-driven stereoselection, we postulated that an epoxide functionality strategically placed in the geraniol fragment would be an effective means of relaying stereochemical information to the C1, C5, and C8 carbon centers, when employed in concert with a truncated, symmetric cyclohexadiene ring.

An analysis of the cyclization of **3** is depicted in Scheme 2. Owing to the symmetry of compound **3**, the quaternary carbon

Scheme 2. Transition-State Analysis of the Cyclization of 3



at C5 is prostereogenic, and the two methyl enol ethers are diastereotopic. Exposure of **3** to Lewis acid activates the epoxide for nucleophilic attack. While either methyl enol ether may engage this activated epoxide, transition state **TS-1** is


avored to yield **7** over its diastereomeric transition state **TS-2**, which must adopt a boat-like conformation containing two severe eclipsing interactions in forming **8**. Additionally, a 6-(*enolendo*)-tet cyclization should be favored over a 5-(*enolendo*)-tet cyclization due to geometric constraints of the substrate.¹⁵ The contributions of these factors culminate in (1) the formation of the hyperforin bicyclo[3.3.1]nonane skeleton; (2) the introduction of stereochemistry at the C5 quaternary position; and (3) the creation of a stereogenic quaternary center at C8.

Our synthesis of hyperforin is depicted in Scheme 3. Oxymercuration followed by reductive workup of epoxygeranyl bromide **6** and subsequent TESCl-mediated silylation gave **9**. Over the course of the synthesis, it became apparent that formal silanolysis of the olefin present in **6** was indispensable. Deprotonation of cyclohexadiene **4** with *t*-BuLi, treatment with freshly prepared BaI_2 ,¹⁶ and alkylation with prenyl chloride **7** yielded diene **11**. In the absence of BaI_2 , non-regioselective prenylation across the pentadienyl anion took place. Coupling of **9** with the anion generated from diene **11** and *s*-BuLi gave cyclization precursor **12**.

Gratifyingly, exposure of **12** to TMSOTf and 2,6-lutidine gave the ketal **13** in 79% yield as the only isolated product. As previously mentioned, the stereochemistry of two key quaternary centers were established as a result of this transformation: at the previously prostereogenic C5 carbon, and at the C8 position. Aside from the construction of the bicyclo[3.3.1]nonane framework, the formation of a cyclic methyl ketal was an unexpected yet fortuitous outcome to this reaction, safeguarding the C7 carbinol during the subsequent allylic oxidation step. The efficiency of this approach was demonstrated by our ability to easily prepare 60 g of **13**.

After extensive optimization, allylic oxidation of **13** was accomplished using TBHP, $\text{PhI}(\text{O}_2\text{CCF}_3)_2$, and O_2 ¹⁷ to afford

Scheme 3. Total Synthesis of Hyperforin (1)^a



^aConditions: (a) $\text{Hg}(\text{OAc})_2$, $\text{Me}_2\text{CO}/\text{H}_2\text{O}$; NaOH , 0°C ; NaBH_4 , 0°C , 91%; (b) TESCl , imid, DMF, 97%; (c) *t*-BuLi, THF, -78°C ; BaI_2 , -78°C ; **5**, -78 to -5°C , 85%; (d) **11**, *s*-BuLi, THF, -78 to -30°C ; **9**, -78 to -40°C , 85%; (e) TMSOTf, 2,6-lut, CH_2Cl_2 , -78°C , 79%; (f) $\text{PhI}(\text{O}_2\text{CCF}_3)_2$, TBHP, Cs_2CO_3 , 4 Å MS, EtOAc, O_2 , -78 to 0°C , 44%; (g) BrBMe_2 , NEt_3 , CH_2Cl_2 , -95°C ; NEt_3 ; sat. aq. NaHCO_3 , 57%; (h) LiTMP, THF, -78 to 0°C , 97%; (i) $\text{ClC}(\text{S})\text{OC}_6\text{F}_5$, *N*-hydroxysuccinimide, pyr, PhMe, 80°C , 82%; (j) allyl-SnBu₃, BEt₃, PhH, air, 72%; (k) **17**, 2-methyl-2-butene, CH_2Cl_2 , 40°C , 86%; (l) LiTMP, TMSOTf, THF, -78 to 0°C , 90%; (m) LiTMP, THF, -78 to 0°C ; *i*-PrC(O)CN, -78 to -30°C , 49%; (n) *p*-TsOH·H₂O, PhMe, HOAc, 2-methyl-2-butene, microwave, 100°C , 65%; (o) LDA, THF, -78°C ; $\text{Li}(2\text{-Th})\text{CuCN}$, -78 to -40°C ; prenyl bromide, -78 to -30°C , 98%; (p) LiCl, DMSO, 120°C , 55%.

vinylous ester **14**.¹⁸ This was an exceptionally challenging transformation given the four allylic sites present in **13** and the steric environment surrounding the desired oxidation site. Hydrolysis of the ketal present in **14** to yield alcohol **15** was performed in two steps: treatment of **14** with BrBMe₂¹⁹ followed by LiTMP-mediated methanol extrusion from the intermediate hemiketal. It was crucial to maintain the temperature of the BrBMe₂-mediated hydrolysis below −90 °C; above this temperature, elimination of the triethylsilyl ether was observed.

After surveying a variety of methods to install the C7 prenyl group from **15**, we pursued a radical Keck allylation approach.²⁰ A radical precursor, thionocarbonate **16**, was generated from the reaction of alcohol **15** with ClC(S)OC₆F₅.²¹ Using BEt₃/air as an initiator, radical allylation of **16** with allyl–SnBu₃ afforded a product containing a C7 allyl group as a single diastereomer. Employing more reactive radical precursor functionality or either photochemical or thermal radical generation conditions gave inferior results for this coupling reaction. An ensuing cross-metathesis with 2-methyl-2-butene catalyzed by Hoveyda–Grubbs second-generation catalyst **17**²² afforded **18** containing the requisite C7 prenyl moiety.

After silylation at the C3 position, sequential bridgehead deprotonation–acylation using LiTMP and *i*-PrC(O)CN¹⁰ⁿ yielded ketone **19**. This direct, one-step bridgehead acylation is noteworthy given that previously reported instances of PPAP bridgehead acylation at the C1 position require multiple steps involving a bridgehead iodide intermediate.^{10b,d,l} One-pot desilylation and elimination to give **20** was accomplished through microwave irradiation of **19** with *p*-TsOH·H₂O. The final C3 prenyl group was installed utilizing a precedented sequence²³ to afford hyperforin methyl ether **21**.²⁴ (1) deprotonation of **20** with LDA, (2) transmetalation with Li(2-Th)CuCN,²⁵ and (3) trapping with prenyl bromide. Finally, hyperforin (**1**) was revealed by heating a DMSO solution of **21** with LiCl.

In summary, we report an enantioselective total synthesis of hyperforin. The synthesis is 18 steps at its longest sequence, starting from geraniol. This approach is also highly scalable; to date, we have prepared over 40 mg of hyperforin. Latent symmetry elements were utilized to quickly access the hyperforin bicyclo[3.3.1]nonane core and to set two key quaternary stereocenters, specifically in the conversion of epoxide **12** to ketal **13**. This practical and modular route is already being exploited to create diverse hyperforin analogues, which we are using to further understand the SAR and underlying mechanisms of hyperforin biological and medicinal activity. Results from these studies will be reported in due course.

■ ASSOCIATED CONTENT

■ Supporting Information

Experimental procedures, spectroscopic data, and ¹H and ¹³C NMR spectra. This material is available free of charge via the Internet at <http://pubs.acs.org>.

■ AUTHOR INFORMATION

Corresponding Author

shair@chemistry.harvard.edu

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

We acknowledge financial support from Chemiderm, Inc. B.A.S. acknowledges a NSF Predoctoral Fellowship.

■ REFERENCES

- (1) (a) Gurevich, A. I.; Dobrynin, V. N.; Kolosov, M. N.; Popravko, S. A.; Ryabova, I. D.; Chernov, B. K.; Derbentseva, N. A.; Aizenman, B. E.; Garagulya, A. D. *Antibiotiki* **1971**, *16*, 510. (b) Bystrov, N. S.; Chernov, B. K.; Dobrynin, V. N.; Kolosov, M. N. *Tetrahedron Lett.* **1975**, *16*, 2791. (c) Brondz, I.; Greibokk, T.; Groth, P. A.; Aasen, A. J. *Tetrahedron Lett.* **1982**, *23*, 1299. (d) Brondz, I.; Greibokk, T.; Groth, P.; Aasen, A. J. *Acta Chem. Scand. A* **1983**, *37*, 263.
- (2) For reviews specific to hyperforin, see: (a) Beerhues, L. *Phytochem.* **2006**, *67*, 2201. (b) Medina, M. A.; Martínez-Poveda, B.; Amores-Sánchez, M. I.; Quesada, A. R. *Life Sci.* **2006**, *79*, 105. (c) Quiney, C.; Billard, C.; Salanoubat, C.; Fourneron, J. D.; Kolb, J. P. *Leukemia* **2006**, *20*, 1519.
- (3) (a) Greeson, J. M.; Sanford, B.; Monti, D. A. *Psychopharmacology* **2001**, *153*, 402. (b) Di Carlo, G.; Borrelli, F.; Ernst, E.; Izzo, A. A. *Trends Pharmacol. Sci.* **2001**, *22*, 292. (c) Müller, W. E. *Pharmacol. Res.* **2003**, *47*, 101. (d) Linde, K. *Forsch. Komplementmed.* **2009**, *16*, 146.
- (4) Chatterjee, S. S.; Bhattacharya, S. K.; Wonnemann, M.; Singer, A.; Müller, W. E. *Life Sci.* **1998**, *63*, 499. Also, see: *Pharmacopsychiatry* **1998**, *31*, Supplementary Issue 1 and *Pharmacopsychiatry* **2001**, *34*, Supplementary Issue.
- (5) Leuner, K.; Kazanski, V.; Müller, M.; Essin, K.; Henke, B.; Gollasch, M.; Harteneck, C.; Müller, W. E. *FASEB J.* **2007**, *21*, 4101.
- (6) (a) Trifunović, S.; Vajs, V.; Macura, S.; Juranić, N.; Djarmati, Z.; Jankov, R.; Milosavljević, S. *Phytochemistry* **1998**, *49*, 1305. (b) Wolfender, J.-L.; Verotta, L.; Belvisi, L.; Fuzzati, N.; Hostettmann, K. *Phytochem. Anal.* **2003**, *14*, 290. (c) Verotta, L.; Lovaglio, E.; Sterner, O.; Appendino, G.; Bombardelli, E. *Eur. J. Org. Chem.* **2004**, 1193. (d) Ang, C. Y. W.; Hu, L.; Heinze, T. M.; Cui, Y.; Freeman, J. P.; Kozak, K.; Luo, W.; Lui, F. F.; Mattia, A.; DiNovi, M. J. *Agric. Food Chem.* **2004**, *52*, 6156. (e) D'Auria, M.; Emanuele, L.; Racioppi, R. *Lett. Org. Chem.* **2008**, *5*, 583.
- (7) (a) Moore, L. B.; Goodwin, B.; Jones, S. A.; Wisely, G. B.; Serabjit-Singh, C. J.; Willson, T. M.; Collins, J. L.; Kliever, S. A. *Proc. Natl. Acad. Sci. U.S.A.* **2000**, *97*, 7500. (b) Watkins, R. E.; Maglich, J. M.; Moore, L. B.; Wisely, G. B.; Noble, S. M.; Davis-Searles, R.; Lambert, M. H.; Kliever, S. A.; Redinbo, M. R. *Biochemistry* **2003**, *42*, 1430.
- (8) For examples, see: (a) Verotta, L.; Appendino, G.; Belloro, E.; Bianchi, F.; Sterner, O.; Lovati, M.; Bombardelli, E. *J. Nat. Prod.* **2002**, *65*, 433. (b) Gartner, M.; Müller, T.; Simon, J. C.; Giannis, A.; Sleeman, J. P. *ChemBioChem* **2005**, *6*, 171.
- (9) For reviews of PPAPs, see: (a) Ciochina, R.; Grossman, R. B. *Chem. Rev.* **2006**, *106*, 3963. (b) Singh, I. P.; Sidana, J.; Bharate, S. B.; Foley, W. J. *Nat. Prod. Rep.* **2010**, *27*, 393. (c) Tsukano, C.; Siegel, D. R.; Danishefsky, S. J. *J. Synth. Org. Chem. Jpn.* **2010**, *68*, 592. (d) Njardarson, J. T. *Tetrahedron* **2011**, *67*, 7631. (e) Richard, P. A.; Pouwer, R. H.; Chen, D. Y.-K. *Angew. Chem., Int. Ed.* **2012**, *51*, 4536.
- (10) (a) Kuramochi, A.; Usuda, H.; Yamatsugu, K.; Kanai, M.; Shibasaki, M. *J. Am. Chem. Soc.* **2005**, *127*, 14200. (b) Siegel, D. R.; Danishefsky, S. J. *J. Am. Chem. Soc.* **2006**, *128*, 1048. (c) Rodeschini, V.; Ahmad, N. M.; Simpkins, N. S. *Org. Lett.* **2006**, *8*, 5283. (d) Tsukano, C.; Siegel, D. R.; Danishefsky, S. J. *Angew. Chem., Int. Ed.* **2007**, *46*, 8840. (e) Qi, J.; Porco, J. A., Jr. *J. Am. Chem. Soc.* **2007**, *129*, 12682. (f) Nuhant, P.; David, M.; Pouplin, T.; Delpech, B.; Marazano, C. *Org. Lett.* **2007**, *9*, 287. (g) Rodeschini, V.; Simpkins, N. S.; Wilson, C. J. *Org. Chem.* **2007**, *72*, 4265. (h) Ahmad, N. M.; Rodeschini, V.; Simpkins, N. S.; Ward, S. E.; Blake, A. J. *J. Org. Chem.* **2007**, *72*, 4803. (i) Qi, J.; Beeler, A. B.; Zhang, Q.; Porco, J. A., Jr. *J. Am. Chem. Soc.* **2010**, *132*, 13642. (j) Zhang, Q.; Mitasev, B.; Qi, J.; Porco, J. A., Jr. *J. Am. Chem. Soc.* **2010**, *132*, 14212. (k) Garnsey, M. R.; Lim, D.; Yost, J. M.; Coltart, D. M. *Org. Lett.* **2010**, *12*, 5234. (l) Simpkins, N. S.; Taylor, J. D.; Weller, M. D.; Hayes, C. J. *Synlett* **2010**, 639. (m) Simpkins, N. S.; Weller, M. D. *Tetrahedron Lett.* **2010**, *51*, 4823.

- (n) Biber, N.; Möws, K.; Plietker, B. *Nature Chem.* **2011**, *3*, 938.
- (o) Garnsey, M. R.; Matous, J. A.; Kwiek, J. J.; Coltart, D. M. *Bioorg. Med. Chem. Lett.* **2011**, *21*, 2406.
- (p) Uwamori, M.; Saito, A.; Nakada, M. *J. Org. Chem.* **2012**, *77*, 5098.
- (11) For examples, see: (a) Kraus, G. A.; Nguyen, T. H.; Jeon, I. *Tetrahedron Lett.* **2003**, *44*, 659. (b) Nicolaou, K. C.; Carenzi, E. A.; Jeso, V. *Angew. Chem., Int. Ed.* **2005**, *44*, 3895. (c) Couladouros, E. A.; Dakanali, M.; Demadis, K. D.; Vidali, V. P. *Org. Lett.* **2009**, *11*, 4430. (d) Uyeda, C.; Rötheli, A. R.; Jacobsen, E. N. *Angew. Chem., Int. Ed.* **2010**, *49*, 9753. (e) Mehta, G.; Dhanbal, T.; Bera, M. K. *Tetrahedron Lett.* **2010**, *51*, 5302. (f) Richard, J.-A.; Chen, D. Y.-K. *Eur. J. Org. Chem.* **2012**, 484 and references therein.
- (12) (a) Usuda, H.; Kuramochi, A.; Kanai, M.; Shibasaki, M. *Org. Lett.* **2004**, *6*, 4387. (b) Shimizu, Y.; Shi, S.-L.; Usuda, H.; Kanai, M.; Shibasaki, M. *Angew. Chem., Int. Ed.* **2010**, *49*, 1103. (c) Shimizu, Y.; Shi, S.-L.; Usuda, H.; Kanai, M.; Shibasaki, M. *Tetrahedron* **2010**, *66*, 6569.
- (13) **4** is synthesized from 1,3-dimethoxybenzene via Birch reduction: Piers, E.; Grierson, J. R. *J. Org. Chem.* **1977**, *42*, 3755. **6** is synthesized from geraniol in three steps via Sharpless epoxidation, mesylation, and bromination: (a) Hanson, R. M.; Sharpless, K. B. *J. Org. Chem.* **1986**, *51*, 1922. (b) Gash, R. C.; MacCorquodale, F.; Walton, J. C. *Tetrahedron* **1989**, *45*, 5531 for more detail.
- (14) (a) Adam, P.; Arigoni, D.; Bacher, A.; Eisenreich, W. *J. Med. Chem.* **2002**, *45*, 4786. (b) Boubakir, Z.; Beuerle, T.; Liu, B.; Beerhues, L. *Phytochemistry* **2005**, *66*, 51. (c) Klingauf, P.; Beuerle, T.; Mellenthin, A.; El-Moghazy, S. A. M.; Boubakir, Z.; Beerhues, L. *Phytochemistry* **2005**, *66*, 139. (d) Karppinen, K.; Hokkanen, J.; Tolonen, A.; Mattila, S.; Hohtola, A. *Phytochemistry* **2007**, *68*, 1038.
- (15) Baldwin, J. E.; Lusch, M. *J. Tetrahedron* **1982**, *38*, 2939.
- (16) For more information on BaI_2 -promoted allylation, see: (a) Yanagisawa, A.; Yasue, K.; Yamamoto, H. *Synlett* **1993**, 686. (b) Yanagisawa, A.; Habaue, S.; Yasue, K.; Yamamoto, H. *J. Am. Chem. Soc.* **1994**, *116*, 6130. (c) Corey, E. J.; Lin, S.; Luo, G. *Tetrahedron Lett.* **1997**, *38*, 5771. (d) Karavathi, S.; Corey, E. J. *J. Am. Chem. Soc.* **2008**, *130*, 8865.
- (17) For examples of similar allylic oxidation systems, see: (a) Zhao, Y.; Yeung, Y.-Y. *Org. Lett.* **2010**, *12*, 2128. (b) Zhao, Y.; Yim, W.-L.; Tan, C. K.; Yeung, Y.-Y. *Org. Lett.* **2011**, *13*, 4308.
- (18) For other examples of allylic oxidation to form the C2–C4 vinylogous ester, see ref 10p and the following: Takagi, R.; Inoue, Y.; Ohkata, K. *J. Org. Chem.* **2008**, *73*, 9320.
- (19) (a) Guidon, Y.; Yoakim, C.; Morton, H. E. *Tetrahedron Lett.* **1983**, *24*, 2969. (b) Guidon, Y.; Yoakim, C.; Morton, H. E. *J. Org. Chem.* **1984**, *49*, 3912.
- (20) (a) Keck, G. E.; Byers, J. H. *J. Org. Chem.* **1985**, *50*, 5442. (b) Keck, G. E.; Enholm, E. J.; Yates, J. B.; Wiley, M. R. *Tetrahedron* **1985**, *41*, 4079. A similar approach to C7 prenylation has been used involving a C7 iodide intermediate. For more information, see refs 10b and d.
- (21) Barton, D. H. R.; Jaszberenyi, J. C. *Tetrahedron Lett.* **1989**, *30*, 2619.
- (22) Garber, S. B.; Kingsbury, J. S.; Gray, B. L.; Hoveyda, A. H. *J. Am. Chem. Soc.* **2000**, *122*, 8168.
- (23) For other examples of C3 metalation–alkylation, see refs 10d, h, l, p, and the following: Ahmad, N. M.; Rodeschini, V.; Simpkins, N. S.; Ward, S. E.; Wilson, C. *Org. Biomol. Chem.* **2007**, *5*, 1924.
- (24) The NMR data for this compound are consistent with those of **21** generated from treatment of isolated hyperforin with diazomethane (ref 8a). See Supporting Information for more details.
- (25) Lipshutz, B. H.; Koerner, M.; Parker, D. A. *Tetrahedron Lett.* **1987**, *28*, 945.