


Highly Functionalized Enantiopure Oxatricyclo[3.3.1.0^{4,6}]nonanes; Selective Opening of an Activated Cyclopropane Ring

Norbert A. Braun and Dietrich Spitzner*


Institut für Chemie, Abt. Organische Chemie, Universität Hohenheim, Garbenstraße 30,
D-70599 Stuttgart, Germany

Abstract: Baeyer-Villiger oxidation of **3** gives solely the lactones **4**. The double-pull-push substituted cyclopropane moiety of **4b** is selectively opened upon hydrogenolysis of the benzyl ether to yield ketone **5**. Copyright © 1996 Elsevier Science Ltd

Functionalized tricyclo[3.2.1.0^{2,7}]octanes are valuable starting materials in the total synthesis of terpenoid natural products such as kaurene, gibberellins and grayanotoxin.¹ Enantiopure tricyclo[3.2.1.0^{2,7}]octanes **3** are obtained by a diastereoselective cascade reaction² of the chiral Michael acceptors (+)-**1** with the dienolates **2** (d. e. >95%). The absolute configuration of **3b**³ was deduced from the CD spectrum [CD (methanol): λ_{max} ($\Theta/\Delta\epsilon$) = 286.5 nm (+ 14984 / + 4.54)] which is similar to that of **3a** [CD (methanol): λ_{max} ($\Theta/\Delta\epsilon$) = 287 nm (+ 13504 / + 4.09)].⁴



In order to extend the range of accessible carbon skeletons starting from **3** we investigated the insertion of additional hetero atoms. Baeyer-Villiger oxidation⁵ of **3** with MCPBA in dichloromethane buffered with solid $KHCO_3$ gave exclusively the lactones **4** in good yields.⁶ The bridge head C-5 of the parent tricyclo[3.2.1.0^{2,7}]octane **3** migrated in both cases. Hydrogenolysis of the benzyl ether of **4a** gave an intermediate double-pull-push activated cyclopropanol⁷ which spontaneously underwent a retro-grade aldol reaction to give the oxabicyclo[3.3.1]nonane **5** (cleavage of the C4-C6 bond).⁸ The alternative opening of the C5-C6 bond was not observed:



ACKNOWLEDGEMENTS: We thank the *Deutsche Forschungsgemeinschaft*, Bonn, the *Fonds der Chemischen Industrie e. V.*, Frankfurt (Main), for financial support and *Chemetall GmbH*, Frankfurt (Main), for generous gifts of chemicals. N.A.B. thanks the *Landesgraduierföderung*, Baden-Württemberg, for a scholarship.

REFERENCES AND NOTES

1. Sagawa, S.; Nagaoka, H.; Yamada, Y. *Tetrahedron Lett.* **1994**, 35, 603-604.
2. a) Tietze, L.F. *Chem. Rev.* **1996**, 96, 115-136, b) Tietze, L.F.; Beifuss, U. *Angew. Chem.* **1993**, 105, 137-170; *Angew. Chem. Int. Ed. Engl.* **1993**, 32, 131, c) Ihara, M.; Fukumoto, K. *Angew. Chem.* **1993**, 105, 1059-1071; *Angew. Chem. Int. Ed.* **1993**, 1010-1022.
3. **3b**: 68 %, oil. - $[\alpha]^{25}_D = +54.0^\circ$ ($c = 1.42$ in CHCl_3). - ^{13}C NMR(CDCl_3): $\delta = 13.93$ (q), 20.35 (q), 23.88 (t, C-4), 24.67 (t, C-3), 25.53 (q), 26.41 (q), 40.44 (s, C-2), 42.24 (s, C-1), 42.52 (d, C-8), 44.15 (d, C-7), 44.73 (d, C-5), 61.02 (t), 70.10 (t, C-5'), 75.60 (d, C-4'), 107.85 (s, C-2'), 169.42 (s), 210.49 (s, C-6).
4. a) Braun, N. A.; Klein, I.; Spitzner, D.; Vogler, B.; Borrmann, H.; Simon, A. *Tetrahedron Lett.* **1995**, 36, 3675-3678, b) Braun, N. A.; Klein, I.; Spitzner, D.; Vogler, B.; Braun, S.; Borrmann, H.; Simon, A. *Liebigs Ann.* **1995**, 2165-2169.
5. Krow, G. R. *Tetrahedron* **1981**, 37, 2697-2774.
6. **4a**: 90 %, white crystals, mp 117-119°C. - $[\alpha]^{24}_D = +38.1^\circ$ ($c = 1.12$ in CHCl_3). - ^{13}C NMR(CDCl_3): $\delta = 21.61$ (t, C-7), 22.12 (t, C-8), 24.67 (q), 26.58 (q), 27.63 (q), 33.53 (d, C-9), 40.17 (d, C-4), 40.40 (s, C-5), 66.72 (s, C-6), 69.98 (t, C-5'), 70.67 (t), 74.35 (d, C-1), 75.65 (d, C-4'), 82.88 (s), 108.83 (s, C-2'), 127.82, 127.93, 128.29 (d), 136.61 (s), 166.01 (s), 168.54 (s, C-3). **4b**: 55 %, oil. - $[\alpha]^{24}_D = +43.8^\circ$ ($c = 1.28$ in CHCl_3). - ^{13}C NMR(CDCl_3): $\delta = 13.75$ (q), 20.69 (t, C-8), 23.64 (q), 24.15 (t, C-7), 24.79 (q), 26.21 (q), 29.62 (s, C-6), 33.70 (d, C-9), 37.14 (s, C-5), 39.08 (d, C-4), 61.30 (t), 69.63 (t, C-5'), 74.11 (d, C-1), 75.68 (d, C-4'), 108.42 (s, C-2'), 168.87 (s, C-3), 169.31 (s).
7. a) Reißig, H.-U. *Top. Curr. Chem.* **1988**, 144, 73-135, b) Reißig, H.-U.; Hirsch, E. *Angew. Chem.* **1980**, 92, 839-840; *Angew. Chem. Int. Ed. Engl.* **1980**, 19, 813, c) Wenkert, E. *Acc. Chem. Res.* **1980**, 13, 27-31.
8. **5**: 96 %, white crystals, mp 123-125°C. - $[\alpha]^{24}_D = +58.1^\circ$ ($c = 1.08$ in CHCl_3). - ^{13}C NMR(CDCl_3): $\delta = 24.59$ (q), 26.50 (q), 27.82 (q), 28.18 (t, C-8), 33.37 (t, C-7), 37.76 (t, C-4), 48.26 (d, C-9), 58.36 (s, C-5), 71.66 (t, C-5'), 73.83 (d, C-1), 75.05 (d, C-4'), 83.57 (s), 108.96 (s, C-2'), 167.84 (s), 168.54 (s, C-3), 205.00 (s, C-6).

(Received in Germany 17 October 1996; accepted 5 November 1996)