ORIGINAL RESEARCH

Antibacterial properties of 5-substituted derivatives of rhodanine-3-carboxyalkyl acids

Waldemar Tejchman¹ · Izabela Korona-Glowniak² · Anna Malm² · Marek Zylewski³ · Piotr Suder^{4,5}

Received: 31 October 2015 / Accepted: 2 March 2017 © The Author(s) 2017; This article is published with open access at Springerlink.com

Abstract A series of rhodanine 3-carboxyalkanoic acid derivatives possessing 4'-(N,N-dialkyl-amino or diphenylamino)-benzylidene moiety as a substituent at the C-5 position were synthesised and their antibacterial activity was screened. All the rhodanine derivatives showed bacteriostatic or bactericidal activity to the reference gram-positive bacterial strains, but lack of activity to the reference Gramnegative bacterial strains and yeast strains was observed.

Keywords Rhodanine · Thiazolidine-4-one · Rhodanine-3acetic acid · Antibacterial activity

⊠ Izabela Korona-Glowniak iza.glowniak@umlub.pl

- ¹ Departament of Chemistry, Institute of Biology, Pedagogical University of Cracow, Podchorazych 2, Kraków 30-084, Poland
- ² Department of Pharmaceutical Microbiology, Medical University of Lublin, Chodzki 1, Lublin 20-093, Poland
- ³ Jagiellonian Center of Innovation, NMR Laboratory, Bobrzyńskiego 14, Kraków 30-348, Poland
- ⁴ Department of Biochemistry and Neurobiology, AGH University of Science and Technology, Mickiewicza 30, Kraków 30–059, Poland
- ⁵ Academic Centre for Materials and Nanotechnology, AGH University of Science and Technology, Mickiewicza 30, Kraków 30–059, Poland

Introduction

The 2-thiazolidine-4-one derivatives traditionally named rhodanine have been known for over 100 years, and due to their fascinating properties they are still examined (Lesyk and Zimenkovsky 2004). These compounds have a broad spectrum of biological effects (Jain et al. 2012). Rhodanine derivatives show antimalarial (Kumar et al. 2007), antitubercular (Alegaon et al. 2012), cytotoxic (Chandrappa et al. 2009), antitumor (Rao et al. 2011; Lesyk et al. 2011), antiviral (Kaminskyy 2015), and antibacterial activity (Bhatti et al. 2013; Kavitha et al. 2006; Song et al. 2014).

The research to obtain new antibacterial compounds is vitally important. Recently, due to excessive and improper use of antibiotics, there has been an increasing rate of antibiotic resistance in the bacterial strains (Woodford 2003), thus new groups of compounds which may be useful as antibacterial agents have been examined. A few reports has been published regarding the rhodanine derivatives with a carboxyalkyl acid moiety at the N-3 position (Xu et al. 2012). Biological activity of hybrid compounds possessing chalcone and rhodanine-3-acetic acid has been also studied (Chen et al. 2010). Such hybrids demonstrated synergistic effect. Antibacterial activity of rhodanine derivatives and their oxygen analogues derived from 2,4-thiazolidinedione was also compared (Zvarec et al. 2012). However, the results of present study suggested that rhodanine derivatives showed greater antibacterial activity than their analogues from the 2,4-thiazolidinedione group having at the C-2 position exocyclic oxygen atom. It was shown that the activity of the rhodanine derivative correlates with the size of the substituent at the C-5 position (Pardasani et al. 2001). The research conducted by Miao et al. (2013) and Patel et al. (2013) indicated that antibacterial activity of the acid derivatives occurred when a major hydrophobic group was



introduced to the arylidene substituent at the C-5 position. The best results were achieved when an aryl group additionally with an electron-withdrawing group was introduced. The rhodanine derivatives possessing a 4-(*N*,*N*dimethylamino)-benzylidene substituent at the C-5 position were also examined. These compounds acted as β -lactamase inhibitors (Grant et al. 2000). Taking into account the data presented by other authors, we decided to synthesise a series of derivatives having carboxyalkyl (acetic, propionic, butyric) acid fragment at N-3 position and benzylidene para-substituent with dimethyloamino, diethylamino, dibuthyloamino or diphenylamino group at C-5 position.

Chemistry

Our initial research proved that the antibacterial activity of the rhodanine derivatives which have carboxyalkyl fragment at N-3 position was more effective than the compounds with a substituent containing an amino group at C-5 position. We synthesised a series of rhodanine derivatives with a carboxyalkyl acid radical at N-3 position (acetic, propionic, butyric, caproic). The synthesis of the 3carboxyalkylrhodanine acids (Scheme 1) was conducted according to the modified procedure proposed by Körner (1908) at the beginning of the 20th century.

The synthesised compounds underwent Knoevenagel condensation with 4-diethylaminobenzoic, 4-dibutylaminobenzoic aldehydes and 4-diphenylaminobenzoic aldehyde with triethylamine as a catalyst. Quaternary ammonium salts, the intermediates obtained during reactions, were not isolated but transformed to appropriate acids with hydrochloric acid (Scheme 2).

Material and methods

All reagents for the synthesis of rhodanine derivatives were purchased from Sigma-Aldrich and used without further purification.

Melting point (uncorrected) has been determined on the Boetius apparatus. The IR spectrum has been recorded with Jasco FT IR-670 Plus spectrophotometer in the KBr disk.

The NMR spectra were obtained in CDCl_3 on the Bruker Avance III HD spectrometer operating at 400.17 MHz (1H) and 100.62 MHz (13C) and the Varian Mercury-VX 300 spectrometer operating at 300.08 MHz (1H) and 75.46 MHz (13C), the chemical shifts (ppm) have been referenced to lock out the signal of the solvent, *J* has been expressed in Hz.

The MS analyses were obtained on the AmaZon ETD mass spectrometer (Bruker Daltonics, Bremen, Germany). Scan parameters: scan range 100–1000 m/z, positive ionisation mode. CID fragmentation were in the ion trap



Scheme 1 . 3-Carboxyalkylrhodanine acids synthesis



Scheme 2 Rhodanine-3-carboxyalkyl acid condensation with aldehydes

analyser with the aid of helium gas. The collision energy was set to ca. 1 eV. The samples were introduced into the mass spectrometer in a $CH_3OH:CHCl_3$ 1:1 solution with 0.1% HCOOH acidification.

General procedure of rhodanine-3-alkanoic acids synthesis

The solution of 11.22 g (0.2 mol) potassium hydroxide in 50 cm^3 of water was added to the suspension of 0.1 mol of the appropriate amino acid (aminoethanoic acid, 3-aminopropanoic acid, 4-aminobutanoic acid and 6-aminohexanoic acid). The resulting solution was cooled to 5 °C and 7.6 g (0.1 mol) carbon disulphide was added. The content of the flask was mixed at 5 °C for 7 h. The cooling bath was removed and mixing was continued in room temperature for 20 h.

The solution of 9.45 g (0.1 mol) chloroacetic acid in 50 cm^3 water was added to the resulting solution. The solution was mixed for 7 h at the temperature below 15 °C. Next, the solution of 60 cm³ hydrochloric acid in 100 cm³ of water was added to the flask content. The resulting mixture was heated to 90 °C and kept at the temperature for 20 min. After cooling, a sediment was received, which was drained and crystallised from water.

General procedure of rhodanine-3-alkanoic acids condensation with aldehydes

0.005 mol of appropriate rhodanine-3-alkanoic acid, 5 g molecular sieves 4 A, 25 cm³ isopropyl alcohol, 0.0055 mol appropriate aldehyde and 2.53 g (0.025 mol) triethylamine were placed in a flask. The mixture was heated under a reflux condenser for 5 h in nitrogen. After heating, the solution was filtered hot. The permeate was cooled and 50 cm³ of 2M hydrochloric acid solution was added. The resulting sediment was filtered using Büchner funnel and crystallised from isopropyl alcohol or glacial acetic acid.

3a/ 5-(4'-N,N-diethylaminobenzylidene)-rhodanine-3-acetic acid

m.p. 239-241 °C, yield 43.48%, MS [M+1]⁺--351.1, IR cm⁻³: 1719.3C=O, 1699.9C=O conj., 1612.2, C=C exo., 1322.9 C–N. 1185.1C=S, $^{1}\mathrm{H}$ NMR(400 MHz, CDCl₃+MeOD), δ ppm, 7.66 (s, 1H,=CH-Ar), 7.37 (d, J = 8.97 Hz, 2H, Ar-H), 6,70 (d, J = 9.00 Hz, 2H, Ar-H), 4.81 (s, 2H, HOOC-CH₂-N), 3.43 (q, 4H, N(CH₂CH₃)₂), 1.20 (t, 6H, $N(CH_2CH_3)_2$) ¹³C NMR (101 MHz, CDCl₃+MeOD), δ ppm, 12.35 (N(CH₂<u>C</u>H₃)₂), 44.61 (CH₂-N), 47.70 (N(CH₂CH₃)₂), 111.67 (Ar-C), 114.09 (Ar-C), 119.93 (Ar-C), 133.78 (Ar-C), 135.65 (=CH-Ar), 150.01 (S-C=CH), 167.68 (N-C=O), 168.23 (HOOC-), 193.16 (S=C-S)

3b/ 5-(4'-N,N-diethylaminobenzylidene)-rhodanine-3propionic acid

m.p. 202–204 °C, yield 54.70%, MS $[M+1]^+$ —365.1, IR cm⁻³: 1716.2C=O, 1700.9C=O conj., 1610.3, C=C exo., 1331.6 C–N, 1199.5C=S, ¹H NMR(400 MHz, CDCl₃), δ ppm, 7.69 (s, 1H,=C<u>H</u>–Ar), 7.40 (d, J = 9.00 Hz, 2H, Ar–H), 6.72 (d, J = 9.09 Hz, 2H, Ar–H), 4.45 (t, 2H, HOOC–CH₂–CH₂–N), 3.46 (q, 4H, N(CH₂CH₃)₂), 2.85 (t, 2H, HOOC–C<u>H</u>₂–CH₂–N), 1.24 (t, 6H, N(CH₂C<u>H₃)₂) ¹³C</u> NMR (101 MHz, CDCl₃), δ ppm, 12.57 (N(CH₂C<u>H₃)₂), 30.97 (CH₂–CH₂–N), 39.45 (CH₂–CH₂–N), 44.70(N (CH₂CH₃)₂), 111.70 (Ar–C), 114.51 (Ar–C), 120.17</u>

(Ar–C), 133.72 (Ar–C), 135.23 (=<u>C</u>H–Ar), 149.82 (S– <u>C</u>=CH), 167.76 (N–<u>C</u>=O), 175.59 (HOO<u>C</u>–), 192.98 (S=<u>C</u>–S)

3c/ 5-(4'-N,N-diethylaminobenzylidene)-rhodanine-3butyric acid

m.p. 155–157 °C, yield 17.06%, MS $[M+1]^+$ —379.1, IR cm⁻³: 1716.3C=O, 1693.2C=O conj., 1610.2C=C exo., 1340.3 C–N, 1194.7C=S, ¹H NMR(400 MHz, CDCl₃), δ ppm, 7.67 (s, 1H,=C<u>H</u>–Ar), 7.39 (d, J=8.84 Hz, 2H, Ar–H), 6,72 (d, J=9.04 Hz, 2H, Ar–H), 4.21 (t, 2H, HOOC–CH₂–CH₂–CH₂–N), 3.45 (q, 4H, N(C<u>H₂CH₃)₂), 2.45 (t, 2H, HOOC–CH₂–CH₂–CH₂–CH₂–CH₂–N), 2.09 (q, 2H, HOOC–CH₂–CH₂–CH₂–CH₂–CH₂–N), 1.26 (t, 6H, N(CH₂C<u>H₃)₂), 22.28 (CH₂–CH₂–CH₂–N), 31.17 (CH₂–CH₂–CH₂–N), 43.43 (CH₂–CH₂–CH₂–N), 44.67 (N(CH₂CH₃)₂), 111.66 (Ar–C), 114.74 (Ar–C), 120.22 (Ar–C), 133.66 (Ar–C), 134.94 (=<u>C</u>H–Ar), 149.75 (S–<u>C</u>=CH), 168.23 (N–<u>C</u>=O), 177.79 (HOO<u>C</u>–), 193.38 (S=<u>C</u>–S)</u></u>

3d/ 5-(4'-N,N-diethylaminobenzylidene)-rhodanine-3caproic acid

m.p. 135–137 °C, yield 32.96%, MS [M+1]⁺--407.1, IR cm⁻³: 1717.3C=O, 1702.8C=O conj., 1615.1C=C exo., 1327.7 C-N, 1195.7C=S, ¹H NMR(400 MHz, CDCl₃), δ ppm,11.25 (br. s HOOC-) 7.66 (s, 1H,=CH-Ar), 7.39 (d, J = 8.97 Hz, 2H, Ar-H), 6,72 (d, J = 8.48 Hz, 2H, CH2-N), 3.45 (q, 4H, N(CH2CH3)2), 2.39 (t, 2H, HOOC-CH2-CH2-CH2-CH2-CH2-N), 1.74 (m, 4H, HOOC-CH2- $CH_2-CH_2-CH_2-N$, 1.25 (t, 6H, N(CH_2CH_3)₂) ¹³C NMR (101 MHz, CDCl₃), δ ppm, 12.58 (N(CH₂CH₃)₂), 24.19 (CH₂-CH₂-CH₂-CH₂-CH₂-N), 26.20 $(CH_2-CH_2-$ CH₂-CH₂-CH₂-N), 26.62 (CH₂-C -N), 44.21((CH₂-CH₂-CH₂-CH₂-CH₂-CH₂-N), 44.73 (N(CH₂ CH₃)₂), 111.72 (Ar-C), 115.10 (Ar-C), 120.30 (Ar-C), 133.59 (Ar-C), 134.64 (=CH-Ar), 149.60 (S-C=CH), 168.14 (N-C=O), 179.47 (HOOC-), 193.33 (S=C-S)

4a/ 5-(4'-N,N-dibutylaminobenzylidene)-rhodanine-3acetic acid

m.p. 191–194 °C, yield 23.0%, MS $[M+1]^+$ —407.1, IR cm⁻³: 1716.3C=O, 1698.0C=O conj., 1636.3C=C exo., 1324.9 C–N, 1184.1C=S, ¹H NMR(400 MHz, CDCl₃), δ ppm, 7.72 (s, 1H,=C<u>H</u>-Ar), 7.40 (d, J = 8.96 Hz, 2H, Ar–H), 6,72 (d, J = 7.64 Hz, 2H, Ar–H), 4.95 (s, 2H, HOOC–C<u>H₂</u>–N), 3.37 (t, 4H, N(C<u>H₂CH₂CH₂CH₃)₂), 1.63 (q, 4H, N(CH₂C<u>H₂CH₂CH₂CH₃)₂), 1.39 (m, 4H, N</u></u>

4b/ 5-(4'-N,N-dibutylaminobenzylidene)-rhodanine-3propionic acid

m.p. 163-165 °C, yield 21.99%, MS [M+1]⁺-421.2, IR cm⁻³: 1727.9C=O, 1698.0C=O conj., 1610.3C=C exo., 1336.4 C–N, 1189.9C=S, ¹H NMR(400 MHz, CDCl₃), δ ppm, 7.68 (s, 1H,=CH-Ar), 7.38 (d, J=9.18 Hz, 2H, Ar-H), 6,69 (d, J = 8.80 Hz, 2H, Ar-H), 4.46 (t, 2H, HOOC-CH₂-CH₂-N), 3.37 (t, 4H, N(CH₂CH₂CH₂CH₃)₂), 2.86 (t, 2H, HOOC-CH2-CH2-N), 1.62 (q, 4H, N (CH₂CH₂CH₂CH₃)₂), 1.39 (q, 4H, N(CH₂CH₂CH₂CH₃)₂), 1.02 (t, 6H, N(CH₂CH₂CH₂CH₃)₂) ¹³C NMR (101 MHz, CDCl₃), δ ppm, 13.94 (N(CH₂CH₂CH₂CH₃)₂), 20.27 (N (CH₂CH₂CH₂CH₃)₂), 29.37 (N(CH₂CH₂CH₂CH₃)₂), 30.95 $(CH_2-CH_2-N),$ $(CH_2 - CH_2 - N),$ 39.45 50.87 (N)(CH₂CH₂CH₂CH₃)₂), 111.82 (Ar–C), 114.41 (Ar–C), 120.06 (Ar-C), 133.64 (Ar-C), 135.23 (=CH-Ar), 150.22 (S-C=CH), 167.77 (N-C=O), 175.49 (HOOC-), 192.96 (S=C-S)

4c/ 5-(4'-N,N-dibutylaminobenzylidene)-rhodanine-3butyric acid

m.p. 134–136 °C, yield 25.23%, MS $[M+1]^+$ –435.1, IR cm⁻³: 1710.5C=O, 1691.3C=O conj., 1637.3C=C exo., 1333.5 C-N, 1193.7C=S, ¹H NMR(400 MHz, CDCl₃), δ ppm, 11.00 (br. s, 1H, HOOC-) 7.66 (s, 1H, =CH-Ar), 7.38 (d, J = 8.97 Hz, 2H, Ar-H), 6.67 (d, J = 9.05 Hz, 2H, Ar-H), 4.21 (t, 2H, HOOC- $CH_2-CH_2-CH_2-N),$ 3.36 (t, 4H, N(CH₂CH₂CH₂ CH₃)₂), 2.45 (t, 2H, HOOC-CH2-CH2-CH2-N), 2.09 (t, 2H, HOOC-CH₂-CH₂-CH₂-N), 1.62 (q, 4H, N(CH₂CH₂CH₂CH₃)₂), 1.39 (m, 4H, N(CH₂CH₂CH₂CH₂) $(CH_3)_2$), $\overline{0.99}$ (t, 6H, $N(CH_2CH_2CH_3)_2$) ¹³C NMR (101 MHz, CDCl₃), δ ppm, 13.94 (N(CH₂CH₂CH₂) CH₃)₂), 20.27 (N(CH₂CH₂CH₂CH₃)₂), 22.27 (CH₂-CH2-CH2-N), 29.38 (N(CH2CH2CH2CH3)2), 31.21 (CH2-CH2-CH2-N), 43.43 (CH2-CH2-CH2-N), 50.85 (N (CH₂CH₂CH₂CH₃)₂), 111.78 (Ar–C), 114.66 (Ar–C), 120.13 (Ar-C), 133.58 (Ar-C), 134.93 (=CH-Ar), 150.15 (S-C=CH), 168.23 (N-C=O), 178.12 (HOOC-), 193.35 (S=C-S)

5a/ 55-(4'-N,N-diphenylaminobenzylidene)-rhodanine-3acetic acid

m.p. 240–242 °C, yield 64.27%, MS $[M+1]^+$ —447.1, IR cm⁻³: 1724.1C=O, 1706.7C=O conj., 1634.4C=C exo., 1329.7 C–N, 1192.7C=S, ¹H NMR(400 MHz, CDCl₃), δ ppm, 7.76 (s, 1H,=C<u>H</u>–Ar), 7.53 (d, J = 8.96 Hz, 2H, Ar–H), 7.44–7.40 (m. 6H Ar), 7.25–7.18 (m, 6H Ar), 4.73 (s, 2H, HOOC–C<u>H₂–N)</u> ¹³C NMR (101 MHz, CDCl₃), δ ppm, 43.46 (–CH₂–N), 39.87 (CH₂–CH₂–N), 117.67, 118.64, 119.67, 124.97, 125.85. 125.94, 125.66, 126.87, 129.05, 130.47, 130.50, 131.76, 133.32, 134.47, 166.92, (Ar–C), 145.98 (=CH–Ar), 150.65 (S–C=CH), 167.63 (N–C=O), 191.02 (HOOC–), 193.33 (S=C–S)

5b/ 5-(4'-N,N-diphenylaminobenzylidene)-rhodanine-3propionic acid

m.p. 212-215 °C, yield 76.09%, MS [M+1]⁺-461.1 IR cm⁻³: 1733.7C=O, 1706.7C=O conj., 1637.3C=C exo., $^{1}\mathrm{H}$ C–N, 1193.7C=S, NMR(400 MHz, 1338.4 CDCl₃+MeOD), δ ppm, 7.61 (s, 1H,=CH-Ar), 7.32-7.23 (m. 6H Ar), 7.14–7.12 (m, 6H Ar), 6,98 (d, J = 8.76 Hz, 2H, Ar-H), 4.37 (t, 2H, HOOC-CH₂-CH₂-N), 2.71 $(t, 2H, HOOC-CH_2-CH_2-N)$ ¹³C NMR (101 MHz, CDCl₃) +MeOD), δ ppm, 30.97 (-CH₂-CH₂-N), 39.87 (CH₂-CH2-N), 118.24, 120.30, 124.97, 125.22, 126.05, 129.66, 132.39, 133.77 (Ar-C), 146.04 (=CH-Ar), 150.46 (S-C=CH), 167.76 (N-C=O), 172.81 (HOOC-), 192.95 (S=C-S)

5c/ 5-(4'-N,N-diphenylaminobenzylidene)-rhodanine-3butyric acid

m.p. 137–140 °C, yield 75.33%, MS $[M+1]^+$ —475.1, IR cm⁻³: 1721.2C=O, 1700.9C=O conj., 1638.23C=C exo., 1330.6 C–N, 1193.7C=S, ¹H NMR(400 MHz, CDCl₃), δ ppm, 10.90 (br. s, <u>H</u>OOC–), 7.67 (s, 1H,= C<u>H</u>–Ar), 7.37–7.33 (m, 6H Ar), 7.19–7.16 (m, 6H Ar), 7.05 (d, *J*=8.80 Hz, 2H, Ar–H), 4.23 (t, 2H, HOOC–CH₂–CH₂–CH₂–N), 2.47 (t, 2H, HOOC–C<u>H</u>₂–CH₂–CH₂–CH₂–CH₂–N), 2.47 (t, 2H, HOOC–C<u>H</u>₂–CH₂–CH₂–CH₂–CH₂–N), 2.10 (q, 2H, HOOC–CH₂–C<u>H₂–CH₂–CH₂–CH₂–CH₂–CH₂–CH₂–N), 31.10 (<u>C</u>H₂–CH₂–CH₂–N), 43.50 (CH₂–CH₂–CH₂–N), 118.50, 120.43, 124.99, 125.41, 126.09, 129.72, 132.40, 133.58 (Ar–C), 146.14 (=<u>C</u>H–Ar), 150.40 (S–C=CH), 168.12 (N–<u>C</u>=O), 177.75 (HOO<u>C</u>–), 193.27 (S=C–S)</u>

Antibacterial activity assay in vitro

The 5-substituted derivatives of rhodanine-3-carboxyalkyl acids were screened for antibacterial and antifungal

activities by micro-dilution broth method using Mueller-Hinton broth and Mueller-Hinton broth with 5% lysed sheep blood for growth of non-fastidious and fastidious bacteria, respectively or Mueller-Hinton broth with 2% glucose for growth of fungi. Minimal inhibitory concentration (MIC) of the tested derivatives were evaluated for the panel of the reference microorganisms from American Type Culture Collection (ATCC), including Gram-negative bacteria (Escherichia coli ATCC 25922, Salmonella typhimurium ATCC14028, Klebsiella pneumoniae ATCC 13883, Pseudomonas aeruginosa ATCC 9027, Proteus mirabilis ATCC 12453), gram-positive bacteria (Staphylococcus aureus ATCC 25923, Staphylococcus aureus ATCC 6538, Staphylococcus epidermidis ATCC 12228, Micrococcus luteus ATCC 10240, Bacillus subtilis ATCC 6633, Bacillus cereus ATCC 10876, Streptococcus pyogenes ATCC 19615, Streptococcus pneumoniae ATCC 49619, Streptococcus mutans ATCC 25175), and fungi (Candida albicans ATCC 10231, Candida parapsilosis ATCC 22019).

derivatives The 5-substituted of rhodanine-3carboxyalkyl acids dissolved in dimethylosulfoxide (DMSO), were first diluted to the concentration (1000 µg/ mL) in an appropriate broth medium recommended for bacteria or yeasts. Then, using the same media, serial twofold dilutions were made in order to obtain final concentrations of the tested derivatives ranged from 0.98 to 1000 µg/mL. The sterile 96-well polystyrene microtitrate plates (Nunc, Denmark) were prepared by dispensing 200 µl of appropriate dilution of the tested derivatives in broth medium per well. The inocula were prepared with fresh microbial cultures in sterile 0.85% NaCl to match the turbidity of 0.5 McFarland standard and 2 µl were added to wells to obtain final density of 1.5×10^6 CFU/ml for bacteria and 5×10^4 CFU/ml for yeasts; CFU—colony forming units. After incubation (bacterial strains-35 °C for 24 h,

yeast strains—30 °C for 48 h), the MICs were assessed visually as the lowest concentration of the 5-substituted derivatives of rhodanine-3-carboxyalkyl acids showing complete growth inhibition of the reference microbial strains. Appropriate DMSO control (at a final concentration of 10%), a positive control (containing inoculum without the tested derivatives) and negative control (containing the tested derivatives without inoculum) were included on each microplate.

Minimal bactericidal concentration (MBC) or minimal fungicidal concentration (MFC) was determined by subculturing 100 μ l of the microbial culture from each well that showed through growth inhibition, from the last positive one and from the growth control onto the recommended agar plates. The plates were incubated at 35 °C for 24 h and the MBC/MFC was defined as the lowest concentration of the 5-substituted rhodanine-3-carboxyalkyl acids without growth of microorganisms. Ciprofloxacin and vancomycin were used as the standard drugs (Table 1). Each experiment was repeated in triplicate. Representative data is presented.

Results and discussion

Chemistry

All the resulting 3-carboxyalkanoic acid derivatives occurred as crystalline solids red in colour. They were characterised by high solubility in polar solvents (alcohols, glacial acetic acid).

The comparison of the condensation reaction yield of all three groups of the compounds /**3a-d**/, /**4a-c**/ and /**5a-c**/ indicated that 4-*N*,*N*-diphenylaminobenzoic aldehyde had the highest activity in condensation reactions among the aldehydes used. 4-dibutylaminobenzoic aldehyde was characterised by the lowest activity.

Microorganisms	Vancomicin			Ciprofloxacin				
	MIC (µg/ mL)	MBC (µg/ mL)	MBC/MIC ratio	MIC (µg/ mL)	MBC (µg/ mL)	MBC/MIC ratio		
S. aureus ATCC6538	0.49	1.95	4	0.24	0.24	1		
S. aureus ATCC25923	0.98	7.81	8	0.49	0.49	1		
S. epidermidis ATCC12228	0.98	0.98	1	0.49	0.49	1		
M. luteus ATCC10240	0.12	0.12	1	0.98	1.95	2		
B. subtilis ATCC6633	0.24	0.49	2	0.03	0.12	4		
B. cereus ATCC10876	0.98	15.6	16	0.12	0.12	1		
S. pyogenes ATCC19615	0.24	0.49	2	-	-	-		
<i>S. pneumoniae</i> ATCC49619	0.24	0.49	2	-	-	-		
S. mutans ATCC25175	0.98	0.98	1	_	-	-		

Table 1 MIC (µg/mL), MBC
(µg/mL) of vancomicin and
ciprofloxacin towards reference
Gram-positive bacterial strains

The characteristic bands deriving from the stretching C=O and C=S groups vibrations were present in the IR spectra of all the researched compounds. The C=O group vibrations ranged from 1727.9 to 1710.5 cm⁻¹, whereas C=S group vibrations ranged from 1195.7 to 1184.1 cm⁻¹.

The MS spectra were very simple. The highest intensity had always the $[M+1]^+$ ion peak. In most cases it reached 100%.

The ¹H NMR spectra contained a very characteristic signal deriving from the proton in =CH-Ar unit. It was a singlet, which was present in the 7.61–7.76 ppm range of chemical shifts. Position of the signal from a methine proton in this range showed that the condensation reaction carried out to Z isomers (Hardej et al. 2010). The ¹³C NMR spectra were characterised by the signal from the carbon atom bound with exocyclic sulphur atom. It was present in the 192.85–193.38 ppm range of chemical shifts.

Antibacterial activity

The antimicrobial activity of rhodanines has been known for over 50 years. The design and synthesis of antibacterial agents based on this heterocycle have been reported in numerous studies (Pardasani et al. 2001; Grant et al. 2000; Gandhe and Gautam 2004: Tomasic and Peterlin Masic 2012). The 5-ylidene-4-thiazolidinones and 4thiazolidinone-3-carboxylic acids are the most studied and promising 4-thiazolidinones in the context of creating new drug-like molecules (Lesyk and Zimenkovsky 2004; Lesyk et al. 2011). It was shown that introduction of substituents (mainly those containing a carboxyl group) in position N3 is the chemical path to the design of new compounds with a significant biological activity and decreased toxicity (Bhat et al. 2004). In present study, the antimicrobial assay of the novel 5-substituted derivatives of rhodanine-3-carboxyalkyl acids was carried out towards reference strains using a serial dilution method to obtain the MIC. None of the tested derivatives had activity against gram-negative bacteria (Escherichia coli ATCC25922, Salmonella typhimurium ATCC14028, Klebsiella pneumoniae ATCC13883, Pseudomonas aeruginosa ATCC9027, Proteus mirabilis ATCC12453), and yeasts (Candida albicans ATCC10231, Candida parapsilosis ATCC22019) (MIC > 1000 µg/mL, data not shown). Tables 2-4 summarised the results obtained for the MICs of the 10 target compounds (3a-d, 4a-c, 5a-c) to the gram-positive bacteria: staphylococi (Staphylococcus aureus ATCC25923, Staphylococcus aureus ATCC6538, Staphylococcus epidermidis ATCC12228); micrococci (Micrococcus luteus ATCC10240), bacilli (Bacillus subtilis ATCC6633, Bacillus cereus ATCC10876) and streptococci (Streptococcus pyogenes ATCC19615, Streptococcus pneumoniae ATCC49619, Streptococcus mutans ATCC25175). Ciprofloxacin and vancomycin were

Microorganisms	3a			3b			3с			3d		
	MIC (µg/ mL)	MBC (µg/ mL)	MBC/MIC ratio	MIC (μg/ mL)	MBC (µg/ mL)	MBC/MIC ratio	MIC (µg/ mL)	MBC (µg/ mL)	MBC/MIC ratio	MIC (µg/ mL)	MBC (µg/ mL)	MBC/MIC ratio
S. aureus ATCC6538	250	1000	4	31.25	>1000	>32	62.5	1000	16	31.25	>1000	>32
S. aureus ATCC25923	125	>1000	>8<	31.25	1000	32	31.25	>1000	>32	15.6	>1000	>64
S. epidermidis ATCC12228	125	1000	8	31.25	>1000	>32	31.25	1000	32	31.25	>1000	>32
M. luteus ATCC10240	125	>1000	>8	15.6	62.5	4	62.5	>1000	>16	15.6	>1000	>64
B. subtilis ATCC6633	62.5	250	4	15.6	>1000	>64	15.6	62.5	4	7.8	500	64
B. cereus ATCC10876	62.5	>1000	16	125	>1000	>4	15.6	>1000	>64	7.8	>1000	>128
S. pyogenes ATCC19615	250	>1000	>4	125	125	1	125	>1000	>4	250	500	2
S. pneunoniae ATCC49619	125	250	7	500	>1000	>2	125	500	4	250	500	2
S. mutans ATCC25175	250	>1000	>4	500	>1000	>2	500	>1000	>2	1000	>1000	PN

Microorganisms	4a			4b			4c		
	MIC (µg/ mL)	MBC (µg/ mL)	MBC/MIC ratio	MIC (µg/ mL)	MBC (µg/ mL)	MBC/MIC ratio	MIC (µg/ mL)	MBC (µg/ mL)	MBC/MIC ratio
S. aureus ATCC6538	7.8	15.6	2	3.9	>1000	>256	3.9	>1000	>256
S. aureus ATCC25923	3.9	62.5	16	3.9	>1000	>256	3.9	>1000	>256
S. epidermidis ATCC12228	3.9	62.5	16	3.9	7.8	2	3.9	>1000	>256
M. luteus ATCC10240	3.9	125	32	3.9	7.8	2	3.9	>1000	>256
B. subtilis ATCC6633	3.9	3.9	1	1.95	3.9	2	1.95	3.9	2
B. cereus ATCC10876	3.9	125	32	3.9	3.9	1	3.9	3.9	1
S. pyogenes ATCC19615	125	>1000	>8	125	500	4	500	>1000	Nd
S. pneumoniae ATCC49619	125	500	4	125	500	4	500	1000	2
S. mutans ATCC25175	>1000	>1000	Nd	1000	>1000	Nd	1000	>1000	Nd

Table 3 MIC (µg/mL), MBC (µg/mL) of 4a, 4b, and 4c derivatives towards reference gram-positive bacterial strains

Table 4 MIC (µg/mL), MBC (µg/mL) of 5a, 5b, and 5c derivatives towards reference gram-positive bacterial strains

Microorganisms	5a			5b			5c		
	MIC (µg/ mL)	MBC (µg/ mL)	MBC/MIC ratio	MIC (µg/ mL)	MBC (µg/ mL)	MBC/ MICratio	MIC (µg/ mL)	MBC (µg/ mL)	MBC/MIC ratio
S. aureus ATCC6538	1.95	7.8	4	1.95	31.5	16	15.6	>1000	>64
S. aureus ATCC25923	1.95	7.8	4	1.95	31.5	16	1.95	125	64
S. epidermidis ATCC12228	1.95	7.8	4	1.95	31.5	16	1.95	125	64
M. luteus ATCC10240	1.95	7.8	4	1.95	15.6	16	3.9	125	32
B. subtilis ATCC6633	1.95	3.9	2	1.95	1.95	1	1.95	1.95	1
B. cereus ATCC10876	1.95	7.8	4	1.95	15.6	8	3.9	62.5	16
S. pyogenes ATCC19615	125	1000	4	125	>1000	>8	125	>1000	>8
S. pneumoniae ATCC49619	250	500	2	125	500	4	125	500	4
S. mutans ATCC25175	500	>1000	Nd	>1000	>1000	Nd	>1000	>1000	Nd

used as positive controls (Table 1). Mild to moderate activity (MIC 125-1000 µg/mL) of the all synthesised derivatives was observed towards streptococci. The new rhodanine compounds showed different activity from moderate to very strong against other tested gram-positive bacteria, i.e., staphylococci, micrococci, and bacilli, depending on the strain and the synthesised compound. The first group of derivatives /3a-d/ was less active towards the tested gram-positive strains (MIC 15.6-250 µg/mL) as compared to the second group /4a-c/ of derivatives (MIC 1.95–7.8 μ g/mL) and the third group /5a–c/ of derivatives (MIC $1.95-15.6 \mu g/mL$). The most active compounds were /5a/ and /5b/ showing very strong bioactivity with MIC 1.95 µg/mL. The low values of MBC/MIC ratio (2-4) for /5a/ suggested its bactericidal power in contrast to higher values (8-16) for /5b/ indicating bacteriostatic activities except for bactericidal activity of /5b/ against B. subtilis ATCC 6633 (MBC/MIC 1). The remaining derivatives showed bactericidal (MBC/MIC \leq 4) or bacteriostatic activity against the tested bacteria (MBC/MIC > 4), depending on the strain and the rhodanine compound.

In the present study, most of synthesised compounds (4a-c and 5a-c) exhibited strong antibacterial activity amongst the tested gram-positive bacteria, although the mechanism of action is not yet clearly understood. However, rhodanines seem to be inhibitors of the bacterial enzyme MurB (Andres et al. 2000). The enzyme MurB, an NADPH dependant enolpyruvyl reductase, is responsible for the second committed step of bacterial peptidoglycan biosynthesis; it means that rhodanines could be expected to be bactericidal. Peptydoglycan is an essential component of the cell wall of both Gram-positive and Gram-negative bacteria and enzyme MurB is found in both of them. It would be expected that rhodanines might possess a broad

spectrum of antibacterial activity. The differences in biological activity of rhodanines to gram-positive and gram-negative bacteria could be explained by the differences in their cell wall structure and thus in the permeability. Peptydoglycan is major component (90%) of the gram-positive cell wall, whereas in Gram-negative bacteria, peptydoglycan, constituting 10% of cell wall, lies between cytoplasmic membrane and the outer lipid byliayer containing lipopolysaccharide, porins, adhesins which create additional barrier to cross by.

In this study, the preliminary remarks of the structure activity dependence can be noted. Comparison the MIC values determined for the newly synthesised rhodanine derivatives allowed to state that the basic factor increasing the activity to prevent bacteria growth is the size of the substituent at the C-5 position. The number of the carbon atoms present in the connector between carboxylic group and the 2,4-thiazolidinedione core is of much less importance. The influence of the connector length on the activity to suppress bacterial growth is noticeable when 5 atoms of carbon are present in the connector. The derivatives which have an acetic, propionic and butyric acid fragment at N-3 position and have the same substituent at C-5 position, demonstrated similar ability to suppress the growth of Gram-positive bacteria. In many cases the activity was identical.

Determining the MBC value allowed to establish the activity to kill bacteria or inhibit its growth. Antimicrobial agent are usually regarded as bactericidal if MBC value is higher no more than four times the MIC value (French 2006).

It was established that /5a/ and /4b/ 5-(4'-dibutylaminobenzylidene)-4-oxo-2-thioxo-3-thiazolidine acetic acid, as well as 5-(4'-diphenylaminobenzylidene)-4-oxo-2-thioxo-3thiazolidine acetic acid and 5-(4'-diphenylaminobenzylidene)-4-oxo-2-thioxo-3-thiazolidine propionic acid had antibacterial effect on the majority of the gram-positive bacteria strains.

It was surprising that increasing the number of carbon atoms in the connector resulted in decreased antibacterial activity of 5-(4'-diphenylaminobenzylidene)-4-oxo-2-thioxo-3-thiazolidine butyric acid.

Conclusions

A series of new rhodanine-3-carboxyalkyl acid derivatives possessing p-N,N-benzylidenedialkylamine moieties and p-N,N-bezylidenediphenylamine as a substituent at the C-5 were synthesised. The compounds were characterised by antibacterial activity on the tested Gram-positive strains of bacteria, however without biological activity on gramnegative bacteria and yeasts. It was observed that there was a dependency between the growth of the substituent size at C-5 position of the rhodanine ring and the antifungal activity growth. The derivatives having a p-N,N-bezylidenediphenylamine fragment at the C-5 position were characterised by the highest antibacterial activity. The increase of activity was probably caused by higher hydrophobicity of the aryl groups in comparison to the alkyl groups, which has been suggested by previous research (Miao et al. 2013; Patel et al. 2013). It was also established that the size of the connector between the carboxylic group and rhodanine ring had a limited influence on the antibacterial activity. The results have indicated the future direction of the research aiming at synthesis of the compounds characterised by higher antibacterial activity.

Compliance with Ethical Standards

Conflict of interest The authors declare that they have no competing interests.

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (http://creativecommons.org/licenses/by/4.0/), which permits use, duplication, adaptation, distribution, and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

References

- Alegaon SG, Alagawadi KR, Sonkusare PV, Chaudhary SM, Dadwe DH, Shah AS (2012) Novel imidazo[2,1-b][1,3,4]thiadiazole carrying rhodanine-3-acetic acid as potential antitubercular agents. Bioorg Med Chem Lett 22:1917–1921
- Andres CJ, Bronson JJ, D'Andrea SV, Deshpande MS, Falk PJ, Grant-Young KA, Harte WE, Ho HT, Misco PF, Robertson JG, Stock D, Sun Y, Walsh AW (2000) 4-Thiazolidinones: novel inhibitors of the bacterial enzyme MurB. Bioorg Med Chem Lett 10:715–717
- Bhat BA, Ponnala S, Sahu DP, Tiwari P, Tripathi BK, Srivastava AK (2004) Synthesis and antihyperglycemic activity profiles of novel thiazolidinedione derivatives. Bioorg Med Chem 12:5857–5864
- Bhatti RS, Shah S, Suresh Krishan P, Sandhu JS (2013) Recent pharmacological developments on rhodanines and 2,4-thiazolidinediones. Int J Med Chem. doi:10.1155/2013/793260
- Chandrappa S, Kavitha CV, Shahabuddin MS, Vinaya K, Kumar CSA, Ranganatha SR, Raghavan SC, Rangappa KS (2009) Synthesis of 2-(5-((5-(4-chlophenyl)furan-2-yl)methylene)-4oxo-2-thioxo-thiazolidin-3-yl)acetic acid derivatives and evaluation of their cytotoxicity and induction of apoptosis in human leukemia cells. Bioorg Med Chem 17:2576–2584
- Chen ZH, Zheng CJ, Sun LP, Piao HR (2010) Synthesis of new chalcone derivatives containing a rhodanine-3-acetic acid moiety with potential anti-bacterial activity. Eur J Med Chem 45:5739–5743
- French GL (2006) Bactericidal agents in the treatment of MRSA infections—the potential role of daptomycin. J Antimicrob Chemother 58:1107–1117

- Gandhe S, Gautam MD (2004) Synthesis and microbial activity of 5-(2-methyl-4-N-cyanoethyl-N-benzenesulphonyl aminobenzylidene)rhodanine and its metal complexes. Asian J Chem 16:261–264
- Grant EB, Guiadeen D, Baum EZ, Foleno BD, Jin H, Montenegro DA, Nelson EA, Bush K, Hlasta DJ (2000) The synthesis and SAR of rhodanines as novel class C β -lactamase inhibitor. Bioorg Med Chem Lett 10:2179–2182
- Hardej D, Ashby Jr CR, Khadtare NS, Kulkarni SS, Singh S, Talele TT (2010) The synthesis of phenylalanine –derived C5substituted rhodanines and their activity against selected methicillin-resistant *Staphylococus aureus* (MRSA) strains. Eur J Med Chem 45:5827–5832
- Jain KA, Vaidya A, Ravihandran V, Kashaw SK, Agrawal RK (2012) Recent developments and biological activities of thiazolidinone derivatives. A review. Bioorg Med Chem 20:3378–3395
- Kaminskyy DV (2015) Screening of the antiviral activity in the range of C5 and N3 substituted 4-thiazolidinone derivatives. J Org Pharmaceut Chem 13:64–69
- Kavitha CV, Basappa B, Swamy SN, Mantelingu K, Doreswamy S, Sridhar MA, Shashidhara Prasad J, Rangappa KS (2006) Synthesis of new bioactive venlafaxine analogs: novel thiazolidin-4-ones as antimicrobials. Bioorg Med Chem 14:2290–2299
- Körner H (1908) Über einige derivate der dithiocarbamino-essigsäure. Ber Dtsch Chem Ges 41:1901–1905
- Kumar G, Parasuraman P, Sharma SK, Banerjee T, Karmodiya K, Surolia N, Surolia A (2007) Discovery of a rhodanine class of compounds as inhibitor of *Plasmodium falciparum* enoyl-acyl carrier protein reductase. J Med Chem 50:2665–2675
- Lesyk RB, Zimenkovsky BS (2004) 4-Thiazolidones: Centenarian history, current status and perspectives for modern organic and medicinal chemistry. Curr Org Chem 8:1547–1577
- Lesyk RB, Zimenkovsky BS, Kaminsky DV, Kryshchyshyn AP, Havryluk DY, Atamanyuk DV, Subtelna IY, Khyluk DV (2011)

Thiazolidinone motif in anticancer drug discovery. Experience of DH LNMU medicinal chemistry scientific group.. Biopolym Cell 27:107–117

- Miao J, Zheng CJ, Sun LP, Song MX, Xu LL, Piao HR (2013) Synthesis and potential antibacterial activity of new rhodanine-3acetic acid derivatives. Med Chem Res 22:4125–4132
- Pardasani RT, Pardasani P, Sherry D, Chaturvedi V (2001) Synthetic and antibacterial studies of rhodanine derivatives with indol-2,3diones. Ind J Chem 40B:1275–1278
- Patel BA, Ashby Jr CR, Hardej D, Talele TT (2013) The synthesis and SAR study of phenylalanine-derived (Z)-5-arylmethylidene rhodanines as anti-methicillin-resistant *Staphylococcus aureus* (MRSA) compounds. Bioorg Med Chem Lett 23:5523–5527
- Rao BA, Rajpoot RS, Naidu VGM, Srinivas K, Ramakrishna S, Rao VJ (2011) Synthesis and *in vitro* antitumor activity of new nicotinyl-rhodanine derivatives. Int J Pharma Bio Sci 2:191–202
- Song MX, Zheng CJ, Deng XQ, Wei ZY, Piao HR (2014) The synthesis and anti-bacterial activities of N-carboxymethyl rhodanines. Med chem 4:441–448
- Tomasic T, Peterlin Masic L (2012) Rhodanine as a scaffold in drug discovery: a critical review of its biological activities and mechanisms of target modulation. Expert Opin Drug Discov 7:549–560
- Woodford N (2003) Review. Novel agents for the treatment of resistant gram-positive infections. Expert Opin Investig Drugs 12:117–137
- Xu LL, Zheng CJ, Sun LP, Miao J, Piao HR (2012) Synthesis of novel 1,3-diaryl pyrazole derivatives bearing rhodanine-3-fatty acid moieties as potential antibacterial agents. Eur J Med Chem 48:174–178
- Zvarec O, Polyak SW, Tieu W, Kuan K, Dai H, Pedersen DS, Morona R, Zhang L, Booker GW, Abell AD (2012) 5-Benzylidenerhodanine and 5-benzylidene-2,4-thiazolidinedione based antibacterials. Bioorg Med Chem Lett 22:2720–2722