1270-1273

Lewis Acid-Mediated Michaelis—Arbuzov Reaction at Room Temperature: A Facile Preparation of Arylmethyl/ Heteroarylmethyl Phosphonates

Ganesan Gobi Rajeshwaran, Meganathan Nandakumar, Radhakrishnan Sureshbabu, and Arasambattu K Mohanakrishnan*

Department of Organic Chemistry, School of Chemistry, University of Madras, Guindy Campus Chennai 600 025, Tamil Nadu, India

mohanakrishnan@unom.ac.in

Received December 9, 2010

ABSTRACT



A facile preparation of arylmethyl and heteroarylmethyl phosphonate esters was achieved involving a Lewis acid mediated Michaelis—Arbuzov reaction at room temperature. Interaction of arylmethyl halides/alcohols with triethyl phosphite in the presence of Lewis acid at room temperature afforded phosphonate esters in good yields.

Organic phosphonates are a key functional group in biological chemistry as well as synthetic organic chemistry.¹ In particular, arylmethyl and heteroarylmethyl phosphonate esters have been widely utilized as prominent precursor for the synthesis of optical materials such as poly(phenyl-vinylenes),² thienylvinylenes,³ trialkylamine tethered vinylenes,⁴ as well as stilbene dendrimers⁵ involving a

Horner–Wadsworth–Emmons reaction⁶ with aldehydes. The required organophosphonate esters have been traditionally obtained through a Michaelis–Arbuzov reaction.⁷

The Michaelis–Arbuzov reaction is initiated with an S_N2 reaction of nucleophilic trialkylphosphite with arylmethyl halides to give a phosphonium intermediate. The displaced halide anion upon a second S_N2 reaction with phosphonium intermediate led to the phosphonate ester along with an alkyl halide. Despite its acceptance, the Michaelis–Arbuzov reaction has two major drawbacks. First, the elevated temperature required for the activation of the reaction, limits the scope of the substrate suitable for the reaction. Second, the reaction generates one equiv of alkyl halide, which can react with the phosphite under reaction conditions to reduce yield and reaction efficiency. The Michaelis–Becker reaction⁸ utilizing dialkylphosphite salt of alkali metals with alkyl halides somewhat

^{(1) (}a) Engel, R.; Cohen, J. I. Synthesis of Carbon-phosphorous Bonds, 2nd ed.; CRC Press: Boca Raton, 2004. (b) Metcalf, W. W.; van der Donk, W. A. Annu. Rev. Biochem. 2009, 78, 65–94. (c) Bock, T.; Moehwald, H.; Muelhaupt, R. Macromol. Chem. Phys. 2007, 208, 1324–1340. (d) Quntar, A.; Baum, O.; Reich, R.; Srebnik, M. Arch. Pharm. 2004, 337, 76–80. (e) Abbas, S.; Bertram, R. D.; Hayes, C. J. Org. Lett. 2001, 3, 3365–3367. (f) Zmudzka, K.; Johansson, T.; Wojcik, M.; Janicka, M.; Nowak, M.; Stawinski, J.; Nawrot, B. New J. Chem. 2003, 27, 1698–1705.

^{(2) (}a) Zhen, L.; Urian, R. C.; Liu, Y.; Jen, A. K.-Y.; Pu, L. *Chem. Mater.* **2000**, *12*, 13–15. (b) Pu, Y.-J.; Soma, M.; Kido, J.; Nishide, H. *Chem. Mater.* **2001**, *13*, 3817–3819. (c) Liao, L.; Pang, Y.; Ding, L.; Karasz, F. E. *Macromolecules* **2002**, *35*, 3819–3824.

^{(3) (}a) Elandaloussi, E. H.; Frère, P.; Richomme, P.; Orduna, J.; Grain, J.; Roncali, J. *J. Am. Chem. Soc.* **1997**, *119*, 10774–10784. (b) Frère, P.; Raimundo, J.-M.; Blanchard, P.; Delaunay, J.; Richomme, P.; Sauvajol, J.-L.; Orduna, J.; Grain, J.; Roncali, J. *J. Org. Chem.* **2003**, *68*, 7254–7265.

^{(4) (}a) Bordeau, G.; Larita, R.; Metge, G.; Fiorini-Debuisschert, C.; Charra, F.; Teulade-Fichou, M.-P. *J. Am. Chem. Soc.* **2008**, *130*, 16836– 16837. (b) Shen, P.; Sang, G.; Lu, J.; Zhao, B.; Wan, M.; Zou, Y.; Li, y.; Tan, S. *Macromolecules* **2008**, *41*, 5716–5722.

⁽⁵⁾ Segura, J. L.; Gomez, R.; Martin, N.; Guldi, D. M. Org. Lett. 2001, 3, 2645–2648.

⁽⁶⁾ Wadsworth, W.; Emmons, W. D. J. Am. Chem. Soc. 1961, 83, 1733–1738.

^{(7) (}a) Michaelis, A.; Kaehne, R. *Chem. Ber.* **1898**, *31*, 1048–1058. (b) Arbuzov, B. A. *Pure Appl. Chem.* **1964**, *9*, 315–370. (c) Bhattacharya,

A. K.; Thyagarajan, G. *Chem. Rev.* **1981**, *81*, 415–430.

⁽⁸⁾ Michaelis, A.; Becker, T. Chem. Ber. 1897, 30, 1033.

overcomes the above-mentioned problems. However, the reaction conditions are often not convenient requiring either the use of strong anhyd base,⁹ or longer reaction time¹⁰ and in some cases resulted in low yield of the phosphonates.¹¹ Salvatore and co-workers achieved a mild and efficient synthesis of phosphonates via interaction of dialkylphosphonates with alkyl halides in the presence of Cs₂CO₃ and TBAI in dry DMF at room temperature.¹²

Recently, an efficient preparation of benzyl phosphonates was achieved involving the Pd(OAc)2-mediated cross-coupling reaction of benzyl halides with H-phosphonate diesters using Xantphos as a supporting ligand.¹³ Peyrottes et al. achieved the synthesis of sugar and nucleoside phosphonate derivatives involving microwave-assisted Michaelis-Arbuzov reaction.¹⁴ Marder and co-workers reported a convenient method for the synthesis of electron-rich benzyl phosphonates by treating the corresponding alcohols with triethyl phosphite in the presence of 1 equiv of I₂ at room temperature.¹⁵ This method proved to be useful for highly electron-rich heteroaryl methyl alcohols. However, in the case of less electron-rich systems, the method requires high temperature. Pei and co-workers achieved the synthesis of alkyl phosphonate esters via interaction of heteroarylmethyl alcohols with triethyl phosphite and I₂ in the presence of DBU at elevated temperature.¹⁶ Very recently, a roomtemperature alternative to the Michaelis-Arbuzov reaction was realized through reductive dehydrogenation of acvl phosphonates.¹⁷ Using this method, a one-pot sequence for the synthesis of alkyl phosphonates was developed from carboxylic acids. An efficient synthesis of α -aminophosphonates via interaction of N-benzyloxycarbonylamino sulfones with triethyl phosphite in the presence of a catalytic amount of InCl₃ in DCM at room temperature was achieved.¹⁸ In continuation of our studies on Lewis acid mediated domino reactions,¹⁹ we report herein the Lewis acid mediated Michaelis-Arbuzov reaction of benzylic bromides 1 with triethyl phosphite at room temperature (Scheme 1).

- (11) (a) Kosolapoff, G. M. J. Am. Chem. Soc. 1945, 67, 2259–2260.
 (b) Weizhen, Y.; Xiugao, L. Synthesis 1985, 10, 986–988.
- (12) Cohen, R. J.; Fox, D. L.; Eubank, J. F.; Salvatore, R. N. Tetrahedron Lett. 2003, 44, 8617–8621.
- (13) (a) Lavén, G.; Stawinski, J. Synlett **2009**, *2*, 225–228. (b) Laven, G.; Kalek, M.; Jezowska, M.; Stawinski, J. New. J. Chem. **2010**, *34*, 967–975.
- (14) Peyrottes, S.; Gallier, F.; Béjaud, J.; Périgaud, C. *Tetrahedron Lett.* **2006**, *47*, 7719–7721.
- (15) Zheng, S.; Barlow, S.; Parker, T. C.; Marder, S. R. *Tetrahedron Lett.* **2003**, *44*, 7989–7992.
- (16) (a) Wang, J.-L.; Tang, Z.-M.; Xiao, Q.; Ma, Y.; Pei, J. Org. Lett. 2008, 10, 4271–4274. (b) Wang, J.-L.; Xiao, Q.; Pei, J. Org. Lett. 2010, 12, 4164–4167.
- (17) Kedrowski, S. M. A.; Dougherty, D. A. Org. Lett. 2010, 12, 3990-3993.
- (18) Das, B.; Damodar, K.; Bhunia, N. J. Org. Chem. 2009, 74, 5607–5609.

Scheme 1. Michaelis-Arbuzov Reaction



Scheme 2. Lewis Acid-Mediated Michaelis-Arbuzov Reaction



In our ongoing program on the synthesis of carbazole alkaloids, we required various types of N-phenylsulfonyl-2/3-indolylmethylphosphonates as starting materials. The accessibility of these phosphonate esters using the traditional method at elevated temperature was problematic due to the unstable nature of the bromomethylindoles. Our recent success with the Lewis acid mediated domino reaction of bromomethylindoles prompted us to explore a Lewis acid mediated Michaelis-Arbuzov reaction of the same. Accordingly, when the reaction of bromomethylindole 3a was performed with triethyl phosphite in the presence of 10 mol % of InBr3 at room temperature for 4 h followed by workup led to the isolation of known phoshponate ester 4a in 80% yield (Scheme 2). Gratifyingly, the reaction could also be carried out at room temperature using less expensive ZnBr2 as a catalyst in almost comparable yield. This unexpected Lewis acidmediated Michaelis-Arbuzov reaction of the bromomethylindole 3a at room temperature prompted us to study the reaction with other benzylic halides.

To our delight, the Lewis acid-mediated Michaelis-Arbuzov reaction of a wide variety of aryl as well as heteroarylmethyl bromides could be successfully performed at room temperature to afford the corresponding phosphonates in good yields (Table 1). The 2-indolylmethanols **3b** and **3d** also underwent a smooth Lewis acid mediated Michaelis-Arbuzov reaction at room temperature in the presence of 1.1 equiv of $ZnBr_2$ (entry 1). Comparatively, 2-chloromethylindole **3a'/3c'** produced the respective phosphonate ester in diminished yield than the 2-bromomethylindole **3a/3c** (entry 1).

With the exception of 3-chloromethylindole 3l', in all other cases the reaction was found to be successful even in the presence of electron-withdrawing functional groups (entries 1–3). The reaction of bromomethyl ketone **30**

⁽⁹⁾ Reaction and Methods of Organic Compound Investigation; Kabachnik, M., Ed.; Goskhimizdat: Moscow, 1953; Vol. 13, p 427.

⁽¹⁰⁾ Kosolapoff, G. M. Organophosphorous Compounds; Wiley: New York, 1950; Chapter 7.

^{(19) (}a) Mohanakrishnan, A. K.; Dhayalan, V.; Arul Clement, J.; Balamurugan, R.; Sureshbabu, R.; Senthil Kumar., N. *Tetrahedron Lett.* **2008**, *49*, 5850–5854. (b) Dhayalan, V.; Arul Clement, J.; Jagan, R.; Mohanakrishnan, A. K. *Eur. J. Org. Chem.* **2009**, *4*, 531–546. (c) Sureshbabu, R.; Saravanan, V.; Dhayalan, V.; Mohanakrishnan, A. K. *Eur. J. Org. Chem.* **2011**, 922–935.



 Table 1. ZnBr₂-Mediated Michaelis-Arbuzov Reaction of Arylmethyl Halides/Alcohols and Aryl Aldehydes

^{*a*} The reaction was performed using bromo compound (1 mmol), $ZnBr_2$ (0.2 mmol), and P(OEt)₃ (1.2 mmol) in dichloromethane. ^{*b*} The reaction was performed using bromo compound (1 mmol), $ZnBr_2$ (0.4 mmol), and P(OEt)₃ (2.5 mmol) in dichloromethane. ^{*c*} The reaction was performed using benzylic alcohol/benzyl chloride (1 mmol), $ZnBr_2$ (1.1 mmol), and P(OEt)₃ (5 mmol). ^{*d*} The reaction was performed using aldehyde (1 mol), $ZnBr_2$ (1.1 mmol), and P(OEt)₃ (1.3 mmol) in DCE.

failed to produce the expected phosphonate ester 40 and instead led to the isolation of the respective debrominated methylindole (entry 5). The Michaelis-Arbuzov reaction was found to be successful with bis(bromomethyl) compounds as well (entries 4 and 6-9). The reaction of 1-bromomethyl-2-nitroarenes 5d-h with triethyl phosphite in the presence of 20 mol % of ZnBr₂ furnished the respective phosphonate esters 6d-h in excellent yields (entry 10). The ZnBr₂-mediated Michaelis-Arbuzov reaction of 1-chloromethyl-2-nitroarene 5d'/5e' as well as 1-hydroxymethyl-2-nitroarene 5d"/5e" was found to be unsuccessful (entry 10). Recently, a facile Lewis-acid mediated arylation of activated benzylic alcohols was achieved in good yields.²⁰ Hence, we propose to do a Lewis acid mediated Michaelis-Arbuzov reaction of benzylic alcohols. Accordingly, the interaction of electron-rich heteroarylmethyl alcohols with triethyl phosphite in the presence of 1.1 equiv of anhyd ZnBr₂ led to the formation of corresponding phosphonate esters in 75-93% yields (entries 11-15). The interaction of indole-3-carboxaldehyde 7 with triethyl phosphite in the presence of 1.1 equiv of ZnBr₂ at room temperature furnished diphosphonate ester 8 in 85% yield (entry 16). Under identical conditions, the reaction of any aldehydes 9a-d led to the isolation of respective α -hydroxyphosphonate esters 10a-d in 80-91% yields (entry 17). Gratifyingly, the interaction of benzhydrol 11 with triethyl phosphite in the presence of 1.1 equiv of ZnBr₂ at room temperature led to the isolation of known phosphonate ester 12 in 89% yield (entry 18). The reaction was found to be successful with secondary benzylic alcohols as well (entry 19). As expected, the reaction of ethyl bromoacetate 15a with triethyl phosphite in the presence of 20 mol % of ZnBr₂ at room temperature led to the isolation of triethyl phosphonoacetate 16 in 93% vield. However, under identical conditions, the reaction of ethyl chloroacetate 15b with triethyl phosphite in the presence of ZnBr₂ failed to produce the phosphonate ester 16 (entry 20). Finally, the Lewis acid mediated Michaelis-Arbuzov reaction was also found to be unsuccessful with *n*-hexyl bromide/*n*-butyl bromide (entry 21).

To verify the mechanism of the Lewis acid mediated Michaelis–Arbuzov reaction, (S)-(–)-1-phenylethanol 19/(R)-(+)-1-phenylbromoethane 21^{21} was reacted with

Scheme 3. Lewis Acid-Mediated Michaelis-Arbuzov Reaction of Chiral Benzylic Alcohol/Bromide



triethyl phosphite in the presence of $ZnBr_2$ (Scheme 3). Usual workup and purification followed by HPLC analysis (see the Supporting Information) indicated the formation of racemic phosphonate esters **20** and **22**. Hence, it may be concluded that the LA-mediated Michaelis-Arbuzov reaction of benzylic bromide/alcohol involves only an S_N1 -type mechanism.

In summary, for the first time, LA-mediated Michaelis–Arbuzov reaction of arylmethyl as well as heteroarylmethyl halides/alcohols was achieved at room temperature. The reaction has been successfully extended to the synthesis of α -hydroxyphosphonate esters as well. The 2-nitroaryl phosphonate esters reported herein may find application in the synthesis of alkaloids.²² The various types of heteroarylmethyl phosphonates reported herein may be regarded as crucial intermediates for the synthesis of conjugated vinylenes.²³

Acknowledgment. Financial assistance from the Department of Science and Technology (DST), New Delhi, is acknowledged. We thank DST-FIST for the high-resolution NMR facility. A.K.M. thanks Dr. V. Kesavan for HPLC data. G.G.R. thanks CSIR, New Delhi, for a SRF fellowship.

Supporting Information Available. Experimental procedure along with characterization data and ¹H and ¹³C NMR spectra of representative phosphonate esters. This material is available free of charge via the Internet at http://pubs.acs.org.

^{(20) (}a) Iovel, I.; Mertins, K.; Kischel, J.; Zapf, A.; Beller, M. Angew. Chem., Int. Ed. 2005, 44, 3913–3917. (b) Mertins, K.; Iovel, I.; Kischel, J.; Zapf, A.; Beller, M. Adv. Synth. Catal. 2006, 348, 691–695. (c) Rueping, M.; Nachtsheim, B. J.; Ieawsuwan, W. Adv. Synth. Catal. 2006, 348, 1033–1037.

⁽²¹⁾ Thakur, V. V.; Sudalai, A. Indian J. Chem., Sect. B 2005, 44, 557–562.

⁽²²⁾ Mohanakrishnan, A. K.; Srinivasan, P. C. J. Org. Chem. 1995, 60, 1939–1946.

^{(23) (}a) Mohanakrishnan, A. K.; Lakshmikantham, M. V.; McDougal, C.; Cava, M. P.; Baldwin, J. W.; Metzger, R. M. J. Org. Chem. **1998**, 63, 3105–3112. (b) Amaladass., P.; Arul Clement, J.; Mohanakrishnan., A. K. Eur. J. Org. Chem. **2008**, 3798–3810.