

View Article Online View Journal

ChemComm

Accepted Manuscript

This article can be cited before page numbers have been issued, to do this please use: H. Pan, T. W. Ng and Y. Zhao, *Chem. Commun.*, 2015, DOI: 10.1039/C5CC03399C.



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about *Accepted Manuscripts* in the **Information for Authors**.

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard <u>Terms & Conditions</u> and the <u>Ethical guidelines</u> still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.



www.rsc.org/chemcomm

Cite this: DOI: 10.1039/c0xx00000x



Iron-Catalyzed Amination of Alcohols Assisted by Lewis Acid

Hui-Jie Pan, Teng Wei Ng and Yu Zhao*

Received (in XXX, XXX) Xth XXXXXXXX 20XX, Accepted Xth XXXXXXXX 20XX DOI: 10.1039/b000000x

⁵ An efficient Lewis acid-assisted, iron-catalyzed amination of alcohols using borrowing hydrogen methodology was developed. In particular, silver fluoride was identified to be a highly effective additive to overcome the low efficiency in the amination of secondary alcohols catalyzed by the Knölker's complex.

Introduction

Owing to the widespread use of amines in pharmaceuticals, agrochemicals, additive and dyes, the development of efficient and environmentally benign methods to prepare amines has long been an important goal in organic synthesis.¹ Among the various strategies for amine synthesis, the amination of alcohols using a 15 borrowing hydrogen methodology² (or hydrogen autotransfer process) has been recognized as a highly atom economical choice (Scheme 1a). In this overall redox-neutral process, the alcohol substrate serves another role as the H₂ donor so no external reductant is needed, and water is generated as the only side 20 product. Various catalytic systems have been developed for this transformation, most of which use catalysts based on precious metals such as Ir,³ Ru⁴ and others.⁵ The development of efficient system using abundant, inexpensive catalytic and environmentally friendly metals (and especially iron) will be 25 highly desired.



Our group has been interested in the application of borrowing ³⁰ hydrogen methodology to chiral amine synthesis and we recently reported the first example of enantioselective amination of alcohols based on cooperative catalysis of a chiral iridium complex and a chiral phosphoric acid.^{7a} The extension of this

75

system to dynamic kinetic amination of α-branched alcohols was 35 also achieved.^{7b} In an effort to discover more efficient economical catalyst for this process, we were particularly attracted to the possibility of using the Knölker's complex 1 (Scheme 1b) for amination of alcohols. The use of 1 to hydrogenation and transfer hydrogenation of carbonyls is well 40 established,⁹ and the Beller group has reported highly efficient and enantioselective hydrogenation of imines catalyzed by 1 and a chiral phosphoric acid.¹⁰ Very recently, the Feringa group and the Wills group reported successful amination of alcohol. catalyzed by iron complex 2 (precursor to Knölker's complex) ⁴⁵ and the analogous complex **3**, respectively (Scheme 1b),¹¹ whic¹ prompted us to report our investigation on related studies as soon as possible. It is noteworthy that in their reports as well as othe early reports on Fe-catalyzed amination reactions through a borrowing hydrogen process, only primary alcohols (and selecte 50 symmetrical secondary alcohols) can be effectively transformed. to achiral amine products. Secondary alcohols, on the other side are in most cases unreactive (consistent with our own findings) In our studies, we have identified silver fluoride as an effective additive that enabled highly efficient amination of a wide range 55 of secondary alcohols catalyzed by 1.

The amination of 2-octanol with *p*-anisidine was chosen as the model reaction (Table 1). When the reaction was carried ov using 10 mol% 1, 6a was formed in only 4% yield after refluxing in toluene for 48 h (entry 1). In the crude reaction mixture, the 60 corresponding ketone could be observed in a substantial amount together with a small amount of imine (<5%), providin experimental support for the borrowing hydrogen process However, the overall efficiency was disappointing by the use of only 1 as the catalyst. In our previous system a chiral Brønsted 65 acid proved essential for facilitating the imine condensation a well as the asymmetric reduction step.^{7a} Inspired by this ¹⁰ y e observation as well as the report from the Beller group, screened various Brønsted acids for this reaction (entries 2 4). Unfortunately no significant improvement was observed in mc. 70 cases, with only formic acid leading to an improved but still poor yield of 29% (entry 2). The use of phosphoric acid proved ineffective (entry 4). Chiral phosphoric acids were als examined; no product formation was observed to ou. disappointment (results not shown).

[journal], [year], **[vol]**, 00–00

This journal is © The Royal Society of Chemistry [year]

Table 1 Screening of Brønsted and Lewis acid additives for the iron-catalyzed amination of secondary alcohol^a

							View Article Online	
	OH Me n-hexyl + MeO	NH ₂	10 mol% 1 additive 0 °C in toluene	HN Me 6a	P-acid: PhO _P O PhO ^P OH OC		.039/C5CC0335	990
Entry	Additive	Time (h)	Yield (%) ^b	Entry	Additive	Time (h)	Yield (%) ^b	Ċ
1	None	48	4	17	20 mol% Ni(OTf) ₂	48	36	
2	20 mol% HCOOH	48	29	18	20 mol% Ni(acac) ₂	48	29	
3	20 mol% TsOH	48	4	19	20 mol% Co(acac) ₂	48	4	
4	20 mol% P-acid	48	<2	20	20 mol% Co(acac) ₃	48	11	
5	20 mol% Cu(OTf) ₂	48	22	21	20 mol% Ti(OiPr) ₄	48	55	
6	20 mol% CuCl ₂	48	<2	22	20 mol% Ag ₂ O	48	6	\mathbf{n}
7	20 mol% CuCl	48	<2	23	20 mol% AgBF ₄	48	9	
8	20 mol% CuBr	48	<2	24	20 mol% AgOTf	48	23	\geq
9	20 mol% CuI	48	<2	25	20 mol% AgSbF ₆	48	39	
10	20 mol% FeCl ₂	48	<2	26	20 mol% Ag ₂ CO ₃	48	8	
11	20 mol% FeBr ₂	48	<2	27	20 mol% AgF	48	67	
12	20 mol% FeCl ₃	48	<2	28	5 mol% AgF	24	18	Ū
13	20 mol% FeBr ₃	48	<2	29	10 mol% AgF	24	46	-
14	20 mol% Fe ₂ O ₃	48	30	30	40 mol% AgF	24	97	C
15	20 mol% Fe(acac) ₃	48	38	31	20 mol% CsF	24	<2	
16	20 mol% NiCl ₂	48	<2	32°	40 mol% AgF	24	70	

s ^aUnless noted otherwise, the reaction was carried out with **5a** (0.2 mmol), **4a** (5 equiv), **1** (10 mol%) and the additive (20 mol%) in refluxing toluene. ^bYield of **6a** was determined by GC analysis using *n*-dodecane as the internal standard. ^c7 was used as the catalyst instead of **1**.

At this stage, it was decided that a wide range of Lewis acids should be examined. Various readily available salts of Cu, Fe, Ni, Co, Ti and Ag were then subjected to the reaction catalyzed by **1**. As summarized in entries 5-27, Table 1, no general trend could be observed, but the strongly Lewis acidic triflate and antimonate 15 salts all resulted in noticeable improvement (entries 5, 17, 24 and 25). Ti(OiPr)₄ and AgF, in particular, proved to be the most effective additives that yielded **6a** in 55% and 67% yield, respectively (entries 21 and 27). Further optimization with Ti(OiPr)₄ was not successful. To our excitement, the test of 20 loading of AgF proved fruitful (entries 28-30): an excellent 97%

23 Ioading of Agr proved Hundri (chines 20-50), an executent 77/6 yield of **6a** was obtained in 24 h when 40 mol% AgF was used. When other F- source such as CsF was used instead, no reaction was obtained (entry 31). Finally, complex **7**,¹² an air-stable precursor of **1**, was tested. Under otherwise identical conditions, a 25 slightly reduced yield of 70% was obtained (entry 32).

As Ag-catalyzed reactions of alcohols and amines via direct nucleophilic substitution have been documented in the literature,¹³ we tested the amination of **4a** using 40 mol% AgF in the absence of **1** (Scheme 2). No product formation was observed ³⁰ under these conditions, further supporting a borrowing hydrogen process catalyzed by **1** instead of Ag-catalyzed nucleophilic substitution mechanism.



35 Scheme 2 Test of AgF-catalyzed amination of alcohol

With the optimized conditions in hand, we moved on to explore the substrate scope. As shown in Scheme 3, a wide range of secondary alcohols underwent reaction to yield the ⁴⁰ corresponding amines in good to excellent yields. In the case of aliphatic alcohols, not only the linear ones such as **4a** and **4b**, bu those with α - and β -substituents could also be converted to the desired products **6c** and **6d** in good yields. Benzylic amines **6e** and **6f** could also be accessed in good yields. Again, no ⁴⁵ conversion to these products was obtained in the absence of AgF. In the Wills' report using **3** as the catalyst it was also noted that the reactions of aniline and secondary benzylic alcohols resulted in only low conversion to impure products. Beside 2-octanol, 3-

This journal is © The Royal Society of Chemistry [year]

^{2 |} Journal Name, [year], [vol], 00-00

octanol bearing an ethyl substituent was also a suitable substrate yielding **6g** in 82% yield, although a longer reaction time of 48 h was needed to achieve high conversion. It is noteworthy that the amination of symmetrical secondary alcohols such as ⁵ cyclohexanol and isopropanol using aniline were reported by the Feringa group; the yield of the corresponding products **6b** and **6h** were only 12-14%, further highlighting the beneficiary effect of AgF. Different anilines could also be used for the amination reaction (**6i-6l**). While aniline worked well to yield **6i** in excellent ¹⁰ yield, electron deficient p-Cl aniline proved to be much less efficient (**6j**). In addition to different anilines, the use of benzylamine such as **5b** for amination reaction was also

examined. Unfortunately, in this case the improvement from AgF was not as significant. Amine **6m** was produced in a 30% yield 15 that will require further optimization.



^aSee Table 1 for general conditions. ^bAll the yields refer to isolated yields of **6**. ^cIsopropyl alcohol as the solvent. ^d48 h ²⁰ reaction.

Scheme 3 Scope of amination of secondary alcohols^{a,b}

The amination of selected primary alcohols that were used in ²⁵ Feringa's and Wills' reports were also tested (Scheme 4). Under the optimized conditions as before, 1-octanol and benzyl alcohol underwent reaction with *p*-anisidine to yield a mixture of secondary amine and tertiary amine products. In this case, the higher reactivity of this system resulted in a partial double ³⁰ amination reaction to yield the tertiary amine side product. A

simple solution was to use close-to-stoichiometric loading of the alcohol. As an example, secondary amine **60** could be obtained in

This journal is © The Royal Society of Chemistry [year]

a high yield of 80% when 1.2 equiv. of **40** was used (Scheme 4b). Under similar conditions, alkene-containing substrate underwer amination smoothly to yield **6p** in excellent yield (Scheme 4c): The alcohols containing other functionalities such as all ester c an ether group, however, failed to yield any product, representing some limitation of the current catalytic system. The use of dio' arin previous work by Feringa worked out well (Scheme 4d) 40 Finally, secondary amine such as piperidine **5g** could underge reaction with **4n** to yield tertiary amine **8d** in a moderate yield of



45 Scheme 4 Amination of primary alcohols



Scheme 5 Proposed catalytic cycle

This reaction is believed to proceed through a borrowing hydrogen mechanism, consistent with previous reports. A controreaction using enantiopure alcohol yielded the amine product in racemic form. The exact nature of activation by AgF is not clea⁵⁵ at this point. Considering the general improvement of the product yield from various Lewis acids examined in Table 1, we believ that AgF serves as the Lewis acid to facilitate imine condensation and activate the imine intermediate towards reduction by the iron hydride species 1 in the catalytic cycle shown in Scheme 5. Computational studies will be carried out to help elucidate the details of this activation by AgF.

⁵ In conclusion, we report herein a highly efficient iron-catalyzed amination of primary as well as secondary alcohols assisted by Lewis acid. The development of an enantioselective variant of this reaction by examining chiral Lewis acids and chiral analogs of Knölker's complex is currently under investigation in our ¹⁰ laboratory.

Notes and references

Department of Chemistry, National University of Singapore, 3 Science Drive 3, 117543, Republic of Singapore. E-mail: zhaoyu@nus.edu.sg

- (a) A. Seayad, M. Ahmed, H. Klein, R. Jackstell, T. Gross, M. Beller, Science, 2002, 297, 1676-1678. (b) S. A. Lawerencem, Amine: Synthesis Properties and Applications, Cambridge University, Cambridge, 2004. (c) B. R. Brown, The Organic Chemistry of Apliphatic Nitrogen Compounds, Cambridge University, Cambridge, 2004.
- For selected reviews on borrowing hydrogen methodology, see: (a) M. H. S. A. Hamid, P. A. Slatford, J. M. J. Williams, *Adv. Synth. Catal.*, 2007, **349**, 1555–1575. (b) T. D. Nixon, M. K. Whittlesey, J.
 M. J. Williams, *Dalton Trans.*, 2009, 753–762. (c) G. E. Dobereiner, R. H. Crabtree, *Chem. Rev.*, 2010, **110**, 681–703. (d) G. Guillena, D. J. Ramón, M. Yus, *Chem. Rev.*, 2010, **110**, 1611–1641. (e) A. J. A.
 Watson, J. M. J. Williams, *Science*, 2010, **329**, 635–636. (f) S. Bähn, S. Imm, L. Neubert, M. Zhang, H. Neumann, M. Beller, *ChemCatChem*, 2011, **3**, 1853–1864. (g) S. Pan, T. Shibata, *ACS Catal.*, 2013, **3**, 704–712. (h) Q. Yang, Q. Wang, Z. Yu, *Chem. Soc. Rev.*, 2015, **44**, 2305-2329.
- 3 For selected examples of the amination of alcohols under Ir catalysis, see: (a) K. Fujita, K. Yamamoto, R. Yamaguchi, Org. Lett., 2002, 4, 2691-2694. (b) K. Fujita, Z. Li, N. Ozeki, R. Yamaguchi, 35 Tetrahedron Lett., 2003, 44, 2687-2690. (c) K. Fujita, T. Fujii, R. Yamaguchi, Org. Lett., 2004, 6, 3525-3528. (d) K. Fujita, Y. Enoki, R. Yamaguchi, Tetrahedron, 2008, 64, 1943-1954. (e) R. Kawahara, K. Fujita, R. Yamaguchi, J. Am. Chem. Soc., 2010, 132, 15108-15111. (f) R. Kawahara, K. Fujita, R. Yamaguchi, Adv. Synth. Catal., 2011, 353, 1161-1168. (g) A. Prades, R. Corberan, M. Poyatos, E. Peris, Chem. Eur. J., 2008, 14, 11474-11479. (h) B. Blank, M. Madalska, R. Kempe, Adv. Synth. Catal., 2008, 350, 749-758. (i) B. Blank, S. Michlik, R. Kempe, Adv. Synth. Catal., 2009, 351, 2903-2911. (j) B. Blank, S. Michlik, R. Kempe, Chem. Eur. J., 2009, 15, 45 3790-3799. (k) S. Michlik, R. Kempe, Chem. Eur. J., 2010, 16,
- 13193–13198. (I) S. Michlik, T. Hille, R. Kempe, *Adv. Synth. Catal.*, 2012, **354**, 847–862. (m) D. Gnanamgari, E. L. O. Sauer, N. D. Schley, C. Butler, C. D. Incarvito, R. H. Crabtree, *Organometallics*, 2009, **28**, 321–325. (n) O. Saidi, A. J. Blacker, M. M. Farah, S. P. Marsden, J. M. J. Williams, *Chem. Commun.*, 2010, **46**, 1541–1543.
- (o) I. Cumpstey, S. Agrawal, E. K. Borbas, B. Martin-Matute, *Chem. Commun.*, 2011, **47**, 7827–7829. (p) S. Agrawal, M. Lenormand, B. Martin-Matute, *Org. Lett.*, 2012, **14**, 1456–1459. (q) A. Bartoszewicz, R. Marcos, S. Sahoo, A. K. Inge, X. Zou, B. Martin-
- Matute, *Chem. Eur. J.*, 2012, **18**, 14510–14519. (r) J.-Q. Li, P. G. Andersson, *Chem. Commun.*, 2013, **49**, 6131–6133.
- For selected examples of the amination of alcohols under Ru catalysis, see: (a) D. Hollmann, A. Tillack, D. Michalik, R. Jackstell,
 M. Beller, *Chem. Asian J.*, 2007, 2, 403–410. (b) A. Tillack, D.
- Hollmann, K. Mevius, D. Michalik, S. Bahn, M. Beller, *Eur. J. Org. Chem.*, 2008, 4745–4750. (c) M. Zhang, S. Imm, S. Bahn, H. Neumann, M. Beller, *Angew. Chem. Int. Ed.*, 2011, **50**, 11197–11201. (d) M. H. S. A. Hamid, J. M. J. Williams, *Chem. Commun.*, 2007, 725–727. (e) M. H. S. A. Hamid, J. M. J. Williams,
- Zoori, 120 (2017), 48, 8263–8265. (f) M. H. S. A. Hamid, C. L. Allen, G.W. Lamb, A. C. Maxwell, H. C. Maytum, A. J. A. Watson, J. M. J. Williams, J. Am. Chem. Soc., 2009, 131, 1766–1774. (g) C. Gunanathan, D. Milstein, Angew. Chem. Int. Ed., 2008, 47, 8661–
- 4 | Journal Name, [year], [vol], 00-00

- 8664. (h) K. Yamaguchi, J. He, T. Oishi, N. Mizuno, *Chem. Eur. J.*, 2010, **16**, 7199–7207. (i) R. N. Monrada, R. Madsen, *Org. Biomol. Chem.*, 2011, **9**, 610–615. View Article Online
- 5 For selected examples of Cu-catalyzed amination of aloobadis, See (a) 90 F. Shi, M. K. Tse, X. Cui, D. Gordes, D. Michalik, K. Thurow, Y.
- Deng, M. Beller, Angew. Chem. Int. Ed., 2009, 48, 5912–5915. (b) A. Martinez-Asencio, D. J. Ramon, M. Yus, Tetrahedron Lett., 2010, 51, 325–327. For Os-catalyzed amination, see: (c) M. Bertoli, A. Choualeb, A. J. Lough, B. Moore, D. Spasyuk, D. G. Gusev, Organometallics, 2011, 30, 3479–3482. For a Fe-catalyzed N-alkylation of amines with alcohols not through a borrowing hydrogen process, see: (d) Y. Zhao, S. W. Food, S. Saito, Angew. Chem. Int. Ed., 2011, 50, 3006–3009. (e) U. Jana, S. Maiti, S. Biswas, Tetrahedron Lett., 2008, 49, 858-862.
- For general reviews and highlights on iron catalysis, see: (a) K.
 Gopalaiah, *Chem. Rev.*, 2013, **113**, 3248–3296. (b) M. Darwish, M.
 Wills, *Catal. Sci. Technol.*, 2012, **2**, 243–255. (c) S. Enthaler, K.
 Junge, M. Beller, *Angew. Chem. Int. Ed.*, 2008, **47**, 3317–3321. (d)
 C. Bolm, J. Legros, J. Le Paih, L. Zani, *Chem. Rev.*, 2004, **104**, 6217–6254.
- 90 7 (a) Y. Zhang, C.-S. Lim, D. S. B. Sim, H.-J. Pan, Y. Zhao, Angew. Chem. Int. Ed., 2014, 53, 1399–1403. (b) Z.-Q. Rong, Y. Zhang, R. H. B. Chua, H.-J. Pan, J. Am. Chem. Soc., 2015, 137, 4944-4947. For other related examples on the use of borrowing hydroge methodology in asymmetric catalysis, see: (c) N. J. Oldenhuis, V. M. Dong, Z. Guan, J. Am. Chem. Soc., 2014, 136, 12548–12551. (d) F. J. Shermer, P. A. Slatford, D. D. Edney, J. M. J. Williams, *Tetrahedron: Asymmetry*, 2007, 18, 2845–2848. (e) A. E. Putra, Y. Oe, T. Ohta, *Eur. J. Org. Chem.*, 2013, 6146–6151. (f) A. Quintard T. Constantieux, J. Rodriguez, Angew. Chem. Int. Ed., 2013, 52, 12883–12887.
- 8 (a) H.-J. Knölker, E. Baum, H. Goesmann, R. Klauss, *Angew. Chem. Int. Ed.*, 1999, 38, 2064–2066. For a highlight of the chemistry using 1, see: (b) A. Chakraborty, H. Guan, *e-Encyclopedia of Reagents for Organic Synthesis*, John Wiley & Sons Ltd., 2014, DOI. 10.1002/047084289X.rn01666.
- 9 (a) C. P. Casey, H. Guan, J. Am. Chem. Soc., 2007, 129, 5816–5817.
 (b) M. G. Coleman, A. N. Brown, B. A. Bolton, H. Guan, Adv. Synth. Catal., 2010, 352, 967–970.
- 10 S. Zhou, S. Fleischer, K. Junge, M. Beller, *Angew. Chem. Int. Ed.*, 2011, **50**, 5120–5124.
 - (a) T. Yan, B. L. Feringa, K. Barta, *Nat. Commun.*, 2014, publish online 26 November 2014. DOI 10.1038/ncomms6602. (b) A. J. Rawlings, L. J. Diorazio, M. Wills, *Org. Lett.*, 2015, **17**, 1086–1089. For earlier examples: (c) X. Cui, F. Shi, Y. Zhang, Y. Deng, *Tetrahedron Lett.*, 2010, **51**, 2048–2051. (d) M. Bala, P. K. Verma, U. Sharma, N. Kumar, B. Singh, *Green Chem.*, 2013, **15**, 1687–1693.
 - 12 T. N. Plank, J. L. Drake, D. K. Kim, T. W. Funk, *Adv. Synth. Catal* 2012, **354**, 597-601.
- 13 (a) B. Sreedhar, P. S. Reddy, M. A. Reddy, B. Neelima, R. Arundhathi, *Tetrahedron Lett.*, 2007, **48**, 8174–8177. (b) S. R. Mothe, P. Kothandaraman, S. J. L. Lauw, S. M. W. Chin, P. W. H. Chan, *Chem. Eur. J.*, 2012, **18**, 6133–6137. (c) D. Susanti, F. Koh, J. A. Kusuma, P. Kothandaraman, P. W. H. Chan, *J. Org. Chem.*, 2012, 77, 7166–7175.