

A Highly Active Phosphine–Borane Organocatalyst for the Reduction of CO₂ to Methanol Using Hydroboranes

Marc-André Courtemanche,[†] Marc-André Légaré,[†] Laurent Maron,^{*,‡} and Frédéric-Georges Fontaine^{*,†}

[†]Département de Chimie, Centre de Catalyse et de Chimie Verte (C3V), Université Laval, Québec G1V 0A6, Canada

[‡]Université de Toulouse, INSA, UPS, LCPNO, CNRS, UMR 5215 CNRS-UPS-INSA, 135 avenue de Rangueil, Toulouse, France

Supporting Information

ABSTRACT: In this work, we report that organocatalyst 1-Bcat-2-PPh₂-C₆H₄ ((1); cat = catechol) acts as an ambiphilic metal-free system for the reduction of carbon dioxide in presence of hydroboranes (HBR₂ = HBcat (catecholborane), HBpin (pinacolborane), 9-BBN (9-borabicyclo[3.3.1]nonane), BH₃·SMe₂ and BH₃·THF) to generate CH₃OBR₂ or (CH₃OBO)₃, products that can be readily hydrolyzed to methanol. The yields can be as high as 99% with exclusive formation of CH₃OBR₂ or (CH₃OBO)₃ with TON (turnover numbers) and TOF (turnover frequencies) reaching >2950 and 853 h⁻¹, respectively. Furthermore, the catalyst exhibits “living” behavior: once the first loading is consumed, it resumes its activity on adding another loading of reagents.

It is widely known that carbon dioxide is a green-house gas and one of the most important contributors to global warming, and several political initiatives have been put forward to reduce carbon dioxide emissions.¹ Most of the current systems known to catalyze the reduction of CO₂ into valuable products use transition metals,² including notably the reverse water–gas shift reaction to generate carbon monoxide which in turn can be transformed into several useful chemicals.³ Recently, some homogeneous organometallic systems have shown promise in generating valuable chemicals.⁴ The most active systems to date for the reduction of CO₂ into high hydrogen content molecules include a ruthenium phosphine complex^{4c} and a nickel pincer complex,^{4d} using respectively H₂ and HBcat (HBcat = catecholborane), to generate MeOH from CO₂, and an iridium catalyst that can reduce CO₂ into methane using hydrosilanes as hydrogen source with turnover numbers (TON) up to 8300.^{4e}


Recently, a variety of transition metal-free systems have emerged for carbon dioxide activation and functionalization. Indeed, it has recently been shown that Lewis acidic Et₂Al⁺ species can catalytically reduce carbon dioxide to methane.⁵ Similarly, silyl cations can catalytically reduce CO₂ to a mixture of benzoic acid, formic acid, and methanol.⁶ However, both systems greatly lack selectivity and generate undesirable alkylation byproducts. An avenue of interest for carbon dioxide activation is the use of “frustrated Lewis pairs” (FLP), work pioneered by Stephan and Erker.⁷ Since this initial discovery, many ambiphilic systems have been shown to be active in the stoichiometric fixation CO₂.⁸ Piers demonstrated elegant use of this concept for the catalytic reduction of CO₂ into methane

using the robust TMP/B(C₆F₅)₃ (TMP = 2,2,6,6-tetramethylpiperidine) system with Et₃SiH, albeit with limited turnovers.⁹ It has been shown that the FLP system consisting of PMes₃/AlX₃, (Mes = mesityl, X = Cl, Br) not only binds CO₂ but also reduces it to methanol using BH₃·NH₃ as hydrogen source.¹⁰ Alternatively, O’Hare and Ashley demonstrated that CO₂ could be hydrogenated using TMP/B(C₆F₅)₃.¹¹ Unfortunately, the last two systems require stoichiometric amounts of FLP. Although interesting in concept, none of the FLP or ambiphilic systems reported to date demonstrate good catalytic activity for carbon dioxide reduction. The only efficient organocatalytic system reported to date for the reduction of CO₂ into methanol use highly Lewis basic N-heterocyclic carbene catalysts and diphenylsilane as hydrogen source with turnover frequencies (TOF) of 25 h⁻¹ at 25 °C.¹²

Our research program targets ambiphilic systems with little “frustrated” character and/or weak Lewis acidity and basicity.¹³ One ambiphilic system of interest is that of aryl bridged phosphine–boranes extensively studied by Bourissou and collaborators.¹⁴ These molecules have been shown to be quite robust, stable, and easy to synthesize. More recently, they have been used in the activation of singlet oxygen¹⁵ and as organocatalysts for the Michael addition reaction,¹⁶ but to our knowledge the activity of these molecules for carbon dioxide reduction has not been investigated. Here we report that the 1-Bcat-2-PPh₂-C₆H₄ ambiphilic system is one of the most active catalysts for the selective catalytic reduction of carbon dioxide to methanol.

Although several ambiphilic phosphine–boranes were prepared by Bourissou,^{14,15} the synthesis of the catecholborane derivative 1-Bcat-2-PPh₂-C₆H₄ (1, Scheme 1) was never reported. The air-stable product is easily synthesized in 80% yield from previously reported o-lithiated triphenylphosphine using a known synthetic pathway (Figure S1, Supporting

Scheme 1. Reduction of CO₂ in Presence of HBcat and catalyst 1



Received: May 7, 2013

Information [SI]).¹⁷ Multinuclear NMR characterization of species **1** demonstrates this molecule to be monomeric in solution having no observable P–B interaction. The $^{31}\text{P}\{^1\text{H}\}$ and $^{11}\text{B}\{^1\text{H}\}$ NMR chemical shifts are respectively of -4.57 and 33.1 ppm. The solid-state structure (see SI Figure S24) does not show any evidence of P–B interaction, the latter distance being quite long (3.28 Å).

Unsurprisingly, exposing **1** to 1 atm of CO_2 at room temperature resulted in no spectroscopic change in solution (^1H , ^{31}P , and ^{11}B NMR spectroscopy). Although no adduct was observed between CO_2 and **1**, the addition of 100 equiv of HBcat to a 9 mM solution of **1** in benzene- d_6 in a J-Young NMR tube under one atmosphere of CO_2 resulted in the formation of a white precipitate after 24 h. This was characterized as catBOBcat on the basis of spectroscopic comparison with the independently synthesized product (see Scheme 1). Monitoring of the solution using ^1H NMR spectroscopy showed the presence of a single new peak at 3.37 ppm attributed to CH_3OBcat by comparison to the independently synthesized product. Hydrolysis of the latter product produces methanol, which was confirmed using GC-FID. As expected, carrying out the same reaction under an atmosphere of $^{13}\text{C}\text{O}_2$ shows the formation of $^{13}\text{CH}_3\text{OBcat}$ with the expected $^1J_{\text{C-H}}$ of 145 Hz.^{4d}

Monitoring the reduction of CO_2 in presence **1** and 100 equiv of HBcat using ^1H and $^{31}\text{P}\{^1\text{H}\}$ NMR spectroscopy showed an induction period of 30 min where no spectroscopic change was observed in the solution. However, after the induction period the reaction started readily, and after 2 h a 34% yield (TON = 34, TOF = 17 h^{-1}) of CH_3OBcat was obtained (Figure 1, \blacklozenge). The rate of the reaction diminished as




Figure 1. Turnover numbers (TON) for the formation of CH_3OBcat from a 9 mM solution of **1** in benzene- d_6 in the presence of 100 equiv of HBcat under one atmosphere of CO_2 . The TONs are based on the number of hydrogen atoms transferred to CO_2 . Reactions were carried out at (\blacklozenge) 23 °C and (\blacksquare) 70 °C.

the reaction progressed, suggesting that conversion is dependent on the concentration of HBcat in solution. Indeed, 50% conversion to CH_3OBcat was obtained in less than 5 h and a yield of 69% of CH_3OBcat was observed after a period of 24 h. The reduction of CO_2 also proceeded in the presence of 100 equiv of $\text{BH}_3\cdot\text{SMe}_2$ to generate $(\text{CH}_3\text{OBO})_3$ but a longer induction period was observed (>2 h; Figure 2, \blacklozenge). Nevertheless, the conversion to the methoxyborane species is rapid once catalysis starts, obtaining respectively 108 and 200 TON at 2 and 5 h after the induction period (respective TOFs of 54 and 40 h^{-1}). After a period of 14 h, a TON of 257 was




Figure 2. Turnover numbers (TON) for the formation of $(\text{CH}_3\text{OBO})_3$ from a 9 mM solution of **1** in benzene- d_6 in the presence of 100 equiv of $\text{BH}_3\cdot\text{SMe}_2$ under one atmosphere of CO_2 . The TONs are based on the number of hydrogen atoms transferred to CO_2 . Reactions were carried out at (\blacklozenge) 23 °C and (\blacksquare) 70 °C.

obtained. The TON numbers being greater than 100 suggests that all hydrogen atoms from $\text{BH}_3\cdot\text{SMe}_2$ are available for the reduction of CO_2 . To our knowledge, it represents the first time that BH_3 is used as a hydrogen source for the catalytic reduction of CO_2 to methanol. It is interesting that the catalyst remains active even if BH_3 is known to coordinate phosphine moieties, which could inhibit catalysis; it is thus logical to presume that the longer induction period is caused by a competition between BH_3 and CO_2 for coordination to the catalyst. BH_3 is of great interest since it has the highest hydrogen content of any hydroborane.

A factor that dramatically increased the efficiency of the catalytic system was temperature. Heating a solution of **1** with 100 equiv of HBcat to 70 °C under one atmosphere of CO_2 generated CH_3OBcat without any observable induction period (Figure 1, \blacksquare). After 36 min, a TON of 86 was observed (TOF = 143 h^{-1}), which increased to 92 after a period of 98 min (Table 1, entries 1–2). After letting the solution rest for a 24-h period, another loading of 100 equiv of HBcat was added and the solution reheated to 70 °C. The catalytic reaction resumed, but with a rate that seemed somewhat slower (an overall TON of 136 after 30 min), possibly due to the presence of a large quantity of precipitate in the solution (catBOBcat) that reduced the homogeneity of the solution. However, 60 min after the addition of the second loading a TON of 185 was measured (Table 1, entries 3–4), a yield similar to that observed in the first run. Such behavior is reminiscent of a durable and “living” catalyst. Under similar conditions, $\text{BH}_3\cdot\text{SMe}_2$ proved to be an excellent hydrogen source, generating 90% yield of $(\text{CH}_3\text{OBO})_n$ in 67 min (TON = 271). A TON of 211 was obtained after only 13 min, representing a TOF of 973 h^{-1} (Figure 2, \blacksquare). The latter result is remarkable since the highest TOF reported for the reduction of CO_2 to a methanol derivative is 495 h^{-1} by a homogeneous nickel catalyst using HBcat as an hydrogen source.^{4d} The reaction was also carried out using other hydroborane sources. The addition of 100 equiv of HBpin to a solution of **1** under one atmosphere of CO_2 at 70 °C generated 60% yield of the desired product in a 3-h period (Table 1, entry 6). The significantly lower activity of the latter borane compared to HBcat is not surprising since it is known that HBpin is less reactive for the hydroboration reaction.¹⁸ Similarly, 9-BBN only showed 34 TON in a 3-h period (entry 7).

Table 1. Reduction of CO₂ with Various Hydroboranes^a

entry	borane	equiv	time (min)	TON ^b	TOF (h ⁻¹)
1	HBcat	100	36	86	143
2	HBcat	100	98	92	56
3	HBcat	100 + 100 ^c	30	136	72
4	HBcat	100 + 100 ^c	60	185	85
5	BH ₃ ·SMe ₂	100	67	271	242
6	HBpin	100	174	60	21
7	9-BBN	50 ^d	174	34	12
8 ^e	HBcat	300	60	145 ^g	145
9 ^f	HBcat	1000	240	664 ^g	166
10 ^e	BH ₃ ·SMe ₂	300	60	853 ^g	853
11 ^f	BH ₃ ·SMe ₂	1000	240	>2,950 ^g	>737
12 ^e	BH ₃ ·THF	300	60	340 ^g	340

^aReaction conditions: Unless noted otherwise, 2.0 mg (0.0053 mmol) of **1** in 0.6 mL of benzene-*d*₆ at 70 °C. ^bBased on mole of B–H consumed per mole of **1**, determined by ¹H NMR integration using hexamethylbenzene as internal standard for entries 1–7, and determined by GC-FID with ¹PrOH as a standard for entries 8–12. ^cA second addition of 100 equiv of HBcat was added 24 h after the first addition. ^dLimited at 50 equiv because of low solubility of 9-BBN. ^e2.0 mg (0.0053 mmol) of **1** in 3 mL of benzene at 70 °C under ~2 atm of CO₂. ^f2.0 mg (0.0053 mmol) of **1** in 9 mL of benzene at 70 °C under ~2 atm of CO₂. ^gQuenched with excess H₂O and analyzed by GC-FID with ¹PrOH as a standard.

Since diffusion problems could limit the rate of the reaction when carried out in NMR conditions, catalytic tests were carried out on a larger scale using Fisher-Porter bottles under ~2 atm of CO₂. The products obtained were hydrolyzed to methanol, and the turnover numbers were calculated on the basis of the concentration of methanol using gas chromatography with a flame ionization detector. As can be observed in Table 1, the activities observed at the NMR scale can be reproduced at larger scale and lower catalyst loading. Indeed, the reduction of CO₂ using 300 equivalents of HBcat and BH₃·SMe₂ gave in one hour methanol in 48% and 95% yield, giving respectively TOF of 145 and 853 h⁻¹ (Table 1, entries 8 and 10). It is notable that the TOF observed under large loading of hydroboranes are consistent with those observed at the NMR scale at low conversion that were 143 and 973 h⁻¹ for HBcat and BH₃·SMe₂, respectively. Catalysis using a 0.1% catalyst loading (1000 equiv of substrate) in a 4-h period also gave impressive results. In the presence of HBcat, a TON of 664 was observed, which indicates that the rate of reaction remains the same during the 4-h period even with a lower catalyst loading and a lower catalyst concentration (Table 1, entry 9, TOF = 166 h⁻¹). In the presence of BH₃·SMe₂, all of the substrate was consumed since the conversion to methanol was quantitative (Table 1, entry 11), once more suggesting that the TOF observed after one hour is conserved over a longer reaction time. The reaction works also with BH₃·THF, albeit less efficiently (Table 1, entry 12, TOF = 340 h⁻¹).

Density functional theory studies at the B3PW91 6-31G** level of theory were performed to obtain further insight at the mechanistic pathway, using HBcat as the hydrogen source. It should be noted that only potential intermediates were considered in the following, and the results are summarized in Figure 3. Also, we did not account for the fact that HBcat is known to degrade in presence of Lewis bases since control experiments have shown this process to be marginal in our system.¹⁹ As observed experimentally, the coordination of CO₂ to **1** to generate intermediate **IM1** is disfavored by 9.9




Figure 3. Enthalpy profile (in kcal·mol⁻¹) for the reduction of CO₂ by **1** and catecholborane.

kcal·mol⁻¹, in line with a weak coordination of carbon dioxide as reflected by its geometry. Indeed, despite the bending of the molecule (indicative of CO₂ activation), the C–O bonds appear to be only slightly elongated compared to free CO₂ (1.28 and 1.21 Å). Nevertheless, this adduct can undergo addition of HBcat to yield a novel species whose formation is favorable by 14.4 kcal·mol⁻¹ compared to **1**. Once the formation of the complex **IM2** is achieved, the second reduction to generate the formaldehyde-1 adduct (**IM3**) and catBOBcat is downhill by 33.3 kcal·mol⁻¹. The third reduction to regenerate the catalyst as well as CH₃OBCat is an even more exothermic process (34.0 kcal·mol⁻¹). To summarize, as soon as the difficult coordination of CO₂ has taken place, the reduction is thermodynamically highly favorable.

In order to confirm these computational results, **1** was reacted with methylformate in an attempt to generate an analogous compound to **IM2**, namely species **IM2mf** (see Figure S22, SI). In line with the DFT results, where the adduct is predicted to be 3.9 kcal·mol⁻¹ higher in energy than **1**, no product could be observed by NMR spectroscopy. However, upon the addition of 3 equiv of catecholborane *without the presence of CO₂*, a 90% conversion to CH₃OBCat was observed after 20 h at room temperature. These latter results suggest that, although the formation of the adduct **IM2mf** is thermodynamically slightly disfavored, the reduction occurs in presence of a hydroborane. It is also interesting to note that the intermediate **IM3** is proposed to be formed in both reduction pathways. A similar formaldehyde intermediate was identified as a key intermediate in previous systems,^{4d,12} but could not be observed experimentally. While monitoring the reduction of CO₂ in the presence of hydroboranes and catalyst **1**, only one resonance in the ¹H NMR spectra, a broad singlet at 5.20 ppm, could not be assigned to the starting materials or products. Running the experiment in the presence of ¹³CO₂ allowed the observation of a ¹J_{C–H} of 151 Hz, suggesting that this species arise from the reduction of CO₂. In the latter experiment, a ¹J_{P–C} of 52 Hz was also observed both in the ¹³C{¹H} and ³¹P{¹H} NMR spectra. The latter species could not be isolated from the catalytic mixture, being in too small concentration in solution. However, when a solution of **1** was reacted with paraformaldehyde and heated at 70 °C for 15 min, the same product was observed to be formed with 74% conversion, as

characterized by multinuclear NMR spectroscopy as **IM3** (see Figure S19, SI).

In summary, we have reported a metal-free system for the reduction of carbon dioxide to methanol using a borane as reducing agent. The system is a robust, living catalytic system and generates TOFs up to 973 h^{-1} and TONs up to 2950 at 70°C under 1 atm of CO_2 , although larger TONs can be expected by additional loadings of hydroboranes. The key aspect of this reported system compared to the other metal-free systems for the activation of CO_2 is the weak interaction between the catalyst and carbon dioxide. Indeed, contrary to most ambiphilic and FLP systems reported to date, no adduct formation is observed between **1** and CO_2 . Nevertheless, CO_2 , being an ambiphilic molecule with its electrophilic carbon atom and nucleophilic oxygen atoms available, does not require significant bonding interaction with an ambiphilic catalyst to undergo reduction with hydroboranes. Once the first reduction has occurred, following reductions occur readily to generate CH_3OBR_2 . Preliminary results demonstrate that the BPin analogue 1-Bpin-2-PPh₂-C₆H₄¹⁵ is an active catalyst for the CO_2 reduction using $\text{BH}_3\cdot\text{SMe}_2$, albeit working less efficiently than **1** (TOF of 24 h^{-1} in conditions similar to those of entry 2 of Table 1). Current work focuses on optimizing the steric and electronic properties at boron and phosphorus centers to obtain optimal catalytic activity. Computational studies to unveil the full reaction mechanism are also well underway.

■ ASSOCIATED CONTENT

Supporting Information

Synthesis and characterization of **1**, catalytic procedures, and DFT details. This material is available free of charge via the Internet at <http://pubs.acs.org>.

■ AUTHOR INFORMATION

Corresponding Author

frederic.fontaine@chm.ulaval.ca

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

This work was supported by the National Sciences and Engineering Research Council of Canada (NSERC, Canada) and the Centre de Catalyse et Chimie Verte (Quebec). M.-A.C. and M.-A.L. thank NSERC and FQRNT for scholarships. We acknowledge W. Bi for the X-ray structure of **1** and R. Lafleur-Lambert with his help setting some catalytic experiments. L.M. is member of the Institut Universitaire de France. Cines and CALMIP are acknowledged for a generous grant of computing time. The Humboldt foundation is also acknowledged for financial support.

■ REFERENCES

- (1) D'Alessandro, D. M.; Smit, B.; Long, J. R. *Angew. Chem., Int. Ed.* **2010**, *49*, 6058–6082.
- (2) (a) Huang, K.; Sun, C.-L.; Shi, Z.-J. *Chem. Soc. Rev.* **2011**, *40*, 2435–2452. (b) Sakakura, T.; Choi, J.-C.; Yasuda, H. *Chem. Rev.* **2007**, *107*, 2365–2387. (c) Aresta, M.; Dibenedetto, A. *Dalton Trans.* **2007**, 2975–2992.
- (3) Olah, G. A.; Goeppert, A.; Surya Prakash, G. K. *J. Org. Chem.* **2009**, *74*, 487–498.
- (4) (a) Balaraman, E.; Gunanathan, C.; Zhang, J.; Shimon, L. J. W.; Milstein, D. *Nat. Chem.* **2011**, *3*, 609–614. (b) Huff, C. A.; Sanford, M. S. *J. Am. Chem. Soc.* **2011**, *133*, 18122–18125. (c) Wesselbaum, S.; von Stein, T.; Klankermayer, J.; Leitner, W. *Angew. Chem., Int. Ed.* **2012**, *51*, 7499–7502. (d) Chakraborty, S.; Zhang, J.; Krause, J. A.; Guan, H. *J. Am. Chem. Soc.* **2010**, *132*, 8872–8873. (e) Park, S.; Bézier, D.; Brookhart, M. *J. Am. Chem. Soc.* **2012**, *134*, 11404–11407. (f) Matsuo, T.; Kawaguchi, H. *J. Am. Chem. Soc.* **2006**, *128*, 12362–12363. (g) Mitton, S. J.; Turculet, L. *Chem.—Eur. J.* **2012**, *48*, 15258–15262. (h) Berkefeld, A.; Piers, W. E.; Parvez, M.; Castro, L.; Maron, L.; Eisenstein, O. *Chem. Sci.* **2013**, *4*, 2152–2162. (i) Schmeier, T. J.; Dobereiner, G. E.; Crabtree, R. H.; Hazari, N. *J. Am. Chem. Soc.* **2011**, *133*, 9274–9277. (j) Zhang, L.; Cheng, J.; Hou, Z. *Chem. Commun.* **2013**, *49*, 4782–4784. (k) Bontemps, S.; Vendier, L.; Sabo-Etienne, S. *Angew. Chem., Int. Ed.* **2012**, *51*, 1671–1674. (l) Cleeborg, C.; Cheung, M. S.; Lin, Z.; Marder, T. B. *J. Am. Chem. Soc.* **2011**, *133*, 19060–19063. (m) Lalrempuia, R.; Iglesias, M.; Polo, V.; Miguel, P. J. S.; Fernandez-Alvarez, F. J.; Pérez-Torrente, J. J.; Oro, L. A. *Angew. Chem., Int. Ed.* **2012**, *51*, 12824–12827. (n) Tominaga, K.; Sasaki, Y.; Kawai, M.; Watanabe, T.; Saito, M. *J. Chem. Soc. Chem. Commun.* **2013**, 629–631.
- (5) Khandelwal, M.; Wehmschulte, R. J. *Angew. Chem., Int. Ed.* **2012**, *51*, 7323–7326.
- (6) Schäfer, A.; Saak, W.; Haase, D.; Müller, T. *Angew. Chem., Int. Ed.* **2012**, *51*, 2981–2984.
- (7) Mömming, C.; Otten, M.; Kehr, E. G.; Fröhlich, R.; Grimme, S.; Stephan, D. W.; Erker, G. *Angew. Chem., Int. Ed.* **2009**, *48*, 6643–6646.
- (8) (a) Boudreau, J.; Courtemanche, M.-A.; Fontaine, F.-G. *Chem. Commun.* **2011**, *47*, 11131–11133. (b) Appelt, C.; Westenberg, H.; Bertini, F.; Ehlers, A. W.; Slootweg, J. C.; Lammertsma, K.; Uhl, W. *Angew. Chem., Int. Ed.* **2011**, *50*, 3925–3928.
- (9) Berkefeld, A.; Piers, W. E.; Parvez, M. *J. Am. Chem. Soc.* **2010**, *132*, 10660–10661.
- (10) Ménard, G.; Stephan, D. W. *J. Am. Chem. Soc.* **2010**, *132*, 1796–1797.
- (11) Ashley, A. E.; Thompson, A. L.; O'Hare, D. *Angew. Chem., Int. Ed.* **2009**, *48*, 9839–9843.
- (12) Riduan, S. N.; Zhang, Y.; Ying, J. Y. *Angew. Chem., Int. Ed.* **2009**, *48*, 3322–3325.
- (13) (a) Fontaine, F.-G.; Boudreau, J.; Thibault, M.-H. *Eur. J. Inorg. Chem.* **2008**, 5439–5454. (b) Boudreau, J.; Fontaine, F.-G. *Organometallics* **2011**, *30*, 511–519. (c) Fontaine, F.-G.; Zargarian, D. *J. Am. Chem. Soc.* **2004**, *126*, 8786–8794. (d) Boudreau, J.; Courtemanche, M.-A.; Marx, V. M.; Burnell, D. J.; Fontaine, F.-G. *Chem. Commun.* **2012**, *48*, 11250–11252.
- (14) (a) Bontemps, S.; Bouhadir, G. D.; Apperley, C.; Dyer, P. W.; Miqueu, K.; Bourissou, D. *Chem.—Asian J.* **2009**, *4*, 428–435. (b) Bontemps, S.; Bouhadir, G.; Miqueu, K.; Bourissou, D. *J. Am. Chem. Soc.* **2006**, *128*, 12056–12057.
- (15) Porcel, S.; Bouhadir, G.; Saffon, N.; Maron, L.; Bourissou, D. *Angew. Chem., Int. Ed.* **2010**, *49*, 6186–6189.
- (16) Baslé, O.; Porcel, S.; Ladeira, S.; Bouhadir, G.; Bourissou, D. *Chem. Commun.* **2012**, *48*, 4495–4497.
- (17) Harder, S.; Brandsma, L.; Kanters, J. A.; Duisenberg, A.; van Lenthe, J. H. *J. Organomet. Chem.* **1991**, *420*, 143.
- (18) Pereira, S.; Srebnik, M. *Organometallics* **1995**, *14*, 3127–3128.
- (19) Westcott, S. A.; Blom, H. P.; Marder, T. B.; Baker, R. T.; Calabrese, J. C. *Inorg. Chem.* **1993**, *32*, 2175–2182.