

Tetrahedron Letters 39 (1998) 5437-5438

TETRAHEDRON LETTERS

On the use of 9-Borabicyclo[3.3.1]nonane as an Initiator for Low-Temperature Free-Radical Reductions

V. Tamara Perchyonok and Carl H. Schiesser*

School of Chemistry, The University of Melbourne, Parkville, Victoria, Australia, 3052

Received 27 March 1998; accepted 19 May 1998

Abstract: 9-Borabicyclo[3.3.1]nonane (9-BBN) (1) is an effective low-temperature initiator for use in stannane mediated free-radical reductions of a wide cross-section of typical radical precursors. © 1998 Elsevier Science Ltd. All rights reserved.

While there exist several classes of initiator for stannane-based free-radical reactions carried out at around room temperature and above, low temperature initiation has largely been restricted to photolysis or the use of triethylborane in the presence of oxygen.¹ Neither of these two techniques are appropriate for reactions involving compounds containing photo-labile or oxygen-sensitive functionalities; there are some suggestions however that reactions involving triethylborane can be initiated by very minor amounts of oxygen.¹

Our recent interest in the development of new reagents for free-radical reactions at -78°C required that we locate a more user-friendly low-temperature initiator. Our requirements included the ability to initiate efficiently at -78° without the need for oxygen while maintaining acceptable yields in reasonable reaction times. We now report that 9-borabicyclo[3.3.1]nonane (9-BBN) $(1)^2$ efficiently initiates the reduction of a wide cross-section of typical radical precursors (2 - 7) by tributyl- and triphenyltin hydride, under nitrogen at both 0° and -78° in good to excellent yields.





We began to explore the use of 9-BBN in radical chemistry through the low-temperature reaction of 1bromooctane with tributyltin hydride. To that end, a toluene solution of tributyltin hydride (1 equiv.) was injected into a deoxygenated solution containing 1-bromooctane and $(4 \text{ mol}\%)^3$ 9-BBN in toluene at the required temperature and the resultant solution (ca. 0.06M in stannane) stirred under nitrogen for 2h at either 0° or -78°. At this time TLC analysis indicated the absence of starting material. Further analysis by GC revealed the formation of octane (by comparison with an authentic sample) in yields of 74% (0°) and 60% (-78°) through integration against an internal standard (decane). Importantly, when the reaction was repeated in the absence of 9-BBN, no reaction was observed after 2h. It is interesting to compare these results with those obtained using triethylborane as radical initiator; the use of 4 mol% Et_aB under nitrogen afforded octane

0040-4039/98/\$19.00 © 1998 Elsevier Science Ltd. All rights reserved. PII: S0040-4039(98)01048-X in 30% (-78°) and 19% (0°) yields after 2h. When these reactions were repeated under an aerobic atmosphere, as expected, improved yields (43% at -78°; 80% at 0°) were observed.

The generality of this procedure was explored by reacting tributyl- as well as triphenyltin hydride with a representative set of radical precursors (2-8);⁴ Table 1 lists the results of these reactions.

RX	Bu ₃ SnH			Pb ₃ SnH		
	9-BBN/N ₂	Et ₃ B/O ₂	Et ₃ B/N ₂	9-BBN/N ₂	Et ₃ B/O ₂	Et ₃ B/N ₂
2	74 (60)	80 (43)	19 (30)	73 (84)	83 (60)	25 (20)
3	70 (61)	76 (60)	14 (17)	100 (100)	90 (71)	22 (10)
4	84 (65)	78 (44)	20 (33)	83 (83)	61 (43)	22 (26)
5	72 (67)	75 (48)	15 (25)	100 (88)	87 (44)	11 (13)
6	78 (68)	46 (56)	0 (15)	86 (64)	74 (57)	0 (12)
7	96 (52)	75 (37)	14 (18)	88 (76)	79 (47)	19(11)
<u> </u>	88 (70)	62 (47)	16 (0)	84 (80)	46 (67)	18 (0)

Table 1. Yields of hydrocarbons (RH) obtained by reaction of radical precursors (RX) with tributyl- and triphenyltin hydride in toluene (ca. 0.06M) initiated by 9-BBN or Et₃B (4mol%) at 0° and -78°(in parentheses).

^aProduct consisted of a mixture of 1-hexene (9), methylcyclopentane (10) and cyclohexane (11).

Inspection of Table 1 reveals that at both temperatures, 9-BBN effectively initiates the reduction of a wide selection of common radical precursors by tributyl- and triphenyltin hydride in good to excellent yield. Importantly, these results are obtained under nitrogen, whereas similar initiation by Et_3B requires at least some oxygen present in the reaction.

In order to establish the intermediacy of radicals in these transformations, we examined the reaction of 6-bromo-1-hexene with $Bu_3SnH/9$ -BBN under similar conditions. GC analysis indicated the formation 1-hexene (9), methylcyclopentane (10) and cyclohexane (11) in the relative yields given in Scheme 1 and in the overall yield listed in Table 1. Application of the appropriate integrated rate equation⁵ leads to values of k_c/k_H of 0.04 (0°) and 0.006 (-78°), in excellent agreement with the expected values of 0.05 and 0.006 based on the well established kinetic equation for the cyclization of the 5-hexenyl radical.⁶ These data leave little doubt that the low-temperature reductions initiated by 9-BBN presented in this report are radical in nature.

We thank the Australian Research Council for financial support.

Notes and References

- Miura, K.; Ichinose, Y.; Nozaki, K.; Fugami, K.; Oshima, K.; Utimoto, K., Bull. Chem. Soc. Jpn., 1989, 62, 143. Curran, D. P.; Nanni, Tetrahedron: Asymmetry, 1996, 7, 2417.
- 9-BBN is an air-stable dimeric solid useful for selective hydroborations. For examples, see: Knights, E. F.; Brown, H. C., J. Am. Chem. Soc., 1968, 90, 5280. Knights, E. F.; Brown, H. C., J. Am. Chem. Soc., 1968, 90, 5281. Brown, H. C.; Krishnamurthy, S.; Yoon, N. M., J. Org. Chem., 1976, 41, 1778.
- 3. Based on monomeric 9-BBN.
- Giese, B., Radicals in Organic Synthesis: Formation of Carbon-Carbon Bonds, 1986, Pergamon Press, Oxford. Regitz, M.; Giese, B., Radicale, Houben-Weyl Methoden der Organischen Chemie, 1989, Vol E 19a, Georg Thieme, Stuttgart. Perkins, M. J., Radical Chemistry, 1994, Ellis-Horwood, New York.
- 5. Rate equation: $d[9]/d[10] = k_c/(k_H[Bu_3SnH])$ where k_c and k_H are the rate constants for the formation of 9 and 10 respectively.
- Johnston, L. J.; Lusztyk, J.; Wayner, D. D. M.; Abeywickrema, A. N.; Beckwith, A. L. J.; Scaiano, J. C.; Ingold, K. U., J. Am. Chem. Soc., 1985, 107, 4594. Chatgilialoglu, C.; Ingold, K. U.; Scaiano, J. C., J. Am. Chem. Soc., 1981, 103, 7739. Dakternieks, D.; Henry, D. J.; Schiesser, C. H., J. Org. Chem., submitted for publication.