



A Journal of



Accepted Article

Title: Direct stereoselective aziridination of cyclohexenols with 3-amino-2-(trifluoromethyl)quinazolin-4(3H)-one in the synthesis of cyclitol aziridine glycosidase inhibitors

Authors: Marta Artola, Shirley Wouters, Sybrin P. Schröder, Casper de Boer, Yurong Chen, Rita Petracca, Adrianus M. C. H. van den Nieuwendijk, Johannes M. F. G. Aerts, Gijsbert A. van der Marel, Jeroen D. C. Codée, and Herman S. Overkleeft

This manuscript has been accepted after peer review and appears as an Accepted Article online prior to editing, proofing, and formal publication of the final Version of Record (VoR). This work is currently citable by using the Digital Object Identifier (DOI) given below. The VoR will be published online in Early View as soon as possible and may be different to this Accepted Article as a result of editing. Readers should obtain the VoR from the journal website shown below when it is published to ensure accuracy of information. The authors are responsible for the content of this Accepted Article.

To be cited as: *Eur. J. Org. Chem.* 10.1002/ejoc.201801703

Link to VoR: <http://dx.doi.org/10.1002/ejoc.201801703>

Supported by



WILEY-VCH

FULL PAPER

Direct stereoselective aziridination of cyclohexenols with 3-amino-2-(trifluoromethyl)quinazolin-4(3*H*)-one in the synthesis of cyclitol aziridine glycosidase inhibitors

Marta Artola,^{*,[a]} Shirley Wouters,^[a] Sybrin P. Schröder,^[a] Casper de Boer,^[a] Yurong Chen,^[a] Rita Petracca,^[a] Adrianus M. C. H. van den Nieuwendijk,^[a] Johannes M. F. G. Aerts,^[b] Gijsbert A. van der Marel,^[a] Jeroen D. C. Codée,^[a] and Herman S. Overkleeft^{*,[a]}

Abstract: Cyclophellitol aziridine and its configurational and functional isomers are powerful covalent inhibitors of retaining glycosidases and find application in fundamental studies on glycosidases, amongst others in relation to inherited lysosomal storage disorders caused by glycosidase malfunctioning. Few direct and stereoselective aziridination methodologies are known for the synthesis of cyclophellitol aziridines. Here, we present our studies on the scope of direct 3-amino-2-(trifluoromethyl)quinazolin-4(3*H*)-one-mediated aziridination on a variety of configurational and functional cyclohexenol isosteres. We demonstrate that the aziridination can be directed by an allylic or homoallylic hydroxyl through H-bonding and that steric hindrance plays a key role in the diastereoselectivity of the reaction.

Introduction

Glycosidases are enzymes involved in the degradation of complex glycoconjugates in nature and are of relevance both in biomedicine and biotechnology.^[1] Many glycosidases follow a two-step Koshland double displacement mechanism, which involves a covalent enzyme-glycoside intermediate.^[2] The active site of such retaining glycosidases is usually composed of an aspartic acid or glutamic acid, termed the catalytic acid/base, and an aspartate/glutamate (or occasionally a tyrosine) termed the nucleophile. In the first step of substrate hydrolysis, the exocyclic oxygen is protonated by the acid/base residue. Next, the catalytic nucleophile attacks at the anomeric carbon and effects an S_N2




Figure 1. A) Koshland double displacement mechanism of retaining β -glucosidases. B) Cyclophellitol aziridines are 4H₃ transition-state mimics and inhibit covalently and irreversibly retaining β -glucosidases.

displacement of the aglycon, yielding a covalent enzyme–glycoside complex with inversion of the anomeric stereochemistry. In the second step, a water molecule is deprotonated by the acid/base carboxylate and hydrolyses the enzyme–substrate intermediate with a second inversion of the anomeric configuration (Figure 1A).^[3] Cyclitol aziridines can mimic the conformation of the oxocarbenium ion transition state and irreversibly inactivate glycosidases by covalently reacting with the nucleophilic carboxylate (Figure 1B). Based on this virtue and the fact, as supported by several studies in recent years from our group^[4], that covalent and irreversible inhibition is often both very effective and highly selective,

^[a] Dr. Marta Artola, Shirley Wouters, Dr. Sybrin P. Schröder, Casper de Boer, Yurong Chen, Dr. Rita Petracca, Adrianus M. C. H. van den Nieuwendijk, Prof. Gijsbert A. van der Marel, Dr. Jeroen D. C. Codée, Prof. Herman S. Overkleeft.
 Department of Bio-organic Synthesis, Leiden Institute of Chemistry, Leiden University, Einsteinweg 55, 2333 CC Leiden, The Netherlands

^[b] Prof. Johannes M. F. G. Aerts.
 Department of Medical Biochemistry, Leiden Institute of Chemistry, Leiden University, Einsteinweg 55, 2333 CC Leiden, The Netherlands

*Corresponding authors E-mails: m.e.artola@lic.leidenuniv.nl; h.s.overkleeft@lic.leidenuniv.nl
 Homepage: <https://www.universiteitleiden.nl/en/science/chemistry/biosyn>

Supporting information and ORCIDs from the authors for this article are available on the WWW under <https://doi.org/>

FULL PAPER

Previous work:




Figure 2. Literature aziridination reactions on glycoside configured cyclohexenes. A) Stereoselective aziridination through cyclic imidates; B) Staudinger-like ring closure of a 1,2-vicinal azido-alcohol obtained after nucleophilic addition of sodium azide to an epoxide intermediate; C) Direct synthesis of unprotected N-H aziridines with DHP and $\text{Rh}_2(\text{esp})_2$ catalyst; D) Stereoselective aziridination by means of Et-Q-NH₂ as nitrogen donor; E) Exploration of the aziridination scope by means of 3-aminoquinazolin-4(3H)-ones (R-Qs) on various glycoside-configured cyclohexenols.

cyclitol aziridines and their corresponding activity-based probes (ABPs) are highly useful tools for chemical glycobiology research.^[5]

An important step in the synthesis of cyclitol aziridine inhibitors and ABPs involves the stereoselective aziridination of a suitable cyclohexene precursor.^[6] In contrast to epoxidations, synthetic methodologies for the direct and stereoselective aziridination of alkenes are scarce.^[7] Our previous work on the synthesis of cyclophellitol aziridines relied mostly on either intramolecular iodocyclization followed by aziridine formation (Figure 2A)^[5a, c, d] or Staudinger-type ring closure of 1,2-azido-alcohols obtained from an epoxide precursor (Figure 2B).^[5b] We also investigated the use of O-(2,4-dinitrophenyl)hydroxylamine (DHP) as nitrogen donor with $\text{Rh}_2(\text{esp})_2$ as catalyst (Figure 2C).^[8] However, some limitations were encountered with these methodologies. Although the iodocyclization/intramolecular substitution sequence gives complete stereochemical control with reasonable to good overall yields, this sequence is not an option when the desired aziridine has the opposite stereochemistry of that of the directing alcohol moiety. The rhodium-catalyzed aziridination methodology appeared to be non-stereoselective and to proceed in relatively low yields,^[8] while in the Staudinger approach, purification can be challenging and over-reduced


amino-alcohols are occasionally generated as side products (although the use of polymer-bound triphenylphosphine alleviates these shortcomings to certain extent^[8b, 9]).

In the late 1960s, Atkinson and co-workers described the lead acetate mediated *in situ* oxidation of diverse hydrazines, amongst which 3-aminobenzoxazolin-2-one was used as aziridinating agent.^[10] They initially proposed that nitrenes could be formed as intermediates^[10-11] which would react with electrophilic and nucleophilic olefins to give the observed aziridines. Atkinson *et al.* extensively re-examined the reactivity of different substituted 3-aminoquinazolin-4(3H)-ones (Q-NH₂s) towards diverse olefins,^[12] and concluded that *N*-acetoxy-aminoquinazolones (Q-NHOAc) are the reactive intermediates, rather than nitrenes.^[13] Recently, the stereospecific aziridination of a partially protected *galacto*-configured cyclohexene has been described by Llebaria and co-workers based on the use of 3-amino-2-ethylquinazolin-4(3H)-one (Et-Q-NH₂) and $\text{PhI}(\text{OAc})_2$ (PIDA). The thus formed β -*galacto*-configured acetylated aziridine was employed in this study to produce *N*-acetoxy-aminoaziridine based irreversible inhibitors (Figure 2D).^[14] Inspired by this work of Llebaria and co-workers we decided to explore the scope of the direct olefin aziridination reaction by investigating the reactivity of diverse aminoquinazolin-4-ones towards differently configured and functionalized cyclohexenol substrates (Figure 2E). As we show here, this reaction proved to be particularly well suited for the synthesis of α -L-*idose*-configured cyclophellitol aziridine, a key intermediate for the synthesis of new α -L-iduronidase inhibitors and ABPs, as we recently reported in a separate body of work.^[15]


Results and Discussion

As the first research objective, D-*gluco*-configured cyclohexene **1a** was used as starting material to screen the most promising aminoquinazolinones described as nitrogen donors in the literature: 3-amino-2-ethylquinazolin-4(3H)-one (Et-Q-NH₂), 3-amino-2-(trifluoromethyl)quinazolin-4(3H)-one (CF_3 -Q-NH₂) and the chiral (S)-3-amino-2-(1-hydroxy-2,2-dimethylpropyl)quinazolin-4(3H)-one (HO-Q-NH₂), which all form the reactive *N*-acetoxy-aminoquinazolinones *in situ* in the presence of PIDA. In line with previous results,^[16] CF_3 -Q-NH₂ gave superior yields (69-75% of **1b**) when using cyclohexene **1a**, PIDA and the quinazolinone in a 1:2:2 ratio respectively and by forming the reactive *N*-acetoxy-aminoquinazolinone at -78 °C prior to addition of the olefin at -23 °C. Aziridine intermediate **1d** was isolated in 54% yield when using HO-Q-NH₂ as aziridination agent, whereas reaction with Et-Q-NH₂ returned starting material only (Scheme 1). Notably, the β -*gluco*-configured aziridine was formed stereoselectively in a 1.5 mmol reaction scale, indicating that hydrogen bonding from the homoallylic alcohol C7-OH guides the incoming Q-NHOAc, in agreement with the mechanistic proposal of Atkinson *et al.* (Scheme 2).^[12a] These results led us to further investigate aziridinations with CF_3 -Q-NH₂ on different cyclohexene substrates.

FULL PAPER




Scheme 1. Stereoselective aziridination of cyclohexene **1a** with 3-amino-2-(trifluoromethyl)quinazolin-4(3*H*)-one ($\text{CF}_3\text{-Q-NH}_2$), chiral (*S*)-3-amino-2-(1-hydroxy-2,2-dimethylpropyl)quinazolin-4(3*H*)-one (HO-Q-NH_2) or 3-amino-2-ethylquinazolin-4(3*H*)-one (Et-Q-NH_2) as nitrogen donor. Reagents and conditions: (i) $\text{PhI}(\text{OAc})_2$, R-Q-NH , DCM, rt, 48 h.




Scheme 2. Proposed reaction transition state driven by H-bonding from the homoallylic or allylic alcohols of *gluco*-cyclohexene **1a** and **2a**, respectively, with *N*-acetoxy-3-amino-2-(trifluoromethyl)quinazolin-4(3*H*)-one ($\text{CF}_3\text{-Q-NHOAc}$, formed *in situ*) during the aziridination reactions, and subsequent deprotection under Birch conditions. Starting material was recovered when the reaction was performed on perbenzylated *gluco*-cyclohexene **3a**. Reagents and conditions: (i) $\text{PhI}(\text{OAc})_2$, $\text{CF}_3\text{-Q-NH}$, DCM, rt, 48 h; (ii) Li , $\text{NH}_3(\text{liq.})$, THF, $-60\text{ }^\circ\text{C}$, 1 h.

When *gluco*-configured cyclohexene **2a** bearing a 4,6-benzylidene acetal and an allylic alcohol at C-2 was used, α -aziridine **2b** was exclusively formed in 55% yield, providing further support for H-bonding guided delivery of the aziridinating reagent (Scheme 2). When perbenzylated *gluco*-cyclohexene **3a** was subjected to the same reaction conditions no conversion

was observed, indicating that the system is not reactive enough without the hydrogen bonding guided delivery and/or that the double bond, with relatively bulky substituents on either side of the alkene, is too crowded to allow for an effective addition. A similar pattern was observed with *galacto*-configured cyclohexenes. *Galacto*-configured cyclohexene **4a** could be stereoselectively transformed into β -aziridine **4b** while cyclohexene **5a** afforded α -aziridine **5b** in 61% yield (Scheme 3). These examples again illustrate the impact of neighboring (homo)allylic alcohol functionalities. Partially protected conduritol **6a** was also amenable to stereoselective aziridination, affording β -aziridine **6b** in 66% yield, whereas the starting material was recovered when the reaction was performed with perbenzylated conduritol **7a** (Scheme 4).




Scheme 3. Proposed reaction transition state driven by H-bonding from the homoallylic or allylic alcohols of *galacto*-cyclohexene **4a** and **5a** respectively, with *N*-acetoxy-3-amino-2-(trifluoromethyl)quinazolin-4(3*H*)-one ($\text{CF}_3\text{-Q-NHOAc}$, formed *in situ*) during the aziridination reactions, and subsequent deprotection under Birch conditions. Reagents and conditions: (i) $\text{PhI}(\text{OAc})_2$, $\text{CF}_3\text{-Q-NH}$, DCM, rt, 48 h; (ii) Li , $\text{NH}_3(\text{liq.})$, THF, $-60\text{ }^\circ\text{C}$, 1 h.



Scheme 4. Proposed reaction transition state driven by H-bonding from the allylic alcohol of *conduritol*-cyclohexene **6a** with *N*-acetoxy-3-amino-2-(trifluoromethyl)quinazolin-4(3*H*)-one ($\text{CF}_3\text{-Q-NHOAc}$, formed *in situ*) during the aziridination reaction, and subsequent deprotection under Birch conditions. Perbenzylated conduritol **7a** does not react under the aziridination conditions. Reagents and conditions: (i) $\text{PhI}(\text{OAc})_2$, $\text{CF}_3\text{-Q-NH}$, DCM, rt, 48 h; (ii) Li , $\text{NH}_3(\text{liq.})$, THF, $-60\text{ }^\circ\text{C}$, 1 h.


FULL PAPER

In order to investigate whether an alcohol further away from the alkene could guide the reagent to one of the diastereotopic faces of the double bond, we examined the aziridination of partially protected *xylo*-configured cyclohexene **8a**. In this case only β -isomer **8b** was obtained, indicating that the 4-OH is too distal for a productive H-bond interaction and that the aziridination takes place on the least hindered face of the double bond, opposite of the C-2-benzyl ether (Scheme 5).



Scheme 5. Proposed reaction transition state driven by steric hindrance of *arabino*-cyclohexene **8a** with *N*-acetoxy-3-amino-2-(trifluoromethyl)quinazolin-4(3*H*)-one ($\text{CF}_3\text{-Q-NHOAc}$, formed *in situ*) during the aziridination reaction, and subsequent deprotection under Birch conditions. Reagents and conditions: (i) $\text{PhI}(\text{OAc})_2$, $\text{CF}_3\text{-Q-NH}$, DCM, rt, 48 h; (ii) Li, NH_3 (liq.), THF, -60°C , 1 h.

We finally explored aziridination of *L-ido*-configured cyclohexenes in order to obtain α -*L-ido*-configured aziridines as potential intermediates for the development of new iduronidase inhibitors.^[15] Partially protected cyclohexene **9a** was not amenable to aziridination, possibly because the primary alcohol directs to the beta side while this region may be hindered by the allylic benzyl ether. Considering that β -*D-xylo*-configured aziridine **8a** was obtained without H-bonding, we postulated H-bonding would not be essential for a satisfactory aziridination in case the double bond is readily accessible. To test this hypothesis, the free alcohols in **9a** were benzylated (benzyl bromide, sodium hydride) to generate cyclohexene **10a**. From this fully protected cyclohexenol, α -*L-ido*-aziridine **10b** was obtained in 43% yield together with 32% recovered starting material after direct aziridination using $\text{CF}_3\text{-Q-NHOAc}$ as the nitrogen donor (Table 1, Scheme 6). The α -configuration of aziridine **10c** was confirmed by comparison of the experimental ^1H NMR coupling constants with the corresponding calculated values acquired from DFT calculations.^[15]



Scheme 6. Partially protected cyclohexene **9a** does not react under the aziridination conditions. Proposed reaction transition state driven by steric hindrance of perbenzylated *ido*-cyclohexene **10a** with *N*-acetoxy-3-amino-2-(trifluoromethyl)quinazolin-4(3*H*)-one ($\text{CF}_3\text{-Q-NHOAc}$, formed *in situ*) during the aziridination reaction, and subsequent deprotection under Birch conditions.

Table 1. Aziridination reaction and subsequent Birch reduction on various configured cyclohexenes **1a-10a**.

Starting material	$\text{CF}_3\text{-Q}$ -mediated aziridination	Birch deprotection
1a	β -D- <i>gluco</i> 1b : 75%	β -D- <i>gluco</i> 1c : 99%
2a	α -D- <i>gluco</i> 2b : 55%	α -D- <i>gluco</i> 2c : 96%
3a	[a]	[a]
4a	β -D- <i>galacto</i> 4b : 48% and 15% S.M.	β -D- <i>galacto</i> 4c : 81%
5a	α -D- <i>galacto</i> 5b : 61%	α -D- <i>galacto</i> 5c : 85%
6a	β -D- <i>conduritol</i> 6b : 66%	<i>conduritol</i> 6c : 99%
7a	[a]	[a]
8a	β -D- <i>xylo</i> 8b : 45% and 37% S.M. ^[a]	β -D- <i>xylo</i> 8c : 96%
9a	[a]	[a]
10a	α -L- <i>ido</i> 10b : 43% and 32% S.M.	α -L- <i>ido</i> 10c : 93%

[a] Starting material (S.M.) recovered only.

In all cases, one step deprotection of the aziridine and hydroxyls in the aforementioned $\text{CF}_3\text{-Q}$ functionalized aziridine intermediates was achieved under Birch conditions using lithium and liquid ammonia at -78°C . The reactions were quenched with H_2O and impurities derived from $\text{CF}_3\text{-Q}$ precipitated and were removed by filtration. The cyclitol aziridines were finally obtained in excellent yields after cation-exchange chromatography with Amberlite H^+ resin to eliminate the lithium hydroxide salts (Table 1, 81-99%).

Conclusions

We have explored direct aziridination of both, partially protected and fully protected, configurational cyclohexenol using different substituted 3-aminoquinazolin-4(3*H*)-ones. From these studies we identified 3-amino-2-(trifluoromethyl)quinazolin-4(3*H*)-one ($\text{CF}_3\text{-Q}$) as the superior aziridinating agent. Using this reagent, direct aziridination reaction can be applied on diverse glycoside configured cyclohexenes, and it appears that aziridination can be directed by allylic or homoallylic hydroxyls through H-bonding and that steric hindrance plays an essential role in the diastereoselectivity of the reaction. With this in mind, one could tune the cyclohexene scaffold depending on the desired configuration of the target aziridine and thus, synthesize diverse glycosidase inhibitors effectively in asymmetric fashion.

Experimental Section

General methods and materials: All reagents were of a commercial grade and were used as received unless stated otherwise. Dichloromethane (DCM), tetrahydrofuran (THF) and *N,N*-dimethylformamide (DMF) were stored over 4 Å molecular sieves, which were dried *in vacuo* before use. All reactions were performed under an argon atmosphere unless stated otherwise. Reactions were monitored by analytical thin-layer chromatography (TLC) using Merck aluminum sheets pre-coated

FULL PAPER

with silica gel 60 with detection by UV absorption (254 nm) and by spraying with a solution of $(\text{NH}_4)_6\text{Mo}_7\text{O}_{24} \cdot \text{H}_2\text{O}$ (25 g/L) and $(\text{NH}_4)_4\text{Ce}(\text{SO}_4)_4 \cdot \text{H}_2\text{O}$ (10 g/L) in 10% sulfuric acid followed by charring at $\sim 150^\circ\text{C}$ or by spraying with an aqueous solution of KMnO_4 (7%) and K_2CO_3 (2%) followed by charring at $\sim 150^\circ\text{C}$. Column chromatography was performed manually or with a Biotage IsoleraTM flash purification system using silica gel cartridges (Screening devices SiliaSep HP, particle size 15–40 μm , 60A) in the indicated solvents. ^1H NMR and ^{13}C NMR spectra were recorded on Bruker AV-400 (400/101 MHz) and Bruker AV-500 (500/126 MHz) spectrometer in the given solvent. Chemical shifts are given in ppm relative to the chloroform residual solvent peak or tetramethylsilane (TMS) as internal standard. Coupling constants are given in Hz. All given ^{13}C spectra are proton decoupled. The following abbreviations are used to describe peak patterns when appropriate: s (singlet), d (doublet), t (triplet), q (quadruplet), m (multiplet), br (broad), Ar (aromatic), C_q (quaternary carbon), Q (quinazolinone). 2D NMR experiments (HSQC, COSY and NOESY) were carried out to assign protons and carbons of the new structures. High-resolution mass spectra (HRMS) of intermediates were recorded with a LTQ Orbitrap (Thermo Finnigan) and final compounds were recorded with an apex-QE instrument (Bruker). Optical rotations were measured on an Anton Paar MCP automatic polarimeter (Sodium D-line, $\lambda = 589\text{ nm}$). LC/MS analysis was performed on an LCQ Advantage Max (Thermo Finnigan) ion-trap spectrometer (ESI+) coupled to a Surveyor HPLC system (Thermo Finnigan) equipped with a C18 column (Gemini, 4.6 mm x 50 mm, 3 μm particle size, Phenomenex) equipped with buffers A: H_2O , B: acetonitrile (MeCN) and C: 1% aqueous TFA, or an Agilent Technologies 1260 Infinity LCMS with a 6120 Quadrupole MS system equipped with buffers A: H_2O , B: acetonitrile (MeCN) and C: 100 mM NH_4OAc .

2,3-Bis-O-benzyl-D-glucocyclohexene (**1a**)^[1], 7,4-O-benzylidene-D-glucocyclohexene (**2a**)^[17], perbenzylated D-glucocyclohexene (**3a**)^[4b], 2,3-Bis-O-benzyl-D-galactocyclohexene (**4a**)^[18], 2,3-Bis-O-benzyl-conduritol cyclohexene (**6a**)^[19], perbenzylated conduritol cyclohexene (**7a**)^[17], 2,3-Bis-O-benzyl-D-xylo-cyclohexene (**8a**)^[8b], 2,3-Bis-O-benzyl-L-ido-cyclohexene (**9a**)^[15] and perbenzylated L-ido-cyclohexene (**10a**)^[15], 3-amino-2-ethylquinazolin-4(3H)-one^[20], (S)-3-amino-2-(1-hydroxy-2,2-dimethylpropyl)quinazolin-4(3H)-one^[21] and 3-amino-2-(trifluoromethyl)-2,3-dihydroquinazolin-4(1H)-one^[20] were synthesized following procedures previously described and their spectroscopic data are in agreement with those previously reported.

Note: Numbering of proton peaks in cyclohexene and cyclitol aziridine derivatives is according to the numbering in Scheme 1.

(3aS,4S,5R,7aS)-5-(hydroxymethyl)-2-phenyl-3a,4,5,7a-tetrahydrobenzo[d][1,3]dioxol-4-ol (5a): (1S,2R,3S,6R)-6-(hydroxymethyl)cyclohex-4-ene-1,2,3-triol^[9, 22] was dissolved in dry DMF (2.0 mL) and dry MeCN (6.0 mL) in an inert atmosphere. CSA (173 mg, 0.74 mmol, 0.2 equiv.) was added to the solution, followed by $\text{PhCH}(\text{OMe})_2$ (838 μL , 5.58 mmol, 1.5 equiv.). The reaction was stirred overnight and then it was quenched with Et_3N (100 μL , 0.23 mmol, 0.2 equiv.). The reaction mixture was extracted with EtOAc (x2) and water. The combined organic layers were washed with brine, dried over Na_2SO_4 , filtered and concentrated *in vacuo*. The product was purified by silica column chromatography (from 30%→70% EtOAc in pentane), affording

cyclohexene **5a** (598 mg, 2.41 mmol, 61%). ^1H NMR (500 MHz, CDCl_3): $\delta = 7.45 - 7.37$ (m, 2H, 2 x CH Ar), 7.33 (dd, $J = 5.1, 2.0$ Hz, 3H, 3 x CH Ar), 5.79 (dt, $J = 10.1, 2.5$ Hz, 1H, CH=CH), 5.55 (dd, $J = 10.1, 2.1$ Hz, 1H, CH=CH), 5.43 (s, 1H, CH-Ph), 4.41 (d, $J = 7.2$ Hz, 1H, CH-2), 4.29 (d, $J = 1.4$ Hz, 1H, CH-4), 4.16 (d, $J = 11.6$ Hz, 1H, CH-7a), 4.02 (dd, $J = 11.7, 3.4$ Hz, 1H, CH-7b), 3.59 (t, $J = 7.3$ Hz, 1H, CH-3), 3.33 – 3.14 (m, 2H, 2 x OH), 2.22 (s, 1H, CH-5). ^{13}C NMR (126 MHz, CDCl_3): δ 138.3 (C_q Ph), 130.8 (C-6), 129.2 (CH Ar), 128.4 (2 x CH Ar), 127.8 (C-1), 126.4 (2 x CH Ar), 101.2 (CHPh), 77.0 (C-4), 75.8 (C-3), 70.6 (C-2), 70.5 (C-7), 35.8 (C-5). ESI-MS (m/z): 248.9 [$\text{M}+\text{H}^+$], HRMS: calcd. for $[\text{C}_{14}\text{H}_{17}\text{O}_4]^+$ 249.11268; found 249.11220.

General procedure for aziridination: A solution of 3-amino-2-(trifluoromethyl)-2,3-dihydroquinazolin-4(1H)-one (2 equiv.) in anhydrous DCM (10 mL/mmol of cyclohexene) was added dropwise over a period of 30 min to a stirred suspension of $\text{PhI}(\text{OAc})_2$ (PIDA) (2 equiv.) in anhydrous DCM (5 mL/mmol of cyclohexene) at -78°C . The resultant mixture was stirred for additional 30 min and then cooled to -23°C and a solution of the corresponding cyclohexene (1 equiv.) in DCM (1 mL/mmol of cyclohexene) was added dropwise over a period of 15 min. The reaction mixture was stirred at -23°C for one hour and then the reaction was allowed to warm to room temperature and stirred for 1–2 days. The mixture was diluted with EtOAc and subsequently washed with 0.5 M aqueous KOH solution and water. The combined water layers were extracted with EtOAc (x2), and the combined organic layers were dried over MgSO_4 , filtered and concentrated *in vacuo*. Purification by column chromatography (from pentane to pentane:EtOAc 1:1) gave the desired aziridines.

β -D-glucocyclitol CF₃-Q-aziridine (1b): Obtained from cyclohexene **1a** (487 mg, 1.40 mmol) as an orange oil in 75% yield (609 mg, 1.07 mmol). ^1H NMR (400 MHz, CDCl_3): $\delta = 8.15$ (dt, $J = 8.1, 1.1$ Hz, 1H, CH Q), 7.86 – 7.80 (m, 2H, 2 x CH Q), 7.60 (ddd, $J = 8.2, 5.0, 3.3$ Hz, 1H, CH Q), 7.40 – 7.30 (m, 7H, 7 x CH Ar), 7.18 (t, $J = 7.7$ Hz, 2H, 2 x CH Ar), 6.99 – 6.91 (m, 1H, CH Ar), 4.97 (d, $J = 11.2$ Hz, 1H, CHHPh), 4.75 (d, $J = 11.4$ Hz, 1H, CHHPh), 4.69 (dd, $J = 11.4, 3.4$ Hz, 2H, 2 x CHHPh), 4.42 (dd, $J = 7.3, 3.4$ Hz, 1H, CH-6), 4.03 (dd, $J = 11.0, 6.7$ Hz, 1H, CH-7a), 3.92 (d, $J = 8.2$ Hz, 1H, CH-2), 3.84 (dd, $J = 11.0, 6.1$ Hz, 1H, CH-7b), 3.70 (d, $J = 7.3$ Hz, 1H, CH-1), 3.50 (t, $J = 10.0$ Hz, 1H, CH-4), 3.38 (dd, $J = 10.0, 8.2$ Hz, 1H, CH-3), 3.16 (br s, 1H, OH), 2.96 (br s, 1H, OH), 2.14 (dtd, $J = 10.0, 6.4, 3.4$ Hz, 1H, CH-5). ^{13}C NMR (101 MHz, CDCl_3): δ 161.0 (C=O), 143.9 (C_q Q), 143.4 (q, $J = 35.0$ Hz, CCF_3), 138.2 (C_q Ph), 137.4 (C_q Ph), 135.1, 129.4, 128.6, 128.6, 128.4, 128.0, 126.5 (10 x CH Ar, 4 x CH Q), 123.2 (C_q Q), 118.2 (q, $J_{\text{CF}} = 277.0$ Hz, CF_3), 84.1 (C-3), 79.8 (C-2), 75.1 (CH_2Ph), 73.3 (CH_2Ph), 69.3 (C-4), 63.7 (C-7), 43.6 (C-5), 40.5, 40.4 (C-1, C-6). HRMS: calcd. for $[\text{C}_{30}\text{H}_{29}\text{F}_3\text{N}_3\text{O}_5]^+$ 568.20593; found 568.20551.

β -D-glucocyclitol OH-Q-aziridine (1d): Obtained from cyclohexene **1a** (507 mg, 1.49 mmol) as an orange oil in 51% yield (443 mg, 0.76 mmol). ^1H NMR (400 MHz, CDCl_3): $\delta = 8.21$ (dd, $J = 8.1$ Hz, 1H, CH Q), 7.81 – 7.73 (m, 1H, CH Q), 7.71 – 7.66 (m, 1H, CH Q), 7.54 – 7.47 (m, 1H, CH Q), 7.47 – 7.42 (m, 2H, 2 x CH Ar), 7.37 – 7.30 (m, 8H, 8 x CH Ar), 5.06 – 5.01 (m, 2H, CHHPh, CH-2), 4.99 (d, $J = 11.1$ Hz, 1H, CHHPh), 4.79 (s, 1H, OH), 4.76 – 4.69 (m, 2H, 2 x CHHPh), 4.27 – 4.06 (m, 3H, CH-7a,b, CH-3), 3.65 – 3.55 (m, 1H, OH), 3.50 – 3.42 (m, 2H, CH-4, Bu-CH-OH), 3.39 (dd, $J = 8.1, 3.5$ Hz, 1H, CH-6), 2.76 (d, $J =$

FULL PAPER

8.0 Hz, 1H, CH-1), 2.22 (ddt, $J = 10.7, 7.4, 3.6$ Hz, 1H, CH-5), 1.02 (s, 9H, 3 x CH₃). ¹³C NMR (101 MHz, CDCl₃): δ 159.4 (C=O), 144.5 (C_q Q), 138.1 (C_q Ph), 137.3 (C_q Ph), 134.6, 128.7, 128.6, 128.5, 128.1, 128.0, 127.9, 127.8, 127.4, 127.0, 126.6 (10 x CH Ar, 4 x CH Q), 121.2 (C_q Q), 84.6 (tBu-CH-OH), 80.0 (C-3), 75.4 (CH₂Ph), 74.9 (C-2), 72.6 (CH₂Ph), 66.3 (C-4), 61.3 (C-7), 51.5 (C-1), 49.2 (C-6), 44.2 (C-5), 38.5 (C(CH₃)₃), 25.9 (3 x CH₃). ESI-MS (m/z): 586.3 [M+H⁺]

α -D-glucocyclitol CF₃-Q-aziridine (2b): Obtained from cyclohexene **2a** (40 mg, 0.161 mmol) as an orange oil in 55% yield (43 mg, 0.089 mmol). ¹H NMR (500 MHz, CDCl₃): δ 8.21 (d, $J = 7.7$ Hz, 1H, CH Q), 7.90 – 7.80 (m, 2H, 2 x CH Q), 7.62 (ddd, $J = 8.2, 6.2, 2.1$ Hz, 1H, CH Q), 7.50 (dd, $J = 7.6, 2.1$ Hz, 2H, 2 x CH Ar), 7.42 – 7.33 (m, 3H, 3 x CH Ar), 5.54 (s, 1H, CHPh), 4.47 (dd, $J = 10.7, 4.7$ Hz, 1H, CH-7a), 4.10 (d, $J = 7.2$ Hz, 1H, CH-2), 4.06 (dd, $J = 7.2, 4.5$ Hz, 1H, CH-1), 3.84 (dd, $J = 11.8, 10.7$ Hz, 1H, CH-7b), 3.73 (dd, $J = 10.3, 7.8$ Hz, 1H, CH-3), 3.66 (br s, 1H, OH), 3.45 (d, $J = 7.2$ Hz, 1H, CH-6), 3.30 (t, $J = 10.5$ Hz, 1H, CH-4), 2.85 (br s, 1H, OH), 2.38 (dddd, $J = 12.0, 10.5, 4.7, 1.8$ Hz, 1H, CH-5). ¹³C NMR (126 MHz, CDCl₃): δ 160.8 (C=O), 143.9 (C_q Q), 142.4 (q, $J = 36.6$ Hz, CCF₃), 137.4 (C_q Ph), 135.4, 129.8, 129.4, 128.8, 128.5, 126.8, 126.3 (10 x CH Ar, 4 x CH Q), 122.7 (C_q Q), 128.2 (q, $J_{CF} = 278.2$ Hz, CF₃), 102.3 (CHPh), 80.0 (C-4), 73.6 (C-3), 72.2 (C-2), 68.8 (C-7), 43.8 (C-1), 41.4 (C-6), 37.4 (C-5). HRMS: calcd. for [C₂₃H₂₁F₃N₃O₅]⁺ 476.1433; found 476.1423; calcd. for [C₂₃H₂₀F₃N₃NaO₅]⁺ 498.1252; found 498.1244.

β -D-galactocyclitol CF₃-Q-aziridine (4b): Obtained from cyclohexene **4a** (470 mg, 2.05 mmol) as an orange oil in 49% yield (225 mg, 1.00 mmol) and 15% starting material recovered. $[\alpha]_D^{20} = +69.4$ ($c = 0.5$, CHCl₃). ¹H NMR (400 MHz, CDCl₃): δ = 8.22 – 8.16 (d, $J = 7.6$ Hz, 1H, CH Q), 7.88 – 7.80 (m, 2H, 2 x CH Q), 7.62 (ddd, $J = 8.2, 6.3, 2.1$ Hz, 1H, CH Q), 7.42 – 7.21 (m, 9H, CH Ar), 7.15 – 7.07 (m, 1H, CH Ar), 4.93 (d, $J = 10.9$ Hz, 1H, CHHPh), 4.78 (d, $J = 11.6$ Hz, 1H, CHHPh), 4.76 (d, $J = 10.8$ Hz, 1H, CHHPh), 4.67 (d, $J = 11.8$ Hz, 1H, CHHPh), 4.29 (d, $J = 8.6$ Hz, 1H, CH-2), 4.23 (dd, $J = 11.6, 8.2$ Hz, 1H, CH-7a), 4.10 (br s, 1H, CH-4), 3.99 (dd, $J = 11.6, 5.8$ Hz, 1H, CH-7b), 3.94 (dd, $J = 7.9, 3.0$ Hz, 1H, CH-6), 3.53 (br s, 1H, OH), 3.45 – 3.36 (m, 2H, CH-1, CH-3), 2.65 (d, $J = 8.0$ Hz, 1H, OH), 2.19 (br s, 1H, CH-5). ¹³C NMR (101 MHz, CDCl₃): δ 160.1 (C=O), 143.5 (C_q Q), 141.4 (q, $J = 35.5$ Hz, CCF₃), 138.0 (C_q Ph), 137.9 (C_q Ph), 135.3, 129.8, 128.6, 128.5, 128.4, 128.3, 128.1, 127.9, 126.9 (10 x CH Ar, 4 x CH Q), 122.8 (C_q Q), 118.1 (q, $J_{CF} = 275.0$ Hz, CF₃), 82.9 (C-3), 77.5 (C-2), 74.8 (CH₂Ph), 71.9 (CH₂Ph), 67.3 (C-4), 61.9 (C-7), 49.1 (C-1), 44.0 (C-6), 41.0 (C-5). HRMS: calcd. for [C₃₀H₂₉F₃N₃O₅]⁺ 568.2059; found 568.2061; calcd. for [C₃₀H₂₈F₃N₃NaO₅]⁺ 590.1879; found 590.1882.

α -D-galactocyclitol CF₃-Q-aziridine (5b): Obtained from cyclohexene **5a** (550 mg, 2.2 mmol) as an orange oil in 61% yield (643 mg, 1.35 mmol). ¹H NMR (400 MHz, CDCl₃): δ 8.18 – 8.09 (m, 1H, CH Q), 7.80 – 7.70 (m, 2H, 2 x CH Q), 7.55 (ddd, $J = 8.3, 6.4, 2.0$ Hz, 1H, CH Q), 7.40 (dq, $J = 5.2, 2.9$ Hz, 2H, 2 x CH Ar), 7.34 – 7.23 (m, 3H, 3 x CH Ar), 5.39 (s, 1H, CHPh), 4.37 – 4.23 (m, 2H, CH-2, CH-7a), 4.14 – 4.03 (m, 2H, CH-4, CH-7b), 3.90 (dd, $J = 7.7, 4.1$ Hz, 1H, CH-1), 3.68 – 3.57 (m, 2H, CH-3, CH-4), 2.08 (s, 1H, CH-5). ¹³C NMR (101 MHz, CDCl₃): δ 160.3 (C=O), 143.6 (C_q Q), 141.8 (q, $J = 36.0$ Hz, CCF₃), 137.9 (C_q Ph), 135.0, 129.5, 129.0, 128.4, 128.1, 126.6, 126.2 (5 x CH Ar, 5 x CH Q), 122.4 (C_q Q), 114.0 (q, $J_{CF} = 277.8$ Hz, CF₃), 101.1 (CHPh), 77.5

(C-4), 72.9 (C-3), 69.9 (C-7), 68.2 (C-2), 47.7 (C-6), 46.5 (C-1), 34.3 (C-5). HRMS: calcd. for [C₂₃H₂₁F₃N₃O₅]⁺ 476.14333; found 476.14255.

β -D-conduritol CF₃-Q-aziridine (6b): Obtained from cyclohexene **6a** (300 mg, 0.72 mmol) as an orange oil in 66% yield (305 mg, 0.47 mmol). ¹H NMR (400 MHz, CDCl₃): δ 8.17 (dd, $J = 7.8, 1.2$ Hz, 1H, CH Q), 7.92 – 7.79 (m, 2H, 2 x CH Q), 7.61 (ddd, $J = 8.3, 5.2, 3.2$ Hz, 1H, CH Q), 7.45 – 7.28 (m, 12H, 12 x CH Ar), 7.28 – 7.14 (m, 2H, 2 x CH Ar), 7.07 – 6.96 (m, 1H, CH Ar), 4.97 (d, $J = 11.3$ Hz, 1H, CHHPh), 4.90 – 4.81 (m, 2H, 2 x CHHPh), 4.79 (d, $J = 11.2$ Hz, 1H, CHHPh), 4.71 (dd, $J = 11.3, 6.6$ Hz, 2H, 2 x CHHPh), 4.41 (dd, $J = 7.2, 3.7$ Hz, 1H, CH-6), 3.99 (td, $J = 7.0, 6.1, 2.9$ Hz, 2H, CH-3, CH-5), 3.93 (d, $J = 7.2$ Hz, 1H, CH-1), 3.55 – 3.42 (m, 2H, CH-2, CH-4), 2.59 (br s, 1H, OH). ¹³C NMR (101 MHz, CDCl₃): δ 161.0 (C=O), 143.5 (q, $J = 34.9$ Hz, CCF₃), 142.9, 138.6, 138.5 (3 x C_q Ph), 137.6, 135.2, 129.5, 128.7, 128.6, 128.5, 128.3, 128.1, 128.0, 127.8, 126.6 (15 x CH Ar, 4 x CH Q), 123.2 (C_q Q), 117.3 (q, $J_{CF} = 277.6$ Hz, CF₃), 83.6, 80.2, 79.8 (3 x CH), 75.8, 75.7, 73.9 (3 x CH₂Ph), 71.4 (CH), 42.6 (CH-6), 40.6 (CH-1). HRMS: calcd. for [C₃₆H₃₃F₃N₃O₅]⁺ 644.2372; found 644.2368; calcd. for [C₃₆H₃₂F₃N₃NaO₅]⁺ 666.2192; found 666.2189.

β -D-xylo-cyclitol CF₃-Q-aziridine (8b): Obtained from cyclohexene **8a** (300 mg, 0.97 mmol) as an orange oil in 45% yield (233 mg, 0.43 mmol) and 37% starting material recovered (112 mg, 0.36 mmol). ¹H NMR (400 MHz, CDCl₃): δ 8.19 (dt, $J = 8.0, 1.1$ Hz, 1H, CH Q), 7.88 – 7.81 (m, 2H, 2 x CH Q), 7.65 – 7.58 (m, 1H, CH Q), 7.43 – 7.29 (m, 7H, 7 x CH Ar), 7.27 – 7.19 (m, 2H, 2 x CH Ar), 7.07 – 7.00 (m, 1H, CH Ar), 4.99 (d, $J = 11.2$ Hz, 1H, CHHPh), 4.80 (d, $J = 2.1$ Hz, 2H, 2 x CHHPh), 4.67 (d, $J = 11.2$ Hz, 1H, CHHPh), 4.03 – 3.95 (m, 2H, CH-2, CH-6), 3.77 (d, $J = 7.2$ Hz, 1H, CH-1), 3.65 (td, $J = 10.5, 5.4$ Hz, 1H, CH-4), 3.30 (dd, $J = 10.0, 8.2$ Hz, 1H, CH-3), 2.65 – 2.54 (m, 2H, CH-5a, OH), 1.74 (ddd, $J = 14.2, 10.9, 3.5$ Hz, 1H, CH-5b). ¹³C NMR (101 MHz, CDCl₃): δ 160.8 (C=O), 144.1 (C_q Q), 143.5 (q, $J = 36.4$ Hz, CCF₃), 138.4, 137.7 (2 x C_q Ph), 135.0, 129.4, 128.7, 128.7, 128.5, 128.1, 128.0, 126.6 (10 x CH Ar, 4 x CH Q), 123.2 (C_q Q), 118.3 (q, $J_{CF} = 275.4$ Hz, CF₃), 84.63 (C-3), 80.4 (C-2), 75.1 (CH₂Ph), 73.0 (CH₂Ph), 65.7 (C-4), 41.4, 41.1 (C-1, C-6), 30.6 (C-5). HRMS: calcd. for [C₂₉H₂₇F₃N₃O₄]⁺ 538.1954; found 538.1960; calcd. for [C₂₉H₂₆F₃N₃NaO₄]⁺ 560.1773; found 560.1779.

α -L-ido-cyclitol CF₃-Q-aziridine (10b): Obtained from cyclohexene **10a** (2.9 g, 5.57 mmol) as an orange oil in 43% yield (1.79 g, 2.40 mmol). $[\alpha]_D^{20} = +1.2$ ($c = 1$, CHCl₃). ¹H NMR (400 MHz, CDCl₃): δ = 8.25 (d, $J = 7.9$ Hz, 1H, CH Ar), 7.87 (d, $J = 3.6$ Hz, 2H, 2 x CH Ar), 7.67 – 7.63 (m, 1H, CH Ar), 7.49 – 7.25 (m, 19H, 19 x CH Ar), 7.14 (t, $J = 7.4$ Hz, 1H, CH Ar), 4.87 (d, $J = 11.1$ Hz, 1H, CHHPh), 4.84 (s, 2H, 2 x CHHPh), 4.78 (d, $J = 11.1$ Hz, 1H, CHHPh), 4.74 (d, $J = 11.6$ Hz, 1H, CHHPh), 4.67 – 4.57 (m, 3H, 3 x CHHPh), 4.22 (d, $J = 7.3$ Hz, 1H, CH-6), 4.06 (d, $J = 7.0$ Hz, 1H, CH-3), 4.00 (d, $J = 7.4$ Hz, 1H, CH-1), 3.98 – 3.96 (m, 1H, CHHPh), 3.78 – 3.69 (m, 3H, CH-2, CH-4, CHHPh), 2.99 – 2.93 (m, 1H, CH-5). ¹³C NMR (101 MHz, CDCl₃): δ = 160.7 (C=O), 144.0 (C_q Ar Q), 143.4 (q, $J = 34.1$ Hz, CCF₃), 139.0, 138.6, 138.3, 138.1 (C_q Ar), 134.9, 129.3, 128.6, 128.5, 128.4, 128.0, 127.9, 127.8, 127.7, 127.6, 126.6 (20 x CH Ar, 4 x CH Q), 123.2 (C_q Q), 118.2 (q, $J = 276.68$ Hz, CF₃), 80.7 (C-4/2), 80.2 (C-3), 76.0 (C-4/2), 75.2, 73.4, 73.2, 73.1 (4 x CH₂Ph), 67.6 (C-7), 44.2 (C-6), 41.2 (C-1), 38.0 (C-5). HRMS: calcd. for [C₄₄H₄₀F₃N₃O₅]⁺

FULL PAPER

748.29983; found 748.29901. HRMS: calcd. for $[C_{44}H_{40}F_3N_3NaO_5]^+$ 770.28123; found 770.28076. Data in agreement with those previously reported.^[15]

General procedure for aziridine deprotection: Ammonia (10 mL/mmol of starting material) was condensed at -60 °C. Lithium (15 equiv.) was added and the mixture was stirred until the lithium was completely dissolved and a bright blue solution was observed. Then, a solution of protected aziridine dissolved in anhydrous THF (5 mL/mmol) was added dropwise. The reaction mixture was stirred for 1 h at -60 °C and subsequently quenched with MeOH and milliQ-H₂O. The solution was allowed to warm to room temperature and stirred until the ammonia was evaporated. The reaction crude was concentrated *in vacuo*, redissolved in MilliQ-H₂O and filtered to remove orange solid impurities from CF₃-Q. The filtrate was neutralized with Amberlite IR-120 H⁺ and the aziridine product bound to the resin was washed with water (3 times) to remove Li salts and subsequently eluted with an aqueous 1M NH₄OH solution, and concentrated under reduced pressure to afford the fully deprotected aziridine.

β-D-glucocyclitol aziridine (1c): Obtained from aziridine **1b** (479 mg, 0.84 mmol) as an oil in 99% yield (146 mg, 0.83 mmol). ¹H NMR (400 MHz, D₂O): δ 3.94 (dd, *J* = 10.9, 4.1 Hz, 1H, CH-7a), 3.70 – 3.58 (m, 2H, CH-7b, CH-2), 3.26 (dd, *J* = 10.2, 8.6 Hz, 1H, CH-3), 3.02 (t, *J* = 10.1 Hz, 1H, CH-4), 2.57 (dd, *J* = 6.1, 3.4 Hz, 1H, CH-6), 2.29 (d, *J* = 6.2 Hz, 1H, CH-1), 2.00 (m, 1H, CH-5). ¹³C NMR (101 MHz, D₂O): δ 76.9 (C-3), 72.1 (C-2), 67.7 (C-4), 61.8 (C-7), 43.1 (C-5), 34.4 (C-1), 32.5 (C-6). Data in agreement with those previously reported.^[23]

α-D-glucocyclitol aziridine (2c): Obtained from aziridine **2b** (104 mg, 0.22 mmol) as an oil in 96% yield (37 mg, 0.21 mmol). ¹H NMR (400 MHz, D₂O): δ 3.87 – 3.77 (m, 2H), 3.69 (dd, *J* = 11.1, 6.4 Hz, 1H), 3.28 (dd, *J* = 10.3, 8.6 Hz, 1H), 3.19 (t, *J* = 10.1 Hz, 1H), 2.52 (dd, *J* = 6.4, 3.6 Hz, 1H), 2.30 (d, *J* = 6.4 Hz, 1H), 1.86 – 1.78 (m, 1H). Data in agreement with those previously reported.^[24]

β-D-galactocyclitol aziridine (4c): Obtained from aziridine **4b** (100 mg, 0.18 mmol) as an orange oil in 81% yield (25 mg, 0.14 mmol). ¹H NMR (500 MHz, D₂O): δ 3.95 (d, *J* = 8.4 Hz, 1H, CH-2), 3.83 (dd, *J* = 7.3, 1.0 Hz, 3H, CH₂OH, CH-4), 3.40 (dd, *J* = 8.3, 2.4 Hz, 1H, CH-3), 2.44 – 2.36 (m, 1H, CH-1), 2.35 – 2.27 (m, 1H, CH-6), 2.22 – 2.11 (m, 1H, CH-5). ¹³C NMR (126 MHz, D₂O): δ 76.0 (C-3), 70.4 (C-2), 70.1 (C-4), 61.2 (CH₂), 39.2 (C-5), 34.4 (C-6), 31.9 (C-1). Data in agreement with those previously reported.^[9]

α-D-galactocyclitol aziridine (5c): Obtained from aziridine **5b** (622 mg, 1.31 mmol) as an oil in 85% yield (195 mg, 1.11 mmol). ¹H NMR (400 MHz, D₂O): δ 4.05 (dd, *J* = 9.0, 4.3 Hz, 1H, CH-2), 3.82 (dt, *J* = 3.2, 1.6 Hz, 1H, CH-4), 3.77 – 3.65 (m, 2H, CH-7a,b), 3.32 (dd, *J* = 9.0, 1.9 Hz, 1H, CH-3), 2.57 (dd, *J* = 6.4, 4.2 Hz, 1H, CH-1), 2.12 (dt, *J* = 6.3, 1.3 Hz, 1H, CH-6), 1.93 (tdd, *J* = 7.5, 3.2, 1.3 Hz, 1H, CH-5). ¹³C NMR (126 MHz, D₂O): δ 72.6 (C-3), 71.4 (C-4), 69.0 (C-2), 61.5 (C-7), 42.6 (C-5), 35.3 (C-1), 31.4 (C-6). Data in agreement with those previously reported.^[5b]

Conduritol aziridine (6c): Obtained from aziridine **6b** (100 mg, 0.16 mmol) as an oil in 99% yield (25 mg, 0.16 mmol). ¹H NMR (500 MHz, D₂O): δ 3.90 – 3.84 (m, 1H, CH-3), 3.71 – 3.67 (m, 1H, CH-5), 3.25 – 3.18 (m, 2H, CH-2, CH-4), 2.61 (dd, *J* = 6.2, 3.6 Hz, 1H, CH-6), 2.34 (d, *J* = 6.2 Hz, 1H, CH-1). ¹³C NMR (126 MHz, D₂O): δ 78.2 (C-2/4), 74.7 (C-5), 74.1 (C-3), 73.5 (C-2/4), 37.9 (C-

6), 37.4 (C-1). Data in agreement with those previously reported.^[25]

β-D-xylo-cyclitol aziridine (8c): Obtained from aziridine **8b** (115 mg, 0.21 mmol) as an oil in 96% yield (30 mg, 0.20 mmol). ¹H NMR (400 MHz, D₂O): δ 3.61 (d, *J* = 8.3 Hz, 1H, CH-2), 3.34 (td, 1H, *J* = 10.5, 5.4 Hz, CH-4), 3.15 (dd, *J* = 10.3, 8.4 Hz, 1H, CH-3), 2.42 (br s, 1H, CH-6), 2.31 (ddd, *J* = 13.8, 5.4, 1.6 Hz, 1H, CH-5a), 2.17 (d, *J* = 6.1 Hz, 1H, CH-1), 1.66 (ddd, *J* = 14.0, 10.7, 3.3 Hz, 1H, CH-5b). ¹³C NMR (101 MHz, D₂O): δ 77.5 (C-3), 72.6 (C-2), 66.2 (C-4), 34.5 (C-1), 31.3 (C-5), 30.8 (C-6). Data in agreement with those previously reported.^[8b]

α-L-ido-cyclitol aziridine (10c): Obtained from aziridine **10b** (1.79 g, 2.39 mmol) as an oil in 93% yield (389 mg, 2.22 mmol). ¹H NMR (400 MHz, D₂O): δ 2.47 (dd, *J* = 11.2, 4.4 Hz, 1H, CH-7a), 2.28 – 2.17 (m, 2H, CH-7b, CH-3), 2.11 (dd, *J* = 10.5, 5.6 Hz, 1H, CH-4), 1.98 (dd, *J* = 10.5, 7.4 Hz, 1H, CH-2), 1.07 (d, *J* = 6.0 Hz, 1H, CH-6), 1.01 (dt, *J* = 5.8, 5.1 Hz, 1H, CH-5), 0.77 (d, *J* = 6.0 Hz, 1H, CH-1). ¹³C NMR (101 MHz, D₂O): δ = 76.0 (C-2), 75.2 (C-3), 70.1 (C-4), 62.2 (C-7), 43.5 (C-5), 36.9 (C-1), 36.3 (C-6). HRMS: calcd. for $[C_7H_{14}NO_4]^+$ 176.09228; found 176.09175. Data in agreement with those previously reported.^[15]

Acknowledgements

We thank The Netherlands Organization for Scientific Research (NWO-CW, ChemThem grant to J.M.F.G.A. and H.S.O.), the European Research Council (ERC-2011-AdG290836 “Chembiosphing”, to H.S.O.) and Sanofi Genzyme (research grant to J.M.F.G.A. and H.S.O. and postdoctoral contract to M.A.) for financial support.

Conflict of interest

The authors declare no conflict of interest.

Keywords: stereoselective aziridination • cyclophellitol aziridines • glycosidases • 3-aminoquinazolin-4(3*H*)-ones • irreversible glycosidase inhibitors

[1] Varki, A.; Cummings, R. D.; Esko, J. D.; Stanley, P.; Hart, G. W.; Aebi, M.; Darvill, A. G.; Kinoshita, T.; Packer, N. H.; Prestegard, J. H.; Schnaar, R. L.; Seeberger, P. H. *Essentials of Glycobiology*, 3rd Edition; Cold Spring Harbor (NY): Cold Spring Harbor Laboratory Press; 2015-2017.

[2] D. E. Koshland, *Biol. Rev.* **1953**, 28, 416-436.

[3] A. Ardevol and C. Rovira, *J. Am. Chem. Soc.* **2015**, 137, 7528-7547.

[4] a) I. Z. Breen, M. Artola, L. Wu, T. J. M. Beenakker, W. A. Offen, J. M. F. G. Aerts, G. J. Davies and H. S. Overkleeft, (January 2018) Competitive and Covalent Inhibitors of Human Lysosomal Retaining Exoglycosidases. In: eLS. John Wiley & Sons, Ltd: Chichester. DOI: 2010.1002/9780470015902.a9780470027591; b) M. Artola, L. Wu, M. J. Ferraz, C. L. Kuo, L. Raich, I. Z. Breen, W. A. Offen, J. D. C. Codee, G. A. van der Marel, C. Rovira, Jmfg Aerts, G. J. Davies and H. S. Overkleeft, *ACS Cent. Sci.* **2017**, 3, 784-793.


[5] a) W. W. Kallemeijn, K. Y. Li, M. D. Witte, A. R. Marques, J. Aten, S. Scheij, J. Jiang, L. I. Willems, T. M. Voorn-Brouwer, C. P. van Roomen, R. Ottenhoff, R. G. Boot, H. van den Elst, M. T. Walvoort, B. I. Florea, J. D. Codée, G. A. van der Marel, J. M. Aerts and H. S. Overkleeft, *Angew. Chem. Int. Ed.* **2012**, 51, 12529-12533. *Angew. Chem.* **2012**, 124, 12697-12701; b) L. I. Willems, T. J. Beenakker, B. Murray, S. Scheij, W. W. Kallemeijn,

FULL PAPER

- R. G. Boot, M. Verhoek, W. E. Donker-Koopman, M. J. Ferraz, E. R. van Rijssel, B. I. Florea, J. D. Codee, G. A. van der Marel, J. M. Aerts and H. S. Overkleeft, *J. Am. Chem. Soc.* **2014**, 136, 11622-11625; c) J. Jiang, C. L. Kuo, L. Wu, C. Franke, W. W. Kallemeijn, B. I. Florea, E. van Meel, G. A. van der Marel, J. D. Codee, R. G. Boot, G. J. Davies, H. S. Overkleeft and J. M. Aerts, *ACS Cent. Sci.* **2016**, 2, 351-358; d) S. P. Schroder, J. W. V. De Sande, W. W. Kallemeijn, C. L. Kuo, M. Artola, E. J. van Rooden, J. B. Jiang, T. J. M. Beenakker, B. I. Florea, W. A. Offen, G. J. Davies, A. J. Minnaard, J. M. F. G. Aerts, J. D. C. Codee, G. A. van der Marel and H. S. Overkleeft, *Chem. Commun.* **2017**, 53, 12528-12531; e) L. Wu, J. B. Jiang, Y. Jin, W. W. Kallemeijn, C. L. Kuo, M. Artola, W. Dai, C. van Elk, M. van Eijk, G. A. van der Marel, J. D. C. Codee, B. I. Florea, J. M. F. G. Aerts, H. S. Overkleeft and G. J. Davies, *Nat. Chem. Biol.* **2017**, 13, 867-873.
- [6] J. B. Jiang, M. Artola, T. J. M. Beenakker, S. P. Schroder, R. Petracca, C. de Boer, J. M. F. G. Aerts, G. A. van der Marel, J. D. C. Codee and H. S. Overkleeft, *Eur. J. Org. Chem.* **2016**, 3671-3678.
- [7] L. Degennaro, P. Trinchera and R. Luisi, *Chem. Rev.* **2014**, 114, 7881-7929.
- [8] a) J. L. Jat, M. P. Paudyal, H. Y. Gao, Q. L. Xu, M. Yousufuddin, D. Devarajan, D. H. Ess, L. Kurti and J. R. Falck, *Science* **2014**, 343, 61-65; b) S. P. Schroder, R. Petracca, H. Minnee, M. Artola, J. M. F. G. Aerts, J. D. C. Codee, G. A. van der Marel and H. S. Overkleeft, *Eur. J. Org. Chem.* **2016**, 4787-4794.
- [9] L. I. Willems, T. J. M. Beenakker, M. Murray, B. Gagestein, H. van den Elst, E. R. van Rijssel, J. D. C. Codee, W. W. Kallemeijn, J. M. F. G. Aerts, G. A. van der Marel and H. S. Overkleeft, *Eur. J. Org. Chem.* **2014**, 6044-6056.
- [10] R. S. Atkinson and C. W. Rees, *J. Chem. Soc. (C)* **1969**, 772-778.
- [11] D. J. Anderson, T. L. Gilchrist, D. C. Horwell and C. W. Rees, *J. Chem. Soc. (C)* **1970**, 0, 576-582.
- [12] a) R. S. Atkinson, M. P. Coogan and C. L. Cornell, *J. Chem. Soc. Chem. Comm.* **1993**, 1215-1216; b) R. S. Atkinson, E. Barker, P. J. Edwards and G. A. Thomson, *J. Chem. Soc. Chem. Comm.* **1995**, 727-728; c) R. S. Atkinson and P. J. Williams, *J. Chem. Soc. Perkin Trans. 2* **1996**, 205-211; d) R. S. Atkinson, M. P. Coogan and I. S. T. Lochrie, *Chem. Commun.* **1996**, 789-790; e) R. S. Atkinson, M. P. Coogan and C. L. Cornell, *J. Chem. Soc. Perkin Trans. 1* **1996**, 157-166; f) R. S. Atkinson, A. P. Ayscough, W. T. Gattrell and T. M. Raynham, *J. Chem. Soc. Perkin Trans. 1* **1998**, 2783-2793.
- [13] R. S. Atkinson, M. J. Grimshire and B. J. Kelly, *Tetrahedron* **1989**, 45, 2875-2886.
- [14] A. Alcaide, A. Trapero, Y. Perez and A. Llebaria, *Org. Biomol. Chem.* **2015**, 13, 5690-5697.
- [15] M. Artola, C. L. Kuo, S. A. McMahon, V. Oehler, T. Hansen, M. van der Lienden, X. He, H. van den Elst, B. I. Florea, A. R. Kermode, T. M. Gloster, G. A. van der Marel, J. D. C. Codee, H. S. Overkleeft and J. M. F. G. Aerts, *Chem. Eur. J.* 10.1002/chem.201804662.
- [16] R. S. Atkinson, M. P. Coogan and C. L. Cornell, *J. Chem. Soc. Perkin Trans. 1* **1996**, 157-166.
- [17] A. Trapero and A. Llebaria, *ACS Med. Chem. Lett.* **2011**, 2, 614-619.
- [18] Y. Harrak, C. M. Barra, A. Delgado, A. R. Castano and A. Llebaria, *J. Am. Chem. Soc.* **2011**, 133, 12079-12084.
- [19] C. Jaramillo, R. F. Delapradilla and M. Martinlomas, *Carbohydr. Res.* **1991**, 209, 296-298.
- [20] S. G. Davies, K. B. Ling, P. M. Roberts, A. J. Russell, J. E. Thomson and P. A. Woods, *Tetrahedron* **2010**, 66, 6806-6813.
- [21] R. S. Atkinson, E. Barker and S. Ulukanli, *J. Chem. Soc. Perkin Trans. 1* **1998**, 583-589.
- [22] F. G. Hansen, E. Bundgaard and R. Madsen, *J. Org. Chem.* **2005**, 70, 10139-10142.
- [23] K. Tatsuta, Y. Niwata, K. Umezawa, K. Toshima and M. Nakata, *J. Antibiot. (Tokyo)* **1991**, 44, 912-914.
- [24] M. Nakata, C. Chong, Y. Niwata, K. Toshima and K. Tatsuta, *J. Antibiot. (Tokyo)* **1993**, 46, 1919-1922.
- [25] G. Caron and S. G. Withers, *Biochem. Biophys. Res. Commun.* **1989**, 163, 495-499.

FULL PAPER

This work describes direct aziridination reactions on differently substituted glycoside configured cyclohexenes with 3-amino-2-(trifluoromethyl)quinazolin-4(3H)-one as nitrogen donor. We show that aziridination can be directed by allylic or homoallylic hydroxyls through H-bonding and that steric hindrance plays an essential role in the diastereoselective outcome of the final cyclitols aziridines.

**Stereoselective aziridination**

Marta Artola,* Shirley Wouters, Sybrin P. Schröder, Casper de Boer, Yurong Chen, Rita Petracca, Adrianus M. C. H. van den Nieuwendijk, Johannes M. F. G. Aerts, Gijsbert A. van der Marel, Jeroen D. C. Codée and Herman S. Overkleef*

Page No. – Page No.

Direct stereoselective aziridination of cyclohexenols with 3-amino-2-(trifluoromethyl)quinazolin-4(3H)-one in the synthesis of cyclitol aziridine glycosidase inhibitors