

Communication

Reactions of HDDA-derived Benzynes with Sulfides: Mechanism, Modes, and Three-component Reactions

Junhua Chen, Vignesh Palani, and Thomas R. Hoye

J. Am. Chem. Soc., Just Accepted Manuscript • DOI: 10.1021/jacs.6b01025 • Publication Date (Web): 24 Mar 2016

Downloaded from http://pubs.acs.org on March 24, 2016

Just Accepted

"Just Accepted" manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides "Just Accepted" as a free service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. "Just Accepted" manuscripts appear in full in PDF format accompanied by an HTML abstract. "Just Accepted" manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are accessible to all readers and citable by the Digital Object Identifier (DOI®). "Just Accepted" is an optional service offered to authors. Therefore, the "Just Accepted" Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the "Just Accepted" Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these "Just Accepted" manuscripts.



Journal of the American Chemical Society is published by the American Chemical Society. 1155 Sixteenth Street N.W., Washington, DC 20036 Published by American Chemical Society. Copyright © American Chemical Society. However, no copyright claim is made to original U.S. Government works, or works produced by employees of any Commonwealth realm Crown government in the course of their duties.

Reactions of HDDA-derived Benzynes with Sulfides: Mechanism, Modes, and Three-component Reactions

Junhua Chen, Vignesh Palani, and Thomas R. Hoye*

Department of Chemistry, 207 Pleasant St. SE, University of Minnesota, Minneapolis, MN 55455

Supporting Information Placeholder

ABSTRACT: We report here reactions of alkyl sulfides with benzynes thermally generated by the hexadehydro-Diels–Alder (HDDA) cycloisomerization. The initially produced 1,3-betaine (*o*-sulfonium/aryl carbanion) undergoes intramolecular proton transfer to generate a more stable S-aryl sulfur ylide. This can react in various manners, including engaging weak acids (HA) in the reaction medium. This can produce transient ion pairs $ArSR_2^+A^-$ that proceed to the products ArSR + RA. When cyclic sulfides are used, A^- opens the ring and is incorporated into the product, an outcome that constitutes a versatile, three-component coupling process.

o-Benzyne (1) is the parent member of one of the most versatile of all reactive intermediates in organic chemistry, largely because of its adaptability as an electrophilic agent capable of capture by many different types of external nucleophiles. Its reaction with sulfide trapping agents, largely by way of ylide intermediates (Figure 1), was studied in a number of laboratories in the period 1962¹-1989² but has lain fallow until the recent report of Xu and coworkers. As they pointed out, "nucleophilic attack of alkyl thioether on benzyne followed by an intramolecular 1,4-proton shift has been almost totally neglected since its debut³."⁴ The majority of those early studies were described in a series of reports from the Nakayama laboratory.^{5a} All of this previous work used simple benzyne derivatives, the vast majority of studies being explored with o-benzyne (1) itself.

$$\begin{bmatrix} & & & & \\ & & & & \\ & & & \\ & & & & & \\ & & & & \\ & & & & \\ & & & & & \\ & & & & \\ & & & & & \\ & & & & & \\ & & & & & \\ & & & & & \\ & & & & & \\ & & & & & \\ & & & & & \\ & & & & & \\ & & & & & \\ & & & & & \\ & & & & & \\ & & & & & \\ & & & & & & \\ & & & & & \\ & & & & & & \\ & & & & & & \\ & & & & & \\ & & & & &$$

Figure 1. Trapping of *o*-benzyne (1) with an alkyl sulfide produces ylide 3, presumably via the zwitterionic betaine 2.

The generation of benzyne intermediates by the thermal cycloisomerization reaction of tethered triynes⁶ via the hexadehydro-Diels–Alder (HDDA) reaction⁷ (e.g., **4** to **8**, Figure 2) produces this important class of reactive species in environments free of the reagents and

byproducts that necessarily accompany all other methods⁸ of benzyne generation. This permits interrogation of inherent benzyne reactivity at a more fundamental level than is often otherwise possible.⁹ In addition, new modes of benzyne trapping reactions^{10,11,12} can be uncovered that would be otherwise incompatible with most if not all of the more traditional methods of benzyne generation. We describe here results of our investigations of the reactivity between HDDA-generated benzyne derivatives and various sulfides. These studies provide both mechanistic insight as well as novel trapping processes, including new three component reactions.^{8f,13}

To probe details of ylide formation, we first explored reactions between the benzyne **8** and thioanisole (Figure 2). Heating tetrayne **4** with PhSCH₃ in the presence of *p*-chlorobenzaldehyde in d_6 -benzene produced the diaryl sulfide **5-h** and epoxide **6** [>60% yield (NMR, see SI)]. Replacing the aldehyde with acetic acid as the additive efficiently produced **5-h** (79% yield following isolation). Both results suggest the intermediacy of ylide **10**; the epoxide would result from a Corey-Chaykovsky-like process⁴ and the outcome with AcOH would arise by protonation of **10-h**₃ to give **11-h**₃ (not shown) and subsequent methylation^{5b} (ca. one equivalent of MeOAc was observed by NMR spectroscopy).

We used labeling studies to probe for the intermediacy of a zwitterion like 9. Tetrayne 4 was heated in C_6D_6



Figure 2. Trapping of HDDA-generated benzyne **8** by thioanisole: epoxidation of *p*-ClPhCHO by ylide $10-h_3$ and alkylation within the sulfonium acetate ion pair $11-d_3$.

with PhSCD₃ and AcOH. This gave, nearly exclusively, the mono-deuterated sulfide **5-d** and dideuterated methyl acetate (**7**, NMR, see SI). Protonation by AcOH could have occurred at the stage of either the initially formed 1,3-zwitterion **9-d**₃ or ylide **10-d**₃. The deuterium labeling pattern of the products **5-d** and **7** clearly indicates the involvement of **11-d**₃ and, importantly, establishes the viability of intramolecular proton transfer within **9** (cf. **2** to **3**). Finally, heating **4**, AcOH, and an equimolar mixture of PhSCH₃ and PhSCD₃ gave nearly equal amounts of **5-h** and **5-d** (as well as AcOCH₃ and AcOCHD₂), suggesting that reversal of formation of adduct **9** is slower than the proton transfer that converts **9** to **10**. This result also argues against a concerted event for conversion of **8** directly to ylide **10** (cf., **1** to **3**).

We next explored HDDA-benzyne trapping with a variety of different types of sulfides (Table 1). These studies demonstrated various modes of trapping processes, some precedented and some not. Consistent with our intent, these results establish the viability of each reaction mode in the context of HDDA-generated benzynes. In each case, benzyne 8 from the symmetrical, etherlinked tetrayne 4 (entries 1-3 and Figure 2) gives a mixture of isomeric products arising from competitive trapping at C6 (major, shown) and C7. Details about the structure assignment of these constitutional isomers and DFT calculations showing the extent of ring distortion of the intermediate benzyne¹⁴ are provided in the SI.

Diallyl sulfide trapped the benzyne **8** to provide the desymmetrized diene **15** (Table 1, entry 1), presumably via **15-int**. This is consistent with the outcome of the reaction of **1** with digeranyl³ or diprenyl^{5c} sulfide. The rearrangement of **15-int** to **15** could, in principle, occur either by a [2,3]-sigmatropic process^{3,15} (green arrows) or by a Stevens rearrangement.¹⁶ We therefore carried out analogous reactions with a propargyl sulfide (entry 2) or dibenzyl sulfide^{5d} (entry 3). The former cleanly gave an allene (**16a** or **16b**), showing that [2,3]-Wittig processes (**16-int**) are viable. The latter gave adduct **17**, the product of a Stevens rearrangement (**17-int**).

Given the considerable generality of the HDDA cascade, we were unsurprised to see that sulfide trapping is not unique to benzyne **8**. Those derived from the tetraynes **12–14** participate as well (entries 4–6). In those instances, we happened to use various cyclic sulfides as the trapping agents. Either thiirane or tetrahydrothiophene gave rise to the same product, the vinyl sulfide **18** (entry 4), likely via the fragmentations shown in **18-int** and **18'-int**, respectively. Similar modes of ring-cleavage have been reported.^{2,5e-f,17} In contrast, use of thietane (entry 5) gave rise to the ring-contracted cyclopropane **19**—an apparently unprecedented mode of reaction. Finally, *N*-methylbenzothiazoline (in the presence of acetic acid) gave the aminophenyl sulfide **20** (entry 6), again a new type of trapping reaction.

We then explored the use of cyclic sulfides in the presence of various protic nucleophiles (H–Nu) with the

Table 1. Various modes of trapping of HDDA-generatedbenzynes by sulfides^a



^a Reaction conditions: Substrate tetrayne in PhH (0.1 M) containing the indicated sulfide (1.5-2 equiv) was heated in a sealed culture tube (see SI for reaction temperatures and times). For entry 5, five equiv of o-NO₂PhCH₂CO₂Me was present. For entry 6, five equiv of AcOH was present. ^b R = 1-propynyl. ^c Benzyne **8** (from **4**) is trapped by sulfides competitively at C6 and C7 to give separable major and minor (yield in parentheses) isomers, respectively.

goal of expanding this chemistry into a three-component process.¹³ We are aware of one report in which such a reaction has been described: namely, generation of *o*-benzyne in the presence of (three) cyclic sulfides and aqueous hydrochloric acid produced chloroalkyl sulfides (e.g., 4-chlorobutyl phenyl sulfide from tetrahydrothiophene).^{5c} The HDDA version, via the ion pair **22**, is portrayed generically in the graphic at the top of Table 2. In the products **23**, the tetrayne-derived fragment and nucleophile are linked in a fashion dictated by the structure

of the sulfide used. This structural motif is relevant to the field of drug discovery.¹⁸

The reaction is general with respect to the type of benzyne used (see 21a-g). Two (21d and 21e) represent a previously unreported type of HDDA substrate. It is notable that HDDA cycloisomerization leading to 21e occurred faster (ca. 5x) than an analog lacking the geminal methyl substituents, presumably a reflection of the bond angle compression that brings the proximal termini of the diyne and diynophile closer to one another.^{12c}

Results of three-component coupling reactions are listed in Table 2. Some highlights are: (i) carbon-, nitro-

gen- and oxygen-centered nucleophiles, including *N*heterocycles (entries 7 and 9), will all participate; (ii) hindered (entry 5) or relatively weakly acidic H–Nu species (Brønsted acids) are functional; (iii) the unsymmetrical sulfonium ion intermediate in entry 6 opens selectively at the allylic center; (iv) additional functionality can be tolerated within the cyclic thioether (entries 6 and 8); (v) a number of these nucleophiles are capable of trapping HDDA benzyne directly—the presence of the polarizable sulfide nucleophile outpaces those processes. One limitation is that thiirane has not been an effective participant in the 3-component reaction to date—

Table 2. Three-component reactions of HDDA benzynes (21, green), cyclic sulfides (blue), and H-nucleophiles (red)



 $^{a}R = 1$ -propynyl.

apparently fragmentation to the vinyl sulfide **18** is so rapid^{5e} that it supersedes protonation by the external protic nucleophile. All told, there is considerable generality to this process. The preparation of arrays of products from any one benzyne precursor, itself selected from among multiple options, can be envisioned.

In conclusion, we have (i) presented evidence supporting the involvement of initial 1,3-zwitterions in reactions of aliphatic sulfides with benzynes (Figure 2), (ii) demonstrated the viability of several new modes of trapping reaction (Table 1), (iii) presented two new types of HDDA-benzynes (**21d** and **21e**, Table 2), and (iv) shown that three-component reactions that engage HDDAbenzynes, cyclic sulfides, and a protic nucleophile have considerable generality.

ASSOCIATED CONTENT

Supporting Information. The Supporting Information is available free of charge on the ACS Publications website.

Experimental details for the preparation of new compounds; spectroscopic data for their characterization, including copies of 1 H and 13 C NMR spectra; and computed (DFT) geometries of benzyne intermediates and transition state for ylide formation (PDF).

AUTHOR INFORMATION

Corresponding Author

* hoye@umn.edu

Notes

The authors declare to have no competing financial interests.

ACKNOWLEDGMENT

Support for this research was provided by the National Institutes of Health (GM65597). V.P. was supported by a Gleysteen-Heisig fellowship for undergraduate students. NMR spectra were recorded using instrumentation purchased with funds provided by the NIH Shared Instrumentation Grant program (S10OD011952). We thank Sean Ross and Juntian Zhang for providing samples of and procedures for preparing tetraynes **13** and precursor to **21c**.

REFERENCES

(1) (a) Hellmann, H.; Eberle, D. Liebigs Ann. Chem. **1963**, 662, 188–201. (b) Franzen, V.; Joschek, H.-I.; Mertz, C. Liebigs Ann. Chem. **1962**, 654, 82–91.

(2) Nakayama, J.; Kumano, Y.; Hoshino, M. Tetrahedron Lett. 1989, 30, 847-850.

(3) Blackburn, G. M.; Ollis, W. D. Chem. Commun. 1968, 1261–1262.

(4) Xu, H.; Cai, M.; He, W.; Hu, W.; Shen, M. RSC Adv. 2014, 4, 7623–7626.

(5) (a) Summarized (pp 400–406) in Nakayama, J. J. Sulfur Chem.
2009, 30, 393–468. (b) Nakayama, J.; Fujita, T.; Hoshino, M. Chem.
Lett. 1982, 1777–1780. (c) Nakayama, J.; Hoshino, K.; Hoshino, M.
Chem Lett. 1985, 677–678. (d) Iwamura, H.; Iwamura, M.; Nishida, T.; Yoshida, M.; Nakayama, J. Tetrahedron Lett. 1971, 12, 63–66. (e) Nakayama, J.; Takeue, S.; Hoshino, M. Tetrahedron Lett. 1984, 25, 2679–2682. (f) Nakayama, J.; Ozasa, H.; Hoshino, M. Heterocycles 1984, 22, 1053–1056.

(6) (a) Bradley, A. Z.; Johnson, R. P. J. Am. Chem. Soc. 1997, 119, 9917-9918.
(b) Miyawaki, K.; Suzuki, R.; Kawano, T.; Ueda, I. Tetrahedron Lett. 1997, 38, 3943-3946.

(7) (a) Hoye, T. R.; Baire, B.; Niu, D. W.; Willoughby, P. H.; Woods, B. P. *Nature* **2012**, *490*, 208-212. (b) Baire, B.; Niu, D.; Willoughby, P. H.; Woods, B. P.; Hoye, T. R. *Nature Protocols* **2013**, *8*, 501–508.

(8) For some recent reviews and commentary on aryne chemistry:
(a) Holden, C.; Greaney, M. F. Angew. Chem Int. Ed. 2014, 53, 5746–5749.
(b) Dubrovskiy, A. V.; Markina, N. A.; Larock, R. C. Org. Biomol. Chem. 2013, 11, 191.
(c) Wu, C.; Shi, F. Asian J. Org. Chem. 2013, 2, 116.
(d) Perez, D.; Peña, D.; Guitian, E. Eur. J. Org. Chem. 2013, 5981.
(e) Tadross, P. M.; Stoltz, B. M. Chem. Rev. 2012, 112, 3550–3577.
(f) Gampe, C. M.; Carreira, E. M. Angew. Chem., Int. Ed. 2012, 51, 3766–3778.
(g) Bhunia, A.; Yetra, S. R.; Biju, A. T. Chem. Soc. Rev. 2012, 41, 3140.
(h) Okuma, K. Heterocycles 2012, 85, 515.
(i) Yoshida, H.; Ohshita, J.; Kunai, A. Bull. Chem. Soc. Jpn. 2010, 83, 199.
(j) Kitamura, T. Aust. J. Chem. 2010, 63, 987–1001.

(9) (a) Niu, D.; Willoughby, P. H.; Baire, B.; Woods, B. P.; Hoye, T. R. *Nature* **2013**, *501*, 531–534. (b) Hoye, T. R.; Baire, B.; Wang, T. *Chem. Sci.* **2014**, *5*, 545–550. (c) Niu, D.; Wang, T.; Woods, B. P.; Hoye, T. R. *Org. Lett.* **2014**, *16*, 254–257. (d) Willoughby, P. H.; Niu, D.; Wang, T.; Haj, M. K.; Cramer, C. J.; Hoye, T. R. *J. Am. Chem. Soc.* **2014**, *136*, 13657–13665. (e) Marell, D. J.; Furan, L. R.; Woods, B. R.; Lei, X.; Bendelsmith, A. J.; Cramer, C. J.; Hoye, T. R.; Kuwata, K. T. *J. Org. Chem.* **2015**, *80*, 11744–11754.

(10) (a) Yun, S. Y.; Wang, K.; Lee, N.; Mamidipalli, P.; Lee D. J. Am. Chem. Soc. **2013**, 135, 4668–4671. (b) Karmakar, R.; Mamidipalli, P.; Yun, S. Y.; Lee, D. Org. Lett. **2013**, 15, 1938–1941. (c) Wang, K.; Yun, S. Y.; Mamidipalli, P.; Lee, D. Chem. Sci. **2013**, 4, 3205–3211. (d) Mamidipalli, P.; Yun, S. Y.; Wang, K.; Zhou, T.; Xia, Y.; Lee, D. Chem. Sci. **2014**, 5, 2362-2367. (e) Lee, N.; Yun, S. Y.; Mamidipalli, P.; Salzman, R. M. Lee, D.; Zhou, T.; Xia, Y. J. Am. Chem. Soc. **2014**, 136, 4363–4368. (f) Karmakar, R.; Yun, S. Y.; Wang, K.; Lee, D. Org. Lett. **2014**, 16, 6-9. (g) Karmakar, R.; Yun, S. Y.; Chen, J.; Xia, Y.; Lee, D. Angew. Chem. Int. Ed. **2015**, 54, 6582-6586. (h) Karmakar, K.; Ghorai, S.; Xia, Y.; Lee, D. Molecules **2015**, 20, 15862-15880.

(11) (a) Vandavasi, J. K.; Hu, W.; Hsiao, C.; Senadia, G. C.; Wang, J. J. *RSC Adv.* **2014**, *4*, 57547–57552. (b) Liang, Y.; Hong, X.; Yu, P.; Houk, K. N. *Org. Lett.* **2014**, *16*, 5702–5705. (c) Nobusue, S.; Yamane, H.; Miyoshi, H.; Tobe, Y. *Org. Lett.* **2014**, *16*, 1940–1943. (d) Kerisit, N.; Toupet, L.; Larini, P.; Perrin, L.; Guillemin, J.; Trolez, Y. *Chem. Eur. J.* **2015**, *21*, 6042–6047. (e) Watanabe, T.; Curran, D. P.; Taniguchi, T. *Org. Lett.* **2015**, *17*, 3450–3453.

(12) (a) Niu, D.; Hoye, T. R. Nature Chem. 2014, 6, 34–40. (b) Chen, J.; Baire, B.; Hoye, T. R. Heterocycles 2014, 88, 1191–1200.
(c) Woods, B. P.; Baire, B.; Hoye, T. R. Org. Lett. 2014, 16, 4578–4581. (d) Woods, B. P.; Hoye, T. R. Org. Lett. 2014, 16, 6370–6373.
(e) Pogula, V. D.; Wang, T.; Hoye, T. R. Org. Lett. 2015, 17, 856–859. (f) Luu Nguyen, Q.; Baire, B.; Hoye, T. R. Tetrahedron Lett. 2015, 56, 3265–3267.

(13) Roy, T.; Baviskar, D. R.; Biju, A. T. J. Org. Chem. **2015**, 80, 11131–11137 and references therein.

(14) (a) Hamura, T.; Ibusuki, Y.; Sato, K.; Matsumoto, T.; Osamura, Y.; Suzuki, K. Org. Lett. 2003, 5, 3551–3554. (b) Cheong, P. H. Y.; Paton, R. S.; Bronner, S. M.; Im, G.-Y. J.; Garg, N. K.; Houk, K. N. J. Am. Chem. Soc. 2010, 132, 1267–1269. (c) Garr, A. N.; Luo, D.; Brown, N.; Cramer, C. J.; Buszek, K. R.; VanderVelde, D. Org. Lett. 2010, 12, 96–99.

(15) Baldwin, J. E.; Hackler, R. E. J. Am. Chem. Soc. 1969, 91, 3646-3647.

(16) Bhakat, S. J. Chem. Pharm. Res. 2011, 3, 115-121.

(17) Brewer, J. P. N.; Heaney, H.; Jablonski, J. M. Tetrahedron Lett. **1968**, *9*, 4455–4456.

(18) E.g., the linked sulfide F15845 has been in Phase I and II clinical trials for treatment of angina: (a) Le Grand, B.; Pignier, C.; Létienne, R.; Cuisiat, F.; Rolland, F.; Mas, A.; Vacher, B. *J. Med.*

Chem. **2008**, *51*, 3856–3866. (b) Pignier, C.; Rougier, J.-S.; Vié, B.; Culié, C.; Verscheure, Y.; Vacher, B.; Abriel, H.; Le Grand, B. *Brit. J. Pharmacol.* **2010**, *161*, 79–91.





