

Tetrahedron 54 (1998) 10999-11010

TETRAHEDRON

# Double Michael Addition Reaction of Oxophorone and Its Derivatives Leading to Bicyclo[2.2.2]octane Compounds

# Hisahiro Hagiwara,<sup>\*a</sup> Yu Yamada,<sup>a</sup> Hitoshi Sakai,<sup>a</sup> Toshio Suzuki,<sup>a</sup> and Masayoshi Ando<sup>b</sup>

Graduate School of Science and Technology,<sup>a</sup> and Faculty of Engineering,<sup>b</sup> Niigata University, 8050, 2 No-cho, Ikarashi, Niigata 950-21, Japan

Received 25 May 1998; accepted 29 June 1998

Abstract: Trimethylsilylenol ethers of oxophorone 1 and its derivatives provided bicyclo[2.2.2]octane compounds 5 in a selective way. © 1998 Elsevier Science Ltd. All rights reserved.

Bicyclo[2.2.2]- or a bicyclo[3.2.1] octane system forms an integral part of several natural products.<sup>1</sup> Normally, bicyclo[2.2.2] octane framework is derived through a successive double Michael addition reaction involving the kinetic enolate of 2-cyclohexenone and an  $\alpha,\beta$ -unsaturated carbonyl compound. This procedure has turned out as a popular synthetic tool to prepare such bicyclic systems,<sup>2</sup> since an alternative procedure employing the Diels-Alder reaction involving cyclohexadiene and an electron deficient dienophile sometimes lacks regio as well as endo/exo selectivities.<sup>3</sup>

As a part of our ongoing research program directed towards the synthesis of polycyclic molecules via a single pot operation by employing nucleophilic domino reactions,<sup>4</sup> we investigated the reaction of trimethylsilylenol ether of oxophorone 1<sup>5</sup> and its derivatives with  $\alpha$ , $\beta$ -unsaturated carbonyl compounds 3 (Scheme 1).



0040-4020/98/\$ - see front matter © 1998 Elsevier Science Ltd. All rights reserved. PII: S0040-4020(98)00646-2 In this direction, we wish to report herein, the synthesis of bicyclo[2.2.2] octane derivatives formed via a double Michael addition reaction of 1 with the respective  $\alpha,\beta$ -unsaturated compounds 3. The advantage of the present reported procedure lies in the fact that only bicyclo[2.2.2] octane derivatives 5 were obtained selectively though there were two options available for the intermediate adduct 4 before it undergoes the successive second addition reaction to provide either 5 or 6 (path a or b).

To start with, the kinetically derived enolate of oxophorone 1 was reacted with methyl acrylate 7 to provide the bicyclo[2.2.2]octane derivative 8 in 23% yield (Table 1, entry 1). No bicyclo[3.2.1]octane derivative 6 was formed under the reaction conditions employed. All attempts to optimize the yield of 8 under the basic conditions employed proved futile in providing any better results and hence the reaction was repeated by following the reaction conditions reported by Fukumoto and Ihara.<sup>6</sup> It was observed that under the reaction conditions, the reaction proceeded to furnish 8 in 42% yield along with the corresponding TMS enol ether 10, a product derived from single Michael addition reaction, which was reacted further on prolonged heating (entry 3).

The relative stereochemistry of the bicyclo[2.2.2] octane derivative **8** was determined as depicted in Fig. 1 by the W-type long range coupling (J 1.3 Hz) between a proton on a carbon bearing methoxycarbonyl group and an *endo* -proton of methylene protons  $\alpha$  to carbonyl group.



Fig. 1 Determination of relative stereochemistries by W-type long range coupling

In an attempt to use mild reaction conditions, the TMS enol ether **2b** of oxophorone **1** was prepared with TMSI and  $(TMS)_2NH^{6.7}$  since an attempt to prepare trimethylsilylenol ether of oxophorone **2b** with LDA/TMSCI failed and provided a complex mixture. The TMS enol ether **2b** was made to react with methyl acrylate **7** employing Et<sub>2</sub>AlCl catalyst<sup>8</sup> to afford stereoselectively, the bicyclic compounds **8**, **9** and single Michael addition adduct **10** (Table 1, entry 4). Other Lewis acid catalyst such as titanium tetrachloride<sup>8</sup> resulted in decomposition of **2b**.



Table 1. Optimization of the Reaction Condition of Double Michael Addition Reactions of Oxophorone with Methyl Acrylate 7.

In order to probe further, the synthetic utility of this procedure, the reaction of 2b and its derivatives such as 24 and 28° with various Michael addition partners 12~15 were investigated and the results obtained are summarized in Table 2. It is noteworthy that the  $\alpha,\beta$ -unsaturated ketones 12~14 reacted smoothly with 2b, 24 or 28 to afford the corresponding bicyclo[2.2.2]octane derivatives, since such substrates are known to be fragile double Michael partners during the reaction with cyclohexenone enolates resulting in either decomposition or forming aldol condensation products. The isomeric silylenol ether 28 was prepared by Lewis acid promoted rearrangement of the silylenol ether 24 (Scheme 2, see Experimental section). The mechanistic aspects of this rearrangement is under investigation and will be reported elsewhere. In the reaction of methyl  $\alpha$ -bromoacrylate 15, cyclopropane derivative 23 was obtained (entry 4) as a result of double Michael-alkylation reaction.<sup>9</sup>

Entry Silvlenol ether **Michael Partner** Products OTMS 0<sup>/</sup> 1 OTMS O ö ő 12 2b 17 30% 16 48% 18 15% OTMS 2 O' 13 19 93% O OTMS OTMS 3 0 0 14 21 20 Total 43% (6:1) MeO OMe B Br 0 OMe 4 0 0 01 15 22 15% 23 28% 0″ C OTMS Bn B Bno Bn OTMS 12 5 П Ó 0 Ö 27 18% Ö 24 26 65% 25 5% 0″ O OTMS Bn Bn OTMS 12 6 Π 0^ Π Bn ö ö 31 8% 30 76% 28

Table 2. Et<sub>2</sub>AICI Promoted Double Michael Addition Reactions of Oxophorone Trimethylsilylenol Ether **2b** and its Derivatives **24** and **27** with Other  $\alpha$ , $\beta$ -Unsaturated Carbonyl Compounds.



Since single Michael addition adduct 11 was formed at low temperature conditions (Table 1, entry 5), the present reaction seems to proceed via two successive silyl transfer Michael pathway. Isolation of a mixture of diastereomers 20 and 21 in the reaction of 3-methyl-3-penten-2-one 14 (Table 2, entry 3) also supports such pathway. Relative stereochemistries of the bicyclo[2.2.2]octanes 16, 17, 19, 25, 26 and 30 were confirmed as depicted in Table 2 by the W-type long range couplings between bridge and *endo*-protons of the corresponding triketones 16, 32, 25, and 33 (Fig 1) derived by acid catalyzed hydrolyses. The relative stereochemistry of 21 was determined by NOESY experiment utilizing triketone 35 formed after acid catalyzed hydrolysis (Fig 1).

Preferential formation of bicyclo[2.2.2]octane derivatives (*via* path a) rather than bicyclo[3.2.1]octane derivative (*via* path b) could be explained on the basis of a larger HOMO coefficient at C-3 than at C-2 (MOPAC, AM1) in case of the silylenol ether **2b**, thereby providing for the second Michael addition to occur concurrently.<sup>10</sup> In the case of the lithium enolate **2a**, the sign of LUMO coefficients of methyl acrylate matches only with those of HOMO at C-3 and C-6. The resultant stereochemical aspect of bicyclo[2.2.2]octane derivatives may be explained by chelation of metal with carbonyl oxygen of Michael partner and oxygen of enolate or enol ether.<sup>4</sup>

In summary, we have demonstrated that the double Michael addition reactions of oxophorone 1, its trimethylsilylenol ether 2b and other derivatives 24 and 28 furnished selectively the bicyclo[2.2.2]octane derivatives in single pot operation.

### Acknowledgement.

We thank Soda Aromatics Co. Ltd. for measurement of mass spectral data.

#### Experimental

All m.p.s were determined with a Yanaco MP hot-stage apparatus and are uncorrected. IR spectra were recorded on a Shimadzu FT/IR-4200 spectrophotometer for solutions in carbon tetrachloride unless otherwise indicated. <sup>1</sup>H-NMR spectra were obtained for solutions in deuteriochloroform with Varian Gemini 200H (200 MHz) and Unity 500 (500 MHz) instruments with tetramethylsilane as internal standard. <sup>13</sup>C-NMR spectra were measured with Varian Gemini 200H (50 MHz) or Unity 500 (125 MHz) instruments. Mass spectral data were run on a Hitachi M-80B spectrometer with M0101 data system. Medium-pressure liquid chromatography (MPLC) were carried out on a JASCO PRC-50 instrument with a silica gel packed column. Microanalyses were carried out in the microanalytical laboratory of Institute for Chemical Reaction Science, Tohoku University.

Methyl (2S\*)-1,5,5-Trimethyl-6,8-dioxobicyclo[2.2.2]octane-2-carboxylate **8**.— To a stirred solution of diisopropylamine (158 µl, 1.2 mmol) in THF (5 ml) was added a solution of n-BuLi (730 µl, 1.2 mmol, 1.6 M solution in n-hexane) at 0 °C under nitrogen atmosphere. After being cooled at -78 °C, a solution of oxophorone **1** (102 µl, 1 mmol) in THF (5 ml) was added and the resulting solution was stirred for 5 min. Subsequently, methyl acrylate (120 µl., 1 mmol) was added and stirring was continued for 45 min. The reaction was quenched by addition of aq. NH<sub>4</sub>Cl and extracted with chloroform twice. The combined organic layers were washed with brine and dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>. Evaporation of the solvent *in vacuo* followed by MPLC purification afforded bicyclic compound **8** (54 mg, 23%) as white needles which had; mp 72-75 °C;  $v_{max}$ /cm<sup>-1</sup> 2973, 1742, 1730, 1203 and 1177; <sup>1</sup>H-NMR  $\delta$  (200 MHz) 1.07 (s, 3H), 1.08 (s, 3H), 1.23 (s, 3H), 2.10 (dd, 1H, J 19.8, 1.6 Hz), 2.18 (ddd, 1H, J 15.4, 7.3, 2.9 Hz), 2.43 (ddd, 1H, J 15.4, 10.5, 3.8 Hz), 2.45 (dd, 1H, J 7.0, 3.8 Hz), 2.74 (ddd, 1H, J 10.5, 7.3, 1.6 Hz), 3.07 (d, 1H, J 19.8 Hz) and 3.73 (s, 3H); <sup>13</sup>C-NMR (50MHz)  $\delta$  (ppm) 17.1 (q), 23.1 (q), 23.7 (t), 25.1 (q), 41.3 (t), 41.8 (d), 44.1 (q), 48.0 (d), 52.1 (s), 55.8 (s), 173.2 (s) and 209.9 (s) (Anal. Calc. for C<sub>13</sub>H<sub>18</sub>O<sub>4</sub>: C, 65.53 %; H, 7.6 %. Found: C, 65.27 %; H, 7.41%).

Methyl 3-(2,2,4-Trimethyl-3,6-dioxo-4-cyclohexenyl) propanoate 11.— To a stirred solution of oxophorone trimethylsilylenol ether 2b (59 mg, 0.26 mmol) in dichloromethane (5 ml) was successively added methyl acrylate (32  $\mu$ l, 0.32 mmol) and diethylaluminum chloride (356  $\mu$ l, 0.32 mmol, 0.9 M solution in n-hexane) at -78 °C under nitrogen atmosphere. The resulting solution was allowed to warm to 0 °C over 6 hr and the reaction was quenched by addition of aq. NaHCO<sub>3</sub>. After extraction with chloroform twice, the organic layer was washed with brine and dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>. Evaporation of the solvent followed by MPLC purification provided single Michael adduct 11 (31 mg, 66%) along with recovered oxophorone 1 (15 mg, 25%).

The single Michael adduct 11 had; mp 90-92 °C;  $v_{max}/cm^1$  2773, 1755, 1714, 1682, 1437, 1267 and 1181; <sup>1</sup>H-NMR  $\delta$  (200MHz) 1.86 (d, 3H, J 1.4 Hz), 1.98 (t, 3H, J 14.1 Hz), 2.33-2.53 (m 2H), 3.34 (t, 1H, J 8 Hz), 3.7 (s, 3H) and 5.88 (q, 1H, J 1.4 Hz) (Anal. Calc. for  $C_{13}H_{18}O_4$ : C, 65.53 %; H, 7.61 %. Found: C, 65.30 %; H, 7.47 %).

Methyl (2S\*)-3,5,5-Trimethyl-7-trimethylsiloxy-4-oxobicyclo[2.2.2]oct-7-ene-2-carboxylate 9.— A solution of oxophorone 1 (20  $\mu$ l, 0.2 mmol), zinc chloride (273 mg, 2 mmol), triethylamine (446  $\mu$ l, 3.2 mmol) and methyl acrylate (39  $\mu$ l, 0.8 mmol) in toluene (2 ml) was heated in a sealed glass tube at 180 °C for 43 hr. After cooling to room temperature, the resulting solution was poured into 1N HCl and extracted with ethyl acetate twice. The organic layer was washed with water and brine and dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>. Evaporation of the solvent followed by MPLC purification provided trimethylsilylenol ether of bicyclic compound 9 (21 mg, 34 %) and bicyclic product 8 (18 mg, 38 %).

The trimethylsilylenol ether **9** had; mp 63-66 °C;  $v_{max}$ /cm<sup>-1</sup> 2973, 1744, 1731, 1368, 1240 and 1204; <sup>1</sup>H-NMR  $\delta$  (200MHz) 0.27 (s, 9H), 1.05 (s, 3H), 1.14 (d, 3H, J 1.7 Hz), 1.15 (s, 3H), 1.87 (ddd, 1H, J 13.2, 7.0, 2.7 Hz), 2.27 (ddd, 1H, J 13.2, 9.4, 2.7 Hz), 2.33 (dd, 1H, J 5.5, 2.7 Hz), 2.57 (dd, 1H, J 9.4, 7.0 Hz), 2.67 (s, 3H) and 4.41 (d, 1H, J 1.7 Hz).

2,6,6-Trimethyl-4-trimethylsiloxy2-cyclohexene-1-one **2b**. — To a stirred solution of 3,5,5-trimethyl-2cyclohexene-1,4-dione (oxophorone) **2a** (204  $\mu$ l, 2 mmol) in 1,2-dichloromethane (10 ml) was added hexamethyldisilazane (630  $\mu$ l, 3 mmol) and trimethylsilyliodide (345  $\mu$ l, 2.4 mmol) at 0 °C under nitrogen atmosphere. The resulting solution was stirred at room temperature for 1 hr and poured into aq. NaHCO<sub>3</sub>. After extraction with chloroform twice, the organic layer was washed with brine and dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>. Evaporation of the solvent *in vacuo* followed by purification with MPLC provided the trimethylsilylenol ether of oxophorone **2b** (298 mg, 67%).

The silylenol ether **2b** had;  $v_{max}/cm^{-1}$  2965, 1692, 1651, 1372, 1254 and 1150; <sup>1</sup>H-NMR  $\delta$  (200 MHz) 0.23 (s, 9H), 1.19 (s, 6H), 1.87 (s, 3H), 5.28 (s, 1H) and 6.63 (s, 1H).

Methyl (2S\*)-1,5,5-Trimethyl-6,8-dioxobicyclo[2.2.2]octane-2-carboxylate **8**, Methyl (2S\*)-3,5,5-Trimethyl-7trimethylsiloxy-4-oxobicyclo[2.2.2]oct-7-ene-2-carboxylate **9**, and Methyl 3-(2,2,4-Trimethyl-4-trimethylsiloxy-3-oxo-4,6-cyclohexa dienyl) propanoate **10**.— To a stirred solution of oxophorone trimethylsilylenol ether **2b** (50 mg, 0.22 mmol) in dichloromethane (4.4 ml) was successively added methyl acrylate (26  $\mu$ l, 0.26 mmol) and diethylaluminum chloride (295  $\mu$ l, 0.26 mmol, 0.9M solution in n-hexane) at -78 °C under nitrogen atmosphere. The resulting solution was stirred at -78 °C for 30 min, 0 °C for 2 hr and then room temperature for 21 hr. The reaction was quenched by addition of aq. NaHCO<sub>3</sub>. After extraction with chloroform twice, the organic layer was washed with brine and dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>. Evaporation of the solvent *in vacuo* followed by MPLC purification gave silylenol ether of bicyclic compound **9** (24 mg, 35%), silylenol ether of single Michael adduct **10** (11 mg, 20%) and bicyclic compound **8** (5 mg, 10%) in the order of elution. The silylenol ether of single Michael adduct **10** had;mp 88-92 °C; v<sub>max</sub>/cm<sup>1</sup> 2955, 1745, 1682, 1253, 1203,

The shylenoi ether of single Michael adduct 10 had; inp 88-92 °C;  $v_{max}$ /Cirl 2933, 1743, 1882, 1233, 1203, 1034 and 914; <sup>1</sup>H-NMR  $\delta$  (200 MHz) 0.27 (s, 9H), 0.92 (s, 3H), 1.07 (s, 3H), 1.86 (d, 3H, J 1.5 Hz), 1.91-2.03 (m, 2H), 2.31-2.48 (m, 1H), 3.36-3.47 (m, 1H), 3.70 (s, 3H) and 5.88 (q, 1H, J 1.5 Hz).

(2S\*)-2-Acetyl-1,5,5-trimethyl-6,8-dioxobicyclo[2.2.2]octane 16, (2S\*)-2-Acetyl-1,5,5-trimethyl-8-trimethylsiloxy-6-oxobicyclo[2.2.2]oct-7-ene 17 and 2,2,6-Trimethyl-3-(3-oxobutyl)-5-cyclohexene-1,4-dione 18.— To a stirred solution of oxophorone trimethylsilylenol ether 2b (62 mg, 0.27 mmol) in dichloromethane (5.5 ml) was successively added 3-pentene-2-one (28  $\mu$ l, 0.34 mmol) and diethylaluminum chloride (370  $\mu$ l, 0.33 mmol, 0.9 M solution in n-hexane) at 0 °C under nitrogen atmosphere. The resulting solution was stirred at 0 °C for 30 min and then room temperature for 40 hr and poured into aq. NaHCO<sub>3</sub>. The products were extracted with chloroform twice, and the organic layer was washed with brine and dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>. Evaporation of the solvent *in vacuo* followed by MPLC purification afforded bicyclic compound 16 (29 mg, 48%), bicyclic silylenol ether 17 (25 mg, 30%), and Michael product 18 (9.3 mg, 15%) in the order of elution.

The bicyclic compound **16** had; mp 67-68 °C;  $v_{max}/cm^{-1}$  2973, 1734, 1458, 1359, 1153 and 1051; <sup>1</sup>H-NMR  $\delta$  (200 MHz) 1.06 (s, 3H), 1.13 (s, 3H), 1.26 (s, 3H), 2.04 (d, 1H, *J* 19.9 Hz), 2.07 (dt, 1H, *J* 3.7, 2.1 Hz), 2.23 (s, 3H), 2.33 (ddd, 1H, *J* 14.1, 10.2, 3.7 Hz), 2.44 (ddd, 1H, *J* 3.7, 2.1, 0.6 Hz), 2.93 (ddd, 1H, *J* 10.2, 7.4, 1.2 Hz) and 3.03 (dd, 1H, *J* 19.9, 0.6 Hz) (Anal. Calc. for C<sub>13</sub>H<sub>18</sub>O<sub>3</sub>: C, 70.25; H, 8.16. Found: C, 70.13; H, 8.05).

The bicyclic silylenol ether 17 had;  $v_{max}/cm^{-1}$  2968, 1721, 1634, 1475, 1356, 1255, 1161 and 878; <sup>1</sup>H-NMR  $\delta$  (200 MHz) 0.27 (s, 9H), 1.06 (s, 3H), 1.14 (s, 3H), 1.18 (s, 3H), 1.75 (ddd, 1H, J 13.1, 7.2, 2.5 Hz), 2.15 (s, 3H), 2.25 (ddd, 1H, J 13.1, 9.5, 3.4 Hz), 2.36 (dd, 1H, J 5.7, 2.5 Hz), 2.69 (dd, 1H, J 9.5, 7.2 Hz) and 4.42 (d, 1H, J 2.5 Hz).

The Michael product **18**;  $v_{max}/cm^{-1}$  2986, 1718, 1699, 1682, 1624, 1462, 1358, 1259 and 868; <sup>1</sup>H-NMR  $\delta$  (200 MHz) 0.92 (s, 3H), 0.98 (s, 3H), 2.04 (d, 3H, J 1.6 Hz), 1.99-2.18 (m, 1H), 2.25 (s, 3H), 2.35 (dt, 2H, J 12.3, 6.4 Hz), 2.54 (dd, 1H, J 7.2, 1.9 Hz), 2.76 (dd, 1H, J 12.3, 6.4 Hz) and 5.76 (q, 1H, J 1.6 Hz); m/z 222 (M<sup>+</sup>, 12), 151 (49), 123 (38), 109 (47), 83 (31), 81 (36), 43 (100) and 41 (58) (Found:M<sup>+</sup>, 222.1221. C<sub>13</sub>H<sub>18</sub>O<sub>3</sub> requires m/z, 222.1256).

(2S\*)-3-[1,5,5-Trimethyl-8-trimethylsiloxy-6-oxo-2-(1-oxopropyl)]bicyclo[2.2.2]oct-7-ene **19**.— To a stirred solution of oxophorone trimethylsilylenol ether **2b** (50 mg, 0.22 mmol) in dichloromethane (4.5 ml) was successively added 4-pentene-3-one (36 µl, 0.27 mmol) and diethylaluminum chloride (300 µl, 0.27 mmol, 0.9 M solution in n-hexane) at 0 °C under nitrogen atmosphere. The resulting solution was stirred at 0 °C for 10 min and then room temperature for 2 hr and poured into aq. NaHCO<sub>3</sub>. The products were extracted with chloroform twice, and the organic layer was washed with brine and dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>. Evaporation of the solvent *in vacuo* followed by MPLC purification afforded bicyclic silylenol ether **19** (64 mg, 93%);  $v_{max}/cm^{-1}$  2871, 1724, 1712, 1634, 1458, 1254, 1145 and 879; <sup>1</sup>H-NMR  $\delta$  (200 MHz) 0.27 (s, 9H), 1.03 (t, 3H, J 7.3 Hz), 1.07 (s, 3H), 1.13 (s, 6H), 1.72 (ddd, 1H, J 13.0, 7.2, 2.4 Hz), 2.22 (ddd, 1H, J 13.0, 9.5, 3.4 Hz), 2.34 (m, 1H), 2.42 (q, 2H, J 7.3 Hz), 2.70 (dd, 1H, J 9.5, 7.2 Hz) and 4.41 (d, 1H, J 2.4 Hz).

(2S\*)-2-Propionyl-1,5,5-trimethyl-6,8-dioxabicyclo[2.2.2]octane **32**.— To a stirred solution of trimethylsilylenol ether **19** (62 mg, 0.20 mmol) in MeOH (2.0 ml) and H<sub>2</sub>O (36 µl, 2.0 mmol) was added PTSA-H<sub>2</sub>O (37.9 mg, 0.20 mmol) and stirring was continued at room temperature for 2.5 hr. The resulting solution was poured into aq. NaHCO<sub>3</sub> and extracted with ethyl acetate twice. The organic layer was washed with water, brine and dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>. Evaporation of the solvent followed by MPLC (EtOAc/Hexane=1/3) purification afforded triketone **32** (34 mg, 65 %) which had;  $v_{max}/cm^{-1}$  2975, 1744, 1726, 1714, 1458 and 1111; <sup>1</sup>H-NMR  $\delta$  (500 MHz) 1.05 (t, 3H, J 7.3 Hz), 1.06 (s, 6H), 2.02 (dt, 1H, J 7.2, 2.2 Hz), 2.03 (dd, 1H, J 19.9, 1.4 Hz), 2.32 (ddd, 1H, J 19.9 Hz); <sup>13</sup>C-NMR  $\delta$  (125 MHz) 7.3 (q), 17.1 (q), 23.2 (q), 23.9 (t), 25.0 (q), 39.2 (t), 40.9 (t), 44 (s), 45.8 (d), 48.5 (s), 56.2 (d), 210 (s), 211.5 (s) and 215.1 (s) (Anal. Calc. for C<sub>14</sub>H<sub>20</sub>O<sub>4</sub>: C, 71.16 %; H, 8.53 %. Found: C, 71.11 %; H, 8.44 %).

 $(2S^*)$ -2-Acetyl-1,2,5,5-tetramethyl-8-trimethylsiloxy-6-oxobicyclo[2.2.2]oct-7-ene **20** and  $(2R^*)$ -2-Acetyl-1,2,5,5-tetramethyl-8-trimethylsiloxy-6-oxobicyclo[2.2.2]oct-7-ene **21**.— To a stirred solution of oxophorone trimethylsilylenol ether **2b** (50 mg, 0.22 mmol) in dichloromethane (4.5 ml) was successively added 3-methyl-3-buten-2-one (26 µl, 0.27 mmol) and diethylaluminum chloride (300 µl, 0.27 mmol, 0.9 M solution in n-hexane) at 0 °C under nitrogen atmosphere. The resulting solution was sitrred at 0 °C for 10 min and then room temperature for 6 hr and poured into aq. NaHCO<sub>3</sub>. The products were extracted with chloroform twice, and the organic layer was washed with brine and dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>. Evaporation of the solvent *in vacuo* followed by MPLC purification afforded bicyclic silylenol ether **20** (25 mg, 37%) and its diastereomer **21** (4 mg, 6%).

The bicyclic silylenol ether 20 had;  $v_{max}/cm^{-1}$  2980, 1746, 1725, 1642, 1455, 1256, 1229 and 1159; <sup>1</sup>H-NMR  $\delta$  (200 MHz) 0.24 (s, 9H), 1.13 (s, 6H), 1.14 (s, 3H), 1.17 (s, 3H), 1.88 (dd, 1H, J 13.9, 2.7 Hz), 2.14 (s, 3H), 2.18 (dd, 1H, J 13.9, 3.1 Hz), 2.30 (t, 1H, J 3.1 Hz) and 4.52 (d, 1H, J 2.7 Hz).

The diastereomer **21** had;  $v_{max}/cm^{-1}$  2970, 1730, 1712, 1459, 1356, 1236, 1151 and 1086; <sup>1</sup>H-NMR  $\delta$  (200 MHz) 0.25 (s, 9H), 1.00 (s, 3H), 1.12 (s, 3H), 1.17 (s, 3H), 1.33 (s, 3H), 1.49 (dd, 1H, J 13.7, 2.8 Hz), 2.17 (s, 3H), 2.08-2.21 (m, 1H), 2.29 (dd, 1H, J 13.7, 2.8 Hz) and 4.33 (d, 1H, J 2.7 Hz).

(2S\*)-2-Acetyl-1,2,5,5-tetramethyl-6,8-dioxabicyclo[2.2.2]octane **34**.— To a stirred solution of trimethylsilylenol ether **20** (25 mg, 0.08 mmol) in MeOH (2 ml) and H<sub>2</sub>O (15 μl, 0.83 mmol) was added PTSA-H<sub>2</sub>O (152 mg, 0.8 mmol) and stirring was continued at room temperature for 2.5 hr. The resulting solution was poured into aq. NaHCO<sub>3</sub> and extracted with ethyl acetate twice. The organic layer was washed with water, brine and dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>. Evaporation of the solvent followed by MPLC (EtOAc/Hexane=1/2) purification afforded triketone **34** (16 mg, 77 %) which had  $v_{max}$ /cm<sup>-1</sup> 2980, 1746, 1726, 1706, 1455, 1354 and 1228; <sup>1</sup>H-NMR δ (500 MHz) 1.07 (s, 3H), 1.10 (s, 3H), 1.24 (s, 3H), 1.29 (s, 3H), 1.88 (dd, 1H, J 14.7, 2.9 Hz), 2.09 (d, 1H, J 20.0 Hz), 2.19 (s, 3H), 2.42 (t, 1H, J 2.9 Hz), 2.48 (dd, 1H, J 14.7, 2.9 Hz) and 2.75 (d, 1H, J 20.0 Hz); <sup>13</sup>C-NMRδ (125 MHz) 15.2 (q), 22.6 (q), 23.8 (q), 26.5 (q), 28.5 (q), 32.1 (t), 42.5 (t), 44.1 (s), 49.2 (s), 52.3 (s), 56.6 (d), 210.1 (s), 210.6 (s) and 215.8 (s) (Anal. Calc. for C<sub>14</sub>H<sub>20</sub>O<sub>3</sub>: C, 71.16; H, 8.53. Found: C, 71.01; H, 8.48).

(2R\*)-2-Acetyl-1,2,5,5-tetramethyl-6,8-dioxabicyclo[2.2.2]octane **35**.— To a stirred solution of trimethylsilylenol ether **21** (41 mg, 0.13 mmol) in MeOH (2 ml) and H<sub>2</sub>O (47 μl, 2.6 mmol) was added PT\$A-H<sub>2</sub>O (247 mg, 1.3 mmol) and stirring was continued at room temperature for 4hr. The resulting solution was poured into aq. NaHCO<sub>3</sub> and extracted with ethyl acetate twice. The organic layer was washed with water, brine and dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>. Evaporation of the solvent followed by MPLC (EtOAc/Hexane=1/2) purification afforded triketone **35** (28 mg, 83 %) which had;  $v_{max}/cm^1$  2980, 1747, 1730, 1712, 1452, 1356 and 1236; <sup>1</sup>H-NMR δ (500 MHz) 1.02 (s, 3H), 1.09 (s, 3H), 1.23 (s, 3H), 1.42 (s, 3H), 1.61 (dd, 1H, J 14.9, 3.2 Hz), 2.11 (d, 1H, J 19.8 Hz), 2.20 (s, 3H), 2.35 (t, 1H, J 3.2 Hz), 2.53 (dd, 1H, J 14.9, 3.2 Hz) and 2.57 (d, 1H, J 19.8 Hz); <sup>13</sup>C-NMRδ (125 MHz) 15.1 (q), 21.0 (q), 22.9 (q), 27.0 (q), 27.6 (q), 33.3 (t), 43.5 (t), 43.7 (s), 51.6 (s), 51.7 (s), 56.1 (d), 211.2 (s), 211.4 (s) and 215.1 (s) ; *m/z* 236 (M<sup>+</sup>, 18), 137 (30), 95 (33), 83 (42), 43 (100) and 41 (65) (Found:M<sup>+</sup>, 236.1373. C<sub>14</sub>H<sub>20</sub>O<sub>3</sub> requires *m/z*, 236.1412).

Methyl (2R\*)-2-Bromo-1,5,5-trimethyl-6,8-dioxobicyclo[2.2.2]octane-2-carboxylate 22 and Methyl 2,4,4-Trimethyl-3,6,dioxotricyclo[3.2.1.0<sup>2.7</sup>]octane-1-carboxylate 23.— To a stirred solution of oxophorone trimethylsilylenol ether 2b (112 mg, 0.5 mmol) in dichloromethane (10 ml) was successively added methyl  $\alpha$ bromoacrylate 15 (65  $\mu$ l, 0.65 mmol) and diethylaluminum chloride (670  $\mu$ l, 0.6 mmol, 0.9M solution in nhexane) at 0 °C under nitrogen atmosphere. The resulting solution was stirred at 0 °C for 30 min and then room temperature for 30 hr. After addition of diethylaluminum chloride (670  $\mu$ l, 0.6 mmol, 0.9M solution in n-hexane), the resulting solution was stirred for 85 hr. The reaction was quenched by addition of aq. NaHCO<sub>3</sub>. After extraction with chloroform twice, the organic layer was washed with brine and dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>. Evaporation of the solvent *in vacuo* followed by MPLC purification gave bicyclic compound 22 (23 mg, 15%) and tricyclic compound 23 (33 mg, 28%) in the order of elution.

The bicyclic compound **22** had;  $v_{max}/cm^{-1}$  (CHCl<sub>3</sub>) 2980, 1744, 1728, 1456, 1400, 1279 and 1197; <sup>1</sup>H-NMR  $\delta$  (200 MHz) 1.13 (s, 3H), 1.21 (s, 3H), 1.27 (s, 3H), 2.28 (d, 1H, J 19.7 Hz), 2.41 (m, 1H), 2.75 (dd, 1H, J 16.5,

3.5 Hz), 3.07 (d, 1H, J 19.7 Hz), 3.39 (dd, 2H, J 16.5, 2.6 Hz) and 3.80 (s, 3H) (Anal. Calc. for C<sub>13</sub>H<sub>17</sub>O<sub>4</sub>Br: C, 49.23 %; H, 5.40 %; Br, 20.18 %. Found: C, 48.99 %; H, 5.27 %; Br, 20.40 %).

The tricyclic compound **23** had;  $v_{max}/cm^{-1}$  (CHCl<sub>3</sub>) 2980, 1738, 1709, 1464, 1439 and 1244; <sup>1</sup>H-NMR  $\delta$  (200 MHz) 1.04 (s, 3H), 1.15 (s, 3H), 1.39 (s, 3H), 2.10 (dd, 1H, J 5.4, 1.8 Hz), 2.38 (d, 1H, J 13.2 Hz), 2.44 (dd, 1H, J 13.2, 5.4 Hz), 2.53 (d, 1H, J 1.8 Hz) and 3.79 (s, 3H); <sup>13</sup>C-NMR (50 MHz, CDCl<sub>3</sub>)  $\delta$  (ppm) 12.5 (q), 22.6 (q), 24.2 (q), 26.6 (s), 41.7 (s), 41.9 (s), 46.2 (s), 47.9 (s), 51.3 (d), 52.7 (q), 168.2 (s), 205.9 (s) and 208.2 (s); *m/z* 236 (M<sup>+</sup>, 44), 121 (60), 70 (59), 41 (100) and 39 (82) (Found:M<sup>+</sup>, 236.0998. C<sub>13</sub>H<sub>16</sub>O<sub>4</sub> requires *m/z*, 236.1049).

(2S\*)-2-Acetyl-4-benzyl-1,5,5-trimethyl-8-trimethylsiloxy-6-oxobicyclo[2.2.2]oct-7-ene 25, 3-Benzyl-2,2,6trimethyl-3-(3-oxoproryl)-5-cyclohexene-1,4-dione 26 and 2-Benzyl-2,2,6-trimethyl-3-(3-oxopropyl)-5cyclohexene-1,4-dione 27.— To a stirred solution of  $\alpha$ -benzyloxophorone trimethylsilylenol ether 24 (115 mg, 0.37 mmol) in dichloromethane (3.7 ml) was successively added 3-penten-2-one (37 µl, 0.44 mmol) and diethylaluminum chloride (490 µl, 0.44 mmol, 0.9M solution in n-hexane) at 0 °C under nitrogen atmosphere. The resulting solution was stirred at 0 °C for 2.5 hr and then at room temperature for 3.5 hr. The reaction was quenched by addition of aq. NaHCO<sub>3</sub>. After extraction with chloroform twice, the organic layer was washed with brine and dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>. Evaporation of the solvent *in vacuo* followed by MPLC purification gave trimethylsilylenol ether of bicyclic compound 26 (91 mg, 65%), bicyclic compound 25 (5 mg, 5%) and single Michael product 27 (20 mg, 18%) in the order of elution.

The trimethylsilylenol ether bicyclic compound **25** had; mp 184-185 °C;  $v_{max}$ /cm<sup>-1</sup> (CHCl<sub>3</sub>) 3048, 1725, 1718, 1495, 1456, 1386, 1363 and 1165; <sup>1</sup>H-NMR  $\delta$  (200 MHz) 1.02 (s, 3H), 1.08 (s, 3H), 1.32 (s, 3H), 1.47 (dd, 1H, J 14.9, 8.5 Hz), 1.97 (dd, 1H, J 14.9, 10.5 Hz), 1.98 (dd, 1H, J 19.6, 1.2 Hz), 2.14 (s, 3H), 2.36 (d, 1H, J 13.5 Hz), 2.88 (ddd, 1H, J 10.5, 8.5, 1.2 Hz), 3.15 (d, 1H, J 19.6 Hz), 3.53 (d, 1H, J 13.5 Hz ) and 7.13-7.29 (m, 5H).

The bicyclic compound **26** had; mp 134-136 °C;  $v_{max}$ /cm<sup>1</sup> 2975, 1150, 1717, 1626, 1456, 1256, 864, 849 and 704; <sup>1</sup>H-NMR  $\delta$  (200 MHz) 0.32 (s, 9H), 1.09 (s, 3H), 1.15 (s, 6H), 1.26 (dd, 1H, *J* 13.6, 7.3 Hz), 1.96 (dd, 1H, *J* 13.6, 9.5 Hz), 2.00 (s, 3H), 2.58 (d, 1H, *J* 13.9 Hz), 2.59 (dd, 1H, *J* 9.5, 7.3 Hz), 3.39 (d, 1H, *J* 13.9 Hz), 4.47 (s, 1H) and 7.14-7.33 (m, 5H).

The single Michael product **27** had; mp 163-165 °C;  $v_{max}/cm^{-1}$  (CHCl<sub>3</sub>) 2939, 1709, 1669, 1495, 1454, 1358, 874 and 689; <sup>1</sup>H-NMR  $\delta$  (200 MHz) 0.66 (s, 3H), 1.21 (s, 3H), 1.76 (dd, 2H, J 12.4, 6.7 Hz), 1.89 (d, 3H, J 1.5 Hz), 1.96-2.28 (m, 1H), 2.17 (s, 3H), 2.98 (d, 1H, J 14.0 Hz), 3.13 (d, 1H, J 14.0 Hz), 3.27 (dd, 1H, J 10.4, 6.7 Hz), 5.96 (d, 1H, J 1.5 Hz) and 7.14-7.24 (m, 5H).

## 3-Benzyl-2,6,6-trimethyl-4-trimethylsiloxy-2,4-cyclohexadienone 28 and 2-benzyl-3,5,5-trimethyl-2-

cyclohexene-1,4-dione 29.— To a solution of trimethylsilylenol ether 24 (678 mg, 2.2 mmol) in dichloromethane (22 ml) was added a solution of diethyl aluminumchloride (2.7 ml, 2.7 mmol, 1M in n-hexane) at 0 °C under nitrogen atmosphere. After stirring for 30 min at 0 °C and then 8 hr at room temperature, the resulting solution was added to aq. NaHCO<sub>3</sub>. Extraction with chloroform three times followed by evaporation of solvent gave a residue which was purified by MPLC to provide trimethylsilylenol ether 28 (370 mg, 55%) and 1,4-dione 29 (184 mg, 35%) in the order of elution.

The trimethylsilylenol ether **28** had;  $v_{max}/cm^{-1}$  2967, 1715, 1682, 1495, 1454, 1381, 1257, 1151, 1084, 845 and 700; <sup>1</sup>H-NMR  $\delta$  (200 MHz) 0.07 (s, 9H), 1.22 (s, 6H), 1.95 (s, 3H), 3.82 (s, 2H), 5.25 (s, 1H) and 7.08-7.32 (m, 5H).

The 1,4-dione **29** had;  $v_{max}/cm^{-1}$  (neat) 2980, 1678, 1495, 1455, 1277, 727 and 700; <sup>1</sup>H-NMR  $\delta$  (200 MHz) 1.23 (s, 6H), 2.05 (s, 3H), 2.75 (s, 2H), 3.82 (s, 2H) and 7.08-7.31 (m, 5H); <sup>13</sup>C-NMR $\delta$  (50 MHz) 14.2 (q), 26.7 (q), 32.9 (t), 45.8 (s), 52.4 (t), 127.0 (d), 129.0 (d), 129.2 (d), 138.4 (s), 145.2 (s), 147.5 (s), 198.0 (s) and 204.0 (s); *m*/z 242 (M<sup>+</sup>, 84), 227 (75), 130 (53), 129 (92), 128 (62), 115 (100), 91 (77), 41 (66) and 39 (59) (Found: M<sup>+</sup>, 242.1277. C<sub>16</sub>H<sub>18</sub>O<sub>2</sub> requires *m*/z, 242.1307).

 $(2S^*)$ -2-Acetyl-7-benzyl-1,5,5-trimethyl-8-trimethylsiloxy-6-oxobicyclo[2.2.2]oct-7-ene **30** and 5-Benzyl-2,2,6trimethyl-3-(3-oxoproryl)-5-cyclohexene-1,4-dione **31**.— To a stirred solution of benzyloxophorone trimethylsilylenol ether **28** (72 mg, 0.23 mmol) in dichloromethane (2.3 ml) was successively added 3-penten-2-one (23 µl, 0.28 mmol) and diethylaluminum chloride (290 µl, 0.28 mmol, 0.9M solution in n-hexane) at 0 °C under nitrogen atmosphere. The resulting solution was stirred at 0 °C for 2 hr and then at room temperature for 30 min. The reaction was quenched by addition of aq. NaHCO<sub>3</sub>. After extraction with chloroform twice, the organic layer was washed with brine and dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>. Evaporation of the solvent *in vacuo* followed by MPLC purification gave trimethylsilylenol ether of bicyclic compound **30** (67 mg, 76%) and single Michael product **31** (6 mg, 8%) in the order of elution.

The trimethylsilylenol ether of bicyclic compound **30** had; $v_{max}$ /cm<sup>-1</sup> 2969, 1717, 1657, 1495, 1455, 1354, 1159, 847 and 696; <sup>1</sup>H-NMR  $\delta$  (200 MHz) 0.26 (s, 9H), 1.07 (s, 6H), 1.21 (s, 3H), 1.70 (ddd, 1H, *J* 13.0, 7.0, 2.7 Hz), 2.13 (s, 3H), 2.28 (ddd, 1H, *J* 13.0, 9.7, 3.2 Hz), 2.44 (dd, 1H, *J* 3.2, 2.7 Hz), 2.70 (dd, 1H, *J* 9.7, 7.0 Hz), 3.02 (d, 1H, *J* 15.4 Hz), 3.88 (d, 1H, *J* 15.4 Hz) and 6.99-7.26 (m, 5H).

The single Michael product **31** had;  $v_{max}$ /cm<sup>-1</sup> 2961, 1694, 1667, 1433, 1358, 1183 and 1154; <sup>1</sup>H-NMR  $\delta$  (200 MHz) 0.90 (s, 3H), 0.91 (s, 3H), 1.95-2.20 (m, 2H), 2.07 (s, 3H), 2.23 (s, 3H), 2.37 (dd, 1H, J 14.4, 7.0 Hz), 2.66 (dd, 2H, J 11.8, 7.0 Hz), 3.58 (d, 1H, J 14.7 Hz), 3.67 (d, 1H, J 14.7 Hz) and 7.09-7.29 (m, 5H); *m/z* 312 (M<sup>+</sup>, 23), 227 (21), 91 (77), 55 (20), 43 (100) and 41 (29) (Found: M<sup>+</sup>, 312.1686. C<sub>20</sub>H<sub>24</sub>O<sub>3</sub> requires *m/z*, 312.1725).

(2S\*)-2-Acetyl-7-benzyl-1,5,5-trimethyl-6,8-dioxobicyclo[2.2.2]octane **33**.— To a stirred solution of trimethylsilylenol ether **30** (47 mg, 0.12 mmol) in acetone (1.2 ml) and H<sub>2</sub>O (22 µl, 1.2 mmol) was added PTSA-H<sub>2</sub>O (23 mg, 0.12 mmol) and stirring was continued at room temperature for 3 hr. The resulting solution was poured into aq. NaHCO<sub>3</sub> and extracted with ethyl acetate twice. The organic layer was washed with water, brine and dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>. Evaporation of the solvent followed by MPLC (EtOAc/Hexane±1/2) purification afforded triketone **33** (16 mg, 44 %) which had;  $v_{max}/cm^{-1}$  (CHCl<sub>3</sub>) 2970, 1779, 1742, 1721, 1456, 1387, 1109 and 1024; <sup>1</sup>H-NMR (500 MHz)  $\delta$  1.17 (s, 3H), 1.24 (s, 3H), 1.43 (s, 3H), 1.57 (s, 3H), 1.74-1.79 (m, 1H), 1.84 (ddd, 1H, J 14.6, 12.2, 4.0 Hz), 2.00 (dd, 1H, J 10.8, 2.7 Hz), 2.32 (ddd, 1H, J 14.6, 10.8, 2.0 Hz), 2.49 (dd, 1H, J 4.0, 2.0 Hz), 2.57 (d, 1H, J 15.3 Hz), 2.93 (d, 1H, J 15.3 Hz) and 7.13-7.31 (m, 5H).

## **References and Notes**

- 1 For example, see: Corey, E. J.; Chen, X-M., In *The logic of chemical synthesis*, John Wiley & Sons, New York, 1989.
- 2 Recent efforts of intermolecular double Michael reactions leading to the bicyclo[2.2.2]octane ring picked up arbitrary: Maiti, S.; Bhaduri, S.; Achari, B.; Banerjee, A. K.; Nayak, N. P.; Mukherjee, A. K. *Tetrahedron Lett.*, **1996**, *37*, 8061; Ley, S. V.; Mynett, D. M.; Koor, W-J. Syn. Lett., **1995**, 1017; Nagaoka, H.; Shibuya, K.; Kobayashi, K.; Miura, I.; Muramatsu, M.; Yamada, Y. *Tetrahedron Lett.*, **1993**, *34*, 4039.
- For example: Loh, T-P.; Pei, J.; Lin, M. J. Chem. Soc. Chem. Commun., 1996, 2315; Hung, S-C.; Liao, C
  C. Tetrahedron Lett., 1991, 32, 4011; Carr, R. V. C.; Williams, R. V.; Paquette, L. A. J. Org. Chem., 1983, 48, 4976.
- Tietze, L. F.; Beifuss, U. Angew. Chem. Int. Ed. Engl. 1993, 32, 131; Ihara, M.; Fukumoto, K. Angew.
  Chem. Int. Ed. Engl. 1993, 32, 1010; Hagiwara, H. J. Synth. Org. Chem. Soc., Jpn. 1992, 50, 713;
  Posner, G. H. Chem. Rev. 1986, 86, 831; Ihara, M.; Fukumoto, K. J. Synth. Org. Chem. Soc., Jpn. 1986, 44, 96.
- 5 Ito, N.; Etoh, T.; Hagiwara, H.; Kato, M. Synthesis, 1997, 153.
- 6 Ihara, M.; Makita, K.; Fujiwara, Y.; Tokunaga, Y.; Fukumoto, K. J. Org. Chem., 1996, 61, 6416.
- 7 Hagiwara, H.; Akama, T., Okano, A.; Uda, H. J. Chem. Soc. Perkin Trans. 1, 1993, 2173.
- 8 Asaoka, M.; Ishibashi, K.; Yanagida, N.; Takei, H. Tetrahedron Lett., 1983, 24, 5127.
- 9 Hagiwara, H.; Abe, F.; Uda, H. J. Chem. Soc. Perkin Trans. 1, 1993, 2651.
- 10 Hagiwara, H.; Okamoto, T.; Harada, N.; Uda, H. Tetrahedron, 1995, 51, 9891.