DOI: 10.1002/ejoc.200600627

Chemoenzymatic Synthesis of 5-Thio-D-xylopyranose

Franck Charmantray,^[a] Philippe Dellis,^[b] Virgil Hélaine,^[a] Soth Samreth,^[c] and Laurence Hecquet^{*[a]}

lase.

Germany, 2006)

Keywords: Biocatalysis / C-C bond formation / Carbohydrates / Isomerisation

5-thio-D-xylopyranose, a synthon used for the preparation of drugs with antithrombotic activity, was synthesised by an enzymatic isomerisation from the corresponding ketose, 5-thio-D-xylulofuranose, with glucose isomerase. This compound was obtained by two different chemoenzymatic routes, the

Introduction

Its sulphur-in-the-ring moiety^[1] lends 5-thio-D-xylopyranose (1) interesting biological properties. This compound is an inhibitor of β -D-xylosidase^[2] and a useful chiral building block for the synthesis of D-xylopyranosides displaying antithrombotic activity (Scheme 1). Current therapies with venous antithrombotic agents use either indirect anticoagulants of the dicoumarol type or a heparin biopolymer. Each of these therapeutic classes has both advantages and limitations. Desirable features of an ideal agent would be rapid onset of action at the target, no side effects, easy monitoring and oral administration. The effects of β -D-xylopyranosides including dicoumarol-like compounds have been studied in animal models by oral administration.^[3,4]



Scheme 1. 5-Thio-D-xylopyranosides derivatives tested as oral venous antithrombotics.

The replacement of the oxygen by sulphur in the sugar ring was found to increase potency. Hence, the 5-thio-D-xylopyranose moiety of 5-thio- β -D-xylopyranosides constituted a good pharmacophore for oral antithrombotic activity (Scheme 2). Clinical studies with some of these compounds are in progress. For instance, odiparcil,^[3,4] (X = S,

- 47 rue de Longvic, 21300 Chenôve, France [c] Fournier Pharma,
- 50 rue de Dijon, 21121 Daix, France

5526

Y = O and R = methylcoumarin in Scheme 1) is currently undergoing Phase II clinical trials.

key step being the stereospecific formation of a C-C bond,

catalysed by transketolase or fructose-1,6-bisphosphate aldo-

(© Wiley-VCH Verlag GmbH & Co. KGaA, 69451 Weinheim,



Scheme 2. The 5-thio-D-xylopyranose moiety.

Various methods for producing these glycosides in a highly stereoselective manner have been described.^[5–7]

Here we report the synthesis of the aldose 5-thio-D-xylopyranose (1). General chemical routes have been described from D-xylopyranose involving protection/deprotection and replacement with nucleophilic sulphur-containing reagents.^[3,8,9] Since the first synthesis of 1 was reported,^[10] Bellamy et al.^[4] have published a five-step synthesis of thioaldose 1 from D-xylose in 36.5% overall yield. More recently, Lalot and co-workers^[11] reported the five-step synthesis of 1 from D-xylono-1,4-lactone in 42% overall yield.

In the field of monosaccharide synthesis, an enzymatic approach is particularly attractive, as enzymes usually exhibit high regio- and stereoselectivity, and the reactions proceed under gentle conditions. The preparation of ketoses and analogues has been largely described with enzymes catalysing stereoselective C-C bond formation such as transketolases and aldolases. Access to aldoses from corresponding ketoses was proposed by Wong et al.^[12] based on enzymatic ketol-aldol isomerisation. The efficiency of this rather general approach was initially reported on for the synthesis of D-glucose derivatives^[13] with commercially available fructose-1,6-bisphosphate aldolase (FruA, EC 4.1.2.13) and glucose isomerase (GlcI, EC 5.3.1.5) and by Fessner and co-workers for the synthesis of L-fucose analogues^[14] with L-fuculose 1-phosphate aldolase (FucA, EC 4.1.2.17) and L-fucose isomerase (FucI, EC 5.3.1.3).

[[]a] Laboratoire SEESIB, UMR 6504 CNRS, Université Blaise Pascal, 24 avenue des Landais, 63177 Aubière, France

E-mail: Laurence.HECQUET@univ-bpclermont.fr [b] Synkem,

In this paper, we investigated two different chemoenzymatic strategies to obtain 5-thio-D-xylopyranose (1) based on ketol-aldol enzymatic isomerisation.

Results and Discussion

Our retrosynthetic approach (Scheme 3) was based on a two-step sequence. An enzymatic ketol-aldol isomerisation of 5-thio-D-xylulofuranose (2) to 5-thio-D-xylopyranose (1) was catalysed by GlcI. Enzymes catalysing C-C bond formation were used to prepare ketose 2. Two enzymes used in organic synthesis are able to create a D-three (3S, 4R)ketose moiety: a transferase, transketolase^[15] (TK, Route A) and a lyase, FruA^[16] (Route B). These enzymes require donor and acceptor substrates. TK transfers a ketol with two carbon units from a donor substrate, hydroxypyruvate (used for synthetic purposes because the reaction becomes irreversible) to an α -hydroxyaldehyde of R configuration. The C2-C3 bond is formed, and the new asymmetric carbon is in an S configuration. Aldolases catalyse an aldol addition reaction. The ones most widely used in organic synthesis are DHAP-aldolases such as fructose-1,6-bisphosphate aldolase (FruA). This enzyme catalyses the aldol condensation of DHAP (Dihydroxy Acetone Phosphate), a ketol with three carbon units, with an α -hydroxyaldehyde, and forms the C3–C4 bond, giving (3*S*,4*R*) stereochemistry.

TK Catalysed the Synthesis of 5-Thio-D-xylulofuranose (Route A)

As TK is generally highly enantioselective towards α -hydroxylated aldehydes as acceptor substrates, compound **3** could be prepared in a racemic form. It was obtained in a three-step sequence starting from commercially available acrolein diethyl acetal **6** (Scheme 4). This underwent epoxidation with hydrogen peroxide as the oxidising agent in 60% yield.^[17] Regioselective ring opening of the epoxide in **5** was carried out with potassium thiolate made in situ by bubbling a stream of H₂S into aqueous KOH.^[18] Under these conditions, compound **4** was obtained in 95% yield. Subsequent hydrolysis of the acetal to the aldehyde was achieved with an acidic resin in water giving **3** in quantitative yield, immediately available for enzymatic condensation without further purification.

As described by Effenberger et al.,^[19] compound **3** was a substrate of TK, and it was possible to run the enzymatic reaction on a preparative scale with concentrations of substrate **3** up to 250 mM. The best yield obtained was 24%



Scheme 3. The 5-thio-D-xylopyranose retrosynthetic pathway; Route A: catalysed by transketolase (TK), Route B: catalysed by fructose-1,6-bisphosphate aldolase (FruA).



Scheme 4. Route A catalysed by TK; i: KHCO₃, PhCN, H_2O_2 , 60%; ii: H_2S , KOH, 95% or AcSK/AcSH, 63% or Na₃SPO₃/ H_2O , 70%; iii: Dowex H⁺, 100%; iv: TK, HOCH₂COCO₂Li, 24%.

FULL PAPER

(based on the racemic acceptor substrate), which is lower than the yield stated by Effenberger et al. on an analytical scale. A free thiol may inactivate TK. Thus, as an alternative to improve the yield, we used compound **3**, protected on its thiol function as a thioacetate or thiophosphate. Epoxide opening with either potassium thioacetate in thioacetic acid or sodium thiophosphate in water proceeded smoothly to give the corresponding compound **4a** or **4b** in 63% or 70% yield, respectively. Unfortunately, the acetal group could not be hydrolysed under various acidic conditions without loss of the protective group, which invariably led back to compound **3**. To improve the yield, we turned to another enzyme able to form a C–C bond with (3S, 4R)stereochemistry, FruA.

FruA Catalysed the Synthesis of 5-Thio-D-xylulofuranose 2 (Route B)

Route B1

Our initial goal was to improve the conditions reported in the literature for the straightforward enzymatic synthesis of 5-thio-D-xylulofuranose (2) catalysed by FruA from DHAP and 1,4-dithiane-2,5-diol (Scheme 5).^[18] The effectiveness of the synthesis was hampered by inhibition of FruA by the starting aldehyde itself. After 1.25 h at 37 °C, FruA activity was halved in the presence of 1,4-dithiane-2,5-diol as the acceptor (10 mm).^[20] As a result, a large amount of FruA (7 U/mL of reaction mixture) was necessary to fully convert the starting aldehyde. Also, the substrate inhibition limited the final concentration of 5-thio-Dxylulofuranose (2) to 10 mm in the reaction mixture. As with TK, substrates bearing a free thiol function may cause irreversible inactivation of FruA. Thus Route B1 was not suitable for large scale synthesis of compound 2. Accordingly, we sought another strategy.



Scheme 5. FruA-catalysed aldolisation reaction from DHAP; i: a: $(C_2H_4SO)_2$, FruA, pH 7.5, b: acid phosphatase (Pase), pH 4.7, 60%;^[18] ii: a: XCH₂CHO, FruA, pH 7.5, b: Pase, pH 4.7, 71% (7) or 55% (8); iii: Na₃SPO₃, 55% (from 7 or 8); or NaSH, 57% (from 7) or 61% (from 8).

Route B2

Our goal in this approach was to avoid the use of an aldehydic acceptor substrate with a free thiol function. We

introduced the thiol after the reaction catalysed by FruA. According to analytical studies reported in the literature, haloaldehydes such as 2-chloro or 2-bromo acetaldehyde lead to high initial velocities with FruA. We investigated the preparation of haloxyluloses that could be further substituted by a thiol with a non-enzymatic procedure (Scheme 5).

Synthesis of Haloxyluloses 7 and 8

In the first step, DHAP and 2-chloro- or 2-bromoacetaldehyde were condensed with FruA. The reactions yielded either 5-chloro-D-xylulose-1-phosphate or 5-bromo-D-xylulose-1-phosphate. These intermediates were not isolated but readily submitted to ester hydrolysis with acid phosphatase type XA (EC 3.1.3.2, from sweet potato, Pase). After purification by column chromatography, 5-chloro-D-xylulose (7) was obtained in 71% yield, whereas 5-bromo-D-xylulose (8) was obtained in 55% yield (Scheme 5).

Halogen Displacement

Two strategies were investigated. We used an efficient method^[21] described for tosylated compounds by adding the thiophosphate under acidic conditions. The phosphorylated thiosugars thus obtained decomposed in situ to give the free thiol sugar. Conversion of 5-chloro-D-xylulose (7) or 5-bromo-D-xylulose (8) into 5-thio-D-xylulofuranose (2) was achieved by nucleophilic substitution with Na₃SPO₃ reagent in water, followed by thioester hydrolysis at pH 4.0. 2 was recovered in 55% yield after purification by column chromatography. One major limitation of this method was the cost of the sodium thiophosphate, which precluded large-scale synthesis. We circumvented this problem by converting 5-halo-D-xyluloses 7 and 8 directly into compound 2 with sodium sulfide as the nucleophilic reagent. After completion of the reaction at pH 9.5 in water, the crude mixture was purified by column chromatography to give compound 2, isolated in 57% and 61% yield from 7 and 8, respectively.

In conclusion, Route B2 appeared more suitable than A or B1. Compound **2** was obtained in two steps in an overall yield of 40% from DHAP. We note that DHAP can be prepared by a chemical route,^[22] and we have recently described an easy, efficient and cheap two-step synthesis of DHAP from 2,3-epoxypropanol (*rac*-glycidol).^[23]

Enzymatic Isomerisation of 5-Thio-D-xylulofuranose (2) to 5-Thio-D-xylopyranose (1)

We converted **2** into **1** by a ketol-aldol enzymatic isomerisation (Scheme 6).^[12–14] This reaction was catalysed by GlcI with a cocktail of divalent cations (Mg^{2+} , Mn^{2+} and Co^{2+} , 1 mM each). In vivo, this enzyme catalyses the reversible isomerisation of D-xylose or D-glucose into D-xylulose or D-fructose (the ratio at equilibrium is in favour of the aldose). This enzyme is widely used in industry in its immobilized form for the large-scale conversion of D-glucose into D-fructose.^[24] The isomerisation of **2** was followed by HPLC with a refractometer as the detector. To increase the ratio in favour of the aldose, we tested different reaction conditions (Table 1). We varied the starting material concentrations from 100 mM to 200 mM along with the frequency of GlcI addition. The ketose/aldose ratio increased in favour of the aldose with time and the quantity of GlcI. The best result (ketose/aldose, 40:60) was obtained for 100 mM after 8 d at 48 °C (the optimum temperature for GlcI) and with 415 U of GlcI added portionwise (45 U every 24 h, Table 1, entry 4).



Scheme 6. Enzymatically catalysed ketol-aldol isomerisation.

Table 1. GlcI-catalysed interconversion of 5-thio-D-xylulofuranose (2) into 5-thio-D-xylopyranose (1).

Experiment	Substrate ketose 2 [mM]	GlcI [U] ^[a]	Reaction time [d]	Ketose 2/aldose 1
1	100	45 ^[b]	4	77:22
2	100	315 ^[c]	4	60:40
3	100	45 ^[b]	8	65:35
4	100	315 ^[c]	8	40:60
5	200	45 ^[b]	4	84:16
6	200	315 ^[c]	4	71:28
7	200	45 ^[b]	8	77:23
8	200	315 ^[c]	8	57:43

[a] 1 U (unit) is defined as the amount of enzyme that converts glucose to fructose at an initial rate of 1 μ mol/min under standard analytical conditions. [b] Addition of all the GlcI at time zero. [c] Portionwise addition of 45 U of GlcI every 24 h.

Under these conditions, compound 1 was isolated in its two isomeric forms (α/β ratio, 87:13) in 60% yield after semi-preparative HPLC.

Conclusions

We performed a new chemoenzymatic synthesis of 5thio-D-xylopyranose based on the GlcI reaction of the corresponding thioketose **2**. To the best of our knowledge, enzymatic isomerisation of a thioketose into a thioaldose on a preparative scale had never been described before.

Two routes to prepare **2**, catalysed by FruA or TK, have been investigated. The use of an acceptor substrate with a free thiol function is rate-limiting for these enzymes. We have developed an efficient enzymatic route catalysed by FruA, in which the acceptor substrate is a haloaldehyde and the donor substrate is DHAP, leading to 5-halo-D-xylulose **7** or **8**. The thiol was introduced after the enzymatic step by displacement of the halogen by NaSH. After isomerisation, the overall yield of **1** was 23% from commercially available 2-chloroacetaldehyde.

This chemoenzymatic strategy offers an attractive alternative to conventional chemical methods because of its stereochemical control, mild conditions and no requirement for a protecting group. To improve the yield of the isomerisation of thioketose 2 into thioaldose 1, recycling of unreacted 2 could be considered, as in the industrial production of D-fructose from D-glucose.

Experimental Section

General Experimental Information: Chemicals and solvents were purchased from Aldrich and Acros and were reagent grade. L-glycerol-3-phosphate oxidase, L-GPO, (EC 1.1.3.21, from Thermophilus bacillus), catalase (EC 1.11.1.6, from bovine liver), fructose-1,6-bisphosphate aldolase (EC 4.1.2.13, from rabbit muscle), acid phosphatase type XA (EC 3.1.3.2, from sweet potato), α -glycerophosphate dehydrogenase type I (EC 1.1.1.8, from rabbit muscle) and peroxidase type I (EC 1.11.1.7, from horseradish) were used as received from Sigma Chem. Co. Immobilised glucose isomerase (EC 5.3.1.5, from Streptomyces murinus, Sweetzyme®), was purchased from Novozyme. TK crude extract comes from the yeast strain H402.^[15e,25] Merck 60 F254 silica gel TLC plates and Merck 60/ 230-400 and 60/40-63 mesh silica gel for column chromatography were used. 1H, 13C and 31P NMR spectra were recorded with a Bruker Avance 400 spectrometer in CDCl₃, D₂O, or CD₃OD. δ values are given in parts per million (ppm) and J values in Hertz. MS and HRMS were recorded with a Micromass Q-Tof spectrometer equipped with an electrospray ionisation source. HPLC analyses of 1 were performed with a μ Bondapak NH₂ column (4.0 mm \times 150 mm) adopting the following conditions: CH₃CN/ H_2O , 95:5 v/v, 1.5 mL min⁻¹, 25 °C and with a refractometer as the detector.

Assay for DHAP Monitoring: DHAP concentration was determined enzymatically with NADH-consuming α -glycerophosphate dehydrogenase.^[26]

Assay for Hydroxypyruvate Monitoring: Hydroxypyruvate concentration was determined enzymatically with NADH-consuming lactate dehydrogenase.^[15e]

Glycidaldehyde Diethyl Acetal (5): To a suspension of KHCO₃ (4.52 g, 45 mmol) in methanol (150 mL), commercially available acrolein diethyl acetal (45 mL, 34.8 g, 395 mmol), benzonitrile (28 mL) and H₂O₂ (35%, 27 mL, 313 mmol) were added. The solution was heated in a water bath at 40 °C for 8 h. H₂O₂ (10 mL) was added, followed by an additional 10 mL 8 h later. The resulting mixture was stirred for 20 h. Water (200 mL) was poured in, and the solution was extracted with CH_2Cl_2 (4 × 50 mL). The combined organic phases were dried on MgSO4, and the solvents were evaporated under vacuum. n-Hexane (150 mL) was added to the residue thus obtained, and the precipitate of benzamide was filtered off. After evaporation of the filtrate under reduced pressure, crude compound 5 was distilled (bp = 86 °C under 50 Torr) to give a pale yellow oil in 60% yield (26 g). ¹H NMR (400 MHz, CDCl₃): δ = 1.12 (2t, J = 6 Hz, 6 H, 2×CH₃), 2.65 (m, 2 H, 3-H), 3.00 (m, 1 H, 2-H), 3.52 (m, 2 H, CH₂), 3.67 (m, 2 H, CH₂), 4.26 (d, J = 4 Hz, 1 H, 1-H) ppm. ¹³C NMR (100 MHz, CDCl₃): δ = 15.0 (CH₃), 43.4 (C³), 51.6 (C²), 62.1 (CH₂), 62.6 (CH₂), 103.3 (C¹) ppm.

3-Thioglyceraldehyde Diethyl Acetal (4): Into a solution of KOH (3.83 g, 68.4 mmol) in methanol (15 mL) at 0 °C was bubbled a stream of H₂S for 1 h. A solution of compound **5** (5 g, 34.2 mmol) in methanol (10 mL) was added dropwise for 30 min under a H₂S atmosphere. The mixture was then stirred until it reached room temperature. The solution was stirred for a further 1 h at room temperature and then poured into ice-water (150 mL). The pH was adjusted to 2 with a H₂SO₄ (5 N) with cooling, and the solution was extracted with CH₂Cl₂ (5 × 50 mL). The combined organic

phases were washed with saturated aq NaHCO₃ (20 mL) and water (20 mL) and dried with MgSO₄. After evaporation under vacuum, a yellow oil was obtained in 95% yield (5.8 g). ¹H NMR (400 MHz, CDCl₃): δ = 1.17 (td, *J* = 7 Hz, 6 H, 2×CH₃), 1.57 (t, *J* = 8 Hz, 1 H, SH), 2.6 (ddd, *J* = 7, 8, 14 Hz, 1 H, 3-H), 2.64 (s, 1 H, OH), 2.75 (ddd, *J* = 4, 8, 14 Hz, 1 H, 3-H), 3.55 (qd, *J* = 7, 10 Hz, 2 H, CH₂), 3.62 (ddd, *J* = 4, 6, 7 Hz, 1 H, 2-H), 3.72 (qd, *J* = 7, 10 Hz, 1 H, CH₂O), 4.43 (d, *J* = 6 Hz, 1 H, 1-H) ppm. ¹³C NMR (100 MHz, CDCl₃): δ = 15.3 (CH₃), 26.7 (C-3), 63.5 (CH₂), 63.8 (CH₂), 72.4 (C-2), 103.5 (C-1) ppm.

3-Acetylthioglyceraldehyde Diethyl Acetal (4a): To a solution of potassium thioacetate (962 mg, 8.3 mmol) in thioacetic acid (5.6 mL) and CH₂Cl₂ (20 mL) was added dropwise compound 3 (2.23 g, 15.9 mmol) at -78 °C under argon. The mixture was stirred at -78 °C for 1 h and for 38 h at room temperature. After filtration, the solution was poured onto ice, and the aqueous phase was extracted with CH_2Cl_2 (3×20 mL). The combined organic phases were washed with saturated aq NaHCO₃ and brine. After drying with MgSO₄ and purifying by column chromatography on silica gel (cyclohexane/ethyl acetate, 8:2), 2.2 g of a yellow oil was obtained (63% yield). ¹H NMR (400 MHz, CDCl₃): δ = 1.15 (t, J = 7 Hz, 6 H, CH₃), 2.26 (s, 3 H, CH₃), 2.65 (s, 1 H, OH), 2.9 (dd, J = 8, 14 Hz, 1 H, 3 -H), 3.18 (dd, J = 4, 14 Hz, 1 H, 3 -H), 3.5 (m, 3.5 H)2 H, CH₂), 3.67 (m, 3 H, CH₂, 2-H), 4.3 (d, J = 5 Hz, 1 H, 1-H) ppm. ¹³C NMR (100 MHz, CDCl₃): δ = 15.2 (CH₃), 30.3 (CH₃), 31.2 (C-3), 62.5 (CH₂), 63.7 (CH₂), 70.9 (C-2), 103.5 (C-1), 195.9 (CO) ppm

3-Thiophosphateglyceraldehyde Diethyl Acetal (4b): Sodium thiophosphate (2.7 g, 6.8 mmol) was added to a solution of compound **3** (1 g, 6.8 mmol) in water (20 mL). The reaction mixture was then stirred for 4 h at room temperature. After evaporation of water under vacuum, 2.1 g of a white solid was obtained (70% yield). ¹H NMR (400 MHz, D₂O): $\delta = 1.26$ (t, J = 7 Hz, 6 H, CH₃), 2.86 (dd, J = 9, 14 Hz, 1 H, 3-H), 3.33 (dd, J = 4, 14 Hz, 1 H, 3-H), 3.73 (m, 2 H, CH₂), 3.84 (m, 2 H, CH₂, 2-H), 4.6 (d, J = 6 Hz, 1 H, 1-H) ppm. ¹³C NMR (100 MHz, D₂O): $\delta = 17.2$ (CH₃), 34.5 (C-3), 66.8 (CH₂), 67.3 (CH₂), 74.8 (C-2), 106.5 (C-1) ppm.

3-Thioglyceraldehyde (3): An aqueous solution of compound **4** (1 g, 5.54 mmol) in water (32 mL) was treated with Dowex 50WX8 resin (H⁺ form, 800 mg) for 24 h. Compound **3** thus obtained (in quantitative yield) was used in the next step without further purification.

5-Thio-D-xylulofuranose (2): In a 100 mL flask, Tris buffer (0.2 M, 1.36 g), lithium hydroxypyruvate (25 mm, 154 mg), compound 3 (175 mM, 16 mL), MgCl₂ (3 mM, 34 mg) and thiamine pyrophosphate (2 mm, 52 mg) were added to water (35 mL). The pH of the solution was adjusted to 7.5 and yeast TK (200 U) was added. The reaction mixture was stirred at 30 °C under argon until complete disappearance of a-hydroxypyruvate was detected (Lactate dehydrogenase/NADH enzymatic assay). Proteins were precipitated with methanol (170 mL) and removed by centrifugation at 8000 rpm for 15 min. Dowex 50WX8 resin (H⁺ form, 10 mL) was added, the mixture was stirred for 30 min, and the resin was removed by filtration. Dowex 1X8 resin (HCO3⁻ form, 10 mL) was added to the filtrate, and the mixture was stirred for 30 min. After filtration of the resin, the solution was evaporated to dryness, and crude compound 2 was purified by flash column chromatography (CH₂Cl₂/MeOH, 8:2) to afford 110 mg of compound 2 (48% yield).

β Isomer: ¹H NMR (400 MHz, CD₃OD): δ = 2.57 (dd, *J* = 9, 10 Hz, 1 H, 5-H), 3.03 (dd, *J* = 8, 10 Hz, 1 H, 5'-H), 3.59 (d, *J* = 13 Hz, 1 H, 1-H), 3.62 (d, *J* = 13 Hz, 1 H, 1'-H), 3.73 (d, *J* = 9 Hz, 1 H, 3-H), 4.23–4.32 (m, 1 H, 4-H) ppm. ¹³C NMR (100 MHz, CD₃OD): δ = 31.2 (C-5), 67.5 (C-1), 77.0 (C-4), 79.8 (C-3), 89.7 (C-2) ppm. **α Isomer:** ¹H NMR (400 MHz, CD₃OD): δ = 2.94 (dd, J = 5, 11 Hz, 1 H, 5-H), 3.03 (dd, J = 5, 11 Hz, 1 H, 5'-H), 3.73 (d, J = 11 Hz, 1 H, 1-H), 3.81 (d, J = 11 Hz, 1 H, 1'-H), 4.03 (d, J = 5 Hz, 1 H, 3-H), 4.23–4.32 (m, 1 H, 4-H) ppm. ¹³C NMR (100 MHz, CD₃OD): δ = 35.9 (C-5), 66.8 (C-1), 79.0 (C-4), 83.9 (C-3), 96.0 (C-2) ppm. HR-ESI-MS calculated for C₅H₁₀O₄NaS [M + Na]⁺: 189.0198; found 189.0192.

5-Chloro-D-xylulose (7): 2-Chloroacetaldehyde (850 µL, mmol) at 50% (w/v) in water was added to an aqueous solution of DHAP (60 mL, 5.7 mmol) at pH 6.8. FruA (200 U) was added. After 1 h an additional amount of 2-chloroacetaldehyde (150 µL) was added to the reaction mixture and allowed to react for 18 h at room temperature. The pH of the water phase was adjusted to 4.7 with HCl (1 N), and acid phosphatase (50 U) was added. The reaction mixture was shaken for a further 48 h at room temperature. After completion of the reaction, the pH was raised to 7.0 with NaOH (1 N), and MeOH (180 mL) was poured into the solution. The resulting precipitate was removed by filtration through celite. The filtrate was concentrated under vacuum. Column chromatography (CH2Cl2/ MeOH, 9:1) yielded 5-chloro-D-xylulose as a pale yellow oil (679 mg, 71% from DHAP). ¹H NMR (400 MHz, CD₃OD): δ = 3.54 (dd, J = 8, 10 Hz, 1 H, 5-H), 3.70 (dd, J = 8, 10 Hz, 1 H, 5'-H), 4.10 (td, J = 2, 8 Hz, 1 H, 4-H), 4.39 (d, J = 2 Hz, 1 H, 3-H), 4.47 (d, J = 19 Hz, 1 H, 1-H), 4.54 (d, J = 19 Hz, 1 H, 1'-H) ppm. ¹³C NMR (100 MHz, CD₃OD): δ = 45.1 (C-5), 67.9 (C-1), 73.6 (C-4), 76.7 (C-3), 212.5 (C-2) ppm. HR-ESI-MS: calculated for $C_5H_9ClNaO_4$ [M + Na]⁺ 191.0087; found 191.0097.

5-Bromo-D-xylulose (8): 2-Bromoacetaldehyde (1.3 mL, 2 mmol) was added to an aqueous solution of DHAP (60 mL, 5.7 mmol) at pH 6.8. FruA (200 U) was added. After 1 h an additional amount of 2-bromoacetaldehyde (150 µL) was added to the reaction mixture and allowed to react for 18 h at room temperature. The pH of the water phase was then adjusted to 4.7 with HCl (1 N) and acid phosphatase (50 U) was added. The reaction mixture was shaken for a further 48 h at room temperature. After completion of the reaction, the pH was raised to 7.0 with NaOH (1 N), and MeOH (180 mL) was poured into the solution. The resulting precipitate was removed by filtration through celite. The filtrate was concentrated under vacuum. Column chromatography (CH2Cl2/MeOH, 9:1) yielded 5-bromo-D-xylulose as a pale yellow oil (666 mg, 55% from DHAP). ¹H NMR (400 MHz, CD₃OD): δ = 3.40 (dd, J = 8, 10 Hz, 1 H, 5-H), 3.58 (dd, J = 8, 10 Hz, 1 H, 5'-H), 4.15 (td, J =2, 8 Hz, 1 H, 4-H), 4.44 (d, J = 2 Hz, 1 H, 3-H), 4.51 (d, J = 19 Hz, 1 H, 1-H), 4.54 (d, J = 19 Hz, 1 H, 1'-H) ppm. ¹³C NMR (100 MHz, CD₃OD): δ = 33.4 (C-5), 67.9 (C-1), 73.7 (C-4), 77.0 (C-3), 213.4 (C-2) ppm. HR-ESI-MS: calculated for C₅H₉BrNaO₄ [M + Na]⁺ 234.9582; found 234.9577.

5-Halo-D-xylulofuranose 7 and 8 (One-Pot Reaction from *rac*-Glycidol): Solid Na₂HPO₄ (3.71 g, 25 mmol) was added to a solution of *rac*-2,3-epoxypropanol (1.9 g, 25 mmol, 50 mL) in distilled water. The mixture was heated at 100 °C for 3 h and assayed for L-glycerol-3-phosphate content. The yield was 61% from (*S*)-2,3-epoxypropanol (30% from *rac*-2,3-epoxypropanol). To a solution containing L-glycerol-3-phosphate (155 mM, 10 mL, based on enzymatic assay) at pH 6.8 was added GPO/catalase mixture (0.1 mL, 45 U/1800 U), FruA (0.07 mL, 20 U) and aldehyde (2-chloroacetaldehyde, 0.13 mL, 2 mmol or 2-bromoacetaldehyde, 545 mg, 2 mmol). The reaction proceeded with stirring at room temperature overnight. The pH was adjusted to 4.7 with HCl (1 N) and acidic phosphatase (50 U) was added. The reaction mixture was shaken for another 24 h at room temperature. The pH was raised to 7.0 with NaOH (1 N), and MeOH (30 mL) was poured into the solu-

tion. The resulting precipitate was removed by filtration through celite. The filtrate was concentrated under vacuum, and the brown residue underwent silica gel chromatography in the same conditions as already described. 5-bromo-D-xylulose was obtained as a light yellow oil in 12% yield. 5-chloro-D-xylulose was recovered as a light yellow oil in 47% yield.

5-Thio-D-xylulofuranose (2) from 5-Halo-D-xylulose 7 or 8

Method A. Na₃SPO₃ as the Nucleophile: To a solution of 5-halo-D-xylulose 7 or 8 (0.62 mmol) in distilled water (2 mL) was added a solution of Na₃SPO₃ (0.65 mmol) in distilled water (2 mL). The pH of the resulting mixture was kept at 10 by adding NaOH (2 N). After 3 h at room temperature and with stirring, the pH was lowered to 4.0 with HCl (1 N), and the reaction was allowed to proceed overnight. The pH was adjusted to 8.2 before the addition of BaCl₂ (1 mL, 0.65 mmol, a solution in water). The white precipitate was isolated by centrifugation at 3500 rpm for 15 min and discarded. The supernatant was concentrated under vacuum and subjected to flash chromatography on silica (CH₂Cl₂/MeOH, 9:1). 5-thio-D-xylulofuranose (2) was recovered as its α and β isomers (α/β , 87:13) as a pale yellow oil in 55% yield, irrespective of whether 5-halo-Dxylulose 7 or 8 was used as the starting material. Analytical data were identical to those previously given.

Method B. NaSH as the Nucleophile: The NaSH.xH₂O reagent was added (0.532 g) to a solution of 5-halo-D-xylulose (4.75 mmol) in distilled water (70 mL). The yellow solution was shaken at room temperature for 3 h and kept at 4 °C overnight. The solvents were eliminated under reduced pressure, and the crude product was purified by flash column chromatography on silica gel. The desired compound **2** was eluted with CH₂Cl₂/MeOH (85:15). It was isolated as its α and β isomers (α/β , 87:13) in 61% (2.9 mmol) and 57% (2.7 mmol) yield starting from 5-chloro- and 5-bromo-D-xylulose, respectively. Analytical data were identical to those previously given.

5-Thio-D-xylopyranose (1): Into a capped vial was introduced an aqueous solution of 5-thio-D-xylulofuranose **2** (100 mM or 200 mM, 4 mL), followed by phosphoric acid (5 μ L, 85%). The pH of the solution was adjusted to 7.5 with NaOH (90 μ L, 2 N). Stock solutions of Co²⁺ (100 mM, 38 μ L), Mg²⁺ (100 mM, 38 μ L) and Mn²⁺ (50 mM, 19 μ L) were then added. Two different processes were investigated. In Process A, the reaction was initiated by pouring GlcI (106 mg, 45 U) into the mixture and letting it react at 48 °C (optimum temperature for GlcI) for 8 d. Aliquots were analysed at intervals by HPLC to monitor the course of the reaction. For Process B, all conditions were the same, except for an additional amount of GlcI (106 mg, 45 U) added to the medium every 24 h.

Starting with 5-thio-D-xylulofuranose (2) at 100 mM and using Process B, the ketose/aldose ratio rose to 40:60 in favour of 5-thio-D-xylopyranose after 8 d at 48 °C. After removal of the protein by ultracentrifugation, the reaction mixture was subjected to semi-preparative HPLC. Purification of 1 was performed with a μ Bondapak NH₂ column (4.0 mm × 150 mm) under the following conditions: CH₃CN/H₂O, 5 mL min⁻¹, 25 °C, injection volume 10 μ L, 4 mg.

Compound 1 was recovered in 60% yield (40 mg). The structure of compound 1 was confirmed by ¹H and ¹³C NMR spectroscopy. The data exactly fit those of the literature.^[11]

Acknowledgments

We thank Professor G. Schneider (Karolinska Institut, Stockholm, Sweden) for the gift of the yeast strain H402 transformed with pTKL1. We thank also Fournier Pharma for financial support.

- a) H. Paulsen, K. Todt, Adv. Carbohydr. Chem. 1968, 23, 115– 232; b) I. Robina, P. Vogel, Z. J. Witczak, Curr. Org. Chem. 2001, 5, 1177–1214; c) I. Robina, P. Vogel, Curr. Org. Chem. 2002, 6, 471–491.
- [2] D. Horton, J. D. Wander, *The Carbohydrate, Chemistry and Biochemistry*; 2nd ed. (Eds.: W. Pigman, D. Horton), vol. IB, Academic Press, New York, **1980**, 799–842.
- [3] S. Samreth, V. Barberousse, F. Bellamy, D. Horton, P. Masson, J. Millet, P. Renaut, C. Sepulchre, J. Thevenaiux, *Actual. Chim. Thér.* 1994, 21, 23–33
- [4] F. Bellamy, D. Horton, J. Millet, F. Picard, S. Samreth, J. B. Chazan, J. Med. Chem. 1993, 36, 898–903.
- [5] H. Paulsen, Angew. Chem. Int. Ed. Engl. 1982, 21, 155–173.
- [6] R. R. Schmidt, Angew. Chem. Int. Ed. Engl. 1986, 25, 212-235.
- [7] H. Kunz, Angew. Chem. Int. Ed. Engl. 1987, 26, 294-308.
- [8] R. L. Whistler, T. Van Es, J. Org. Chem. 1963, 28, 2303-2304.
- [9] D. L. Ingles, R. L. Whistler, J. Org. Chem. 1962, 27, 3896-3898.
- [10] a) J. C. Schwarz, K. C. Yule, Proc. Chem. Soc. London 1961, 417; b) T. J. Adley, L. N. Owen, Proc. Chem. Soc. London 1961, 418.
- [11] J. Lalot, I. Stasik, G. Demailly, D. Beaupère, *Carbohydr. Res.* 2003, 338, 2241–2245.
- [12] J. R. Durrwachter, D. G. Drueckhammer, K. Nozaki, H. M. Sweers, C.-H. Wong, J. Am. Chem. Soc. 1986, 108, 7812–7818.
- [13] W.-C. Chou, L. Chen, J.-M. Fang, C.-H. Wong, J. Am. Chem. Soc. 1994, 116, 6191–6194.
- [14] W.-D. Fessner, C. Gosse, G. Jaeschke, O. Eyrisch, Eur. J. Org. Chem. 2000, 125–132.
- [15] a) T. Ziegler, A. Straub, F. Effenberger, Angew. Chem. Int. Ed. Engl. 1988, 27, 716–721; b) Y. Kobori, D. C. Myles, G. M. Whitesides, J. Org. Chem. 1992, 57, 5899–5907; c) K. G. Morris, M. E. B. Smith, N. J. Turner, Tetrahedron: Asymmetry 1996, 7, 2185–2188; d) F. T. Zimmermann, A. Schneider, Y. Schörken, G. A. Sprenger, W. D. Fessner, Tetrahedron: Asymmetry 1999, 10, 1643–1646; e) L. Hecquet, J. Bolte, C. Demunyck, Tetrahedron 1996, 52, 8223–8232; f) C. Guérard, V. Alphand, A. Archelas, C. Demuynck, L. Hecquet, R. Furstoss, J. Bolte, Eur. J. Org. Chem. 1999, 1, 3399–3402; g) D. Crestia, C. Guérard, H. Veschambre, L. Hecquet, C. Demuynck, J. Bolte, Tetrahedron: Asymmetry 2001, 12, 869–876.
- [16] a) M. D. Bednarski, E. S. Simon, N. Bischofberger, W. D. Fessner, M. J. Kim, W. Lees, T. Saito, H. Waldmann, G. M. Whitesides, J. Am. Chem. Soc. 1989, 111, 627–635; b) C. Andre, C. Guerard, L. Hecquet, C. Demuynck, J. Bolte, J. Mol. Catal. B: Enzym. 1998, 5, 459–466; c) L. Espelt, T. Parella, J. Bujons, C. Solans, J. Joglar, A. Delgado, P. Clapes, Chem. Eur. J. 2003, 9, 4887–4899; d) L. El Blidi, D. Crestia, E. Gallienne, C. Demuynck, J. Bolte, M. Lemaire, Tetrahedron: Asymmetry 2004, 15, 2951–2954.
- [17] Z. Liu, L. M. Sayre, Chem. Res. Toxicol. 2003, 16, 232-241.
- [18] F. Effenberger, A. Straub, V. Null, *Liebigs Ann. Chem.* 1992, 12, 1297–1301.
- [19] F. Effenberger, V. Null, T. Ziegler, *Tetrahedron Lett.* 1992, 33, 5157–5160.
- [20] As determined enzymatically through DHAP monitoring by NADH-consuming glycerol-3-phosphate dehydrogenase in the presence of triose phosphate isomerase.
- [21] W. B. Knight, D. S. Sem, K. Smith, H. M. Miziorko, A. Rendina, W. W. Cleland, *Biochemistry* **1991**, *30*, 4977–4979.
- [22] a) O. Meyer, M. Rohmer, C. Grosdemange-Billiard, *Tetrahedron Lett.* 2004, 45, 7921–7923; b) N. Ouwerkerk, M. Steenweg, M. De Ruijter, J. Brouwer, J. H. Van Boom, J. Lugtenburg, J. Raap, J. Org. Chem. 2002, 67, 1480–1489; c) E. L. Ferroni, V. DiTella, N. Ghanayem, R. Jeske, C. Jodlowski, M. O'Connell, J. Styrski, R. Svodoba, A. Venkataraman, B. M. Winkler, J. Org. Chem. 1999, 64, 4943–4945; d) T. Gefflaut, M. Valentin, M.-L. Lemaire, J. Bolte, J. Org. Chem. 1997, 62, 5920–5922; e) M.-L. Valentin, J. Bolte, Bull. Soc. Chim. Fr. 1995, 132, 1167–

1171; f) S.-H. Jung, J.-H. Jeong, P. Miller, C.-H. Wong, J. Org. Chem. 1994, 59, 7182–7184.

- [23] F. Charmantray, P. Dellis, S. Samreth, L. Hecquet, *Tetrahedron Lett.* **2006**, *47*, 3261–3263.
- [24] S. H. Bhosale, M. B. Rao, V. V. Deshpande, *Microbiol. Rev.* 1996, 60, 280–300.
- [25] L. Wikner, U. Meshalkina, M. Nikkola, Y. Lindqvist, G. Schneider, J. Biol. Chem. 1994, 269, 32144–32150.
- [26] H. U. Bergmeyer, Methods in Enzymatic Analysis, 34th ed., Verlag Chemie, 1984, 11, 146–147.

Received: July 20, 2006 Published Online: October 12, 2006