Effects of the Hydrogen-bonded Interactions on the Thermal *cis* to *trans* Isomerization of 3,3'-Diacetylazobenzene

Jacqueline Rosengaus and Itamar Willner*

Institute of Chemistry and the Farkas Center for Light Induced Processes, The Hebrew University of Jerusalem, Jerusalem 91904, Israel

The thermal isomerization of 3,3'-diacetyl-*cis*-azobenzene **1b** is retarded in the presence of 2,4-bis(cyclohexylamino)-6-methoxy-1,3,5-triazine due to the formation of an intermolecular hydrogen-bonded assembly.

Extensive research activity is directed towards the design of supramolecular host-guest assemblies originating from complementary hydrogen-bonded interactions.^{1,2} Of particular interest is the influence of hydrogen-bonded interactions on chemical transformations; *e.g.* self-replicating systems by complementary hydrogen-bonds,³ phosphoryl transfer processes⁴ and control of pericyclic reactions⁵ represent chemical transformations in hydrogen-bonded supramole-cular assemblies.

Recently, the effects of hydrogen-bonded interactions on the photophysical properties of a pyrene substituted receptor have been studied by Shinkai.⁶ In this system the fluorescence of the pyrene units is affected by intermolecular hydrogenbonding of the receptor with barbiturate host compounds.⁷ In a previous study we have demonstrated the formation of hydrogen-bonded assemblies between di- and tri-amino-triazines and bemegride.⁸ We have shown that these compounds provide three sites for the formation of complementary hydrogen-bonded interactions with guest substates.

Azobenzenes undergo reversible photochemical *trans* \rightleftharpoons *cis* and thermal *cis* \rightarrow *trans* isomerization reactions.⁹ Here we report the influence of hydrogen bond interactions on the thermal isomerization of 3,3'-diacetyl-*cis*-azobenzene, **1b**. We describe the influence of the hydrogen-bonded assembly formed between 2,4-bis(cyclohexylamino)-6-methoxy-1,3,5-triazine, **2**, and **1b** on the isomerization process [eqn. (1)].



3; $R^1 = OMe$, $R^2 = c-C_6H_{11}NH$



Fig. 1 Rates of isomerization of **1b**, 8.36×10^{-5} mol dm⁻³ in the presence of **2**. Concentrations of **2**: (a) 0; (b) 2.09×10^{-5} ; (c) 4.18×10^{-5} ; (d) 8.36×10^{-5} ; (e) 4.18×10^{-4} mol dm⁻³; (f) rate of isomerization of **1b** 8.36×10^{-5} mol dm⁻³, in the presence of 2.09×10^{-5} mol dm⁻³ of **3**.

Molecular mechanics calculations show that a stable complex between the *cis* isomer **1b**, and the triazine receptor **2**, is formed through two complementary hydrogen bonds.† The resulting intermolecular complex 1b-2 exhibits minimum energy. The calculated 1b-2 complex shows intermolecular hydrogen bond distances of 2.055 Å between the amine groups and the complementary carbonyl functionalities. No stable complex is formed with the *trans* isomer 1a or with 2,2'- or 2,3'-diacetylazobenzenes. The formation of the hydrogenbonded assembly 1b-2 is evidenced by ¹H NMR spectroscopy. The amine protons of **2** are shifted downfield in the presence of 1b as expected for a hydrogen-bonded assembly.⁸ Formation of the complex is anticipated to retard the thermal isomerization of 1b to 1a by the stabilization of the cisazobenzene isomer in the complex structure, eqn. (1). 3,3'-Diacetyl-trans-azobenzene, 1a, was prepared as described by Nakagawa.¹⁰[±] 3,3'-Diacetyl-cis-azobenzene, 1b, was prepared by monochromatic light irradiation, $\lambda = 355$ nm, of 1a dissolved in a dry dichloroethane solution. Sealed cells containing 1b and different concentrations of 2 were prepared, and the thermal isomerization of 1b to 1a was followed spectroscopically at 50 °C. Fig. 1 shows the rates of isomerization of 1b in the presence of different concentrations of 2. Evidently, the rate of isomerization is slowed down as the concentration of the triazine receptor 2 is increased. The inhibition of the isomerization process of 1b to 1a is attributed to the formation of the hydrogen-bonded intermolecular complex 1b-2. In this structure the cis-configuration is stabilized and the structural isomerization is retarded. Increase of the concentration of 2 results in higher concentration of the complex and, therefore, a decrease in the isomerization rates. The association constant for the complex

1b–2, has been derived from the kinetic studies§ and corresponds to $K_a = 1.2 \times 10^6 \text{ dm}^3 \text{ mol}^{-1}$.

Further support that the inhibition of the $cis \rightarrow trans$ isomerization reaction is due to the formation of a hydrogenbonded assembly between the triazine receptor 2 and the azo compound 1b has been obtained by a control experiment where the monodentate receptor 2-cyclohexylamino-4,6dimethoxy-1,3,5-triazine 3 replaces 2 as additive. The monoaminotriazine is unable to form two complementary hydrogen-bonded interactions with 1b. Fig. 1 compares the rates of isomerization of 1b in the presence of the triazine receptors 2 and 3 respectively, at a molar ratio corresponding to 1:0.25. No retarding effect of the rate of isomerization is observed in the presence of the monodentate receptor 3. In fact, addition of 3 results in even a slight enhancement in the isomerization of 1b.

This research was supported by the Basic Research Foundation administered by the Israel Academy of Sciences and Humanities.

Received, 1st March 1993; Com. 3/01175E

References

- For general reviews, see J.-M. Lehn, *Science*, 1985, 227, 849; D. J. Cram, *Science*, 1983, 219, 1177; H.-J. Schneider, *Angew. Chem.*, *Int. Ed. Engl.*, 1991, 30, 1471; J. S. Lindsey, *New J. Chem.*, 1991, 15, 153.
- 2 For hydrogen bond reviews, see J. Rebek, Jr., Acc. Chem. Res., 1990, 23, 399; Angew. Chem., Int. Ed. Engl., 1990, 29, 245; M. C. Etter, Acc. Chem. Res., 1990, 23, 120; G. R. Desiraju, Acc. Chem. Res., 1991, 24, 290.
- 3 T. Tjivikua, P. Ballester and J. Rebek, Jr., J. Am. Chem. Soc., 1990, **112**, 1249; G. Kiedrowski, B. Wlotzka, J. Heliberg, M. Matzen and S. Jordan, Angew. Chem., Int. Ed. Engl., 1991, **30**, 423.
- 4 P. Tecilla, S.-K. Chang and A. D. Hamilton, J. Am. Chem. Soc., 1990, 112, 9586.
- 5 S. C. Hirst and A. D. Hamilton, J. Am. Chem. Soc., 1991, 113, 382.
- 6 I. Aoki, T. Harada, T. Sakaki, Y. Kawahara and S. Shinkai, J. Chem. Soc., Chem. Commun., 1992, 1341.
- 7 S.-K. Chang and A. D. Hamilton, J. Am. Chem. Soc., 1988, 110, 1318.
- 8 I. Willner, J. Rosengaus and Y. Eichen, J. Phys. Org. Chem., 1993, 6, 29.
- 9 H. Rau, in *Photochromism-Molecules and Systems*, eds. H. Dürr and H. Bouas-Laurent, Elsevier, Amsterdam, 1989, p. 165.
- 10 K. Nakawa and T. Tsuji, Chem. Pharm. Bull., 1963, 11, 296.

§ The association constant of **1b–2** is calculated from the kinetic data using eqn. (2) and eqn. (3), where k_1 corresponds to the isomerization rate constants of **1b** without added host, k_2 is the isomerization rate constant in the complex structure, $[H]^0$ and $[G]^0$ are the added concentrations of **2** and **1** respectively and [HG] is the complex concentration at a given concentration of **2** (cf. A. K. Colter, S. S. Wang, G. H. Megerle and P. S. Ossip, J. Am. Chem. Soc., 1984, **86**, 3106). K_a has been derived by an iterative calculation initiated by assuming [HG] = 0 and plotting $[H]^{0/k}_{obs} - k_1 vs.$ ($[H]^0 + [G]^0 -$ [HG]). The resulting slope has been introduced into eqn. (3) and the resulting concentrations of [HG] have been resubstituted into eqn. (2). The process has been repeated until the slope converges. The association constant is derived from the slope and intercept of the converged plot.

$$\frac{[H]^0}{k_{\rm obs} - k_1} = \frac{1}{K_{\rm a}(k_2 - k_1)} + ([H]^0 + [G]^0 - [HG])\frac{1}{(k_2 - k_1)}$$
(2)

$$[HG] = \frac{k_{obs} - k_1}{k_2 - k_1} [G]^0$$
(3)

[†] The program PCMPI Ver. 4.2, Serena Software, Bloomington, IN. has been applied (cf. N. L. Allinger, J. Am. Chem. Soc., 1977, **99**, 8127).

 $[\]ddagger 3,3'$ -Diacetyl-*trans*-azobenzene, **1a**, was prepared as described by oxidation of 3-aminoacetophenone with nickel peroxide followed by SiO₂ column chromatography using light petroleum-diethyl ether (3:1) as eluent.