

Communication

**Catalytic Hydroxycyclopropanol Ring-Opening
 Carbonylative Lactonization to Fused Bicyclic Lactones**

XINPEI CAI, Weida Liang, Mingxin Liu, Xiating Li, and Mingji Dai

J. Am. Chem. Soc., **Just Accepted Manuscript** • DOI: 10.1021/jacs.0c06179 • Publication Date (Web): 20 Jul 2020

Downloaded from pubs.acs.org on July 21, 2020

Just Accepted

“Just Accepted” manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides “Just Accepted” as a service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. “Just Accepted” manuscripts appear in full in PDF format accompanied by an HTML abstract. “Just Accepted” manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are citable by the Digital Object Identifier (DOI®). “Just Accepted” is an optional service offered to authors. Therefore, the “Just Accepted” Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the “Just Accepted” Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these “Just Accepted” manuscripts.

Catalytic Hydroxycyclopropanol Ring-Opening Carbonylative Lactonization to Fused Bicyclic Lactones

Xinpei Cai, Weida Liang, Mingxin Liu, Xiating Li, and Mingji Dai*

Department of Chemistry and Center for Cancer Research, Purdue University, West Lafayette, Indiana 47907, United States

Supporting Information Placeholder

ABSTRACT: A novel palladium-catalyzed ring opening carbonylative lactonization of readily available hydroxycyclopropanols was developed to efficiently synthesize tetrahydrofuran (THF) or tetrahydropyran (THP)-fused bicyclic γ -lactones, two privileged scaffolds often found in natural products. The reaction features mild reaction conditions, good functional group tolerability, and scalability. Its application was demonstrated in a short total synthesis of (\pm)-paeonilide. The fused bicyclic γ -lactone products can be easily diversified to other medicinally important scaffolds, which further broadens the application of this new carbonylation method.

Nature has been constantly creating natural products with novel, diverse, and complex structures. Many of these natural products or their derivatives have been used as medicines to treat human diseases or are lead compounds for drug discovery.¹ While various biosynthetic pathways and different organisms are involved in natural product production, privileged structural scaffolds shared by a large number of natural products can often be identified.^{1c} *Cis*-2,8-dioxabicyclo[3.3.0]octan-3-one (a THF-fused bicyclic γ -lactone, also called *cis*-tetrahydro[2,3-*b*]furan-2(3*H*)-one) is one such privileged structural scaffold, which has been found in over 100 natural products from various origins (Figure 1).² In these molecules, the THF-fused γ -lactone forges diverse connections with the rest of the molecular structures. Part of them such as ginkgolide B (**1a**),³ gracilin B (**1b**),⁴ paracaseolide A (**1c**),⁵ and darwinolide (**1d**),⁶ have the bicyclic γ -lactone moiety embedded in a large polycyclic ring system. Another portion of them including norrisolide (**1e**)⁷ and cheloviolene A (**1f**)⁸ have the bicyclic γ -lactone connected with a second (poly)cyclic ring system via a C-C single bond. Why and how nature produces such fused bicyclic skeletons are yet to be understood. Natural products with such a bicyclic γ -lactone motif have demonstrated a wide range of biological activity and possess great potential in novel therapeutic development. For example, the ginkgolides can act as anti-platelet-activating factors; gracilins exhibit immunosuppressive and neuroprotective properties;

paracaseolide A inhibits phosphatase CDC25B, an emerging and important target for anticancer drug discovery; the marine spongian diterpenoid norrisolide and its analogs demonstrated unique Golgi-modifying activity; darwinolide is a biofilm-penetrating anti-MRSA agent. Additionally, the related *cis*-2,9-dioxabicyclo[4.3.0]nonan-8-one (a THP-fused bicyclic γ -lactone, also called *cis*-tetrahydro-4*H*-furo[2,3-*b*]pyran-2(3*H*)-one) has been found in natural products,⁹ including applanatumol B (**1h**),^{9a} which showed activity against renal fibrosis.




Figure 1. Selected natural products with a THF/THP-fused bicyclic γ -lactone

Due to their unique and challenging chemical structures as well as their diverse biological applications, natural products with a THF/THP-fused bicyclic γ -lactone moiety have been popular targets in total synthesis practices.¹⁰ Notably, the Overman group recently accomplished elegant total syntheses of several marine spongian diterpenoids including cheloviolene A (**1f**, Figure 1) with a THF-fused γ -lactone.¹¹ So far, most of these total syntheses utilized the hydroxy-acid-aldehyde (or their masked forms) condensation reactions to construct the desired THF/THP-fused γ -lactone moiety in the target

molecules (eq. A-1, Figure 2A). In these processes, tedious functional group manipulations, protections, and deprotections are often involved, which added extra steps to the entire synthesis. Surprisingly, synthetic methodology development in this area is extremely barren.¹² Notably, Trost and Toste developed a Pd-catalyzed asymmetric nucleophilic substitution of 5-acetoxy-2-(5*H*)-furanone followed by reductive intramolecular Heck reaction to build THF-fused γ -lactones,^{12a} but this method is limited to benzo-fused tricyclic systems (eq. A-2). Theodorakis^{12b} and Reiser^{12c} independently reported elegant activated cyclopropane rearrangements to THF-fused γ -lactones (eq. A-3). In the Reiser's case, the activated cyclopropanes were generated via Rh-catalyzed enantioselective cyclopropanation of furans. In general, these prior arts are limited to 5,5-fused γ -lactones and hydrogen atom substitution at the ring junction carbons. Thus, versatile, modular, and catalytic methods are highly desirable.




Figure 2. Synthetic methodology design


We recently developed a Pd-catalyzed hydroxycyclopropanol ring opening carbonylative lactonization to synthesize oxaspirolactones, another privileged scaffold frequently appears in natural products (2→5, Figure 2B).¹³ This new synthetic capability significantly facilitated our total syntheses of C₁₂ oxygenated diterpenes and *Stemona* alkaloids.¹⁴ With this success, we wondered the possibility of developing a catalytic carbonylation chemistry to

prepare THF/THP-fused γ -lactones with various substitution patterns (Figure 2C). To achieve this goal, cyclopropanols such as **6** would be required. They can be convergently assembled from readily available ester **7** and homoallylic alcohol **8** using the cyclopropanol synthesis¹⁵ protocol developed by Cha and co-workers.¹⁶ We envisioned a process involving Pd-catalyzed β -carbon elimination to generate Pd-homoenolate **9**. Ideally, we hoped for an acetal formation followed by carbonylative lactonization to convert **9** to desired fused bicyclic γ -lactone **11**, presumably via an intermediate like **10**. In principle, several processes could compete with the desired one. For instance, β -hydride elimination would decompose **9** to an enone; the tethered secondary alcohol could attack the Pd center and a following C-O reductive elimination would give THF product **12**. Additionally, carbonylative lactonization with the same tethered alcohol would give δ -lactone **14** (**9**→**13**→**14**). At the planning stage, we didn't know which pathway would dominate, but it would be ideal to develop conditions to access **11**, **12**, or **14** at wish. We were particularly attracted by the conversion of hydroxycyclopropanol **6** to fused lactone **11**. The established modular assembly of **6** would enable the introduction of various substitutions in product **11**, especially at the ring junction carbon where the R group resides. Meanwhile, we were hoping that the use of bishomoallylic alcohols may eventually result in THF-fused γ -lactones. Given the prevalence of THF/THP-fused γ -lactones in biologically active natural products, this chemistry is expected to find broad application in facilitating the chemical syntheses of those natural products and their analogs. Herein, we report such a Pd-catalyzed hydroxycyclopropanol ring opening carbonylative lactonization to the aforementioned bicyclic γ -lactones and its application in a short total synthesis of (\pm)-paeonilide (**1g**) and other medicinally important bicyclic and polycyclic scaffolds.

Our investigation started with known hydroxycyclopropanol **15a** (Table 1).¹⁶ When it was subjected to the carbonylative spiro-lactonization conditions we developed before with ([Pd(neoc)(OAc)]₂(OTf)₂ as catalyst and 1,4-benzoquinone (BQ) as oxidant,¹³ the formation of **16a** was not observed (entry 1). Switching [Pd(neoc)(OAc)]₂(OTf)₂ to palladium(II) trifluoroacetate (Pd(TFA)₂) was not fruitful either (entry 2). The breakthrough came by using Cu(OTf)₂ to replace BQ as the oxidant, which resulted in the formation of **16a** in 6% yield (entry 3). The structure of **16a** was unambiguously confirmed by X-ray crystallographic analysis.¹⁷ The use of 2,3-dichloro-5,6-dicyano-1,4-benzoquinone (DDQ) as oxidant increased the yield to 18% (entry 4). Further investigation of palladium catalysts led to the identification of Pd(OAc)₂ as an optimal choice (entry 5-6). Other 1,4-benzoquinone derivatives including 2,5-DMBQ and 2,6-DMBQ were not effective (entry 8, 9). THF (entry 7),

toluene (entry 10), and benzene (entry 14) were suitable solvents with benzene giving slightly better yield (also see **16e** and **16n**, Table 2). Elevating reaction temperature (40 °C, entry 11) or increasing the amount of DDQ (entry 12) led to reduced reaction yield. Finally, reducing the reaction concentration from 0.03 M to 0.01 M was beneficial and product **16a** was obtained in 74% yield (entry 14). The reaction is also scalable. Scaling up the reaction from 0.2 mmol to 2.92 mmol led to the formation of **16a** in 65% yield. Moreover, when enantioenriched homoallylic alcohol (1-phenylbut-3-en-1-ol, 92% ee) was used, **16a** was obtained in 92% ee (73% yield) after a sequence of cyclopropanol formation and carbonylative lactonization. Given the ready availability of enantioenriched homoallylic alcohols, our method offers an efficient asymmetric alternative to THF-fused γ -lactones.

Table 1. Reaction condition optimization^a



entry	reaction conditions	yield ^b
1	[Pd(neoc)(OAc)] ₂ (OTf) ₂ , BQ, DCE (0.03), rt	0%
2	Pd(TFA) ₂ , BQ, DCE (0.03), rt	0%
3	Pd(TFA) ₂ , Cu(OTf) ₂ , DCE (0.03), rt	6%
4	Pd(TFA) ₂ , DDQ, DCE (0.03), rt	18%
5	PdCl ₂ , DDQ, DCE (0.03), rt	0%
6	Pd(dppf)Cl ₂ , DDQ, DCE (0.03), rt	trace
7	Pd(OAc) ₂ , DDQ, THF (0.03), rt	58%
8	Pd(OAc) ₂ , 2,5-DMBQ, THF (0.03), rt	trace
9	Pd(OAc) ₂ , 2,6-DMBQ, THF (0.03), rt	trace
10	Pd(OAc) ₂ , DDQ, toluene (0.03), rt	65%
11	Pd(OAc) ₂ , DDQ, toluene (0.03), 40 °C	51%
12	Pd(OAc) ₂ , DDQ (3.0 equiv.), toluene (0.03), rt	58%
13	Pd(OAc) ₂ , DDQ, toluene (0.01), rt	70%
14	Pd(OAc) ₂ , DDQ, benzene (0.01), rt	74% ^{c,d}

^[a]General reaction conditions: To a stirred solution of **15a** (1.0 equiv., 0.2 mmol) and DDQ (2.0 equiv.) in benzene (0.01 M) under carbon monoxide (balloon) was added Pd(OAc)₂ (0.1 equiv.) in one portion. The resulting solution was stirred at room temperature until no more starting material left. The reaction process was monitored by thin-layer chromatography; ^[b]Isolated yield; ^[c]65% for 0.56-gram scale (2.92 mmol); ^[d] 73% yield and 92% ee when **15a** was prepared from the corresponding enantioenriched homoallylic alcohol with 92% ee; 2,5-DMBQ: 2,5-dimethylbenzoquinone; 2,6-DMBQ: 2,6-dimethylbenzoquinone.

We then evaluated the scope and generality of this new Pd-catalyzed hydroxycyclopropanol ring opening carbonylative lactonization. The reaction proved to be very robust. A variety of hydroxycyclopropanol substrates could be transformed into the corresponding *cis*-fused bicyclic γ -lactones. THF-fused γ -lactone products with either an aryl or an alkyl group at the C5-position could be obtained. Notably, tosylate (**16e**), TBS-ether group (**16k**, **16l**, **16s**), and heteroaromatics including indole (**16g**), furan (**16h**), thiophene (**16i**), and pyrrole (**16j**) are all tolerated under the mild reaction conditions. Substrates containing sterically hindered secondary (**16m**) and tertiary alcohols (**16n**, **16o**, **16p**, **16q**) were effective as well. *Most*





Table 2. Substrate scope


*of the previously reported methods are limited to have hydrogen atom at C6 of the ring junction. In our case, the substituent at this position can be altered from a methyl group to a hydrogen (**16w**) or other bulkier alkyl groups including cyclohexyl and isopropyl groups (**16r**, **16s**, **16t**, **16u**, **16v**, **16x**, and **16y**). Variation of the stereochemistry of the secondary alcohol on the side chain was tolerated*

as well (cf. **16x** vs **16y**). Similar reaction yields were obtained, but the reaction of **16y** took longer time. Furthermore, THP-fused γ -lactones could be produced in modest to good yield (**16za**, **16zb**, **16zc**, and **16zd**). Additionally, slow addition of DDQ to the reaction mixture could be beneficial to improve the reaction yield (cf. **16c** and **16d**), but was not employed to all the substrates due to the operational inconvenience.




Scheme 1. A “competition” case

We also evaluated hydroxycyclopropanol **17**, which could undergo ring opening carbonylation to form either fused γ -lactone **21** or spiro lactone **22** presumably via intermediate **19** or **20**, respectively (Scheme 1). Both **19** and **20** could be derived from the same palladium-homoenolate **18**. When **17** was subjected to the optimized carbonylative conditions (entry 14, Table 1), fused γ -lactone **21** was obtained in 47% yield without any spiro lactone **22** formation. Interestingly, even when the carbonylative spiro lactonization conditions were used ($[\text{Pd}(\text{neoc})(\text{OAc})]_2(\text{OTf})_2$ and BQ),¹³ fused lactone was still the dominant product, but in lower yield.



Scheme 2. Total synthesis of (±)-paeonilide

We next decided to test this new carbonylative lactonization method in natural product total synthesis and chose paeonilide (**1g**) as our initial target. Paeonilide is a monoterpene isolated from *Paeonia delavayi* by Liu and co-workers.¹⁸ It selectively inhibits platelet aggregation induced by platelet activating factor and has been previously synthesized by the groups of Zhang (chiral pool synthesis starting from (*R*)-(-)-carvone, 14 LLS steps,^{19a} racemic, 17 LLS steps^{19b}), Du (racemic, 6 LLS steps),^{19c} Reiser (enantioselective, 12 LLS steps),^{19d} and Argade (racemic, 11 LLS steps)^{19e} from commercially available starting materials. Our synthesis started with known compounds **23**²⁰ and **24**,²¹ which were prepared in one and three steps, respectively, from commercially available starting materials. They were united via a Kulinkovich reaction to give desired hydroxycyclopropanol **25**, albeit in low yield due to the existence of the ketal, free alcohol, and primary TBS ether. When **25** was subjected to the carbonylative lactonization conditions, bicyclic γ -lactone **26** was produced in 42% yield (99% yield brsm). Removal of the TBS and ketal protecting groups under acidic



Scheme 3. Synthetic diversifications.

conditions followed by benzoate formation completed the total synthesis (\pm)-paeonilide in 4 LLS steps in 3.6% yield from known compounds **23** and **24** or 7 LLS steps from commercially available starting materials.

Additionally, the fused bicyclic lactone product can be diversified to other medicinally important structural scaffolds (Scheme 3). α -Methylene γ -butyrolactone is another privileged scaffold frequently found in natural products.²² Treatment of **16a** with the Bredereck's reagent followed by DIBAL-H reduction²³ could convert **16a** to α -methylene γ -butyrolactone containing bicyclic product **28** in high yield. The lactone moiety of **16a** could also be converted to a lactam (cf. **30**) via the treatment of MeNH₂ followed by acid-promoted cyclization. Aldol reaction of **16a** with benzaldehyde gave a 3:1 mixture of diastereomers in 82% yield. The relative stereochemistry of the major product **31** was established by X-ray crystallographic analysis of its *p*-bromobenzoate derivative **32**.¹⁷ Hexahydrofuro[2,3-*b*]furan is an important scaffold in protease inhibitors, including the FDA-approved anti-HIV drug Darunavir (**34**).²⁴ Bicyclic lactone **16a** could be readily converted to hexahydrofuro[2,3-*b*]furan-containing product **33** via a sequential DIBAL-H and triethylsilane reduction process. Furthermore, **16a** could also be elaborated to tricyclic products **36** and **37** via a series of standard functional group manipulations.²⁵ These two tricyclic scaffolds have been found in natural products such as gracilin B (**1b**) and potent protease inhibitors including GRL-0519A.²⁶ These diversification pathways demonstrate the potential broad application of the newly developed carbonylative lactonization method.

In summary, we have developed a novel Pd-catalyzed hydroxycyclopropanol ring opening carbonylative lactonization to synthesize THF/THP-fused bicyclic γ -lactones. This method features mild reaction conditions, good functional group tolerability, and scalability. Its application was demonstrated in a short total synthesis of racemic paeonilide. The bicyclic lactone products could be further elaborated to other valuable structures.

ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website.

Experimental procedures and spectra data (PDF file)
Cystallographic data for **16a** and **32** (cif file)

AUTHOR INFORMATION

Corresponding Author

* Email: mjdai@purdue.edu

ORCID

Mingji Dai: 0000-0001-7956-6426

Xinpei Cai: 0000-0002-2633-1882

Weida Liang: 0000-0001-5205-0374

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENT

This work was financially supported by NSF CAREER Award 1553820. The NIH P30 CA023168 is acknowledged for supporting shared NMR resources to Purdue Center for Cancer Research. The XRD data is collected on a new single crystal X-ray diffractometer supported by the NSF through the Major Research Instrumentation Program under Grant No. CHE 1625543.

REFERENCES

- (1) (a) Wilson, R. M.; Danishefsky, S. J. Small molecule natural products in the discovery of therapeutic agents: the synthesis connection. *J. Org. Chem.* **2006**, *71*, 8329-8351. (b) Newman, D. J.; Cragg, G. M. Natural products as sources of new drugs over the 30 years from 1981 to 2010. *J. Nat. Prod.* **2012**, *75*, 311-335. (c) Davison, E. K.; Brimble, M. A. Natural Product Derived Privileged Scaffolds in Drug Discovery. *Curr. Opin. in Chem. Biol.* **2019**, *52*, 1-8. (d) Rodrigues, T.; Reker, D.; Schneider, P.; Schneider, G. Counting on natural products for drug design. *Nat. Chem.* **2016**, *8*, 531-541.
- (2) Garnsey, M. R.; Slutskyy, Y.; Jamison, C. R.; Zhao, P.; Lee, J.; Rhee, Y. H.; Overman, L. E. Short Enantioselective Total Syntheses of Cheloviolenes A and B and Dendrillolide C via Convergent Fragment Coupling Using a Tertiary Carbon Radical. *J. Org. Chem.* **2018**, *83*, 6958-6976.
- (3) Sakabe, N.; Takada, S.; Okabe, K. The structure of ginkgolide A, a novel diterpenoid trilactone. *J. Chem. Soc., Chem. Commun.* **1967**, 259-261.
- (4) Mayol, L.; Piccialli, V.; Sica, D. Minor Bisnorditerpenes from the Marine Sponge *Spongionella gracilis* and Revision of the Δ^6 Configuration of Gracilin B. *J. Nat. Prod.* **1986**, *49*, 823-828.
- (5) Chen, X.-L.; Liu, H.-L.; Li, J.; Xin, G.-R.; Guo, Y.-W. Paracaseolide A, First α -Alkylbutenolide Dimer with an Unusual Tetraquinane Oxa-Cage Bislactone Skeleton from Chinese Mangrove *Sonneratia Paracaseolaris*. *Org. Lett.* **2011**, *13*, 5032-5035.
- (6) von Salm, J. L.; Witowski, C. G.; Fleeman, R. M.; McClintock, J. B.; Amsler, C. D.; Shaw, L. N.; Baker, B. J. Darwinolide, a New Diterpene Scaffold That Inhibits Methicillin-Resistant *Staphylococcus aureus* Biofilm from the Antarctic Sponge *Dendrilla membranosa*. *Org. Lett.* **2016**, *18*, 2596-2599.
- (7) (a) Hochlowski, J.; Faulkner, D. J.; Matsumoto, G.; Clardy, J. Synthetic Studies on Norrisolide: Enantioselective Synthesis of the Norrisane Side Chain. *J. Org. Chem.* **1983**, *48*, 1141-1142. (b) Guizzunti, G.; Brady, T. P.; Fischer, D.; Malhotra, V.; Theodorakis, E. A. Chemical biology studies on norrisolide. *Bioorg. Med. Chem.* **2010**, *18*, 2115-2122.
- (8) Buckleton, J. S.; Cambie, R. C.; Clark, G. R. Structures of cheloviolene A from the sponge *Chelonaphysilla violacea*. *Acta Crystallogr., Sect. C: Cryst. Struct. Commun.* **1991**, *47*, 1438-1440.
- (9) (a) Luo, Q.; Di, L.; Yang, X. H.; Cheng, Y. X. Applanatumols A and B, meroterpenoids with unprecedented skeletons from *Ganoderma applanatum*. *RSC Adv.* **2016**, *6*, 45963-45967. (b) Qi, B.; Liu, X.; Mo, T.; Zhu, Z.; Li, J.; Wang, J.; Shi, X.; Zeng, K.; Wang, X.; Tu, P.; Abe, I.; Shi, S. 3,5-Dimethylorsellinic Acid Derived Meroterpenoids from

1 *Penicillium chrysogenum* MT-12, an Endophytic Fungus Isolated from
2 *Huperzia serrata*. *J. Nat. Prod.* **2017**, *80*, 2699–2707. (c) Tan, S.-J.;
3 Lim, J.-L.; Low, Y.-Y.; Sim, K.-S.; Lim, S.-H.; Kam, T.-S. Oxidized
4 Derivatives of Macroline, Sarpagine, and Pleiocarpamine Alkaloids
5 from *Alstonia angustifolia*. *J. Nat. Prod.* **2014**, *77*, 2068–2080. (d) Li,
6 T. X.; Liu, R. H.; Wang, X. B.; Luo, J.; Luo, J. G.; Kong, L. Y.; Yang,
7 M. H. Hypoxia-protective azaphilone adducts from *Peyronella glom-*
8 *erata*. *J. Nat. Prod.* **2018**, *81*, 1148–1153. (e) Aziz, A. N.; Ismail, N.
9 H.; Halim, S. N. A.; Looi, C. Y.; Anouar, E. H.; Langat, M. K.; Mul-
10 holland, D.; Awang, K. Laevifins A–G, clerodane diterpenoids from
11 the Bark of *Croton oblongus* Burm.f. *Phytochemistry* **2018**, *156*,
12 193–200.

13 (10) (a) Corey, E. J.; Kang, M. C.; Desai, M. C.; Ghosh, A. K.;
14 Houppis, I. N. Total synthesis of (+)-ginkgolide B. *J. Am. Chem. Soc.*
15 **1988**, *110*, 649–651. (b) Corey, E. J.; Letavic, M. A. Enantioselective
16 Total Synthesis of Gracilins B and C Using Catalytic Asymmetric
17 Diels-Alder Methodology. *J. Am. Chem. Soc.* **1995**, *117*, 9616–9617.
18 (c) Crimmins, M. T.; Pace, J. M.; Nantermet, P. G.; Kim-Meade, A. S.;
19 Thomas, J. B.; Watterson, S. H.; Wagman, A. S. The Total Synthesis
20 of (±)-Ginkgolide B. *J. Am. Chem. Soc.* **2000**, *122*, 8453–8463. (d)
21 Brady, T. P.; Kim, S. H.; Wen, K.; Theodorakis, E. A. Stereoselective
22 Total Synthesis of (+)-Norrisolide. *Angew. Chem., Int. Ed.* **2004**, *43*,
23 739–742 (e) Shi, H.; Fang, L.; Tan, C.; Shi, L.; Zhang, W.; Li, C. C.;
24 Luo, T.; Yang, Z. Total Syntheses of Drimane-Type Sesquiterpenoids
25 Enabled by a Gold-Catalyzed Tandem Reaction. *J. Am. Chem. Soc.*
26 **2011**, *133*, 14944–14947. (f) Guney, T.; Kraus, G. A. Total Synthesis
27 of Paracaseolide A. *Org. Lett.* **2013**, *15*, 613–615. (g) Wang, L.; Wang,
28 H.; Li, Y.; Tang, P. Total Synthesis of Schilancitrilactones B and C.
29 *Angew. Chem., Int. Ed.* **2015**, *54*, 5732–5735. (h) Yuan, Z.; Hu, X.;
30 Zhang, H.; Liu, L.; Chen, P.; He, M.; Xie, X.; Wang X.; She, X. Total
31 synthesis of conosilane A via a site-selective C–H functionalization
32 strategy. *Chem. Commun.*, **2018**, *54*, 912–915. (i) Okamura, H.; Mori,
33 N.; Watanabe, H.; Takikawa, H. Synthesis of both enantiomers of
34 conosilane A. *Tetrahedron Lett.* **2018**, *59*, 4397–4400. (j) Siemon, T.;
35 Steinhauer, S.; Christmann, M. Synthesis of (+) - Darwinolide, a Bio-
36 film - Penetrating Anti - MRSA Agent. *Angew. Chem., Int. Ed.* **2019**,
37 *58*, 1120–1122. (k) Li, C.; Quan, T.; Xue, Y.; Cao, Y.; Chen, S. C.; Luo,
38 T. Synthesis of 17-Deacetoxy Chromodorolide B Based on a Gold-
39 Catalyzed Alkoxy cyclization Reaction. *Org. Lett.*, **2020**, *22*, 1655-
40 1658. (l) Uchida, K.; Kawamoto, Y.; Kobayashi, T.; Ito, H. (+/–)-Lu-
41 cidumone, a COX-2 Inhibitory Caged Fungal Meroterpenoid from
42 *Ganoderma lucidum*. *Org. Lett.* **2019**, *21*, 6199–6201. (m) Li, Y.;
43 Zhang, Q.; Wang, H.; Cheng, B.; Zhai, H. Bioinspired Total Synthesis
44 of (±)-Chaetophenol C Enabled by a Pd-Catalyzed Cascade Cycliza-
45 tion. *Org. Lett.* **2017**, *19*, 4387–4390.

46 (11) (a) Allred, T. K.; Dieskau, A. P.; Zhao, P.; Lackner, G. L.; Over-
47 man, L. E. Enantioselective Total Synthesis of Macfarlandin C, a
48 Spongian Diterpenoid Harboring a Concave - Substituted cis - Dioxabi-
49 cyclo [3.3.0] octanone Fragment. *Angew. Chem., Int. Ed.* **2020**, *59*, 1-
50 6. (b) Tao, D. J.; Slutskyy, Y.; Muuronen, M.; Le, A.; Kohler, P.; Over-
51 man, L. E. Total synthesis of (–)-chromodorolide B by a computa-
52 tionally guided radical addition/cyclization/fragmentation cascade. *J. Am.*
53 *Chem. Soc.* **2018**, *140*, 3091–3102. (c) Slutskyy, Y.; Jamison, C. R.;
54 Zhao, P.; Lee, J.; Rhee, Y. H.; Overman, L. E. Versatile Construction
55 of 6-Substituted Cis-2,8- Dioxabicyclo[3.3.0]Octan-3-Ones: Short En-
56 antioselective Total Syntheses of Cheloviolenes A and B and Dendrilo-
57 lide C. *J. Am. Chem. Soc.* **2017**, *139*, 7192–7195. (d) Tao, D. J.; Slut-
58 skyy, Y.; Overman, L. E. Total synthesis of (–)-chromodorolide B. *J.*
59 *Am. Chem. Soc.* **2016**, *138*, 2186–2189.

(12) (a) Trost, B. M.; Toste, F. D. Palladium-Catalyzed Kinetic and
Dynamic Kinetic Asymmetric Transformation of 5-Acyloxy-2-(5H)-
furanone. Enantioselective Synthesis of (–)-Aflatoxin B Lactone. *J. Am.*
Chem. Soc. **1999**, *121*, 3543–3544. (b) Kim, C.; Hoang, R.; Theodorakis,
E. A. Synthetic studies on norrisolide: enantioselective synthesis
of the norrisane side chain. *Org. Lett.* **1999**, *1*, 1295–1297. (c) Weisser,
R.; Yue, W.; Reiser, O. Enantioselective Synthesis of Furo[2,3b]furan,
a Spongian Diterpenoid Substructure. *Org. Lett.* **2005**, *7*,
5353–5356. (d) Schnermann, M. J.; Beaudry, C. M.; Genung, N. E.;
Canham, S. M.; Untiedt, N. L.; Karanikolas, B. D. W.; Sütterlin, C.;
Overman, L. E. Divergent Synthesis and Chemical Reactivity of Bicy-
clic Lactone Fragments of Complex Rearranged Spongian Diterpenes.
J. Am. Chem. Soc. **2011**, *133*, 17494–17503. (e) Yousuf, S. K.; Taneja,
S. C.; Mukherjee, D. Highly Regio- and Stereoselective One-Pot Syn-
thesis of Carbohydrate-Based Butyrolactones. *Org. Lett.* **2011**, *13*, 576-
579. (f) Zhao, W.- W.; Liu, Y.-K. Enantio- and diastereoselective syn-
thesis of tetrahydrofuro[2,3-b]furan-2(3H)-one derivatives and related
oxygen heterocycles via an asymmetric organocatalytic cascade pro-
cess. *Org. Chem. Front.* **2017**, *4*, 2358–2363. (g) Thorat, S. S.; Kataria,
P.; Kontham, R. Synthesis of Furo[2,3-b]pyran-2-ones through Ag(I)-
or Ag(I)–Au(I)-Catalyzed Cascade Annulation of Alkynols and α -Ke-
toesters. *Org. Lett.* **2018**, *20*, 872–875. (h) Zhang, T.; Shimizu, Y.; Fu-
kaya, S.; Sawa, T.; Maekawa, H. Construction of a CF₃-Containing
Benzofurofuranone Skeleton from Coumarins via Reductive Coupling
and Acid-Mediated Ring Contraction. *J. Org. Chem.* **2019**, *84*, 12165-
12171.

(13) Davis, D. C.; Walker, K. L.; Hu, C.; Zare, R. N.; Waymouth, R.
M.; Dai, M. J. Catalytic Carbonylative Spirolactonization of Hy-
droxycyclopropanols. *J. Am. Chem. Soc.* **2016**, *138*, 10693–10699.

(14) (a) Ma, K.; Yin, X.; Dai, M. J. Total Syntheses of Bisdehydro-
neostemoninine and Bisdehydrostemoninine by Catalytic Carbonylative
Spirolactonization. *Angew. Chem., Int. Ed.* **2018**, *57*,
15209–15212. (b) Yin, X.; Ma, K.; Dong, Y.; Dai, M. J. Pyrrole Strat-
egy for the γ -Lactam-Containing *Stemona* Alkaloids: (±)Stemoamide,
(±)Tuberostemoamide, and (±)Sessilifoliamide A. *Org. Lett.* **2020**, *22*,
5001–5004.

(15) For reviews: (a) Kulinkovich, O. G. The Chemistry of Cyclo-
propanols. *Chem. Rev.* **2003**, *103*, 2597–2632. (b) Mack, D. J.; Njardar-
son, J. T. Recent Advances in the Metal-Catalyzed Ring Expansions of
Three and Four-Membered Rings. *ACS Catal.* **2013**, *3*, 272–286. (c)
Nikolaev, A.; Orellana, A. Transition-Metal-Catalyzed C–C and C–X
Bond-Forming Reactions Using Cyclopropanols. *Synthesis* **2016**, *48*,
1741–1768. (d) Mills, L. R.; Rousseaux, S. A. L. Modern Develop-
ments in the Chemistry of Homo-enolates. *Eur. J. Org. Chem.* **2019**, *8*-
26. (e) Cai, X.; Liang, W.; Dai, M. Total Syntheses via Cyclopropanols.
Tetrahedron **2019**, *75*, 193–208.

(16) (a) Lee, J.; Kim, H.; Cha, J. K. A New Variant of Kulinkovich
Hydroxycyclopropanation. Reductive Coupling of Carboxylic Esters
with Terminal Olefins. *J. Am. Chem. Soc.* **1996**, *118*, 4198–4199. (b)
Quan, L. G.; Kim, S.-H.; Lee, J. C.; Cha, J. K. Diastereoselective Syn-
thesis of trans-1,2-Dialkylcyclopropanols by the Kulinkovich Hy-
droxycyclopropanation of Homoallylic Alcohols. *Angew. Chem. Int.*
Ed. **2002**, *41*, 2160–2162.

(17) CCDC 1997763 and 2016932 contain the supplementary crys-
tallographic data for **16a** and **32**.

(18) Liu, J. K.; Ma, Y. B.; Wu, D. G.; Lu, Y.; Shen, Z. Q.; Zheng,
Q. T.; Chen, Z. H. Paeonilide, a Novel Anti-PAF-active Monoterpe-
noid-derived Metabolite from *Paeonia delavayi*. *Biosci. Biotechnol.*
Biochem. **2000**, *64*, 1511–1514.

1 (19) (a) Wang, C.; Zhang, H.; Liu, J.; Ji, Y.; Shao, Z.; Li, L. Stereoselective Synthesis of (+)-Paeonilide and Confirmation of its Absolute Configuration. *Synlett* **2006**, 1051-1054. (b) Wang, C.; Liu, J.; Ji, Y.; Zhao, J.; Li, L.; Zhang, H. Total Synthesis of (±)-Paeonilide. *Org. Lett.* **2006**, *8*, 2479-2481. (c) Du, Y.; Liu, J.; Linhardt, R. J. Facile Synthesis of (±)-Paeonilide. *J. Org. Chem.* **2007**, *72*, 3952-3954. (d) Harrar, K.; Reiser, O. Enantioselective synthesis of (-)-paeonilide. *Chem. Commun.* **2012**, *48*, 3457-3759. (e) Deore, P. S.; Argade, N. P. Reactivity Umpolung in Intramolecular Ring Closure of 3,4-Disubstituted Butenolides: Diastereoselective Total Synthesis of Paeonilide. *Org. Lett.* **2013**, *15*, 5826-5829.

2 (20) Ueda, Y.; Abe, H.; Iguchi, K.; Ito, H. Synthetic Study of Yonanolide: Stereoselective Construction of the Tricyclic Core. *Tetrahedron Lett.* **2011**, *52*, 3379-3381.

3 (21) Lohse-Fraefel, N.; Carreira, E. M. Polyketide Building Blocks via Diastereoselective Nitrile Oxide Cycloadditions with Homoallylic Alcohols and Monoprotected Homoallylic Diols. *Chem. Eur. J.* **2009**, *15*, 12065-12081.

4 (22) Jackson, P. A.; Widen, J. C.; Harki, D. A.; Brummond, K. M. Covalent Modifiers: A Chemical Perspective on the Reactivity of α,β -

Unsaturated Carbonyls with Thiols via Hetero-Michael Addition Reactions. *J. Med. Chem.* **2017**, *60*, 839-885.

5 (23) Barthel, A.; Kaden, F.; Jager, A.; Metz, P. Enantioselective Synthesis of Guaianolides in the Osmitopsin Family by Domino Metathesis. *Org. Lett.* **2016**, *18*, 3298-3301.

6 (24) Ghosh, A. K.; Chapsal, B. D.; Weber, I. T.; Mitsuya, H. Design of HIV Protease Inhibitors Targeting Protein Backbone: An Effective Strategy for Combating Drug Resistance. *Acc. Chem. Res.* **2008**, *41*, 78-86.

7 (25) Rashid, S.; Bhat, B. A.; Mehta, G. Regenerative γ -Lactone Annulations: A Modular, Iterative Approach to Oligo-tetrahydrofuran Molecular Stairs and Related Frameworks. *Org. Lett.* **2015**, *17*, 3604-3607.

8 (26) Amano, M.; Koh, Y.; Das, D.; Li, J.; Leschenko, S.; Wang, Y. F.; Boross, P. I.; Weber, I. T.; Ghosh, A. K.; Mitsuya, H. A novel bis-tetrahydrofuranylurethane-containing nonpeptidic protease inhibitor (PI), GRL-98065, is potent against multiple-PI-resistant human immunodeficiency virus in vitro. *Antimicrob. Agents Chemother.* **2007**, *51*, 2143-2155.

Table of Contents

