

Direct Hydroxylation of Benzene to Phenol Using Hydrogen Peroxide Catalyzed by Nickel Complexes Supported by Pyridylalkylamine Ligands

Yuma Morimoto, Shuji Bunno, Nobutaka Fujieda, Hideki Sugimoto, and Shinobu Itoh*

Department of Material and Life Science, Division of Advanced Science and Biotechnology, Graduate School of Engineering, Osaka University, 2-1 Yamadaoka, Suita, Osaka 565-0871, Japan

Supporting Information

ABSTRACT: Selective hydroxylation of benzene to phenol has been achieved using H₂O₂ in the presence of a catalytic amount of the nickel complex $[Ni^{II}(tepa)]^{2+}(2)$ (tepa = tris[2-(pyridin-2-yl)ethyl]amine) at 60 °C. The maximum yield of phenol was 21% based on benzene without the formation of quinone or diphenol. In an endurance test of the catalyst, complex 2 showed a turnover number (TON) of 749, which is the highest value reported to date for molecular catalysts in benzene hydroxylation with H2O2. When toluene was employed as a substrate instead of benzene, cresol was obtained as the major product with 90% selectivity. When $H_2^{18}O_2$ was utilized as the oxidant, ¹⁸O-labeled phenol was predominantly obtained. The reaction rate for fully deuterated benzene was nearly identical to that of benzene (kinetic isotope effect = 1.0). On the basis of these results, the reaction mechanism is discussed.

Phenol and its derivatives are an important class of chemicals as precursors of dyes, pharmaceuticals, polymers, etc.¹ Phenol and cresol are currently produced in industry by the cumene process, which consists of three chemical steps: propylation of benzene or toluene, autoxidation to the cumene hydroperoxide derivative, and Hock rearrangement.^{1,2} These reactions are carried out under high temperature, high pressure, and strongly acidic conditions, and an equimolar amount of acetone is produced as a byproduct in the final step. In addition, the overall yield of phenol from benzene is very low (5%). Thus, a much simpler and more efficient process that proceeds under mild conditions is desired. In particular, a direct aromatic oxygenation reaction with an economically and environmentally benign oxidant such as O_2 or H_2O_2 has remained a focal point for extensive research efforts.³⁻¹⁵ However, the direct introduction of hydroxyl functionality into the aromatic C–H bond with such oxidants is challenging because of the notoriously low reactivity of aromatic C-H bonds and the low product selectivity caused by overoxidation of phenols.^{16–18}

In the previous studies of transition-metal complexes of active oxygen species, dinuclear complexes such as dicopper(II) μ - $\eta^2:\eta^2$ -peroxide, dicopper(III) bis(μ -oxo), and dinickel(III) bis(μ -oxo) complexes supported by pyridylalkylamine ligands have been demonstrated to exhibit hydroxylation reactivity toward aromatic groups embedded in their supporting ligands.¹⁹⁻²⁵ To the best of our knowledge, however, there has been no report to

date of *intermolecular* hydroxylation of externally added benzene derivatives with transition-metal complexes supported by pyridylalkylamine ligands, although a large number of active-oxygen transition-metal complexes have been developed to date.^{26–29} We herein report a benzene hydroxylation reaction employing nickel complexes supported by pyridylalkylamine ligands as the catalysts and H₂O₂ as the oxidant under mild conditions.

We first examined the catalytic activity of $[Ni^{II}(tmpa)]^{2+}$ (1) (tmpa = tris[2-(pyridin-2-yl)methyl]amine), $[Ni^{II}(tepa)]^{2+}$ (2) (tepa = tris[2-(pyridin-2-yl)ethyl]amine), and $[Ni^{II}(bepa)]^{2+}$ (3) (bepa = *N*,*N*-bis[2-(pyridin-2-yl)ethyl]-2-phenylethylamine) in the benzene hydroxylation reaction with H₂O₂ (for the ligand structures, see Chart 1).^{30,31,25} Treatment of benzene

Chart 1. Ligands for Ni-Based Hydroxylation Catalysis



(5.0 μ mol) with H₂O₂ (2.5 mmol) in the presence of a catalytic amount of **1**, **2**, or **3** (0.50 μ mol, 10 mol %) and triethylamine (TEA) (5.0 μ mol) afforded phenol in 18%, 21%, or 2% yield based on benzene, respectively, after 5 h (Table 1, entries 1–3).³² This is the first example of hydroxylation of benzene with H₂O₂ catalyzed by nickel complexes. When **1** was employed as the catalyst, the overoxidation product benzoquinone (BQ) was also obtained in 9% yield. In contrast, only a negligible amount (<1% yield) of BQ was generated with catalyst **2** (Figure S1 in the Supporting Information (SI)). The 21% yield of phenol without formation of the overoxidation product (<1%) (entry 2) is the best result reported to date for homogeneous systems using H₂O₂ (Scheme 1; for other systems, see Table S1 in the SI).^{33,34}

Received: February 18, 2015

Table 1. Catalytic Activity of Ni(II) Complexes

entry	cat	conv. (%)	yield (%)	selectivity (%)
1	1	27	18	66
2	2	23	21	>99
3	3	25	2	8
4	4	7	0	-
5	5	10	0	_

Conditions: benzene (5.0 $\mu mol),~H_2O_2$ (2.5 mmol), catalyst (0.50 $\mu mol)$, and TEA (5.0 $\mu mol)$ in MeCN for 5 h at 60 °C.

Scheme 1. Catalytic Hydroxylation of Benzene with 2



The reaction also proceeded at room temperature, but the yield was lower (14% in 5 h). When phenol was employed as the substrate (5.0 mol) with 2 under the same conditions, the hydroxylation reaction hardly took place (phenol was recovered almost quantitatively).

Nickel complexes supported by the tetraaza macrocyclic ligands such as $[Ni^{II}(12-tmc)]^{2+}$ (4) (12-tmc = 1,4,7,10-tetraatexcyclododecane) and $[Ni^{II}(14-tmc)]^{2+}$ (5) (14-tmc = 1,4,8,11-tetramethyl-1,4,8,11-tetraatexcyclotetradecane) did not show any catalytic ability for the reaction (entries 4 and 5).^{35,36} Nickel(II) acetate alone or with 2,2'-bipyridine (bpy) as the ligand also did not work as the catalyst.

Since complex 2 showed the highest activity and selectivity, an endurance test was performed using 0.01 mol % 2, 0.10 mol % TEA, and 5.0 equiv of H_2O_2 .³⁷ In this case, the turnover number (TON) of 2 reached 749 with a 7.5% yield of phenol after 216 h without the formation of substantial amounts of the over-oxidation products (Figure 1).³⁸ This is the highest TON



Figure 1. Time course of the turnover number in the catalytic oxidation of benzene (5.0 mmol) by H_2O_2 (25 mmol) in the presence of **2** (0.50 μ mol) and TEA (5.0 μ mol) in 3.0 mL of MeCN at 60 °C.

reported for benzene hydroxylation reactions using transitionmetal complexes as molecular catalysts; among molecular catalysts showing high selectivity (>95%), the polyoxometalate (POM) catalyst Py_1 – $PMo_{10}V_2O_{40}$ showed a TON of 21, which was the highest number reported previously (Table S1).

The system could also be applied to the hydroxylation of toluene. In the presence of 5 equiv of H_2O_2 and 0.01 mol % 2, the TON reached 96 after 24 h to give cresol as the major product without the formation of quinone derivatives (Table 2, entry 1).

Table 2. Product Distributions in Catalytic Oxidation of Alkylbenzenes with 2



Conditions: substrate (5.0 mmol), H_2O_2 (5 equiv), 2 (0.01 mol %), and TEA (0.10 mol %) in CH₃CN at 60 °C for 24 h. ^{*a*}Ratio of products given by aromatic hydroxylation (sp²), alkyl-substituent oxygenation (sp³), and quinone derivatives (q). ^{*b*}Ratio of products of hydroxylation at the ortho, meta, para, and ipso positions. ^{*c*}Selectivity for aromatic hydroxylation products except phenol. ^{*d*}Calculated on the basis of the amount of aromatic hydroxylation products, except for phenol, which was calculated on the basis of the substrate.

It should be noted that the aromatic sp^2 carbons were preferentially oxygenated with a high selectivity of 90% (Scheme S1 in the SI) despite the fact that the bond dissociation energy (BDE) of the C–H bond of a benzylic sp^3 carbon (90 kcal mol⁻¹) is much lower than that of an aromatic C–H bond (112 kcal mol⁻¹).³⁹

Mizuno and co-workers reported selective aromatic oxygenation of toluene with H_2O_2 using the POM catalyst [γ - $PW_{10}O_{38}V_2(\mu-OH)_2]^{3-}$, where a high aromatic carbon to benzylic carbon selectivity of 98:2 was obtained. However, the total selectivity for cresol decreased to 86% as a result of overoxidation of cresol to methyl-*p*-quinone. Thus, the present system showed the highest selectivity for cresol formation from toluene with H_2O_2 .⁴⁰ The regioselectivity of the present system was also different from that of Mizuno's system: with nickel catalyst 2, the ortho position was mainly oxygenated, whereas the para position of toluene was mainly oxygenated with Mizuno's system. Stimulated by this result, we examined the oxygenation of other alkylbenzene derivatives with more reactive benzylic C-H bonds, namely, ethylbenzene (BDE = 87 kcal mol⁻¹) and cumene (BDE = $84.5 \text{ kcal mol}^{-1}$), in the presence of a catalytic amount of 2 (0.01 mol %).³⁹ To our delight, oxygenation of the sp² carbons of those substrates also proceeded, although the selectivity for sp² carbons slightly decreased (Table 2, entries 2 and 3; for a comparison with previous reports, see Tables S3-S5). In these cases as well, no overoxidation products such as hydroquinone or quinone derivatives were obtained (Schemes S2 and S3). In the oxygenations of ethylbenzene and cumene, the ortho position was mainly attacked, as in the case of toluene oxygenation. However, the selectivity slightly decreased with increasing bulkiness of the substituent, and the selectivity for the ipso position to produce phenol increased.

To gain insight into the reaction mechanism, the kinetic isotope effect (KIE) was examined using a 1:1 mixture of C_6D_6 and C_6H_6 . From the ratio of phenol to phenol- d_5 , the KIE value was determined to be 1.0, which is close to the KIE values for intramolecular aromatic hydroxylation of the dicopper(II) μ -

Journal of the American Chemical Society

 $\eta^{2}:\eta^{2}$ -peroxide, dicopper(III) bis(μ -oxo), and dinickel(III) bis(μ -oxo) complexes (0.9–1.0).^{21–23} Such a small or negligible KIE value is typical in aromatic hydroxylation by metal—active oxygen species via the electrophilic aromatic substitution mechanism.⁴¹ Such a KIE value also suggests that a Fenton-type reaction is not likely to be involved, as the KIE value for the Fenton reaction was reported to be 1.7.⁴² Metal insertion is not a plausible mechanism either, as the KIE value for C–H bond activation via Pd(0) insertion was reported to be 2.9–4.8.⁴³

To verify the origin of the oxygen atom, benzene oxygenation was carried out using $H_2^{18}O_2$ with 90% atom purity in $H_2^{16}O$. In this case, ¹⁸O was incorporated into the phenol product nearly quantitatively (92%), whereas no ¹⁸O atom was incorporated into the product when the reaction was carried out in $H_2^{18}O$ using $H_2^{16}O_2$. These results unambiguously indicated that H_2O_2 is the oxygen source in the present reaction.⁴⁴

A number of nickel complexes supported by pyridylalkylamine ligands, including *N*,*N*-bis[(6-phenylpyridin-2-yl)methyl]-*N*-[(pyridin-2-yl)methyl]amine (Ph₂tmpa) and bepa have been demonstrated to generate dinickel(III) bis(μ -oxo) complexes in the reaction with H₂O₂. Moreover the Suzuki group and our group have independently reported intramolecular aromatic oxygenation in the dinickel(III) bis(μ -oxo) supported by pyridylalkylamine ligands under low-temperature conditions (Scheme 2).⁴⁵ In fact, a nickel(II) complex supported by

Scheme 2. Intramolecular Aromatic Hydroxylation by a Dinickel(III) $Bis(\mu$ -oxo) Complex²⁵



Ph₂tmpa could oxygenate benzene under the same reaction conditions as shown in Table 1, although the yield of phenol was only 3%. The low yield could be attributed to the competitive intramolecular reaction, as the product of intramolecular oxygenation of the phenyl group of the ligand was also found in the electrospray ionization (ESI) mass spectrum of the final reaction solution (Figure S2). Hydroxylation of aliphatic C-H bonds has also been demonstrated to be a characteristic feature of dinickel(III) bis(μ -oxo) complexes supported by pyridylalkylamine ligands, whereas aliphatic hydroxylated products are minor products in the present alkylbenzene oxygenation.^{46–49} By contrast, nickel(II) complexes supported by the tetraaza macrocyclic ligands 12-tmc and 14-tmc have been shown to provide a mononuclear nickel(III) peroxide complex and a nickel(II) superoxide complex, respectively, in the reaction with H_2O_2 ; neither Ni(III) peroxide nor Ni(II) superoxide species are known to have H-atom transfer reactivity.^{35,36,50–52} It should be also noted that the reaction of ethylbenzene and mchloroperoxybenzoic acid catalyzed by complex 2 exclusively provided 1-phenylethanol (sp³-carbon oxygenation product) rather than ethylphenol, where we proposed that a Ni(II) oxyl radical-type species is involved as the key reactive intermediate.³¹ On the basis of these results, we propose that a dinickel(III) bis(μ -oxo) complex is the most plausible reactive intermediate in the present reaction, as illustrated in Scheme 3.

In summary, we have demonstrated the catalytic ability of nickel complexes in direct benzene hydroxylation with H_2O_2 for the first time. In particular, the nickel complex 2 supported by





tepa exhibited a prominent ability to produce phenol in 21% yield based on benzene without the formation of any overoxidation products. The total turnover number of the nickel complex supported by tepa reached 749, which is the highest value ever reported for molecular catalysts. The aromatic hydroxylation method has been shown to be applicable to oxygenation of alkylbenzene derivatives. The selectivity for cresol in the toluene hydroxylation reached 90%, which is also the highest reported value for direct hydroxylation of toluene to cresol with H_2O_2 and a molecular catalyst. Details of the reaction mechanism and the reason for the bizarre selectivity are now under investigation.⁵³

ASSOCIATED CONTENT

Supporting Information

Experimental details. The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/ jacs.5b01814.

AUTHOR INFORMATION

Corresponding Author

*shinobu@mls.eng.osaka-u.ac.jp

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENTS

This work was partly supported by a JSPS Fellowship for Young Scientists (Y.M.) and a Grant for Scientific Research on Innovative Areas (22105007 to S.I.) from MEXT of Japan. Y.M. and S.I. express their special thanks to the JSPS Japanese– German Graduate Externship Program on "Environmentally Benign Bio and Chemical Processes" for financial support of the stay at RWTH Aachen. The authors also express their gratitude to Prof. Takashi Hayashi and Dr. Koji Oohora at Osaka University for their help with ESI mass spectrometry.

REFERENCES

(1) Weber, M.; Weber, M.; Kleine-Boymann, M. In *Ullmann's Encyclopedia of Industrial Chemistry*; Wiley-VCH: Weinheim, Germany, 2000.

(2) Schmidt, R. J. Appl. Catal., A 2005, 280, 89-103.

(3) Mimoun, H.; Saussine, L.; Daire, E.; Postel, M.; Fischer, J.; Weiss, R. J. Am. Chem. Soc. **1983**, 105, 3101–3110.

(4) Kimura, E.; Machida, R. J. Chem. Soc., Chem. Commun. 1984, 499–500.

(5) Hata, E.; Takai, T.; Yamada, T.; Mukaiyama, T. Chem. Lett. 1994, 1849–1852.

(6) Bianchi, D.; Bortolo, R.; Tassinari, R.; Ricci, M.; Vignola, R. *Angew. Chem., Int. Ed.* **2000**, *39*, 4321–4323.

(7) Niwa, S.-i.; Eswaramoorthy, M.; Nair, J.; Raj, A.; Itoh, N.; Shoji, H.; Namba, T.; Mizukami, F. *Science* **2002**, *295*, 105–107.

(8) Tani, M.; Sakamoto, T.; Mita, S.; Sakaguchi, S.; Ishii, Y. Angew. Chem., Int. Ed. 2005, 44, 2586–2588.

Journal of the American Chemical Society

(9) Dong, T.; Li, J.; Huang, F.; Wang, L.; Tu, J.; Torimoto, Y.; Sadakata, M.; Li, Q. *Chem. Commun.* **2005**, 2724–2726.

(10) Tang, Y.; Zhang, J. Transition Met. Chem. 2006, 31, 299-305.

(11) Thibon, A.; Bartoli, J.-F.; Guillot, R.; Sainton, J.; Martinho, M.; Mansuy, D.; Banse, F. J. Mol. Catal. A: Chem. **2008**, 287, 115–120.

(12) Conde, A.; Diaz-Requejo, M. M.; Perez, P. J. Chem. Commun. 2011, 47, 8154-8156.

(13) Shoji, O.; Kunimatsu, T.; Kawakami, N.; Watanabe, Y. Angew. Chem., Int. Ed. 2013, 52, 6606–6610.

(14) Shul'pin, G. B.; Kozlov, Y. N.; Shul'pina, L. S.; Carvalho, W. A.; Mandelli, D. RSC Adv. 2013, 3, 15065–15074.

(15) Raba, A.; Cokoja, M.; Herrmann, W. A.; Kuhn, F. E. Chem. Commun. 2014, 50, 11454–11457.

(16) Sheldon, R. A.; Kochi, J. K. In *Metal-Catalyzed Oxidations of Organic Compounds*; Sheldon, R. A., Kochi, J. K., Eds.; Academic Press: New York, 1981; pp 315–339.

(17) Catalytic Oxidations with Hydrogen Peroxide as Oxidant; Strukul, G., Ed.; Springer: Dordrecht, The Netherlands, 1992.

(18) Haggin, J. Chem. Eng. News 1993, 71 (22), 23-27.

(19) Cruse, R. W.; Kaderli, S.; Karlin, K. D.; Zuberbuehler, A. D. J. Am. Chem. Soc. **1988**, 110, 6882–6883.

(20) Nasir, M. S.; Cohen, B. I.; Karlin, K. D. J. Am. Chem. Soc. 1992, 114, 2482-2494.

(21) Karlin, K. D.; Nasir, M. S.; Cohen, B. I.; Cruse, R. W.; Kaderli, S.; Zuberbuehler, A. D. *J. Am. Chem. Soc.* **1994**, *116*, 1324–1336.

(22) Holland, P. L.; Rodgers, K. R.; Tolman, W. B. Angew. Chem., Int. Ed. **1999**, 38, 1139–1142.

(23) Honda, K.; Cho, J.; Matsumoto, T.; Roh, J.; Furutachi, H.; Tosha, T.; Kubo, M.; Fujinami, S.; Ogura, T.; Kitagawa, T.; Suzuki, M. *Angew. Chem., Int. Ed.* **2009**, *48*, 3304–3307.

(24) Kunishita, A.; Doi, Y.; Kubo, M.; Ogura, T.; Sugimoto, H.; Itoh, S. *Inorg. Chem.* **2009**, *48*, 4997–5004.

(25) Tano, T.; Doi, Y.; Inosako, M.; Kunishita, A.; Kubo, M.; Ishimaru, H.; Ogura, T.; Sugimoto, H.; Itoh, S. *Bull. Chem. Soc. Jpn.* **2010**, *83*, 530–538.

(26) Que, J. L.; Tolman, W. B. Angew. Chem., Int. Ed. 2002, 41, 1114–1137.

(27) Mirica, L. M.; Ottenwaelder, X.; Stack, T. D. P. Chem. Rev. 2004, 104, 1013–1046.

(28) Lewis, E. A.; Tolman, W. B. Chem. Rev. 2004, 104, 1047–1076.
(29) Suzuki, M. Acc. Chem. Res. 2007, 40, 609–617.

(30) Itoh, S.; Bandoh, H.; Nagatomo, S.; Kitagawa, T.; Fukuzumi, S. J. Am. Chem. Soc. **1999**, 121, 8945–8946.

(31) Nagataki, T.; Ishii, K.; Tachi, Y.; Itoh, S. Dalton Trans. 2007, 1120-1128.

(32) The conversion of benzene was not identical to the yield because of evaporation of benzene (\sim 10% in 5 h). This was confirmed by a blank experiment.

(33) A large part of the H₂O₂ (44%) remained after the reaction.

(34) For catalyst 3, we observed ligand hydroxylation at the benzylic position of the phenethyl arm in about 10% yield by 1 H NMR analysis, whereas we did not detect such a ligand-hydroxylated product with catalyst 2. For details, see the SI.

(35) Cho, J.; Sarangi, R.; Annaraj, J.; Kim, S. Y.; Kubo, M.; Ogura, T.; Solomon, E. I.; Nam, W. *Nat. Chem.* **2009**, *1*, 568–572.

(36) Kieber-Emmons, M. T.; Annaraj, J.; Seo, M. S.; Van Heuvelen, K. M.; Tosha, T.; Kitagawa, T.; Brunold, T. C.; Nam, W.; Riordan, C. G. *J. Am. Chem. Soc.* **2006**, *128*, 14230–14231.

(37) The concentrations of benzene, H_2O_2 , and TEA were optimized (see Table S2).

(38) After the reaction, 4.3 mmol of $\rm H_2O_2$ (17% of the initial amount) remained.

(39) Luo, Y.-R. Comprehensive Handbook of Chemical Bond Energies; CRC Press: Boca Raton, FL, 2007.

(40) Kamata, K.; Yamaura, T.; Mizuno, N. Angew. Chem., Int. Ed. 2012, 51, 7275–7278.

(41) In aromatic oxygenation reactions, a shift of the deuteron on the attacked carbon to the neighboring carbon (a so-called NIH shift) is

often observed. In this system, however, the NIH shift was not observed in the oxygenation reaction of toluene-4-*d*₁. For details, see Scheme S4. (42) Augusti, R.; Dias, A. O.; Rocha, L. L.; Lago, R. M. *J. Phys. Chem. A* **1998**, *102*, 10723–10727.

(43) Emmert, M. H.; Cook, A. K.; Xie, Y. J.; Sanford, M. S. Angew. Chem., Int. Ed. 2011, 50, 9409–9412.

(44) Ohkubo, K.; Kobayashi, T.; Fukuzumi, S. Angew. Chem., Int. Ed. 2011, 50, 8652–8655.

(45) Unfortunately, we could not detect any spectral change upon addition of H_2O_2 to a solution of catalyst 2 at low temperature (below -60 °C), under which conditions catalyst 3, supported by the less hindered bepa ligand, generates a dinickel(III) bis(μ -oxo) species (see ref 30).

(46) Shiren, K.; Ogo, S.; Fujinami, S.; Hayashi, H.; Suzuki, M.; Uehara, A.; Watanabe, Y.; Moro-oka, Y. J. Am. Chem. Soc. **2000**, 122, 254–262.

(47) Cho, J.; Furutachi, H.; Fujinami, S.; Tosha, T.; Ohtsu, H.; Ikeda, O.; Suzuki, A.; Nomura, M.; Uruga, T.; Tanida, H.; Kawai, T.; Tanaka,

K.; Kitagawa, T.; Suzuki, M. *Inorg. Chem.* **2006**, *45*, 2873–2885. (48) Hikichi, S.; Yoshizawa, M.; Sasakura, Y.; Komatsuzaki, H.; Moro-

oka, Y.; Akita, M. *Chem.—Eur. J.* **2001**, *7*, 5011–5028.

(49) Itoh, S.; Bandoh, H.; Nakagawa, M.; Nagatomo, S.; Kitagawa, T.;
Karlin, K. D.; Fukuzumi, S. J. Am. Chem. Soc. 2001, 123, 11168–11178.
(50) Cho, J.; Sarangi, R.; Nam, W. Acc. Chem. Res. 2012, 45, 1321–1330.

(51) 14-tmc has also been reported to be able to stabilize dinickel(II) peroxide species generated by the reaction between the corresponding Ni(I) complex and molecular oxygen (see ref 52).

(52) Kieber-Emmons, M. T.; Schenker, R.; Yap, G. P. A.; Brunold, T. C.; Riordan, C. G. Angew. Chem., Int. Ed. **2004**, 43, 6716–6718.

(53) The oxygenation efficiency of benzene derivatives with catalyst **2** was found to decrease upon the introduction of electron-donating substituents. This unusual selectivity of benzene oxygenation over phenol oxygenation may be due to this tendency.