

# On reduction of $\alpha,\beta$ -unsaturated ketones and the respective allylic alcohols, bearing a phenylsulfonyl or phenylsulfanyl group in the $\alpha$ position. Hydroxy group-controlled stereoselective reduction of $3\alpha$ - and $3\beta$ -hydroxy-4-(phenylsulfonyl)cholest-4-ene

Karol Michalak, Waczesław Stepanenko and Jerzy Wicha \*

Institute of Organic Chemistry, Polish Academy of Sciences, POB 58, 01-224 Warsaw, Poland

Received (in Cambridge, UK) 21st January 2000, Accepted 17th March 2000

Published on the Web 19th April 2000

Reduction reactions of cholest-4-en-3-one derivatives bearing the phenylsulfonyl or phenylsulfanyl group at C-4 with various metal hydrides are studied. Lithium aluminumhydride reduction of 4-(phenylsulfonyl)cholest-4-en-3 $\beta$ -ol **13a** and 4-(phenylsulfonyl)cholest-4-en-3 $\alpha$ -ol **15a** occurs with saturation of the double bond and deoxygenation to give 4 $\beta$ -phenylsulfonyl-5 $\beta$ -cholestane **8** and 4 $\alpha$ -phenylsulfonyl-5 $\alpha$ -cholestane **7a**, respectively. Reduction of 4-(phenylsulfonyl)cholest-4-en-3-one **2** with lithium aluminumhydride yields compound **8**. Reduction of compounds **2**, **13a** and **15a** with other metal hydrides affords mixtures of diastereomeric products. Metal hydride reductions of 4-(phenylsulfanyl)cholest-4-en-3-one **1** affect the carbonyl group only. Catalytic hydrogenation of compound **2** gives a mixture of 5 $\alpha$ - and 5 $\beta$ - dihydro derivatives. Mechanistic and stereochemical aspects of the reduction reactions are discussed.


## Introduction

Metal hydride reduction reactions of  $\alpha,\beta$ -unsaturated ketones,<sup>1,2</sup>  $\alpha$ -phenylsulfonyl ketones<sup>3</sup> and vinyl sulfones<sup>4-6</sup> have been extensively studied. However, there are no reports on the reduction of  $\alpha,\beta$ -unsaturated ketones bearing in the  $\alpha$ -position a phenylsulfonyl or a phenylsulfanyl group. In such derivatives relative rates of reduction of the carbonyl group and the double bond may differ from those in the parent compound, and, consequently, the steric outcome of the reaction may be different. Some consecutive reactions due to the presence of the sulfur-containing group may also occur. Recently, we have shown<sup>7,8</sup> that the complex hydrindane derivative **i** with oxo and vinyl sulfone functions, in reaction with lithium aluminumhydride was transformed stereoselectively into product **iii** with *trans*-fused hydrindane rings. We have also noted that the steric course of the carbon-carbon double-bond reduction doesn't depend upon the orientation of the hydroxy group ( $\alpha$  or  $\beta$ ) in the intermediate **ii**. It was thought of interest to examine in some detail the reduction of easily available  $\alpha,\beta$ -unsaturated ketones **1** and **2**, and the corresponding 3-hydroxy derivatives that are likely immediate products of the carbonyl compounds' reduction. It



was expected that we would gain insight into the multistep reduction mechanism and, eventually, develop stereoselective approaches to *trans*- and *cis*-decalin derivatives.

## Results and discussion

Sulfide **1** was prepared by treatment of 4 $\beta$ ,5-epoxy-5 $\beta$ -cholestan-4-one<sup>9</sup> with thiophenol and potassium hydroxide in ethanol.<sup>10</sup> Sulfone **2** was obtained by oxidation of sulfide **1** with MCPBA. In order to prepare some reference compounds, the known<sup>11,12</sup> 3 $\alpha$ ,4 $\alpha$ -epoxy-5 $\alpha$ -cholestane **3** (Scheme 1) was



R = oxygen substituent or modified steroid side chain




**Scheme 1** Reagents and yields: (a) PhSH, EtOH, 83% yield; (b) MsCl, Et<sub>3</sub>N, CH<sub>2</sub>Cl<sub>2</sub>; (c) MCPBA, 100%; (d) DBU, PhH, 95% in 2 steps; (e) Et<sub>3</sub>BHLi, 99%; (f) *tert*-BuOK-*tert*-BuOH, 100%.

allowed to react with sodium thiophenolate in ethanol to give diaxial hydroxy sulfide **4**, by analogy to the procedure developed for related hydrindane derivatives.<sup>13</sup> Compound **4** was treated consecutively with mesyl chloride, MCPBA, and DBU to afford vinylic sulfone **6**. Reduction of **6** with Et<sub>3</sub>BHLi and chromatographic separation of the products afforded

4 $\alpha$ - and 4 $\beta$ -phenylsulfonyl derivatives **7a** and **7b** in 79 and 20% yield, respectively. Alternatively, oxidation of hydroxy sulfide **4** with MCPBA provided sulfone **5**, which on treatment with mesyl chloride and then with Et<sub>3</sub>BHLi gave a mixture of **7a** and **7b**. Compound **7b** with an axially oriented phenylsulfonyl group was transformed into its epimer **7a** quantitatively with potassium *tert*-butoxide in *tert*-butyl alcohol.<sup>14</sup>

Reduction of sulfone **2** with an excess of LiAlH<sub>4</sub> in THF at room temperature afforded saturated sulfone **8** with *cis* configuration at the ring junction, isolated in 78% yield. The structure of compound **8** was confirmed by its reduction with sodium amalgam, which afforded 5 $\beta$ -cholestane.

Reduction of compound **2** with NaBH<sub>4</sub> in methanol afforded a mixture of hydroxy sulfones **9** (5 $\alpha$ -H) and **10** (5 $\beta$ -H) (83% yield). The mixture could not be separated by either chromatography or crystallization; however, a sample of pure compound **9** was isolated by precipitation. Oxidation of the crude mixture of **9** and **10** with Jones reagent followed by chromatography afforded oxo sulfones **11** and **12** in 58% and 41% yield, respectively. Configuration at C-5 in compounds **11** and **12** was established by their reduction to 5 $\alpha$ - and 5 $\beta$ -cholestan-3-one, respectively.



The configuration around C-3 and C-4 in compound **9** follows from the results of nuclear Overhauser effects (NOE) experiments involving C-19, C-4 and C-3 protons (see Experimental section). Configuration around C-3 and C-4 in **10** was deduced from the <sup>1</sup>H NMR spectrum of the mixture of **9** and **10** in which the signals of the C-3 proton ( $\delta$  3.91, broad singlet) and the C-4 proton ( $\delta$  3.15, doublet of doublets, *J* = 12.1 and 2.0 Hz) in **10** could be seen. It was assumed that the value *J* = 2 Hz corresponds to C-3–C-4 proton coupling (*cis*) and that the value *J* = 12.1 Hz reflects coupling of C-4 and C-5 protons in a *trans* configuration (4 $\alpha$ ,5 $\beta$ ). The assignment of structures of **9** and **10** was corroborated by the <sup>1</sup>H NMR spectrum of previously prepared hydroxy sulfone epimer **5** which differs from that of **9** in the orientation of the phenylsulfonyl group. It is noteworthy that in both alcohols **9** and **10** the hydroxy group occupies an axial position whereas the sulfonyl group is in an equatorial position.

Next, reduction of **2** with NaBH<sub>4</sub> in THF in the presence of DMPU as a polar co-solvent was examined. Treatment of **2** with NaBH<sub>4</sub> in THF–DMPU (a suspension) at room temperature afforded a mixture of hydroxy sulfones **9** and **10** in 74% yield in a ratio of  $\approx$ 5.5:1, and oxo sulfone **12** in 23% yield. Oxo sulfone **12** resisted further reduction under the reaction conditions. It is noteworthy that isomeric oxo sulfone **11** was

smoothly reduced under analogous conditions to give alcohol **9**. This experiment shows that the reduction reaction occurs along two mechanistic paths: (1) with reduction of the carbonyl group preceding the reduction of the carbon–carbon double bond and (2) with reduction of the carbon–carbon double bond occurring first, followed by reduction of the carbonyl group (or providing a saturated ketone **12**). In practical terms, selectivity with regard to the ring-junction configuration was poor, providing, in total, 5 $\alpha$  and 5 $\beta$  isomers in the ratio 65:35.

Reduction of **2** with a system, CuCN–*n*-BuLi–DIBALH, that is known to reduce steroidal and related 4-en-3-ones selectively to the corresponding 5 $\alpha$ -H saturated ketones,<sup>15,16</sup> afforded a mixture of ketones **11** (39%) and **12** (22%) and unsaturated alcohol **13a** (10%). Reduction of **2** with DIBALH in methylene dichloride affected the carbonyl group only and yielded quantitatively alcohol **13a**.

Allylic alcohol **13a** was thought to be a likely intermediate in reduction of phenylsulfonyl enone **2** to sulfone **8** by means of LiAlH<sub>4</sub>. Since a directive effect of the hydroxy group appeared to be possible it was of interest to compare the steric courses of reduction of epimeric alcohols **13a** and **15a**. Accordingly, alcohol **13a** was subjected to the Mitsunobu inversion which afforded 3 $\alpha$ -hydroxy sulfone **15a** in 64% overall yield, *via* benzoate **15b**. In parallel experiments **15a** was prepared from 3 $\beta$ -hydroxy sulfide **14** (see below) *via* **16b** and **16a** in a somewhat better overall yield (68%).

Reduction of 3 $\beta$ -hydroxy sulfone **13a** with LiAlH<sub>4</sub> afforded 5 $\beta$ -H sulfone **8** in 78% yield. Reduction of 3 $\alpha$ -hydroxy sulfone **15a** with LiAlH<sub>4</sub> afforded 5 $\alpha$ -H sulfone **7a** in 82% yield.

In light of the clear directing effect exerted by the hydroxy group in the reduction of hydroxy sulfone **13a**, it was of interest to examine the effect of an alkoxy group. To this end, the methoxymethyl derivative **13b**, prepared from **13a** in the usual way, was treated with LiAlH<sub>4</sub>. Three products, **8**, **7a** and **7b**, were obtained; a ratio of 5 $\alpha$ - to 5 $\beta$ -H products was estimated as  $\approx$ 1:1. The above results for the reduction reactions of vinylic sulfones **2**, **13a** and **15a** are compiled in Table 1.

Reduction of sulfide **1** with various reducing agents afforded mixtures of 3-hydroxy-4-(phenylsulfonyl)cholest-4-enes **14** and **16a**. Some results are shown in Table 2. No carbon–carbon double-bond reduction was observed. Only Luche reduction<sup>17</sup> (NaBH<sub>4</sub>–CeCl<sub>3</sub>) was virtually selective with respect to 3 $\beta$ -hydroxy derivative **14**.

Catalytic hydrogenation of sulfone **2** in the presence of 10% palladium on carbon was examined for completeness of our study. Compound **2** was practically unchanged under hydrogenation conditions in EtOH at room temperature. However, at reflux temperature the reaction was complete in *ca.* 5 h to yield *trans* and *cis* products, **11** and **12** in the ratio 6.5:1. Hydrogenation of **2** in EtOH at room temperature in the presence of CF<sub>3</sub>COOH afforded **11** and **12** in the ratio 5.5:1. Our attempts to achieve stereoselective reduction of the carbon–carbon double bond in **2** by varying the solvent in the catalytic hydrogenation reaction failed.

The above described experiments allow for some mechanistic comments. It is likely that reduction of **2** to 5 $\beta$ -H saturated sulfone **8** commences with hydride ion addition to the carbonyl group from the  $\alpha$ -side to form intermediate **iv**, Scheme 2. Further reduction occurs with intramolecular hydride ion delivery and elimination of the metal-bonded oxygen atom. The intermediate vinylic sulfone **v** with *cis* ring junction undergoes further reduction to **8**. This mechanism is corroborated by stereoselective reduction of the allylic alcohol with  $\beta$ -oriented hydroxy group, **13a** (Table 1, entry 2). Intramolecular hydride anion delivery presumably occurs in a similar fashion in reduction of 3 $\alpha$ -hydroxy derivative **15a** (*via* intermediate **vi**) to yield 5 $\alpha$ -H sulfone **7a** (Table 1, entry 3). The importance of the hydroxy group for stereoselective introduction of the hydrogen at C-5 is also shown by the reduction of the MOM ether **13b** in which a mixture of 5 $\alpha$ - and 5 $\beta$ -H products was formed


**Table 1** Reduction of oxo sulfone **2** and hydroxy sulfones **13a** and **15a** with metal hydrides

Entry	Compound	Reducing system	Product (yield %), ratio	5 $\alpha$ :5 $\beta$
1	<b>2</b>	LiAlH <sub>4</sub> -THF	<b>8</b> (78)	5 $\beta$ -H only
2	<b>13a</b>	LiAlH <sub>4</sub> -THF	<b>8</b> (78)	5 $\beta$ -H only
3	<b>15a</b>	LiAlH <sub>4</sub> -THF	<b>7a</b> (82)	5 $\alpha$ -H only
4	<b>13b</b>	LiAlH <sub>4</sub> -THF	<b>8</b> , <b>7a</b> , <b>7b</b>	1:1
5	<b>2</b>	NaBH <sub>4</sub> -MeOH	<b>9</b> , <b>10</b> (83)	3:2
6	<b>2</b>	NaBH <sub>4</sub> -THF-DMPU	<b>9</b> , <b>10</b> (74) and <b>12</b> (23)	2:1
7	<b>2</b>	CuCN- <i>n</i> -BuLi-DIBAH	<b>11</b> (39), <b>12</b> (22), <b>13a</b> (10)	2:1
8	<b>2</b>	DIBAH-CH <sub>2</sub> Cl <sub>2</sub>	<b>13a</b> (100)	

**Table 2** Reduction of oxo sulfide **1** with metal hydrides

Entry	Reducing agent	Solvent	Temp.	Product (yield %, ratio)
1	DIBAH	CH <sub>2</sub> Cl <sub>2</sub> -hexane	-78 °C	<b>14</b> (69) and <b>16a</b> (21)
2	L-Selectride	THF	-78 °C	<b>14</b> and <b>16a</b> (79, 82:18) <sup>a</sup>
3	NaBH <sub>4</sub> -CeCl <sub>3</sub>	THF-MeOH	-78 °C to rt	<b>14</b> (91) <sup>b</sup>
4	LiAlH <sub>4</sub>	THF	rt	<b>14</b> and <b>16a</b> (91, 84:16)

<sup>a</sup> Ratio of products was determined by <sup>1</sup>H NMR spectroscopy. <sup>b</sup> Some **15a**, less than 5%, could be detected by <sup>1</sup>H NMR analysis.

**Scheme 2**

(Table 1, entry 4). The hydroxy group-directing effect has not been observed in the reduction of a related hydrindane system.<sup>7</sup> This difference suggests that in the hydrindane system reduction of the vinyl sulfone unit occurs by intermolecular hydride anion delivery. It is noteworthy that LiAlH<sub>4</sub> reduction of the carbon-carbon double bond at the ring junction in a hydrindane-related vinylic sulfone lacking an oxygen function afforded the *trans*-hydrindane derivative only.<sup>6</sup> NaBH<sub>4</sub>-methanol reduction of **2** affording a mixture of **9** and **10** shows little selectivity with respect to the configuration at the ring junction (Table 1, entry 5).

In conclusion, it was shown that readily available sulfonyl ketone **2** may be stereoselectively transformed into 5 $\alpha$ - or 5 $\beta$ -cholestane derivatives, **7** or **8**. Reduction of the sulfur-containing  $\alpha,\beta$ -unsaturated ketones **1** and **2**, and of the respective alcohols, has been scrutinized. Some mechanistic

suggestions regarding the multistep reduction have been made.

## Experimental

Mps were determined on a Kofler hot-stage melting point apparatus and are uncorrected. <sup>1</sup>H and <sup>13</sup>C NMR spectra were performed for samples in CDCl<sub>3</sub> on a Varian Gemini 200 MHz spectrometer using residual CHCl<sub>3</sub> as an internal standard ( $\delta$  7.26 and 77.0, respectively). Signal multiplicities in <sup>13</sup>C spectra were assigned using DEPT sequence and are given in brackets. *J*-Values are given in Hz. Mass spectra were taken at an ionizing voltage of 70 eV; peak (*m/z*) relative intensities are given in parentheses. Air-sensitive reactions were performed in oven- or flame-dried glassware under argon. Bulk solution of LiAlH<sub>4</sub> (1 M) was prepared using freshly purchased reagent (Aldrich) and Na/K alloy-dried THF. Organic extracts were dried over Na<sub>2</sub>SO<sub>4</sub> and the solvents were evaporated on a rotary evaporator.

### 4-(Phenylsulfonyl)cholest-4-en-3-one **1**

This compound was prepared from 4 $\beta$ ,5-epoxy-5 $\beta$ -cholestan-3-one<sup>9</sup> following the described procedure;<sup>10</sup> mp 105 °C (Found: C, 80.62; H, 9.70. Calc. for C<sub>33</sub>H<sub>48</sub>OS: C, 80.43; H, 9.82%);  $\lambda_{\text{max}}$ (EtOH)/nm 321.3, 250.4, 202.5;  $\nu_{\text{max}}$ (KBr)/cm<sup>-1</sup> 1679, 1581, 1557;  $\delta_{\text{H}}$  7.24–7.02 (5 H, m, ArH), 3.62 (1 H, dtd, *J* 14.5, 3.0, 0.5), 2.54 (2 H, dd, *J* 9.9, 4.5), 2.23 (1 H, td, *J* 14.2, 5.3), 1.29 (3 H, s, 19-H<sub>3</sub>), 0.91 (3 H, d, *J*<sub>20,21</sub> 6.5, 21-H<sub>3</sub>), 0.86 (6 H, d, *J*<sub>25,26</sub> and 25,27 6.6, 26- and 27-H<sub>3</sub>), 0.71 (3 H, s, 18-H<sub>3</sub>);  $\delta_{\text{C}}$  194.1, 177.9, 137.23, 128.7, 126.9, 125.1, 56.0, 55.7, 54.2, 42.3, 41.6, 39.5, 39.4, 36.0, 35.6, 35.2, 34.6, 34.4, 32.0, 31.0, 28.1, 27.9, 24.0, 23.7, 22.7, 22.5, 21.1, 18.6, 18.3, 11.9. Physical constants were in agreement with those described, with the exception of <sup>1</sup>H NMR signal at  $\delta$  3.62 (1 H) which has been not recorded in the original characterization.

### 4-(Phenylsulfonyl)cholest-4-en-3-one **2**

To a stirred solution of sulfide **1** (604 mg, 1.23 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (10 cm<sup>3</sup>) was added MCPBA (60%; 734 mg, 2.55 mmol) in portions. After 2 h, the mixture was partitioned between aq. Na<sub>2</sub>SO<sub>3</sub> and CH<sub>2</sub>Cl<sub>2</sub>. The organic solution was washed with water and the solvent was evaporated. The residue was crystallized from acetone-hexane to give *sulfone 2* (528 mg, 82%), mp 146–148 °C;  $\lambda_{\text{max}}$ (EtOH)/nm 253.3, 221.7, 199.6;  $\nu_{\text{max}}$ (KBr)/cm<sup>-1</sup> 1688;  $\delta_{\text{H}}$  8.02–7.92 (2 H, m, ArH), 7.58–7.42 (3 H, m, ArH), 4.32 (1 H, dt, *J* 14.1, 2.8), 2.38–0.82 (26 H, m) over-

lapping with 1.25 (3 H, s, 19-H<sub>3</sub>), 0.89 (3 H, d,  $J_{20,21}$  7.0, 21-H<sub>3</sub>), 0.85 (6 H, d,  $J_{25,26}$  and  $J_{25,27}$  7.0, 26- and 27-H<sub>3</sub>), 0.70 (3 H, s, 18-H<sub>3</sub>);  $\delta_C$  192.0 (C-3), 181.2 (C-4 or -5), 142.6 (C-5 or -4), 136.1 (C-*ipso*), 132.7 (C-*o*), 128.4 (C-*m*), 127.7 (C-*p*), 55.9, 55.5, 54.7, 42.8, 42.3, 39.4, 39.3, 36.0, 35.6, 35.1, 34.1, 33.4, 32.6, 28.1, 27.9, 27.6, 23.9, 23.7, 22.7, 22.5, 21.3, 18.5, 18.4, 11.9 (C-18);  $m/z$  (EI) 524.332 34 ( $M^+$ , 22%, C<sub>33</sub>H<sub>48</sub>O<sub>3</sub>S requires  $M$ , 524.332 41), 509 [( $M$  - CH<sub>3</sub>)<sup>+</sup>, 2], 460 [( $M$  - SO<sub>2</sub>)<sup>+</sup>, 100], 445 (13), 383 [( $M$  - SO<sub>2</sub>Ph)<sup>+</sup>, 83].

#### 4β-Phenylsulfanyl-5α-cholestan-3α-ol 4

To anhydrous EtOH (30 cm<sup>3</sup>) was added sodium (214 mg, 9.3 mg-atom). After the reaction was complete, thiophenol (0.94 cm<sup>3</sup>, 9.15 mmol) was added, followed by 3α,4α-epoxy-5α-cholestan-3<sup>11,12</sup> (1.80 g, 4.6 mmol). The mixture was heated under reflux for 24 h, cooled and partitioned between water and benzene. The organic extract was washed with water and evaporated. The residue was chromatographed on SiO<sub>2</sub> (10 g; hexane–EtOAc, 9:1) to give **compound 4** (1.93 g, 83%), mp 133 °C (from EtOH);  $\delta_H$  7.41–7.12 (5 H, m, ArH), 4.02 (1 H, br d,  $J_{3,4}$  2.2, 3-H<sub>3</sub>), 3.17 (1 H, br dd,  $J_{3,4}$  2.2,  $J_{4,5}$  2.2, 4-H<sub>3</sub>), 0.97 (3 H, s, 19-H<sub>3</sub>), 0.91 (3 H, d,  $J_{20,21}$  7.1, 21-H<sub>3</sub>) overlapping with 0.87 (6 H, d,  $J_{25,26}$  and  $J_{25,27}$  6.7, 26- and 27-H<sub>3</sub>), 0.66 (3 H, s, 18-H<sub>3</sub>);  $\delta_C$  137.9 (C-*i*), 130.5 (C-*o*), 120.9 (C-*m*), 126.3 (C-*p*), 70.2 (C-3), 56.7, 56.5, 56.2, 55.1, 42.8, 42.5, 39.9, 39.5, 36.5, 36.2, 35.8, 35.3, 32.3, 31.6, 28.2, 28.0, 27.5, 24.3, 24.2, 23.9, 22.8, 22.6, 20.5, 18.7, 13.7 (C-19), 12.0 (C-18);  $m/z$  (EI) 496.373 12 ( $M^+$ , 84%, C<sub>33</sub>H<sub>52</sub>OS requires  $M$ , 496.373 89), 386 (14), 369 (100), 353 (5), 287 (8), 273 (8), 257 (7), 243 (10), 229 (7), 215 (11), 189 (7), 175 (10), 161 (24), 147 (12), 135 (18), 121 (10), 107 (15).

#### 4β-Phenylsulfanyl-5α-cholestan-3α-ol 5

To a solution of sulfide **4** (598 mg, 1.2 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (20 cm<sup>3</sup>) was added MCPBA (60%; 710 mg, 2.47 mmol). The mixture was stirred for 2 h and then partitioned between aq. Na<sub>2</sub>SO<sub>3</sub> and CH<sub>2</sub>Cl<sub>2</sub>. The product was isolated in the usual way to give **sulfone 5** (627 mg, 98%),  $\delta_H$  7.91–7.82 (2 H, m, ArH), 7.65–7.45 (3 H, m, ArH), 4.13 (1 H, br s, 3-H), 3.26 (1 H, d,  $J$  5.3, 4-H), 1.19 (3 H, s, 19-H<sub>3</sub>), 0.90 (3 H, d,  $J_{20,21}$  7.3, 21-H<sub>3</sub>) overlapping with 0.86 (6 H, d,  $J_{25,26}$  and  $J_{25,27}$  6.7, 26- and 27-H<sub>3</sub>), 0.66 (3 H, s, 18-H<sub>3</sub>);  $\delta_C$  141.6 (C-*i*), 133.2 (C-*o*), 129.1 (C-*m*), 127.9 (C-*p*), 70.6 (C-3), 65.2 (C-4), 56.4, 56.2, 55.6, 45.2, 42.4, 39.9, 39.5, 36.2, 35.8, 35.4, 33.3, 31.9, 28.2, 28.0, 27.3, 25.7, 24.2, 23.8, 22.8, 22.6, 20.9, 18.7, 13.4 (C-19), 12.0 (C-18);  $m/z$  (EI) 528.363 67 ( $M^+$ , 5%, C<sub>33</sub>H<sub>52</sub>O<sub>3</sub>S requires  $M$ , 528.363 71), 510 [( $M$  - H<sub>2</sub>O), 8], 495 (4), 445 (3), 387 (23), 369 (100), 355 (17), 341 (3), 313 (2), 287 (6), 257 (6), 247 (10), 229 (23), 215 (33), 201 (8), 175 (10), 161 (20), 147 (21), 135 (27), 121 (23), 107 (28), 95 (38), 81 (47).

#### 4-Phenylsulfanyl-5α-cholest-3-ene 6

To a solution of hydroxy sulfide **5** (306 mg, 0.62 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (10 cm<sup>3</sup>) was added Et<sub>3</sub>N (0.26 cm<sup>3</sup>, 1.9 mmol) followed by MsCl (0.09 cm<sup>3</sup>, 1.2 mmol). The mixture was set aside for 4 h and then partitioned between hexane and water. The hexane layer was washed consecutively with 3% HCl and water. The solvent was evaporated to give 3ξ-chloro-4β-phenylsulfanyl-5α-cholestan-3 (316 mg, 100%),  $\delta_H$  7.38–7.20 (5 H, m, ArH), 4.45 (1 H, br d,  $J_{3,4}$  2.2, 3-H<sub>3</sub>), 3.43 (1 H, br dd,  $J_{3,4}$  2.2,  $J_{4,5}$  2.2, 4-H<sub>3</sub>), 2.59–2.38 (1 H, m), 2.29–2.16 (1 H, m), 1.00 (3 H, s, 19-H<sub>3</sub>), 0.92 (3 H, d,  $J_{20,21}$  7.5, 21-H<sub>3</sub>), 0.89 (6 H, d,  $J_{25,26}$  and  $J_{25,27}$  6.8, 26- and 27-H<sub>3</sub>), 0.67 (3 H, s, 18-H<sub>3</sub>);  $\delta_C$  136.9 (C-*i*), 130.3 (C-*o*), 129.1 (C-*m*), 126.7 (C-*p*), 62.4 (C-3), 57.1, 56.5, 56.2, 54.8, 42.5, 42.3, 39.8, 39.5, 36.6, 36.2, 35.8, 35.3, 32.1, 31.8, 28.2, 28.0, 27.4, 25.0, 24.2, 23.9, 22.8, 22.6, 20.5, 18.7, 14.3 (C-19), 12.0 (C-18).

To a solution of the crude chloride (226 mg, 0.44 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (10 cm<sup>3</sup>) was added MCPBA (60%; 256 mg, 0.89

mmol). After 8 h the mixture was partitioned between benzene and aq. Na<sub>2</sub>SO<sub>3</sub>. The organic layer was washed successively with aq. Na<sub>2</sub>SO<sub>3</sub> and water. Evaporation of the solvent gave 3ξ-chloro-4β-phenylsulfanyl-5α-cholestan-3 (239 mg, 100%),  $\delta_H$  7.94–7.86 (2 H, m, ArH), 7.69–7.51 (3 H, m, ArH), 4.53 (1 H, br s, 3-H), 3.42 (1 H, br d,  $J_{4,5}$  4.3, 4-H), 2.74–2.56 (1 H, m), 2.41–2.23 (2 H, m), 1.21 (3 H, s, 19-H<sub>3</sub>), 0.90 (3 H, d,  $J_{20,21}$  8.1, 21-H<sub>3</sub>), overlapping with 0.86 (6 H, d,  $J_{25,26}$  and  $J_{25,27}$  6.6, 26- and 27-H<sub>3</sub>), 0.67 (3 H, s, 18-H<sub>3</sub>);  $\delta_C$  141.1 (C-*i*)(0), 133.6 (C-*o*)(1), 129.3 (C-*m*)(1), 127.7 (C-*p*)(1), 71.4 (C-3)(1), 56.3 (1), 56.2 (1), 55.3 (1), 44.7 (1), 42.4 (0), 39.8 (2), 39.5 (2), 36.1 (2), 35.9 (1), 35.7 (1), 35.4 (1), 33.2 (2), 32.1 (2), 28.2 (2), 28.0 (1), 27.1 (2), 26.6 (2), 24.1 (2), 23.8 (2), 22.8 (C-26 or -27)(3), 22.5 (C-27 or -26)(3), 20.9 (2), 18.6 (C-21)(3), 13.6 (C-19)(3), 12.0 (C-18)(3).

A solution of the later product (239 mg, 0.44 mmol) in benzene (10 cm<sup>3</sup>) containing DBU (0.07 cm<sup>3</sup>, 0.46 mmol) was heated under reflux for 30 min and then cooled and partitioned between benzene and 3% HCl. The organic layer was washed with water and the solvent was evaporated to give **vinyl sulfone 6** (220 mg, 99%),  $\delta_H$  7.83–7.76 (2 H, m, ArH), 7.60–7.44 (3 H, m, ArH), 7.14 (1 H, br d,  $J_{2,3}$  3.1, 3-H<sub>3</sub>), 0.87 (3 H, d,  $J_{20,21}$  5.0, 21-H<sub>3</sub>) overlapping with 0.84 (6 H, d,  $J_{25,26}$  and  $J_{25,27}$  6.3, 26- and 27-H<sub>3</sub>), 0.70 (3 H, s, 19-H<sub>3</sub>), 0.59 (3 H, s, 18-H<sub>3</sub>);  $\delta_C$  141.7 (C-4 or -*i*)(0), 141.1 (C-*i* or -4)(0), 140.3 (C-3)(1), 132.6 (C-*o*)(1), 128.8 (C-*m*)(1), 127.1 (C-*p*)(1), 56.1 (1), 56.1 (1), 52.8 (1), 45.8 (1), 42.4 (0), 39.9 (2), 39.4 (2), 36.1 (2), 35.7 (1), 34.8 (1), 32.8 (2), 31.4 (2), 28.2 (2), 27.9 (1), 23.9 (2), 23.8 (2), 23.6 (2), 23.4 (2), 22.8 (C-26 or -27)(3), 22.5 (C-27 or -26)(3), 21.4 (2), 18.6 (C-21)(3), 12.4 (C-19)(3), 12.0 (C-18)(3);  $m/z$  (EI) 510.353 64 ( $M^+$ , 63%, C<sub>33</sub>H<sub>50</sub>O<sub>2</sub>S requires  $M$ , 510.353 15), 495 (23), 462 (3), 445 (12), 397 (3), 369 (42), 355 (100), 341 (20), 287 (10), 275 (22), 259 (7), 235 (10), 229 (17), 213 (22), 199 (5), 171 (6), 161 (8), 147 (13), 135 (23), 121 (35), 107 (28), 95 (35), 81 (65).

#### 4α-Phenylsulfanyl-5α-cholestan-7α and 4β-phenylsulfanyl-5α-cholestan-7β

(a) To a stirred solution of vinyl sulfone **6** (100 mg, 0.2 mmol) in THF (5 cm<sup>3</sup>) was added a solution of LiEt<sub>3</sub>BH (0.5 M in THF; 0.25 cm<sup>3</sup>, 0.125 mmol). The mixture was set aside for 2 h and then partitioned between CH<sub>2</sub>Cl<sub>2</sub> and water. The organic solution was evaporated and the residue was chromatographed on SiO<sub>2</sub> (6 g; hexane–AcOEt, 9:1) to give consecutively **7b** (20 mg, 20%) and then **7a** (79 mg, 79%).

(b) To a solution of hydroxy sulfone **5** (108 mg, 0.2 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (5 cm<sup>3</sup>), stirred under argon at –20 °C, was added Et<sub>3</sub>N (0.09 cm<sup>3</sup>, 0.65 mmol) followed by MsCl (0.03 cm<sup>3</sup>, 0.39 mmol). After 1 h, water (5 cm<sup>3</sup>) was added. The mixture was allowed to warm to room temperature and then was extracted with CH<sub>2</sub>Cl<sub>2</sub>. The organic extract was dried and evaporated to give the crude mesyl ester. The later product was dissolved in THF (5 cm<sup>3</sup>) and treated with LiEtBH<sub>3</sub> (0.5 M in THF; 0.7 cm<sup>3</sup>, 0.35 mmol). The mixture was stirred at room temperature for 0.5 h, and then was poured into water and extracted with CH<sub>2</sub>Cl<sub>2</sub>. The organic extract was dried and concentrated. The residue was chromatographed on SiO<sub>2</sub> (10 g; hexane–ethyl acetate, 9:1) to give **7b** (19 mg, 18%), **7a** (80 mg, 76%) and a fraction containing both these components (5 mg, 5%).

Compound **7b** had  $\delta_H$  7.96–7.87 (2 H, m, ArH), 7.63–7.48 (3 H, m, ArH), 3.21 (1 H, br t,  $J$  5.2, 4-H), 1.23 (3 H, s, 19-H<sub>3</sub>), 0.89 (3 H, d,  $J_{20,21}$  7.0, 21-H<sub>3</sub>) overlapping with 0.86 (6 H, d,  $J_{25,26}$  and  $J_{25,27}$  7.0, 26- and 27-H<sub>3</sub>), 0.66 (3 H, s, 18-H<sub>3</sub>) [lit.<sup>14</sup>  $\delta_H$  3.22 (4-H), 1.24 (19-H<sub>3</sub>), 0.68 (18-H<sub>3</sub>)].

Compound **7a** had  $\delta_H$  7.87–7.78 (2 H, m, ArH), 7.66–7.47 (3 H, m, ArH), 3.04 (1 H, br t,  $J$  10.7, 4-H), 2.43 (1 H, dd,  $J$  13.6, 2.4), 0.89 (3 H, d,  $J_{20,21}$  5.2, 21-H<sub>3</sub>) overlapping with 0.85 (6 H, d,  $J_{25,26}$  and  $J_{25,27}$  7.1, 26- and 27-H<sub>3</sub>) overlapping with 0.83 (3 H, s, 19-H<sub>3</sub>), 0.63 (3 H, s, 18-H<sub>3</sub>);  $\delta_C$  139.1 (C-*i*)(0), 133.1 (C-*o*)(1), 128.9 (C-*m*)(1), 128.5 (C-*p*)(1), 64.5 (C-4)(1), 56.3 (1), 56.1 (1), 54.4 (1), 46.7 (1), 42.3 (0), 39.9 (2), 39.5 (2), 37.5 (0),



37.4 (2), 36.1 (2), 35.7 (1), 34.6 (1), 31.4 (2), 28.8 (2), 28.2 (2), 28.0 (1), 25.4 (2), 24.0 (2), 23.8 (2), 22.8 (C-26 or -27)(3), 22.5 (C-27 or -26)(3), 21.1 (2), 20.6 (2), 18.6 (C-21)(3), 13.3 (C-19)(3), 12.0 (C-18)(3) [lit.,<sup>14</sup>  $\delta_{\text{H}}$  2.96 (4-H), 1.27 (19-H<sub>3</sub>), 0.65 (18-H<sub>3</sub>)].

#### Reduction of **2** with LiAlH<sub>4</sub>. 4 $\beta$ -Phenylsulfonyl-5 $\beta$ -cholestane **8**

To a stirred solution of compound **2** (120 mg, 0.229 mmol) in THF (5 cm<sup>3</sup>) was added LiAlH<sub>4</sub> (1 M in THF; 3 cm<sup>3</sup>, 3.00 mmol) dropwise. The reaction was quenched with MeOH<sup>18</sup> (1 cm<sup>3</sup>) and the mixture was poured into 10% aq. tartaric acid. The product was extracted with CH<sub>2</sub>Cl<sub>2</sub>. The solvent was evaporated and the residue was chromatographed on SiO<sub>2</sub> (6 g; hexane–EtOAc, 50:1, 10:1 and 5:1) to give compound **8** (92 mg, 78%),  $\delta_{\text{H}}$  7.88–7.78 (2 H, m, ArH), 7.66–7.46 (3 H, m, ArH), 3.55 (1 H, td,  $J$  11.1, 3.7, 4-H<sub>3</sub>), 2.51–2.37 (1 H, m), 0.99 (3 H, s, 19-H<sub>3</sub>), 0.88 (3 H, d,  $J_{20,21}$  6.3, 21-H<sub>3</sub>), 0.85 (3 H, d,  $J_{25,26}$  6.6, 26-H<sub>3</sub>), 0.85 (3 H, d,  $J_{25,27}$  6.6, 27-H<sub>3</sub>), 0.63 (3 H, s, 18-H<sub>3</sub>);  $\delta_{\text{C}}$  138.9 (C-*i*)(0), 133.1 (C-*o*)(1), 128.8 (C-*m*)(1), 128.5 (C-*p*)(1), 61.1 (C-4)(1), 56.5 (1), 56.3 (1), 43.6 (1), 42.6 (0), 42.1 (1), 40.1 (2), 39.4 (2), 37.1 (0), 36.5 (2), 36.1 (2), 35.7 (1), 35.7 (1), 28.9 (2), 28.3 (2), 28.0 (1), 26.1 (2), 24.1 (C-21)(3), 23.8 (2), 23.1 (2), 22.8 (C-26)(3), 22.5 (C-27)(3), 20.8 (2), 19.5 (2), 18.6 (C-19)(3), 12.0 (C-18)(3);  $m/z$  (EI) 512.368 99 (M<sup>+</sup>, 0.5%, C<sub>33</sub>H<sub>52</sub>O<sub>2</sub>S requires  $M$ , 512.368 80), 510 (1), 495 (0.3), 447 (0.3), 397 (0.4), 371 (100), 355 (7), 245 (14), 163 (23), 149 (35), 135 (30), 109 (41), 95 (55), 81 (45).

#### Desulfuration of **8**

To a stirred solution of sulfone **8** (30 mg, 0.06 mmol) in THF (2 cm<sup>3</sup>) and MeOH (10 cm<sup>3</sup>) was added 6% sodium amalgam (310 mg) in portions during 12 h. The mixture was partitioned between hexane and water, and the product was isolated in the usual way. 5 $\beta$ -Cholestane was obtained (21 mg, 96%),  $\delta_{\text{H}}$  0.91 (3 H, s, 19-H<sub>3</sub>) overlapping with 0.90 (3 H, d,  $J_{20,21}$  6.5, 21-H<sub>3</sub>) overlapping with 0.86 (6 H, d,  $J_{25,26}$  and  $J_{25,27}$  6.7, 26- and 27-H<sub>3</sub>), 0.64 (3 H, s, 18-H<sub>3</sub>);  $\delta_{\text{C}}$  56.7, 56.4, 43.8, 42.7, 40.6, 40.3, 39.5, 37.6, 36.2, 35.9, 35.8, 35.4, 28.3, 28.0, 27.6, 27.3, 27.1, 26.6, 24.3, 23.8, 22.8, 22.6, 21.4, 20.9, 18.7, 12.1 (lit.,<sup>19</sup>  $\delta_{\text{C}}$  56.6, 56.3, 43.7, 42.7, 40.5, 40.3, 39.5, 37.6, 36.2, 35.9, 35.8, 35.3, 28.3, 28.0, 27.5, 27.2, 27.0, 26.6, 24.3, 23.8, 22.8, 22.5, 21.3, 20.8, 18.6, 12.0).

#### Reduction of **2** with NaBH<sub>4</sub> in MeOH. 4 $\alpha$ -Phenylsulfonyl-5 $\alpha$ -cholestan-3 $\alpha$ -ol **9** and 4 $\beta$ -phenylsulfonyl-5 $\beta$ -cholestan-3 $\beta$ -ol **10**

To a stirred solution of ketone **2** (201 mg, 0.38 mmol) in MeOH (20 cm<sup>3</sup>) was added NaBH<sub>4</sub> (161 mg, 4.2 mmol) in portions during 7 h. The mixture was set aside for 16 h and then was poured into water. The product was extracted with CH<sub>2</sub>Cl<sub>2</sub>. The extract was evaporated. The residue was chromatographed on SiO<sub>2</sub> (7 g; benzene–EtOAc, 85:15). A mixture of alcohols **9** and **10** was obtained (168 mg, 83%). A sample of pure isomer **9** was obtained by dissolution of the mixture in CH<sub>2</sub>Cl<sub>2</sub> and precipitation of an amorphous solid with pentane;  $\delta_{\text{H}}$  7.96–7.88 (2 H, m, ArH), 7.70–7.51 (3 H, m, ArH), 3.75 (1 H, br s, 3-H<sub>3</sub>), 3.10 (1 H, dd,  $J_{4,5}$  11.5,  $J_{3,4}$  1.8, 4-H), 2.38–2.20 (2 H, m), 2.01–0.81 (29 H, m) overlapping with 0.89 (3 H, d,  $J_{20,21}$  6.2, 21-H<sub>3</sub>), 0.85 (6 H, d,  $J_{25,26}$  and  $J_{25,27}$  7.3, 26- and 27-H<sub>3</sub>) overlapping with 0.84 (3 H, s, 19-H<sub>3</sub>), 0.64 (3 H, s, 18-H<sub>3</sub>); NOE experiments: Irradiation at  $\delta$  0.84 ppm (19-H<sub>3</sub>) led to an increase of intensity of the signal at  $\delta$  3.1 (3-H) by 1.4% being observed. Irradiation at  $\delta$  3.10 (4-H) led to an increase of signals at  $\delta$  0.84 by 1.4% and at  $\delta$  3.75 ppm (3-H) by 5.2%. Upon irradiation at  $\delta$  3.75 an increase of the signal at  $\delta$  3.10 by 5.0% was observed;  $\delta_{\text{C}}$  139.5 (C-*i*)(0), 133.5 (C-*o*)(1), 129.1 (C-*m*)(1), 128.1 (C-*p*)(1), 67.8 (C-3 or -4)(1), 65.1 (C-4 or 3)(1), 56.2 (1), 56.1 (1), 54.1 (1), 42.3 (0), 40.3 (1), 39.9 (2), 39.5 (2), 37.7 (0), 36.1 (2), 35.8 (1), 34.6 (1), 31.5 (2), 31.2 (2), 28.3 (2), 28.2 (2), 28.0 (1), 25.2 (2), 24.0 (2), 23.8 (2), 22.8 (3), 22.5 (3), 21.2 (2), 18.6 (3), 12.4 (C-19)(3),

12.0 (C-18)(3);  $m/z$  (EI) 528.363 68 (M<sup>+</sup>, 8%, C<sub>33</sub>H<sub>52</sub>O<sub>3</sub>S requires  $M$ , 528.363 71), 463 (3), 445 (8), 387 (86), 369 (100), 353 (5), 329 (3), 287 (5), 261 (8), 243 (10), 231 (21), 215 (12), 175 (11), 161 (14), 135 (16), 107 (7).

Compound **10**:  $\delta_{\text{H}}$  selected signals: 3.91 (1 H, br s), 3.15 (1 H, dd,  $J$  12.1, 2.0), 1.03 (3 H, s, 19-H<sub>3</sub>).

On attempted reduction of **2** with NaBH<sub>4</sub> in MeOH at the reflux temp., the enol of **2** [4-(phenylsulfonyl)cholesta-3,5-dien-3-ol] was obtained in 85% yield,  $\lambda_{\text{max}}$ (EtOH)/nm 201.3, 222.5, 248.9, 286.8;  $\delta_{\text{H}}$  11.25 (1 H, s, OH), 7.84–7.77 (2 H, m, ArH), 7.67–7.43 (3 H, m, ArH), 6.08 (1 H, dd,  $J_{6,7a}$  5.5,  $J_{6,7b}$  2.7, 6-H), 2.50–2.38 (2 H, m), 0.88 (3 H, d,  $J_{20,21}$  6.0, 21-H<sub>3</sub>) overlapping with 0.85 (6 H, d,  $J_{25,26}$  and  $J_{25,27}$  6.2, 26- and 27-H), 0.62 (3 H, s, 19-H<sub>3</sub>), 0.59 (3 H, s, 18-H<sub>3</sub>);  $\delta_{\text{C}}$  165.4, 140.8 (C-*i*), 133.1 (C-*o*), 131.8, 128.6 (C-*m*), 127.0 (C-*p*), 122.0, 106.6, 56.6, 56.0, 48.7, 39.6, 39.5, 42.2, 39.6, 39.5, 36.1, 35.9, 35.7, 32.1, 31.8, 30.5, 28.2, 28.0, 27.5, 24.1, 23.8, 22.8, 22.5, 21.1, 18.6, 17.3, 11.8 (C-18);  $m/z$  (EI) 524 (M<sup>+</sup>, 19%), 509 (2), 460 (100), 445 (15), 427 (3), 383 (43), 369 (3), 365 (4), 271 (10), 257 (8), 247 (10), 229 (15), 213 (8), 197 (16), 173 (15), 147 (15), 135 (18), 121 (19), 105 (22), 95 (35).

#### 4 $\alpha$ -Phenylsulfonyl-5 $\alpha$ -cholestan-3-one **11** and 4 $\beta$ -phenylsulfonyl-5 $\beta$ -cholestan-3-one **12**

A mixture of **9** and **10**, as described above, (95 mg, 0.18 mmol) was dissolved in acetone (5 cm<sup>3</sup>) and treated with Jones reagent until the brown colour persisted (2.7 M; 0.1 cm<sup>3</sup>, 0.27 mmol), then some propan-2-ol was added and the solution was partitioned between water and CH<sub>2</sub>Cl<sub>2</sub>. The organic extract was evaporated. The residue was chromatographed on SiO<sub>2</sub> (10 g; hexane–EtOAc, 92:8) to give consecutively ketones **12** (39 mg, 41%) and **11** (55 mg, 58%).

Compound **12** showed  $\delta_{\text{H}}$  7.82–7.75 (2 H, m, ArH), 7.71–7.50 (3 H, m, ArH), 3.63 (1 H, d,  $J_{4,5}$  6.3, 4-H), 2.76 (1 H, br s), 2.55 (1 H, dd,  $J$  19.6, 7.1), 2.35–2.13 (1 H, m), 1.12 (3 H, s, 19-H<sub>3</sub>), 0.87 (3 H, d,  $J_{20,21}$  6.4, 21-H<sub>3</sub>), 0.84 (6 H, d,  $J_{25,26}$  and  $J_{25,27}$  6.0, 26- and 27-H<sub>3</sub>), 0.63 (3 H, s, 18-H<sub>3</sub>);  $\delta_{\text{C}}$  204.3 (C-3), 137.6 (C-*i*), 134.0 (C-*o*), 129.0 (C-*m*), 128.9 (C-*p*), 76.5 (C-4), 56.1, 55.9, 43.5, 42.8, 42.5, 39.5, 39.4, 36.1, 35.7, 35.0, 34.8, 34.6, 32.1, 28.2, 28.0, 26.3, 24.0, 23.8, 22.8, 22.5, 22.3, 21.4, 18.6, 11.9 (C-18);  $m/z$  (EI) 526.348 00 (M<sup>+</sup>, 2%, C<sub>33</sub>H<sub>50</sub>O<sub>3</sub>S requires  $M$ , 526.348 06), 511 (1), 462 (7), 385 (100), 367 (20), 315 (13), 275 (5), 245 (12), 231 (22), 213 (8), 161 (12), 135 (12), 121 (22), 95 (32).

Compound **11** showed  $\delta_{\text{H}}$  7.82–7.74 (2 H, m, ArH), 7.69–7.48 (3 H, m, ArH), 3.49 (1 H, dd,  $J_{4,5}$  9.1, 1.3, 4-H), 2.87–2.53 (3 H, m), 2.45–2.25 (2 H, m), 0.90 (3 H, d,  $J_{20,21}$  6.6, 21-H<sub>3</sub>) overlapping with 0.86 (6 H, d,  $J_{25,26}$  and  $J_{25,27}$  7.1, 26- and 27-H<sub>3</sub>), 0.74 (3 H, s, 19-H<sub>3</sub>), 0.64 (3 H, s, 18-H<sub>3</sub>);  $\delta_{\text{C}}$  204.5 (C-3), 138.0 (C-*i*), 133.9 (C-*o*), 128.9 (C-*m*), 128.8 (C-*p*), 76.2 (C-4), 58.1, 56.0, 53.8, 45.2, 42.5, 39.8, 39.5, 36.1, 35.9, 35.7, 35.6, 35.0, 34.5, 31.2, 29.1, 28.2, 28.0, 24.1, 23.8, 22.8, 22.5, 21.4, 18.6, 13.7 (C-19), 12.0 (C-18);  $m/z$  (EI) 526.348 00 (M<sup>+</sup>, 8%), 462 (28), 444 (2), 392 (12), 385 (53), 369 (27), 353 (3), 315 (5), 271 (7), 245 (20), 229 (100), 215 (22), 187 (10), 161 (10).

#### Desulfuration of **11**

To a mixture of sulfone **11** (45 mg, 0.09 mmol), THF (2 cm<sup>3</sup>) and MeOH (10 cm<sup>3</sup>) was added 6% sodium amalgam (1.2 g) in portions until starting material was consumed (TLC, 5 days). The mixture was partitioned between hexane and water. The product was isolated in the usual way. 5 $\alpha$ -Cholestan-3-one was obtained (26 mg, 79%), identical in all respects with an authentic sample.

#### Desulfuration of **12**

Sulfone **12** (48 mg, 0.09 mmol) treated with sodium amalgam under conditions analogous to those described above afforded 5 $\beta$ -cholestan-3-one (28 mg, 80%).

### Reduction of **2** with NaBH<sub>4</sub> in a mixture of THF and DMPU

To a stirred mixture of NaBH<sub>4</sub> (10 mg, 0.26 mmol), THF (2 cm<sup>3</sup>) and 1,3-dimethylperhydropyrimidin-2(1*H*)-one (DMPU) (0.3 cm<sup>3</sup>) was added ketone **2** (105 mg, 0.2 mmol) in THF (2 cm<sup>3</sup>) at 0 °C. After 1 h (at 0 °C), the mixture was allowed to warm to room temp. and was partitioned between benzene and 3% HCl. The organic extract was washed successively with water and brine, and the solvent was evaporated. The residue was chromatographed on SiO<sub>2</sub> (5g; hexane–EtOAc, 95:5) to give compound **12** (23%) and a mixture of alcohols **9** and **10** (74%) in the ratio 5.5:1 (determined by <sup>1</sup>H NMR after oxidation of the mixture to a mixture of ketones **11** and **12**). In an analogous reaction carried out at –45 °C, ketone **12** (24%) and alcohols **9** and **10** (73%, in the ratio 4.5:1, respectively) were obtained.

### Reduction of **11** with NaBH<sub>4</sub> in THF–DMPU

A mixture of **11** (48 mg, 0.09 mmol), THF (1 cm<sup>3</sup>), DMPU (0.3 cm<sup>3</sup>) and NaBH<sub>4</sub> (4 mg, 0.1 mmol) was stirred at 0 °C for 1 h, and then worked up in an analogous manner to that described above. Alcohol **9** (47 mg, 97%) was obtained.

Attempted reduction of **12** under analogous conditions failed.

### Reduction of **2** with CuCN–*n*-BuLi–DIBAH system<sup>16,20</sup>

To a suspension of CuCN (46 mg, 0.513 mmol) in THF (5 cm<sup>3</sup>) was added *n*-BuLi (1.3 M in hexane; 0.39 cm<sup>3</sup>, 0.513 mmol) at –20 °C. After 30 min, the mixture was cooled to –50 °C and DIBAH (0.7 M in hexane; 1.833 cm<sup>3</sup>, 1.283 mmol) and HMPA (1 cm<sup>3</sup>) were added consecutively. Stirring at –50 °C was continued for 2 h, and then ketone **2** (52.5 mg, 0.100 mmol) in THF (1 cm<sup>3</sup>) was added. After an additional 3 h, the mixture was allowed to warm to room temp. The reaction was quenched with MeOH (2 cm<sup>3</sup>) and the mixture was poured into 10% aq. tartaric acid. The product was isolated with CH<sub>2</sub>Cl<sub>2</sub> and chromatographed on SiO<sub>2</sub> (6 g; hexane–EtOAc, 40:1, then 20:1) to give 4-(phenylsulfonyl)cholest-4-en-3β-ol **13a** (5 mg, 10%), **12** (11.5 mg, 22%) and **11** (20.5 mg, 39%). Compound **13a** showed  $\delta_{\text{H}}$  7.98–7.90 (2 H, m, ArH), 7.62–7.44 (3 H, m, ArH), 4.75 (1 H, t, *J*<sub>2,3</sub> 5.5, 3-H<sub>3</sub>), 3.05 (1 H, dt, *J* 13.9, 1.8), 1.06 (3 H, s, 19-H<sub>3</sub>), 0.86 (3 H, d, *J*<sub>20,21</sub> 4.6, 21-H<sub>3</sub>), 0.84 (6 H, d, *J*<sub>25,26</sub> and 25,27 6.6, 26- and 27-H<sub>3</sub>), 0.62 (3 H, s, 18-H<sub>3</sub>);  $\delta_{\text{C}}$  164.0 (C-4), 143.4 (C-5), 136.0 (C-*i*), 132.7 (C-*o*), 128.9 (C-*m*), 128.4 (C-*p*), 65.7 (C-3), 55.9, 55.6, 52.3, 42.4, 41.4, 39.6, 39.4, 36.0, 35.6, 35.2, 32.1, 31.0, 28.1, 27.9, 27.7, 26.6, 23.8, 23.7, 22.8, 22.5, 21.5, 20.5, 18.5, 11.9 (C-18); *m/z* (LSIMS) 549.337 77 (M<sup>+</sup> + Na. C<sub>33</sub>H<sub>50</sub>NaO<sub>3</sub>S requires *m/z*, 549.337 84).

### Reduction of **2** with DIBAH. 4-(Phenylsulfonyl)cholest-4-en-3β-ol **13a**

To a solution of **2** (210 mg, 0.38 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (10 cm<sup>3</sup>), stirred at –20 °C, was added DIBAH (0.7 M in hexane; 0.65 cm<sup>3</sup>, 0.46 mmol). After 1 h (at –20 °C) the reaction was quenched with MeOH (0.1 cm<sup>3</sup>). The mixture was warmed to room temp. and partitioned between 3% HCl and CH<sub>2</sub>Cl<sub>2</sub>. The product was isolated in the usual way. Alcohol **13a** (212 mg, 100%) was obtained.

### Reduction of **13a** with LiAlH<sub>4</sub>

A mixture of **13a** (98 mg, 0.19 mmol), THF (3 cm<sup>3</sup>) and LiAlH<sub>4</sub> (10 mg, 0.26 mmol) was stirred for 2 h at room temp. Work-up with saturated aq. Na<sub>2</sub>SO<sub>4</sub> and the usual isolation of product gave **8** (70 mg, 78%).

### 3β-Methoxymethyl-4-(phenylsulfonyl)cholest-4-ene **13b**

To mixture of **13a** (102 mg, 0.19 mmol) in THF (5 cm<sup>3</sup>) and HMPA (0.25 cm<sup>3</sup>) was added NaH (55% in mineral oil; 51 mg,

1.17 mmol), followed by CH<sub>3</sub>OCH<sub>2</sub>Cl (90%; 0.06 cm<sup>3</sup>, 0.74 mmol) and KI (0.1 mg). The mixture was heated under reflux for 24 h and then cooled and partitioned between water and benzene. The organic layer was washed with water and the solvent was evaporated. The residue was chromatographed on SiO<sub>2</sub> (5 g; hexane–EtOAc, 93:7) to give **13b** (106 mg, 96%),  $\delta_{\text{H}}$  7.98–7.88 (2 H, m, ArH), 7.56–7.40 (3 H, m, ArH), 4.95 (1 H, d, *J* 7.0, 3-H), 4.68 (2 H, d, *J* 6.8, OCH<sub>2</sub>O), 3.39 (3 H, s, CH<sub>3</sub>O), 3.07 (1 H, br d, *J* 12.9), 1.01 (3 H, s, 19-H<sub>3</sub>), 0.86 (3 H, d, *J*<sub>20,21</sub> 5.4, 21-H<sub>3</sub>) overlapping with 0.83 (6 H, d, *J*<sub>25,26</sub> and 25,27 6.5, 26- and 27-H<sub>3</sub>), 0.62 (3 H, s, 18-H<sub>3</sub>);  $\delta_{\text{C}}$  165.2 (C-4)(0), 143.9 (C-5)(0), 133.8 (C-*i*)(0), 132.3 (C-*o*)(1), 128.6 (C-*m*)(1), 126.6 (C-*p*)(1), 97.7 (OCH<sub>2</sub>O)(2), 71.5 (C-3)(1), 55.9 (1), 55.8 (1), 55.7 (1), 50.5 (CH<sub>3</sub>O)(3), 42.5 (0), 41.1 (0), 39.6 (2), 39.4 (2), 36.0 (2), 35.6 (1), 35.3 (1), 32.8 (2), 28.5 (2), 28.2 (2), 27.9 (3), 27.6 (2), 25.6 (2), 23.9 (2), 23.7 (2), 22.7 (3), 22.5 (3), 21.9 (2), 21.5 (3), 18.5 (3), 12.0 (C-18)(3).

### Reduction of **13b** with LiAlH<sub>4</sub>

A mixture of ether **13b** (102 mg, 0.18 mmol), THF (5 cm<sup>3</sup>) and LiAlH<sub>4</sub> (35 mg, 0.92 mmol) was heated under reflux for 20 min and cooled. The reagent excess was decomposed with water and the mixture was partitioned between 3% HCl and benzene. Product was isolated in the usual way and chromatographed on SiO<sub>2</sub> (5 g; hexane–EtOAc, 9:1). A mixture of sulfones **8**, **7a** and **7b** was obtained (70 mg, 76%) in proportions 5:4:1 by <sup>1</sup>H NMR. Rechromatography of the mixture gave pure products **8** and **7a** and **7b**.

### 4-(Phenylsulfonyl)cholest-4-en-3α-ol **15a**

(a) **From 13a.** To a solution of alcohol **13a** (113 mg, 0.21 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (3 cm<sup>3</sup>) was added Ph<sub>3</sub>P (265 mg, 1.1 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (1 cm<sup>3</sup>) at –78 °C. The mixture was stirred for 2 h and then benzoic acid (135 mg, 1.1 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (2 cm<sup>3</sup>) and diethyl azodicarboxylate (0.175 cm<sup>3</sup>, 1.1 mmol) were added. The mixture was set aside at room temp. for 4 h and then partitioned between saturated aq. NaHCO<sub>3</sub> and benzene. The organic phase was washed with water and the solvent was evaporated. The residue was chromatographed on SiO<sub>2</sub> (5 g; hexane–EtOAc, 95:5) to give benzoate **15b** (80 mg). This crude product was dissolved in 5% methanolic KOH (3 cm<sup>3</sup>) and set aside for 16 h. The usual work-up and chromatography of the product on SiO<sub>2</sub> (5 g; hexane–EtOAc, 8:2) gave alcohol **15a** (72 mg, 64%).

(b) **From 16a.** To a solution of sulfide **16a** (see below) (59 mg, 0.119 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (3 cm<sup>3</sup>) was added MCPBA (60%; 70 mg, 0.243 mmol) at 0 °C. The mixture was stirred at room temp. for 1 h and poured into aq. Na<sub>2</sub>CO<sub>3</sub>. The product was extracted with CH<sub>2</sub>Cl<sub>2</sub>. The extract was washed successively with aq. Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub> and water, and the solvent was evaporated. The residue was chromatographed on SiO<sub>2</sub> (2 g; hexane–EtOAc, 50:1, 25:1, 10:1) to give alcohol **15a** (61 mg, 97%),  $\delta_{\text{H}}$  7.97–7.90 (2 H, m, ArH), 7.63–7.46 (3 H, m, ArH), 4.77 (1 H, s, 3-H<sub>3</sub>), 3.15 (1 H, dt, *J* 14.5, 3.3), 1.03 (3 H, s, 19-H<sub>3</sub>), 0.87 (3 H, d, *J*<sub>20,21</sub> 4.7, 21-H<sub>3</sub>) overlapping with 0.85 (6 H, d, *J*<sub>25,26</sub> and 25,27 5.7, 26- and 27-H<sub>3</sub>), 0.63 (3 H, s, 18-H<sub>3</sub>);  $\delta_{\text{C}}$  163.0 (C-4)(0), 143.1 (C-5)(0), 135.6 (C-*i*)(0), 132.7 (C-*o*)(1), 128.9 (C-*m*)(1), 126.7 (C-*p*)(1), 62.9 (C-3)(1), 55.8 (1), 55.4 (1), 54.5 (1), 42.2 (0), 40.9 (0), 39.5 (2), 39.3 (2), 35.9 (1), 35.6 (1), 34.9 (1), 31.2 (2), 30.8 (2), 28.0 (2), 27.9 (1), 27.2 (2), 26.5 (2), 23.8 (2), 23.6 (2), 22.7 (3), 22.4 (2), 21.6 (2), 18.4 (2C)(3), 11.8 (C-18)(3); *m/z* (EI) 525 (M<sup>+</sup>, 0.3%), 508.3373 [(M – H<sub>2</sub>O)<sup>+</sup>, 99. C<sub>33</sub>H<sub>48</sub>O<sub>2</sub>S requires *m/z*, 508.3375], 493 (19), 444 (54), 429 (9), 395 (44), 385 (8), 367 (82), 351 (8), 261 (81), 247 (62), 159 (33), 147 (80), 135 (71), 119 (42), 109 (54), 105 (100), 95 (82), 91 (52), 81 (71), 69 (45), 55 (50), 43 (45).

### Reduction of **15a** with LiAlH<sub>4</sub>

To a solution of alcohol **15a** (70 mg, 0.13 mmol) in THF (2 cm<sup>3</sup>)

was added  $\text{LiAlH}_4$  (10 mg, 0.26 mmol) and the mixture was stirred for 2 h. The reagent excess was decomposed with saturated aq.  $\text{Na}_2\text{SO}_4$ , and the mixture was diluted with  $\text{CH}_2\text{Cl}_2$  and dried with  $\text{Na}_2\text{SO}_4$ . The solid was filtered off and the solvent was evaporated. The residue was chromatographed on  $\text{SiO}_2$  (5 g; hexane–EtOAc, 9:1) to give compound **7a** (56 mg, 82%).

#### Reduction of **1** with DIBAH. 4-(Phenylsulfanyl)cholest-4-en-3 $\beta$ -ol **14** and 4-(phenylsulfanyl)cholest-4-en-3 $\alpha$ -ol **16a**

To a solution of **1** (450 mg, 0.91 mmol) in  $\text{CH}_2\text{Cl}_2$  (10  $\text{cm}^3$ ) was added DIBAH (0.7 M in hexane; 4.4  $\text{cm}^3$ , 3.08 mmol) at  $-78^\circ\text{C}$ . The mixture was stirred for 1 h at  $-78^\circ\text{C}$  and allowed to warm to room temp. The reaction was quenched with MeOH (2  $\text{cm}^3$ ) and the mixture was poured into 10% aq. tartaric acid. The product was isolated with  $\text{CH}_2\text{Cl}_2$  and chromatographed on  $\text{SiO}_2$  (45 g; hexane–EtOAc, 100:1 and 75:1) to give alcohols **14** (312 mg, 69%) and **16a** (96 mg, 21%).

Compound **14** showed  $\delta_{\text{H}}$  7.31–7.08 (5 H, m, ArH), 4.08–3.86 (1 H, m, 3-H), 3.31–3.18 (1 H, m, 8 lines, 6- $\alpha$ H), 2.90 (1 H, br s, 3-OH), 1.17 (3 H, s, 19- $\text{H}_3$ ), 0.92 (3 H, d,  $J_{20,21}$  6.6, 21- $\text{H}_3$ ), 0.87 (6 H, d,  $J_{25,26}$  and  $J_{25,27}$  6.6, 26- and 27- $\text{H}_3$ ), 0.70 (3 H, s, 18- $\text{H}_3$ );  $\delta_{\text{C}}$  157.6 (C-5), 136.2 (C-*i*), 128.8 (C-*o*), 127.0 (C-*m*), 125.7 (C-4), 125.3 (C-*p*), 68.0 (C-3), 56.0, 56.0, 54.4, 42.3 (C-13), 40.6 (C-10), 39.7, 39.4, 36.0, 35.6, 35.4, 33.7, 32.7, 29.1, 28.1, 27.9, 27.3, 24.0, 23.7, 22.7, 22.4, 21.2, 19.8, 18.5, 11.9;  $m/z$  (EI) 494.358 35 ( $\text{M}^+$ , 100%,  $\text{C}_{33}\text{H}_{50}\text{OS}$  requires  $M$ , 494.358 24), 476 [( $\text{M} - \text{H}_2\text{O}$ ) $^+$ , 94], 461 [( $\text{M} - \text{CH}_3 - \text{H}_2\text{O}$ ) $^+$ , 13], 399 [( $\text{M} - \text{Ph} - \text{H}_2\text{O}$ ) $^+$ , 6], 385 (19), 367 (65), 105 (29), 95 (34), 57 (25).

Compound **16a** showed  $\delta_{\text{H}}$  7.31–7.06 (5 H, m, ArH), 4.05–3.98 (1 H, m, 3-H), 3.37–3.23 (1 H, m, 6 lines, 6- $\alpha$ H), 2.27 (1 H, br s, 3-OH), 1.14 (3 H, s, 19- $\text{H}_3$ ), 0.90 (3 H, d,  $J_{20,21}$  8.6, 21- $\text{H}_3$ ), 0.87 (6 H, d,  $J_{25,26}$  and  $J_{25,27}$  6.6, 26- and 27- $\text{H}_3$ ), 0.70 (3 H, s, 18- $\text{H}_3$ );  $\delta_{\text{C}}$  157.7 (C-5), 136.7 (C-*i*), 128.9 (C-*o*), 127.5 (C-*m*), 125.3 (C-4), 124.4 (C-*p*), 67.5 (C-3), 56.0, 55.9, 54.4, 42.3 (C-13), 40.6 (C-10), 39.8, 39.4, 36.0, 35.7, 35.6, 32.2, 31.5, 28.5, 28.1, 27.9, 26.9, 24.0, 23.7, 22.7, 22.5, 21.6, 18.6, 18.5, 11.9;  $m/z$  (EI) 494.358 41 ( $\text{M}^+$ , 38%), 476.347 53 [( $\text{M} - \text{H}_2\text{O}$ ) $^+$ , 100],  $\text{C}_{33}\text{H}_{48}\text{S}$  requires  $m/z$ , 476.347 67], 461 [( $\text{M} - \text{CH}_3 - \text{H}_2\text{O}$ ) $^+$ , 12], 399 [( $\text{M} - \text{Ph} - \text{H}_2\text{O}$ ) $^+$ , 6], 385 (5), 367 (30), 109 [(SPh) $^+$ , 14], 105 (36), 95 (24), 57 (19).

#### Reduction of **1** with $\text{LiAlH}_4$

To a stirred solution of  $\text{LiAlH}_4$  in THF (1.5 M; 5  $\text{cm}^3$ ) was added a solution of **1** (150 mg, 0.305 mmol) in THF (3  $\text{cm}^3$ ) in one portion. After 20 min the reaction was quenched with MeOH (1  $\text{cm}^3$ ) and the mixture was poured into 10% aq. tartaric acid. The product was extracted with  $\text{CH}_2\text{Cl}_2$ . The extract was washed with water and evaporated. The residue was filtered through  $\text{SiO}_2$  (15 g; hexane–EtOAc, 40:1) to give a mixture of **14** and **16a** (137 mg, 91%) in the ratio 84:16, respectively, by  $^1\text{H}$  NMR.

#### Reduction of **1** with L-Selectride<sup>®</sup>

To a solution of **1** (105 mg, 0.213 mmol) in THF (7  $\text{cm}^3$ ) was added L-Selectride<sup>®</sup> (1 M in THF; 0.25  $\text{cm}^3$ , 0.25 mmol) at  $-78^\circ\text{C}$ . The mixture was stirred at  $-78^\circ\text{C}$  for 1 h. The reaction was quenched with MeOH (0.5  $\text{cm}^3$ ). The product was isolated as in the above described experiments to give a mixture of **14** and **16a** (83 mg 79%) in the ratio 82:18, respectively, by  $^1\text{H}$  NMR.

#### Reduction of **1** with $\text{NaBH}_4$ – $\text{CeCl}_3$ system<sup>17</sup>

To a mixture of **1** (100 mg, 0.203 mmol),  $\text{CeCl}_3 \cdot 7\text{H}_2\text{O}$  (41 mg, 0.110 mmol), THF (2  $\text{cm}^3$ ) and MeOH (3  $\text{cm}^3$ ) was added  $\text{NaBH}_4$  (8.5 mg, 0.224 mmol) at  $-78^\circ\text{C}$ . After 2 h at  $-78^\circ\text{C}$ , the mixture was allowed to warm to room temp. and then was poured into 10% aq. tartaric acid. The product was isolated

with  $\text{CH}_2\text{Cl}_2$  and chromatographed on  $\text{SiO}_2$  (10 g; hexane–EtOAc, 60:1 and 40:1) to give **14** (91 mg, 91%). Inspection of the  $^1\text{H}$  NMR spectrum of this product showed the presence of traces ( $\approx 3\%$ ) of the isomer **16a**.

#### 4-(Phenylsulfanyl)cholest-4-en-3 $\alpha$ -yl benzoate **16b**

To a solution of alcohol **14** (86 mg, 0.174 mmol),  $\text{Ph}_3\text{P}$  (182 mg, 0.695 mmol) and benzoic acid (78 mg, 0.639 mmol) in THF (1.5  $\text{cm}^3$ ) was added DEAD (0.11  $\text{cm}^3$ , 0.575 mmol) in THF (0.5  $\text{cm}^3$ ) at  $-78^\circ\text{C}$ . The mixture set aside at room temp. for 2 h and the solvent was evaporated. The residue was chromatographed on  $\text{SiO}_2$  (5 g; hexane–EtOAc, 100:1, 50:1 and 20:1) to give the title benzoate **16b** (77 mg, 76%),  $\delta_{\text{H}}$  7.89–7.00 (10 H, m, ArH), 5.51 (1 H, br s, 3-H), 3.51–3.36 (1 H, m, 6 lines, 6- $\alpha$ H), 1.19 (3 H, s, 19- $\text{H}_3$ ), 0.94 (3 H, d,  $J_{20,21}$  6.4, 21- $\text{H}_3$ ), 0.89 (6 H, d,  $J_{25,26}$  and  $J_{25,27}$  6.6, 26- and 27- $\text{H}_3$ ), 0.73 (3 H, s, 18- $\text{H}_3$ );  $\delta_{\text{C}}$  165.9 (C=O), 160.8 (C-5), 137.0 (C-*i*, SPh), 132.4 (C-*p*, Bz), 130.6 (C-*i*, Bz), 129.5 (C-*o*, Bz), 128.6 (C-*o*, SPh), 128.0 (C-*m*, Bz), 127.6 (C-*m*, SPh), 125.1 (C-*p*, SPh), 120.7 (C-4), 72.1 (C-3), 56.1, 55.9, 54.7, 42.3 (C-13), 40.4 (C-10), 39.7, 39.4, 36.0, 35.6, 35.5, 32.2, 32.1, 28.5, 28.1, 27.9, 25.7, 24.0, 23.7, 22.7, 22.5, 21.6, 18.5, 18.5, 11.9;  $m/z$  (EI) 598.384 14 ( $\text{M}^+$ , 11%,  $\text{C}_{40}\text{H}_{54}\text{O}_2\text{S}$  requires  $M$ , 598.384 45), 476.347 80 ( $\text{M}^+ - \text{C}_6\text{H}_5\text{CO}_2\text{H}$ , 100),  $\text{C}_{33}\text{H}_{48}\text{S}$  requires  $m/z$ , 476.347 67], 461 [( $\text{M} - \text{CH}_3 - \text{BzOH}$ ) $^+$ , 21], 367 (24), 122 ( $\text{BzOH}^+$ , 18), 105 ( $\text{C}_6\text{H}_5\text{CO}^+$ , 100).

#### 4-(Phenylsulfanyl)cholest-4-en-3 $\alpha$ -ol **16a** from **16b**

To a solution of benzoate **16b** (77 mg, 0.129 mmol) in  $\text{CH}_2\text{Cl}_2$  (3  $\text{cm}^3$ ) was added DIBAH (0.7 M in hexane; 0.4  $\text{cm}^3$ , 0.280 mmol) at  $-78^\circ\text{C}$ . The mixture was stirred for 1 h before being allowed to warm to room temp., treated with MeOH (0.5  $\text{cm}^3$ ), and poured into 10% aq. tartaric acid. The product was isolated with  $\text{CH}_2\text{Cl}_2$  and chromatographed on  $\text{SiO}_2$  (1.5 g; hexane–EtOAc, 100:1 and 50:1) to give **16a** (59 mg, 92%).

#### Reduction of **15a** with $\text{LiAlH}_4$

To a solution of hydroxy sulfone **15a** (61 mg, 0.116 mmol) in THF (4  $\text{cm}^3$ ) was added  $\text{LiAlH}_4$  (0.25 M in THF; 0.6  $\text{cm}^3$ , 0.150 mmol). The mixture was heated under reflux for 20 min. After cooling, the reaction mixture was quenched with MeOH (0.5  $\text{cm}^3$ ) and poured into 10% aq. tartaric acid. The product was isolated with  $\text{CH}_2\text{Cl}_2$  and chromatographed on  $\text{SiO}_2$  (2 g; hexane–EtOAc 50:1, 10:1 and 5:1) to give sulfone **7a** (49 mg, 82%).

#### Catalytic hydrogenation of **2**

A solution of **2** (60 mg, 0.11 mmol) in EtOH (4.5  $\text{cm}^3$ ), containing 10% palladium on carbon (10 mg), was stirred under hydrogen at the reflux temperature for 5 h. The mixture was cooled and diluted with  $\text{CH}_2\text{Cl}_2$ . The solid was filtered off and the solvent was evaporated. A mixture of compounds **11** and **12** in the ratio 6.5:1 ( $^1\text{H}$  NMR) was obtained.

In an analogous reaction, a solution of **2** (60 mg, 0.11 mmol) in EtOH (4.5  $\text{cm}^3$ ), containing 10% palladium on carbon (10 mg), was treated with  $\text{CF}_3\text{COOH}$  (0.5 mmol) and the mixture was stirred at room temp. for 5 days. A mixture of **11** and **12** in the ratio 5.5:1 ( $^1\text{H}$  NMR) was obtained.

#### References

- 1 J. Seyden-Penne, *Reduction by the Alumino- and Borohydrides in Organic Synthesis*, Wiley-VCH, New York, 1997.
- 2 A. P. Davis, in *Formation of C–C Bonds by the Reduction of Carbonyl Groups with Metal Hydrides*, ed. G. Helmchen, R. W. Hoffmann, J. Mulzer and E. Schaumann, Thieme, Stuttgart, 1996.
- 3 M. Julia, M. Launay, J.-F. Stacino and J.-N. Verpeaux, *Tetrahedron Lett.*, 1982, **23**, 2465.
- 4 J. S. Grossert, in *Reduction of Sulphoxides and Sulphones*, ed. S. Patai, Z. Rappoport and C. Stirling, Wiley, Chichester, 1988.



- 5 P. S. Jones, S. V. Ley, N. S. Simpkins and A. J. White, *Tetrahedron*, 1986, **42**, 6519.
- 6 M. C. Clasby and D. Craig, *Synth. Commun.*, 1994, **24**, 481.
- 7 K. Michalak, W. Stepanenko and J. Wicha, *Tetrahedron Lett.*, 1996, **37**, 7657.
- 8 A. Przewdziecka, W. Stepanenko and J. Wicha, *Tetrahedron: Asymmetry*, 1999, **10**, 1589.
- 9 H. B. Henbest and W. R. Jackson, *J. Chem. Soc. C*, 1967, 2459.
- 10 A. G. Schultz, W. Y. Fu, R. D. Lucci, B. G. Kurr, K. M. Lo and M. Boxer, *J. Am. Chem. Soc.*, 1978, **100**, 2140.
- 11 A. Furst and R. Scotoni, *Helv. Chim. Acta*, 1953, **36**, 1332.
- 12 A. Kasal, *Collect. Czech. Chem. Commun.*, 1976, **41**, 140.
- 13 P. J. Kocienski, B. Lythgoe and S. Ruston, *J. Chem. Soc., Perkin Trans. 1*, 1979, 1290.
- 14 N. Jones and M. J. Green, *J. Chem. Soc. C*, 1967, 532.
- 15 T. Tsuda, T. Hayashi, H. Satomi, T. Kawamoto and T. Saegusa, *J. Org. Chem.*, 1986, **51**, 537.
- 16 A. R. Daniewski and J. Kiegel, *Synth. Commun.*, 1988, **18**, 115.
- 17 J.-L. Luche, L. Rodrigez-Hahn and P. Crabbe, *J. Chem. Soc., Chem. Commun.*, 1978, 601.
- 18 After quenching of the reaction with MeOH only one epimer of sulfone was obtained (with equatorially oriented phenylsulfonyl group). The quench with excess of saturated aq. Na<sub>2</sub>SO<sub>4</sub> leads to a mixture of axial and equatorial sulfones (≈1:9, respectively).
- 19 D. Zeigan, R. Radgelia and G. Engelhardt, *J. Prakt. Chem.*, 1983, **325**, 651.
- 20 T. Tsuda, T. Kawamoto, Y. Kumamoto and T. Saegusa, *Synth. Commun.*, 1986, **16**, 639.