

Note

Mild and Catalyst-Free Petasis/Decarboxylative Domino Reaction: Chemoselective Synthesis of N-Benzyl Propargylamines

Huangdi Feng, Huihui Jia, and Zhihua Sun

J. Org. Chem., **Just Accepted Manuscript** • DOI: 10.1021/jo502349a • Publication Date (Web): 19 Nov 2014

Downloaded from <http://pubs.acs.org> on November 21, 2014

Just Accepted

"Just Accepted" manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides "Just Accepted" as a free service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. "Just Accepted" manuscripts appear in full in PDF format accompanied by an HTML abstract. "Just Accepted" manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are accessible to all readers and citable by the Digital Object Identifier (DOI®). "Just Accepted" is an optional service offered to authors. Therefore, the "Just Accepted" Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the "Just Accepted" Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these "Just Accepted" manuscripts.



ACS Publications
High quality. High impact.

The Journal of Organic Chemistry is published by the American Chemical Society, 1155 Sixteenth Street N.W., Washington, DC 20036
Published by American Chemical Society. Copyright © American Chemical Society. However, no copyright claim is made to original U.S. Government works, or works produced by employees of any Commonwealth realm Crown government in the course of their duties.


Mild and Catalyst-Free Petasis/Decarboxylative Domino
Reaction: Chemoselective Synthesis of *N*-Benzyl
Propargylamines

Huangdi Feng, Huihui Jia and Zhihua Sun*

College of Chemistry and Chemical Engineering, Shanghai University of Engineering
Science, 333 Longteng Road, Shanghai 201620, China

*E-mail: sungaris@gmail.com

Table of Contents Graphic:



ABSTRACT

Multicomponent domino reactions are attractive for assembling functionalized compounds. To this end, a one-pot catalyst-free chemoselective synthesis of *N*-benzyl propargylamines is reported with good functional group compatibility. This mild process involves *in situ* formation of an active amine through Petasis reaction of primary amines, formaldehyde solution and boronic acids, which reacts with propiolic acids to give product in up to 94% yield via decarboxylative coupling reaction.

Propargylamines can be used as important building blocks for the generation of functionalized five-¹, six-², seven-³ and eight-membered⁴ nitrogen heterocyclic frameworks. In addition, they have been found within highly potent and irreversible selective monoamine oxidase type-B inhibitors⁵. A common route to the synthesis of propargylamines involves the reaction of an aldehyde with an amine and an alkyne, known as the A³-coupling reaction⁶. However, examples for the constitution of *N*-benzyl propargylamines have been rarely studied or reported owing to most of the *N*-substituted benzylamines are not commercially available. As such, the development of ~~greener and~~ more efficient approaches for the preparation of structurally diverse propargylamines is highly desirable for synthetic organic chemistry.

Currently, the transition-metal-catalyzed decarboxylative couplings have been successfully used for C–C bond formation under relatively neutral conditions compared to preformed organometallic reagents^{7, 8}. In this regard, copper-catalyzed decarboxylative couplings have been used as a valuable synthetic strategy for the formation of propargylamines, employing amino acid⁹, 2-oxoacetic acid¹⁰, glyoxylic acid¹¹, propiolic acid¹² as starting materials. However, those copper-catalyzed methods usually suffer from some drawbacks in terms of environmental pollution and complication in the purification processes. Recently, the group of Sunwoo Lee reported a metal-free decarboxylative coupling reaction of propiolic acid with secondary amine and paraformaldehyde¹³, similar to A³-coupling reaction. Although this method provided a facile approach to propargylamines, no example with a primary amine as starting material was reported due to primary amines are considered as being difficult substrates ~~in the A³-type coupling reaction~~¹⁴. Meanwhile, there are a limited number of literature examples in which primary amines were employed in the decarboxylative A³-coupling reaction¹⁵. To expand the decarboxylative coupling reaction for the generation of propargylamines to a broad range of starting amines, we envisioned a scenario wherein a primary amine might participate in a Petasis reaction with an aldehyde and a boronic acid¹⁶, to form an active amine *in situ*, followed by metal-free decarboxylative A³-coupling reaction of a propiolic acid (Figure 1).





Figure 1. Petasis and decarboxylative domino reaction

Key to the success of this proposed reaction would be the addition of the aryl boronic acid into imine **I** to produce *N*-benzyl hemiaminal **II**¹⁷, which in the presence of a propiolic acid results in the formation of iminium salt **III**. It is noteworthy that the related addition of another boronic acid into activated iminium salt **III** has been described to obtain amine **5**¹⁸. In this note, the alkynyl carbon next to the carboxylate attacks iminium salt **III** to generate intermediate **IV** instead of the above process. Subsequent extrusion of CO_2 through decarboxylation pathway affords the desired propargylamine in a process similar to the Eschweiler-Clarke methylation¹⁹. Herein, we report the successful development of such a new catalyst-free one-pot chemoselective synthesis of propargylamines through the Petasis/decarboxylative domino reaction of a primary amine, a formaldehyde solution, an aryl boronic acid and a propiolic acid.

We initiated our investigation using benzylamine (**1a**), formaldehyde solution, phenylboronic acid (**2a**), and 3-phenylpropionic acid (**3a**) in toluene at 60 °C for 10 h (Table 1). To our delight, the desired product **4{1}** was obtained in 31% yield (Table 1, entry 1). To improve the yield, various ratios of the four starting materials were examined (Table 1, entries 2 and 3). Among them, 3.2 equiv of formaldehyde solution provided the best yield (Table 1, entry 2). When paraformaldehyde was used instead of formaldehyde solution (Table 1, entries 4 and 5), only 46% and 40% of products

were isolated, which is much lower than using formaldehyde solution (Table 1, entry 2). Subsequently, solvent screening showed that this reaction was sensitive to the nature of solvents (Table 1, entries 6–12), 1,2-dichloroethane (DCE) provided an improved isolated yield of 83% for **4{1}** (Table 1, entry 6), whereas ethyl acetate, water, acetonitrile, ethanol, dichloromethane and tetrahydrofuran gave lower yields of the desired product. Eventually, different reaction time and temperatures were also tested, but did not give much improvement of the yield (Table 1, entries 13–18). Remarkably, using the reaction temperature of 45 °C was slightly more efficient than 60 °C to afford the optimal result (Table 1, entry 16). When the reaction was run at room temperature for 10 h, a yield of 73% was obtained (Table 1, entry 17). Prolonging the reaction time to 26 h, only produced a minor increase of yield (Table 1, entry 18). In addition, the reaction yield under the optimal conditions can be decreased from 84% to 65% after adding an acid to the reaction (Table 1, entry 19), which could be explained by the adverse effect of an acid to the decarboxylative pathway shown in Figure 1. When ethyl phenylpropiolate was used instead of phenylboronic acid (Table 1, entry 20), no desired product was formed in the presence of 1.5 equiv of acetic acid.

Table 1: Optimization of Reaction Parameters^a

					
entry	ratio	solvent	temp [°C]	t [h]	yield [%] ^b
1	1.0/2.2/1.1/1.2	Toluene	60	10	31
2	1.0/3.2/1.1/1.2	Toluene	60	10	69
3	1.0/4.0/1.1/1.2	Toluene	60	10	54
4 ^c	1.0/3.2/1.1/1.2	Toluene	60	10	46
5 ^c	1.0/3.2/1.1/1.2	H ₂ O/Toluene (1:15)	60	10	40
6	1.0/3.2/1.1/1.2	DCE	60	10	83
7	1.0/3.2/1.1/1.2	AcOEt	60	10	8
8	1.0/3.2/1.1/1.2	H ₂ O	60	10	33
9	1.0/3.2/1.1/1.2	CH ₃ CN	60	10	14

10	1.0/3.2/1.1/1.2	EtOH	60	10	6
11	1.0/3.2/1.1/1.2	DCM	45	10	68
12	1.0/3.2/1.1/1.2	THF	60	10	trace
13	1.0/3.2/1.1/1.2	DCE	60	24	77
14	1.0/3.2/1.1/1.2	DCE	80	5	68
15	1.0/3.2/1.1/1.2	DCE	80	10	65
16	1.0/3.2/1.1/1.2	DCE	45	10	84
17	1.0/3.2/1.1/1.2	DCE	rt	10	73
18	1.0/3.2/1.1/1.2	DCE	rt	26	74
19 ^d	1.0/3.2/1.1/1.2	DCE	45	10	65
20 ^e	1.0/3.2/1.1/1.2	DCE	45	10	0


^aReactions conditions: **1a** (0.5 mmol), formaldehyde solution (1.1-2 mmol), **2a** (0.55 mmol), **3a** (0.6 mmol) and solvent (1.5 mL) in a sealed tube. ^bIsolated yield. ^cParaformaldehyde was used instead of formaldehyde solution. ^d1.5 equiv of acetic acid was added. ^eEthyl phenylpropionate instead of phenylpropionic acid with 1.5 equiv of acetic acid

A variety of primary amines were subjected to the optimized conditions to evaluate the scope of the Patisis/decarboxylative domino reaction. For substituted benzylamines, the reaction afforded the corresponding products **4** in moderate to good yield bearing electron-rich and electron-deficient groups (Table 2, **4{2}–4{12}**). In general, halo-substitution led to the cleanest and highest yielding reactions (Table 2, **4{2}–4{5}**), and cyano-substitution also gave good yield (Table 2, **4{6}**). Methoxy-substitution usually afforded lower yield of the products compared to methyl-substitution (Table 2, **4{7}–4{12}**). The use of aliphatic amines also gave the desired products in nice yields (Table 2, **4{13}–4{15}**). Then, a number of boronic acids were investigated. Both electron-poor and -rich aryl-substituted boronic acids were effective, producing the desired compounds in 59–87% yields (Table 2, **4{16}–4{19}**). In addition, the reactions of aromatic naphthalen-1-ylboronic acid, naphthalen-2-ylboronic acid and thiophen-3-ylboronic acid were also examined. The corresponding coupling products were obtained in high yield (Table 2, **4{20}–4{22}**). It is noteworthy that 4-fluorophenyl boronic acid (**2i**), 4-tolylboronic acid (**2g**), 2-methoxyphenyl boronic acid (**2k**), and 4-methoxyphenyl boronic acid (**2l**) also reacted well under standard conditions to give excellent yields in the formation of

4{4}[#], 4{9}[#], 4{10}[#], and 4{11}[#] when compared to the same structures 4{4}, 4{9}, 4{10}, and 4{11} from alternative starting materials. Finally, our investigations were focused on the use of various 3-substituted propiolic acids **3** (Table 2, 4{23}–4{30}). Both aromatic and aliphatic propiolic acids were explored as reaction partners, and the results clearly showed that aromatic propiolic acids worked well (Table 2, 4{23}–4{27}) and alkyl propiolic acids afforded the target compounds only in low to moderate yields (Table 2, 4{28}–4{30}). It seemed that substituents on the phenyl ring of the aromatic propiolic acids played important roles in this process, as electron-donating substituents (Table 2, 4{25}–4{27}) led to higher yields than electron-withdrawing groups (Table 2, 4{23}–4{24}).

Table 2: Substrate scope and limitations^a


$ \begin{array}{c} \text{R}_1\text{-NH}_2 + \text{H}-\text{C}(=\text{O})-\text{H} + \text{R}_2\text{-B(OH)}_2 + \text{R}_3\text{-C}\equiv\text{C}-\text{COOH} \\ \text{1} \qquad \qquad \qquad \text{2} \qquad \qquad \qquad \text{3} \end{array} \xrightarrow[45\text{ }^\circ\text{C, 10 h}]{\text{DCE}} \begin{array}{c} \text{R}_3\text{-C}\equiv\text{C}-\text{CH}_2\text{-N(R}_1\text{)(R}_2\text{)} \\ \text{4} \end{array} $			
	84% 4{1}		93% 4{2}
	90% 4{3}		78% 4{4}
	88% 4{5}		76% 4{6}
	91% 4{7}		86% 4{8}
	65% 4{9}		60% 4{10}
	63% 4{11}		68% 4{12}
	49% 4{13}		54% 4{14}
	73% 4{15}		59% 4{16} ^b



^aReactions conditions: amine (0.5 mmol), formaldehyde solution (1.6 mmol), boronic acid (0.55 mmol), propionic acid (0.6 mmol) and DCE (1.5 mL) in a sealed tube; reaction was monitored by TLC plate. ^bThis reaction was carried out at 60 °C for 18 h. ^cThis reaction was carried out for 26 h.

Not surprisingly, when isobutylboronic acid was used in this protocol, dipropargylamine **6** was obtained in 73% yield (Scheme 1). Only trace amounts of the expected tertiary amine product was observed as indicated by ¹H NMR spectroscopy and LC–MS of the crude reaction mixtures since the Petasis reaction typically did not work with inactive alkyl-boronic acid.

Scheme 1. Selective coupling reaction toward dipropargylamine



In summary, we have described a catalyst free method for the synthesis of *N*-benzyl propargylamines by multicomponent domino reaction of primary amines with formaldehyde solution, boronic acids and propiolic acids. The reaction uses commercially available reagents, and affords significant advantages over the related transition-metal-catalyzed A^3 -coupling or decarboxylative A^3 -coupling, in that it proceeds under much milder reaction conditions and displays a broad substrate scope and high chemoselectivity.

EXPERIMENTAL SECTION

General Information. All commercially available reagents were used without purification unless otherwise noted. Column chromatography was performed using silica gel (100–200 mesh). Visualization of the compounds was accomplished with UV light (254 nm) and iodine. ^1H NMR and ^{13}C NMR spectra were recorded in CDCl_3 operating at 400 MHz and 100 MHz, respectively. Proton chemical shifts are reported relative to the residual proton signals of the deuterated solvent CDCl_3 (7.26 ppm) or TMS. Carbon chemical shifts were internally referenced to the deuterated solvent signals in CDCl_3 (77.00 ppm) and reported in δ (parts per million) values. Coupling constants J are reported in Hz. Proton coupling patterns were described as singlet (s), doublet (d), triplet (t), quartet (q), and multiple (m). High-resolution mass spectra were recorded on a Liquid Chromatograph Mass Spectrometer (LCMS-IT-TOF).

General Procedure for the Synthesis of Propargylamines 4. In a vial equipped with a stirring bar, a mixture of amine (0.5 mmol) and formaldehyde solution (1.6 mmol) was dissolved in DCE (1.5 mL). Then boronic acid (0.55 mmol) and propiolic acid (0.6 mmol) were added. The reaction vessel was sealed and heated in an oil-bath for 10 h at a temperature of 45 $^\circ\text{C}$. The resulting reaction mixture was loaded on a silica gel column and eluted with 2–5% ethyl acetate in hexane to afford the desired product

4 as light yellow oil.

N,N-dibenzyl-3-phenylprop-2-yn-1-amine (**4{1}**). light yellow oil (131.1 mg, 84% yield); ^1H NMR (400 MHz, CDCl_3) δ 7.57–7.54 (m, 2H), 7.50–7.48 (m, 4H), 7.40–7.35 (m, 7H), 7.33–7.28 (m, 2H), 3.817 (s, 4H), 3.53 (t, 2H); ^{13}C NMR (77 MHz, CDCl_3) δ 139.0, 131.8, 129.1, 128.3, 128.3, 128.0, 127.1, 123.5, 85.90, 84.8, 84.5, 57.8, 42.1; MS (ESI) m/z : 312.2 $[\text{M} + \text{H}]^+$.

N-benzyl-*N*-(4-chlorobenzyl)-3-phenylprop-2-yn-1-amine (**4{2}**). light yellow oil (160.9 mg, 93% yield); ^1H NMR (400 MHz, CDCl_3) δ 7.55–7.45 (m, 2H), 7.42–7.40 (m, 2H), 7.38–7.33 (m, 10H), 3.79 (s, 2H), 3.77 (s, 2H), 3.50 (s, 2H); ^{13}C NMR (77 MHz, CDCl_3) δ 138.7, 137.5, 132.9, 131.8, 130.4, 129.1, 128.6, 128.5, 128.4, 128.3, 128.2, 128.1, 127.2, 123.3, 86.0, 84.2, 57.8, 57.0, 42.1; HRMS (ESI) m/z calcd for $\text{C}_{23}\text{H}_{20}\text{ClN}$ $[\text{M} + \text{H}]^+$ 346.1357, found 346.1347.

N-benzyl-*N*-(2-chlorobenzyl)-3-phenylprop-2-yn-1-amine (**4{3}**). light yellow oil (155.7 mg, 90% yield); ^1H NMR (400 MHz, CDCl_3) δ 7.65 (d, $J = 7.2$ Hz, 1H), 7.57–7.55 (m, 2H), 7.50–7.48 (m, 2H), 7.42–7.36 (m, 6H), 7.32–7.28 (m, 2H), 7.26–7.24 (m, 1H), 3.96 (s, 2H), 3.87 (s, 2H), 3.54 (s, 2H); ^{13}C NMR (77 MHz, CDCl_3) δ 138.8, 136.5, 134.7, 131.8, 130.9, 130.0, 129.1, 128.3, 128.0, 127.2, 126.6, 123.4, 86.0, 84.4, 57.9, 54.9, 42.1; HRMS (ESI) m/z calcd for $\text{C}_{23}\text{H}_{20}\text{ClN}$ $[\text{M} + \text{H}]^+$ 346.1357, found 346.1350.

N-benzyl-*N*-(4-fluorobenzyl)-3-phenylprop-2-yn-1-amine (**4{4}**). light yellow oil (128.8 mg, 78% yield); ^1H NMR (400 MHz, CDCl_3) δ 7.61–7.58 (m, 2H), 7.53–7.41 (m, 9H), 7.37–7.33 (m, 1H) 7.13–7.10 (m, 2H), 3.84 (s, 2H), 3.81 (s, 2H), 3.55 (s, 2H); ^{13}C NMR (77 MHz, CDCl_3) δ 163.4, 161.0, 138.9, 134.7, 131.9, 130.7, 130.6, 129.1, 128.4, 128.1, 127.3, 123.5, 115.3, 115.1, 86.1, 84.3, 57.8, 57.0, 42.0; HRMS (ESI) m/z calcd for $\text{C}_{23}\text{H}_{20}\text{FN}$ $[\text{M} + \text{H}]^+$ 330.1653, found 330.1651.

N-benzyl-*N*-(4-bromobenzyl)-3-phenylprop-2-yn-1-amine (**4{5}**). light yellow oil (171.7mg, 88% yield); ^1H NMR (400 MHz, CDCl_3) δ 7.60–7.53 (m, 6H), 7.45–7.34 (m, 8H), 3.83 (s, 2H), 3.79 (s, 2H), 3.55 (s, 2H); ^{13}C NMR (77 MHz, CDCl_3) δ 138.8, 138.1, 131.9, 131.5, 130.8, 129.1, 128.5, 128.4, 128.2, 127.3, 123.4, 121.0, 86.1, 84.2, 57.8, 57.1, 42.1; HRMS (ESI) m/z calcd for $\text{C}_{23}\text{H}_{20}\text{BrN}$ $[\text{M} + \text{H}]^+$ 390.0852, found

390.0840.

4-((benzyl(3-phenylprop-2-yn-1-yl)amino)methyl)benzonitrile (**4{6}**). light yellow oil (127.8mg, 76% yield); ^1H NMR (400 MHz, CDCl_3) δ 7.67 (d, J = 8.4 Hz, 2H), 7.62–7.57 (m, 4H), 7.50 (d, J = 6.8Hz, 2H), 7.44–7.33 (m, 6H), 3.87 (s, 2H), 3.84 (s, 2H), 3.54 (s, 2H); ^{13}C NMR (77 MHz, CDCl_3) δ 144.9, 138.4, 132.3, 131.9, 129.6, 129.1, 128.5, 128.3, 127.5, 123.2, 119.1, 111.0, 86.3, 83.8, 58.0, 57.3, 42.4; HRMS (ESI) m/z calcd for $\text{C}_{24}\text{H}_{20}\text{N}_2$ $[\text{M} + \text{H}]^+$ 337.1699, found 337.1689.

N-benzyl-*N*-(2-methylbenzyl)-3-phenylprop-2-yn-1-amine (**4{7}**). light yellow oil (148.4 mg, 91% yield); ^1H NMR (400 MHz, CDCl_3) δ 7.45 (d, J = 7.2 Hz, 2H), 7.41 (d, J = 4.8 Hz, 3H), 7.39–7.26 (m, 6H), 7.28–7.21 (m, 3H), 3.8 (s, 4H), 3.5 (s, 2H), 2.5 (s, 3H); ^{13}C NMR (77 MHz, CDCl_3) δ 139.0, 137.8, 136.8, 131.8, 130.4, 130.2, 129.2, 128.4, 128.3, 128.0, 127.3, 127.1, 125.6, 123.6, 86.0, 84.6, 58.0, 56.0, 41.7, 19.3; HRMS (ESI) m/z calcd for $\text{C}_{24}\text{H}_{23}\text{N}$ $[\text{M} + \text{H}]^+$ 326.1903, found 326.1894.

N-benzyl-*N*-(3-methylbenzyl)-3-phenylprop-2-yn-1-amine (**4{8}**). light yellow oil (140.3 mg, 86% yield); ^1H NMR (400 MHz, CDCl_3) δ 7.61–7.50 (m, 4H), 7.45–7.29 (m, 9H), 7.18–7.16 (m, 1H), 3.86 (s, 2H), 3.83 (s, 2H), 3.58 (s, 2H), 2.46 (s, 3H); ^{13}C NMR (77 MHz, CDCl_3) δ 139.0, 138.9, 137.9, 131.9, 129.9, 129.2, 128.4, 128.3, 128.0, 127.9, 127.2, 126.2, 123.6, 86.0, 84.6, 57.9, 57.8, 42.2, 21.5; HRMS (ESI) m/z calcd for $\text{C}_{24}\text{H}_{23}\text{N}$ $[\text{M} + \text{H}]^+$ 326.1903, found 326.1896.

N-benzyl-*N*-(4-methylbenzyl)-3-phenylprop-2-yn-1-amine (**4{9}**). light yellow oil (106.0 mg, 65% yield); ^1H NMR (400 MHz, CDCl_3) δ 7.56–7.54 (m, 2H), 7.49 (d, J = 7.2 Hz, 2H), 7.40–7.37 (m, 7H), 7.32–7.28 (m, 1H), 7.20 (d, J = 7.6 Hz, 2H), 3.81 (s, 2H), 3.78 (s, 2H), 3.53 (s, 2H), 2.39 (s, 3H); ^{13}C NMR (77 MHz, CDCl_3) δ 139.0, 136.7, 135.8, 131.8, 129.1, 129.0, 128.3, 128.0, 127.12, 123.5, 86.0, 84.5, 57.7, 57.5, 42.0, 21.1; HRMS (ESI) m/z calcd for $\text{C}_{24}\text{H}_{23}\text{N}$ $[\text{M} + \text{H}]^+$ 326.1903, found 326.1893.

N-benzyl-*N*-(2-methoxybenzyl)-3-phenylprop-2-yn-1-amine (**4{10}**). light yellow oil (102.7 mg, 60% yield); ^1H NMR (400 MHz, CDCl_3) δ 7.58–7.50 (m, 5H), 7.40–7.36 (m, 5H), 7.32–7.27 (m, 2H), 7.01 (t, J = 7.4 Hz, 1H), 6.92 (d, J = 8.4 Hz, 1H), 3.87 (s, 3H), 3.86 (s, 4H), 3.59 (s, 2H); ^{13}C NMR (77 MHz, CDCl_3) δ 158.1, 139.2, 131.8, 130.5, 129.2, 128.3, 128.2, 128.1, 127.9, 127.2, 127.0, 123.7, 120.5, 110.7, 85.7, 85.2,

58.1, 55.5, 51.5, 42.5; HRMS (ESI) m/z calcd for $C_{24}H_{23}NO$ $[M + H]^+$ 342.1852, found 342.1837.

N-benzyl-*N*-(4-methoxybenzyl)-3-phenylprop-2-yn-1-amine (**4{11}**). light yellow oil (107.8 mg, 63% yield); 1H NMR (400 MHz, $CDCl_3$) δ 7.58–7.55 (m, 2H), 7.49 (d, J = 7.2 Hz, 2H), 7.42–7.38 (m, 7H), 7.34–7.28 (m, 1H), 6.94 (d, J = 8.4 Hz, 2H), 3.86 (s, 3H), 3.81 (s, 2H), 3.77 (s, 2H), 3.53 (s, 2H); ^{13}C NMR (77 MHz, $CDCl_3$) δ 158.9, 139.1, 131.8, 131.0, 130.3, 129.1, 128.3, 128.0, 127.1, 123.5, 113.8, 85.9, 84.6, 57.7, 57.1, 55.3, 41.9; MS (ESI) m/z : 342.2 $[M + H]^+$.

N-benzyl-*N*-(3,4-dimethoxybenzyl)-3-phenylprop-2-yn-1-amine (**4{12}**). light yellow oil (126.5 mg, 68% yield); 1H NMR (400 MHz, $CDCl_3$) δ 7.57–7.55 (m, 2H), 7.49 (d, J = 7.2 Hz, 2H), 7.41–7.28 (m, 5H), 7.07 (s, 1H), 7.02 (d, J = 8.4 Hz, 2H), 6.88 (d, J = 8.4 Hz, 1H), 3.95 (s, 3H), 3.92 (s, 3H), 3.80 (s, 2H), 3.77 (s, 2H), 3.55 (s, 2H); ^{13}C NMR (77 MHz, $CDCl_3$) δ 149.1, 148.3, 139.0, 131.8, 131.6, 129.1, 128.4, 128.0, 127.2, 123.5, 121.3, 112.4, 111.1, 85.9, 84.6, 57.7, 57.5, 56.0, 55.9, 42.1; HRMS (ESI) m/z calcd for $C_{25}H_{25}NO_2$ $[M + H]^+$ 372.1958, found 372.1942.

N-benzyl-*N*-(cyclohexylmethyl)-3-phenylprop-2-yn-1-amine (**4{13}**). light yellow oil (88.0 mg, 49% yield); 1H NMR (400 MHz, $CDCl_3$) δ 7.52–7.50 (m, 2H), 7.48–7.43 (m, 2H), 7.38–7.28 (m, 6H), 3.74 (s, 2H), 3.55 (s, 2H), 2.47 (d, J = 7.2 Hz, 2H), 1.93–1.90 (m, 2H), 1.78–1.70 (m, 4H), 1.65–1.48 (m, 4H), 1.40–1.29 (m, 2H); ^{13}C NMR (77 MHz, $CDCl_3$) δ 149.1, 148.3, 139.0, 131.8, 131.6, 129.1, 128.4, 128.0, 127.2, 123.5, 121.3, 112.4, 111.1, 85.9, 84.6, 57.7, 57.5, 56.0, 55.9, 42.1; HRMS (ESI) m/z calcd for $C_{23}H_{27}N$ $[M + H]^+$ 318.2216, found 318.2203.

N-benzyl-*N*-(3-methoxypropyl)-3-phenylprop-2-yn-1-amine (**4{14}**). light yellow oil (79.4 mg, 54% yield); 1H NMR (400 MHz, $CDCl_3$) δ 7.51–7.49 (m, 2H), 7.42 (d, J = 7.2 Hz, 2H), 7.38–7.34 (m, 5H), 7.31–7.28 (m, 1H), 3.74 (s, 2H), 3.57 (s, 2H), 3.51 (t, J = 6.4 Hz, 2H), 3.37 (s, 3H), 2.75 (t, J = 7.2 Hz, 2H), 1.91–1.85 (m, 2H); ^{13}C NMR (77 MHz, $CDCl_3$) δ 139.0, 131.8, 129.1, 128.3, 128.0, 127.1, 123.4, 85.6, 84.5, 71.0, 58.6, 58.1, 50.1, 42.3, 27.9; HRMS (ESI) m/z calcd for $C_{20}H_{23}NO$ $[M + H]^+$ 294.1852, found 294.1843.

N-benzyl-*N*-phenethyl-3-phenylprop-2-yn-1-amine (**4{15}**). light yellow oil (119.1 mg,

73% yield); ^1H NMR (400 MHz, CDCl_3) δ 7.56–7.55 (m, 2H), 7.46–7.28 (m, 13H), 3.85 (s, 2H), 3.69 (s, 2H), 2.98 (s, 4H); ^{13}C NMR (77 MHz, CDCl_3) δ 140.5, 138.8, 131.8, 129.2, 128.8, 128.5, 128.4, 128.0, 127.2, 126.0, 123.5, 85.7, 84.5, 58.2, 55.4, 42.5, 34.3; MS (ESI) m/z : 326.2 $[\text{M} + \text{H}]^+$.

N-benzyl-*N*-(3-chlorobenzyl)-3-phenylprop-2-yn-1-amine (**4{16}**). light yellow oil (102.1 mg, 59% yield); ^1H NMR (400 MHz, CDCl_3) δ 7.55–7.53 (m, 2H), 7.47 (d, J = 8.4 Hz, 3H), 7.40–7.28 (m, 9H), 3.80 (s, 2H), 3.77(s, 2H), 3.52(s, 2H); ^{13}C NMR (77 MHz, CDCl_3) δ 141.2, 138.6, 134.3, 131.8, 129.6, 129.1, 128.4, 128.3, 128.1, 127.4, 127.3, 127.2, 123.3, 86.1, 84.1, 57.8, 57.2, 42.1; HRMS (ESI) m/z calcd for $\text{C}_{23}\text{H}_{20}\text{ClN}$ $[\text{M} + \text{H}]^+$ 346.1357, found 346.1341.

N-benzyl-*N*-(3-fluorobenzyl)-3-phenylprop-2-yn-1-amine (**4{17}**). light yellow oil (112.3 mg, 68% yield); ^1H NMR (400 MHz, CDCl_3) δ 7.56–7.54 (m, 4H), 7.50–7.02 (m, 9H), 7.00–6.98 (m, 1H), 3.81 (s, 2H), 3.80 (s, 2H), 3.53(s, 2H); ^{13}C NMR (77 MHz, CDCl_3) δ 164.3, 161.9, 141.8, 141.7, 138.7, 131.8, 129.8, 129.7, 129.4, 129.1, 128.4, 128.3, 128.1, 127.3, 124.6, 124.5, 123.4, 115.8, 115.6, 114.2, 113.9, 86.1, 84.2, 57.8, 57.2, 42.1; HRMS (ESI) m/z calcd for $\text{C}_{23}\text{H}_{20}\text{FN}$ $[\text{M} + \text{H}]^+$ 330.1653, found 330.1645.

N-([1,1'-biphenyl]-4-ylmethyl)-*N*-benzyl-3-phenylprop-2-yn-1-amine (**4{18}**). light yellow oil (147.5 mg, 76% yield); ^1H NMR (400 MHz, CDCl_3) δ 7.70–7.50 (m, 13H), 7.46–7.34 (m, 6H), 3.90 (s, 2H), 3.89 (s, 2H), 3.61(s, 2H); ^{13}C NMR (77 MHz, CDCl_3) δ 164.3, 161.9, 141.8, 141.7, 138.7, 131.8, 129.8, 129.7, 129.4, 129.1, 128.4, 128.3, 128.1, 127.3, 124.6, 124.5, 123.4, 115.8, 115.6, 114.2, 113.9, 86.1, 84.2, 57.8, 57.2, 42.1; HRMS (ESI) m/z calcd for $\text{C}_{29}\text{H}_{26}\text{N}$ $[\text{M} + \text{H}]^+$ 388.2060, found 388.2054.

N-benzyl-*N*-(3-methoxybenzyl)-3-phenylprop-2-yn-1-amine (**4{19}**). light yellow oil (147.1 mg, 86% yield); ^1H NMR (400 MHz, CDCl_3) δ 7.59–7.57 (m, 2H), 7.53–7.51 (m, 2H), 7.43–7.39 (m, 5H), 7.35–7.28 (m, 2H), 7.11–7.10 (m, 2H), 6.88 (d, J = 7.2Hz, 1H), 3.88 (s, 3H), 3.84 (s, 2H), 3.83 (s, 2H), 3.57(s, 2H); ^{13}C NMR (77 MHz, CDCl_3) δ 159.8, 140.7, 139.0, 131.8, 129.3, 129.1, 128.4, 128.1, 127.2, 123.5, 121.5, 114.6, 112.7, 86.0, 84.5, 57.8, 57.7, 55.2, 42.2; HRMS (ESI) m/z calcd for $\text{C}_{24}\text{H}_{23}\text{NO}$ $[\text{M} + \text{H}]^+$ 342.1852, found 342.1835.

N-benzyl-*N*-(naphthalen-1-ylmethyl)-3-phenylprop-2-yn-1-amine (**4{20}**). light yellow oil (157.6 mg, 87% yield); ¹H NMR (400 MHz, CDCl₃) δ 8.51–8.50 (d, *J* = 7.2 Hz, 1H), 8.00–7.93 (m, 2H), 7.77–7.40 (m, 14H), 4.37 (s, 2H), 4.02 (s, 2H), 3.63 (s, 2H); ¹³C NMR (77 MHz, CDCl₃) δ 139.0, 134.5, 134.2, 132.9, 132.0, 129.5, 129.4, 128.6, 128.5, 128.4, 128.3, 128.2, 128.1, 128.0, 127.4, 125.9, 125.8, 125.3, 125.2, 123.7, 86.4, 84.7, 58.3, 56.4, 42.0; HRMS (ESI) *m/z* calcd for C₂₇H₂₃N [M + H]⁺ 362.1903, found 362.1888.

N-benzyl-*N*-(naphthalen-2-ylmethyl)-3-phenylprop-2-yn-1-amine (**4{21}**). light yellow oil (161.2 mg, 89% yield); ¹H NMR (400 MHz, CDCl₃) δ 7.99–7.94 (m, 4H), 7.76–7.55 (m, 7H), 7.50–7.38 (m, 6H), 4.06 (s, 2H), 3.95 (s, 2H), 3.66 (s, 2H); ¹³C NMR (77 MHz, CDCl₃) 139.0, 136.7, 133.6, 133.0, 131.9, 129.3, 128.5, 128.1, 127.9, 127.8, 127.5, 127.3, 126.0, 125.7, 123.6, 86.1, 84.6, 58.1, 57.9, 42.3; HRMS (ESI) *m/z* calcd for C₂₇H₂₃N [M + H]⁺ 362.1903, found 362.1886.

N-benzyl-3-phenyl-*N*-(thiophen-3-ylmethyl)prop-2-yn-1-amine (**4{22}**). light yellow oil (135.2 mg, 85% yield); ¹H NMR (400 MHz, CDCl₃) δ 7.62–7.54 (m, 4H), 7.47–7.34 (m, 8H), 7.28–7.27 (m, 1H), 3.91 (s, 2H), 3.86 (s, 2H), 3.61 (s, 2H); ¹³C NMR (77 MHz, CDCl₃) 139.9, 139.0, 131.9, 129.2, 128.6, 128.4, 128.1, 127.3, 125.6, 123.5, 123.0, 86.0, 84.6, 57.7, 52.7, 42.3; HRMS (ESI) *m/z* calcd for C₂₁H₁₉NS [M + H]⁺ 318.1311, found 318.1306.

N-benzyl-*N*-(4-fluorobenzyl)-3-phenylprop-2-yn-1-amine (**4{4}**[#]). light yellow oil (148.6 mg, 90% yield); ¹H NMR (400 MHz, CDCl₃) δ 7.58–7.55 (m, 2H), 7.50–7.28 (m, 10H), 7.10–7.06 (m, 2H), 3.82 (s, 2H), 3.78 (s, 2H), 3.53 (s, 2H); ¹³C NMR (77 MHz, CDCl₃) δ 163.4, 161.0, 138.9, 134.7, 131.9, 130.7, 130.6, 129.1, 128.4, 128.1, 127.3, 123.5, 115.3, 115.1, 86.1, 84.3, 57.8, 57.0, 42.0; HRMS (ESI) *m/z* calcd for C₂₃H₂₀FN [M + H]⁺ 330.1653, found 330.1640.

N-benzyl-*N*-(4-methylbenzyl)-3-phenylprop-2-yn-1-amine (**4{9}**[#]). light yellow oil (153.3 mg, 94% yield); ¹H NMR (400 MHz, CDCl₃) δ 7.60–7.51 (m, 4H), 7.42–7.32 (m, 8H), 7.23 (d, *J* = 7.6 Hz, 2H), 3.84 (s, 2H), 3.82 (s, 2H), 3.56 (s, 2H), 2.43 (s, 3H); ¹³C NMR (77 MHz, CDCl₃) δ 139.0, 136.7, 135.8, 131.8, 129.1, 129.0, 128.3, 128.0, 127.12, 123.5, 86.0, 84.5, 57.7, 57.5, 42.0, 21.1; HRMS (EI) *m/z* calcd for C₂₄H₂₃N

[M]⁺ 326.1903, found 326.1889.

N-benzyl-*N*-(2-methoxybenzyl)-3-phenylprop-2-yn-1-amine (**4{10}**[#]). light yellow oil (147.1 mg, 86% yield); ¹H NMR (400 MHz, CDCl₃) δ 7.60–7.52 (m, 5H), 7.42–7.29 (m, 7H), 7.05–7.01 (m, 1H), 6.94 (d, *J* = 8.4 Hz, 1H), 3.89 (s, 3H), 3.88 (s, 4H), 3.60 (s, 2H); ¹³C NMR (77 MHz, CDCl₃) δ 158.1, 139.2, 131.8, 130.5, 129.2, 128.3, 128.2, 128.1, 127.9, 127.2, 127.0, 123.7, 120.5, 110.7, 85.7, 85.2, 58.1, 55.5, 51.5, 42.5; HRMS (ESI) *m/z* calcd for C₂₄H₂₃NO [M + H]⁺ 342.1852, found 342.1842.

N-benzyl-*N*-(4-methoxybenzyl)-3-phenylprop-2-yn-1-amine (**4{11}**[#]). light yellow oil (160.9 mg, 94% yield); ¹H NMR (400 MHz, CDCl₃) δ 7.55–7.53 (m, 2H), 7.48–7.46 (m, 2H), 7.40–7.30 (m, 8H), 6.92 (d, *J* = 8.8 Hz, 2H), 3.84 (s, 3H), 3.83 (s, 2H), 3.78 (s, 2H), 3.50 (s, 2H); ¹³C NMR (77 MHz, CDCl₃) δ 158.9, 139.0, 131.8, 131.0, 130.3, 129.1, 128.3, 128.0, 127.1, 123.5, 113.8, 85.9, 84.6, 57.6, 57.1, 55.3, 41.9; MS (ESI) *m/z* 342.2 [M + H]⁺.

N,N-dibenzyl-3-(3-fluorophenyl)prop-2-yn-1-amine (**4{23}**). light yellow oil (125.5 mg, 76% yield); ¹H NMR (400 MHz, CDCl₃) δ 7.54–7.50 (m, 4H), 7.45–7.27 (m, 9H), 7.14–7.09 (m, 1H), 3.85 (s, 4H), 3.57 (s, 2H); ¹³C NMR (77 MHz, CDCl₃) δ 163.7, 161.3, 138.8, 130.0, 129.9, 129.2, 128.8, 128.4, 128.3, 127.8, 127.7, 127.3, 125.4, 125.3, 118.8, 118.6, 115.6, 115.4, 85.7, 84.8, 57.8, 42.0; HRMS (ESI) *m/z* calcd for C₂₃H₂₀FN [M + H]⁺ 330.1653, found 330.1638.

N,N-dibenzyl-3-(4-(trifluoromethyl)phenyl)prop-2-yn-1-amine (**4{24}**). light yellow oil (120.0 mg, 63% yield); ¹H NMR (400 MHz, CDCl₃) δ 7.67 (s, 4H), 7.50–7.47 (m, 4H), 7.42–7.31 (m, 6H), 3.83 (s, 4H), 3.56 (s, 2H); ¹³C NMR (77 MHz, CDCl₃) δ 138.7, 132.1, 129.1, 128.4, 127.3, 125.4, 125.3, 125.2, 87.4, 84.7, 57.9, 42.0; HRMS (ESI) *m/z* calcd for C₂₄H₂₀F₃N [M + H]⁺ 380.1621, found 380.1609.

N,N-dibenzyl-3-(4-methoxyphenyl)prop-2-yn-1-amine (**4{25}**). light yellow oil (147.1 mg, 86% yield); ¹H NMR (400 MHz, CDCl₃) δ 7.51 (d, *J* = 8.4 Hz, 4H), 7.42–7.31 (m, 4H), 6.93 (d, *J* = 8.8 Hz, 6H), 3.88 (s, 3H), 3.83 (s, 4H), 3.53 (s, 2H); ¹³C NMR (77 MHz, CDCl₃) δ 159.5, 139.0, 133.2, 129.2, 128.4, 127.2, 115.6, 114.0, 85.8, 82.8, 57.7, 55.4, 42.2; HRMS (ESI) *m/z* calcd for C₂₄H₂₃NO [M + H]⁺ 342.1852, found 342.1845.

3-([1,1'-biphenyl]-4-yl)-*N,N*-dibenzylprop-2-yn-1-amine (**4{26}**). light yellow oil (170.8 mg, 88% yield); ^1H NMR (400 MHz, CDCl_3) δ 7.74–7.69 (m, 6H), 7.61–7.54 (m, 6H), 7.50–7.45 (m, 5H), 7.42–7.39 (m, 2H), 3.93 (s, 4H), 3.65 (s, 2H); ^{13}C NMR (77 MHz, CDCl_3) δ 140.9, 140.5, 139.1, 132.4, 129.3, 129.0, 128.5, 128.4, 127.8, 127.3, 127.2, 122.5, 85.9, 85.3, 57.9, 42.3; MS (ESI) m/z 388.2 $[\text{M} + \text{H}]^+$.

N,N-dibenzyl-3-(4-(*tert*-butyl)phenyl)prop-2-yn-1-amine (**4{27}**). light yellow oil (151.0 mg, 82% yield); ^1H NMR (400 MHz, CDCl_3) δ 7.57–7.50 (m, 6H), 7.47–7.41 (m, 6H), 7.37–7.33 (m, 2H), 3.86 (s, 4H), 3.57 (s, 2H), 1.43 (s, 9H); ^{13}C NMR (77 MHz, CDCl_3) δ 151.3, 139.0, 131.6, 129.2, 128.4, 128.3, 127.2, 125.4, 125.3, 120.5, 86.1, 83.7, 57.8, 42.1, 34.8, 31.3; HRMS (ESI) m/z calcd for $\text{C}_{27}\text{H}_{29}\text{N}$ $[\text{M} + \text{H}]^+$ 368.2373, found 368.2360.

N,N-dibenzylhept-2-yn-1-amine (**4{28}**). light yellow oil (68.7 mg, 47% yield); ^1H NMR (400 MHz, CDCl_3) δ 7.43 (d, $J = 7.2\text{Hz}$, 4H), 7.35 (t, $J = 7.2\text{Hz}$, 4H), 7.30–7.26 (m, 2H), 3.27–3.26 (m, 4H), 2.33–2.30 (m, 2H), 1.64–1.30 (m, 2H), 1.01 (t, $J = 7.2\text{Hz}$, 2H); ^{13}C NMR (77 MHz, CDCl_3) δ 139.1, 129.1, 128.2, 127.0, 86.0, 74.3, 57.5, 41.7, 31.2, 22.0, 18.5, 13.7; MS (ESI) m/z : 292.2 $[\text{M} + \text{H}]^+$.

N,N-dibenzylhex-2-yn-1-amine (**4{29}**). light yellow oil (73.7 mg, 53% yield); ^1H NMR (400 MHz, CDCl_3) δ 7.46 (d, $J = 7.2\text{Hz}$, 4H), 7.37 (t, $J = 7.4\text{Hz}$, 4H), 7.31–7.28 (m, 2H), 3.73 (s, 4H), 3.30 (s, 2H), 2.34–2.30 (m, 2H), 1.67–1.64 (m, $J = 7.2\text{Hz}$, 2H), 1.12 (t, $J = 7.2\text{Hz}$, 3H); ^{13}C NMR (77 MHz, CDCl_3) δ 139.2, 129.1, 128.8, 128.2, 127.0, 85.7, 74.7, 57.6, 41.8, 22.6, 20.9, 13.6; HRMS (ESI) m/z calcd for $\text{C}_{20}\text{H}_{23}\text{N}$ $[\text{M} + \text{H}]^+$ 278.1903, found 278.1885.

N,N-dibenzylbut-2-yn-1-amine (**4{30}**). light yellow oil (38.8 mg, 31% yield); ^1H NMR (400 MHz, CDCl_3) δ 7.43 (d, $J = 7.2\text{Hz}$, 4H), 7.35 (t, $J = 7.4\text{Hz}$, 4H), 7.29–7.26 (m, 2H), 3.70 (s, 4H), 3.25 (s, 2H), 1.94 (s, 3H); ^{13}C NMR (77 MHz, CDCl_3) δ 139.1, 129.1, 128.4, 128.2, 127.0, 81.0, 73.8, 57.5, 41.8, 29.7; MS (ESI) m/z : 250.2 $[\text{M} + \text{H}]^+$.

N-benzyl-3-phenyl-*N*-(3-phenylprop-2-yn-1-yl)prop-2-yn-1-amine (**6**). light yellow oil (122.7 mg, 73% yield); ^1H NMR (400 MHz, CDCl_3) δ 7.52–7.48 (m, 6H), 7.41–7.31 (m, 9H), 3.89 (s, 2H), 3.74 (s, 4H); ^{13}C NMR (77 MHz, CDCl_3) δ 138.0, 131.8, 129.4,

128.4, 128.3, 128.1, 127.4, 123.2, 85.5, 84.7, 57.4, 43.1; MS (ESI) m/z : 336.2 $[M + H]^+$.

ACKNOWLEDGMENTS

The authors thank Professor Erkang Fan ~~at the University of Washington~~ for helpful discussions, and are grateful for financial support from Shanghai Municipal Education Commission (No.14ZZ159), Shanghai Municipal Science and Technology Commission (No. 12430501300), Shanghai University of Engineering Science (nhrc-2014-05), ~~and the Program for Special Appointment (Eastern Scholar) at Shanghai Institutions of Higher Education.~~

Supporting Information Available. NMR (1H and ^{13}C) spectra of all compounds.

This material is available free of charge via the Internet at <http://pubs.acs.org>.

REFERENCES

- (a) Merkul, E.; Boersch, C.; Frank, W.; Muller, T. J. *J. Org. Lett.* **2009**, *11*, 2269. (b) Yamamoto, Y.; Hayashi, H.; Saigoku, T.; Nishiyama, H. *J. Am. Chem. Soc.* **2005**, *127*, 10804. (c) Campbell, M. J.; Toste, F. D. *Chem. Sci.* **2011**, *2*, 1369. (d) Chen, J. S.; Properzi, R.; Uccello, D. P.; Young, J. A.; Dushin, R. G.; Starr, J. T. *Org. Lett.* **2014**, *16*, 4146. (e) Ermolat'ev, D. S.; Bariwal, J. B.; Steenackers, H. P. L.; De Keersmaecker, S. C. J.; Van der Eycken, E. V. *Angew. Chem. Int. Ed.* **2010**, *49*, 9465. (f) Bhonde, V. R.; Looper, R. E. *J. Am. Chem. Soc.* **2011**, *133*, 20172. (g) Robles-Machin, R.; Adrio, J.; Carretero, J. C.; *J. Org. Chem.* **2006**, *71*, 5023. (h) Yan, B.; Liu, Y. *Org. Lett.* **2007**, *9*, 4323. (i) Liu, J. X.; Zhang, Y. D.; Li, G. S.; Roschangar, F.; Farina, V.; Senanayake, C. H.; Lu, B. Z. *Adv. Synth. Catal.* **2010**, *352*, 2667.
- (a) Gainer, M. J.; Bennett, N. R.; Takahashi Y.; Looper R. E. *Angew. Chem. Int. Ed.* **2011**, *50*, 684. (b) Pereshivko, O. P.; Peshkov, V. A.; Peshkov, A. A.; Jacobs, J.; Van Meervelt L.; Van der Eycken, E. V. *Org. Biomol. Chem.* **2014**, *12*, 1741. (c) M. Bhandari, R.; Yousufuddin, M.; Lovely, C. J. *Org. Lett.* **2011**, *13*, 1382. (d) Ohta,

- Y.; Kubota, Y.; Watabe, T.; Chiba, H.; Oishi, S.; Fujii, N.; Ohno, H. *J. Org. Chem.* **2009**, *74*, 6299. (e) Xia, Y.Y.; Chen, L. Y.; Lv, S.; Sun, Z. H.; Wang, B. *J. Org. Chem.* **2014**, *79*, 9818.
3. (a) Nguyen, H. H.; Palazzo, T. A.; Kurth, M. J. *Org. Lett.* **2013**, *15*, 4492. (b) Donald, J. R.; Martin, S. F. *Org. Lett.* **2011**, *13*, 852. (c) Donald, J. R.; Wood, R. R.; Martin, S. F. *ACS Comb. Sci.* **2012**, *14*, 135. (d) Buysse, K.; Farard, J.; Nikolaou, A.; Van derheyden, P.; Vauquelin, G.; Pedersen, D. S.; Tourwe, D.; Ballet, S. *Org. Lett.* **2011**, *13*, 6468.
4. Mont, N.; Mehta, V. P.; Appukkuttan, P.; Beryozkina, T.; Toppet, S.; Van Hecke, K.; Van Meervelt, L.; Voet, A.; DeMaeyer, M.; Van der Eycken, E. V.; *J. Org. Chem.* **2008**, *73*, 7509.
5. Yu, P. H.; Bruce, A. D.; Boulton, A. A. *J. Med. Chem.* **1992**, *35*, 3705.
6. For a recent review of the A^3 -coupling reaction, see: (a) Peshkov, V. A.; Pereshivko, O. P.; Van der Eycken, E. V. *Chem. Soc. Rev.* **2012**, *41*, 3790. (b) Yoo, W. J.; Zhao, L.; Li, C. J. *Aldrichim. Acta* **2011**, *44*, 43. (c) Seidel, D. *Org. Chem. Front.* **2014**, *1*, 426.
7. For a recent review of the decarboxylation reaction (a) Rodriguez, N.; Gooßen, L. *J. Chem. Soc. Rev.* **2011**, *40*, 5030. (b) Weaver, J. D.; Recio, A. III.; Grenning, A. J.; Tung, J. A. *Chem. Rev.* **2011**, *111*, 1846. (c) Park, K.; Lee, S. *RSC Adv.* **2013**, *3*, 14165. (d) Dzik, W. I.; Lange, P. P.; Gooßen, L. J. *Chem. Sci.* **2012**, *3*, 2671.
8. (a) Myers, A. G.; Tanaka, D.; Mannion, M. R. *J. Am. Chem. Soc.* **2002**, *124*, 11250. (b) Tanaka, D.; Romeril, S. P.; Myers, A. G. *J. Am. Chem. Soc.* **2005**, *127*, 10323. (c) Gooßen, L. J.; Deng, G.; Levy, L. M. *Science* **2006**, *313*, 662. (d) Streuff, J.; White, D.; Virgil, S.; Stoltz, B. *Nat. Chem.* **2010**, *2*, 192. (e) Park, A.; Park, K.; Kim, Y.; Lee, S. *Org. Lett.* **2011**, *13*, 944. (f) Gooßen, L. J.; Zimmermann, B.; Knauber, T. *Angew. Chem. Int. Ed.* **2008**, *47*, 7103. (g) Liu, X.; Wang, Z.; Cheng, X.; Li, C. J. *J. Am. Chem. Soc.* **2012**, *134*, 14330. (h) Feng, Y. S.; Xu, Z. Q.; Mao, L.; Zhang, F. F.; Xu, H. J. *Org. Lett.* **2013**, *15*, 1472.
9. (a) Bi, H. P.; Zhao, L.; Liang, Y. M.; Li, C. J. *Angew. Chem, Int. Ed.* **2009**, *48*, 792. (b) Bi, H. P.; Teng, Q.; Guan, M.; Chen, W. W.; Liang, Y. M.; Yao, X.; Li, C. J. *J.*

- Org. Chem.* **2010**, *75*, 783. (c) Zhang, C.; Seidel, D. *J. Am. Chem. Soc.* **2010**, *132*, 1798.
10. Feng, H. D.; Ermolat'ev, D. S.; Song, G. H.; Van der Eycken, E. V. *J. Org. Chem.* **2011**, *76*, 7608.
11. Feng, H. D.; Ermolat'ev, D. S.; Song, G. H.; Van der Eycken, E. V. *Org. Lett.* **2012**, *14*, 1942.
12. (a) Feng, H. D.; Ermolat'ev, D. S.; Song, G. H.; Van der Eycken, E. V. *J. Org. Chem.* **2012**, *77*, 5149. (b) Ermolat'ev, D. S.; Feng, H. D.; Song, G. H.; Van der Eycken, E. V. *Eur. J. Org. Chem.* **2014**, 5346. (c) Lim, J.; Park, K.; Byeun, A.; Lee, S. *Tetrahedron Lett.* **2014**, *55*, 4875.
13. Park, K.; Heo, Y.; Lee, S. *Org. Lett.* **2013**, *15*, 3322.
14. Giles, R. L.; Sullivan, J. D.; Steiner, A. M.; Looper, R. E. *Angew. Chem, Int. Ed.* **2009**, *48*, 3116.
15. (a) Feng, H. D.; Ermolat'ev, D. S.; Song, G. H.; Van der Eycken, E. V. *Adv. Synth. Catal.* **2012**, *354*, 505. (b) Palani, T.; Park, K.; Kumar, M. R.; Jung, H. M.; Lee, S. *Eur. J. Org. Chem.* **2014**, 26.
16. (a) Petasis, N. A.; Akritopoulou, I. *Tetrahedron Lett.* **1993**, *34*, 583. (b) Petasis, N. A.; Zavialov, I. A. *J. Am. Chem. Soc.* **1997**, *119*, 445. (c) Morin, M. S. T.; Lu, Y.; Black, D. A.; Arndtsen, B. A. *J. Org. Chem.* **2012**, *77*, 2013. (d) Muncipinto, G.; Moquist, P. N.; Schreiber, S. L.; Schaus, S. E. *Angew. Chem. Int. Ed.* **2011**, *50*, 8172. (e) Frauenlob, R.; García, C.; Bradshaw, G. A.; Burke, H. M.; Bergin, E. *J. Org. Chem.* **2012**, *77*, 4445. (f) Wang, J. Y.; Shen, Q. Y.; Li, P. Z.; Peng, Y. Q.; Song, G. H. *Org. Biomol. Chem.* **2014**, *12*, 5597.
17. Candeias, N. R.; Montalbano, F.; Cal P. M. S. D.; Gois, P. M. P. *Chem. Rev.* **2010**, *110*, 6169.
18. Wang, J. Y.; Li, P. Z.; Shen, Q. Y.; Song, G. H. *Tetrahedron Lett.* **2014**, *55*, 3888
19. (a) Eschweiler, W. *Ber.* **1905**, *38*, 880. (b) Gibson, H. W. *Chem. Rev.* **1969**, *69*, 673.