

Gold Catalysis

International Edition: DOI: 10.1002/anie.201603938 German Edition: DOI: 10.1002/ange.201603938

Well-Defined Dinuclear Gold Complexes for Preorganization-Induced Selective Dual Gold Catalysis

Vincent Vreeken, Daniël L. J. Broere, Anne C. H. Jans, Marianne Lankelma, Joost N. H. Reek, Maxime A. Siegler, and Jarl Ivar van der Vlugt*

Abstract: The synthesis, reactivity, and potential of welldefined dinuclear gold complexes as precursors for dual gold catalysis are explored. Using the preorganizing abilities of the ditopic $PN^{H}P^{iPr}(L^{H})$ ligand, dinuclear $Au^{I}-Au^{I}$ complex 1 and mixed-valent Au¹-Au¹¹¹ complex 2 provide access to structurally characterized chlorido-bridged cationic species 3 and 4 upon halide abstraction. For 2, this transformation involves unprecedented two-electron oxidation of the redoxactive ligand, generating a highly rigidified environment for the Au_2 core. Facile reaction with phenylacetylene affords the σ,π -activated phenylacetylide complex 5. When applied in the dual gold heterocycloaddition of a urea-functionalized alkyne, well-defined precatalyst 3 provides high regioselectivities for the anti-Markovnikov product, even at low catalyst loadings, and outperforms common mononuclear Au¹ systems. This proof-of-concept demonstrates the benefit of preorganization of two gold centers to enforce selective non-classical σ , π -activation with bifunctional substrates.

Gold catalysis has flourished over the past 15 years, enabling a wide range of transformations.^[1] More recently, dual gold catalysis has been developed successfully.^[2] Whereas "traditional" mono-gold catalysis relies on π -activation of a substrate by a cationic Au^I center, dual gold catalysis typically involves both σ - and π -activation (of one or two functionalities, such as C=C bonds) by two gold centers. The prevailing strategy utilizes mononuclear Au^I complexes to induce this mode of activation (Scheme 1).^[3] However, this strategy offers no handles to induce preorganization of both gold centers to specifically target welldefined σ,π -activation of, for example, unsaturated hydrocarbon C-C multiple bonds whilst avoiding π - or $\sigma + \pi\text{-coordination}.$ Furthermore, no control over the selective binding of bifunctional substrates (such as heterocyclizations) can be achieved in this manner.

The synthesis, coordination chemistry and catalytic applications of dinuclear Au^I complexes are well-developed.^[4,5]



Scheme 1. Top: coordination of C=C bonds on mononuclear Au¹ species and a strategy to induce selective σ,π -activation using well-defined diphosphine-supported dinuclear Au¹–Au¹. Bottom: known (left) and novel (right) coinage metal coordination chemistry of **PN^HP^{iPr}** ligand L^H, including unprecedented ligand-based reactivity.

The proximity of both gold centers has occasionally been credited to enhance reactivity.^[6] However, to the best of our knowledge the competence of well-defined dinuclear σ -, π -alkynide complexes in dual gold catalysis has never been reported, despite the potential benefits of two preorganized gold centers with respect to chemoselectivity and activity for this type of reaction. Additionally, facile access to dinuclear mixed-valent gold complexes might be an interesting target in the context of cascade and Au^I/Au^{III} cooperative catalysis.^[7]

The ditopic tridentate ligand $\mathbf{PN^{H}P^{iPr}}$ (L^H; Scheme 1) displays versatile coordination chemistry with respect to a wide range of transition metals,^[8] including Cu^[9] and Ag.^[10] Strikingly, no single complex of gold with this type of ditopic framework is known to date. Furthermore, chemistry related to the redox-active nature^[11] of $\mathbf{L}^{\mathbf{H}}$ has been well-established in nickel, manganese, and rhenium complexes,^[12] but ligand redox-activity with Group 11 metals is limited to one example with Cu^I, leading to dimerization on one of the *para*-positions of the L backbone.^[9a] New avenues for gold coordination chemistry and catalysis may become accessible by developing strategies to preorganize and stabilize multiple gold centers on suitable ligand platforms and to selectively bind and activate bifunctional substrates. To address these challenges, we herein 1) uncover the versatile coordination chemistry of gold using the redox-active ditopic **PN^HP^{iPr}** ligand, 2) disclose the interesting ligand and metal-based reactivity of (mixedvalent) dinuclear species, and 3) establish well-defined precatalyst systems ideally suited for σ,π -activation of alkynes and demonstrate their use in highly selective dual gold catalysis.

© 2016 Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim

Wiley Online Library

^[*] V. Vreeken, Dr. D. L. J. Broere, A. C. H. Jans, M. Lankelma, Prof. Dr. J. N. H. Reek, Dr. J. I. van der Vlugt Homogeneous, Bioinspired and Supramolecular Catalysis, van 't Hoff Institute for Molecular Sciences, University of Amsterdam Science Park 904, 1098 XH Amsterdam (The Netherlands) E-mail: j.i.vandervlugt@uva.nl
Dr. M. A. Siegler Small Molecule X-ray Crystallography Facility, Department of Chemistry, John Hopkins University 3400 N. Charles St., Baltimore, MD 21218 (USA)
Supporting information for this article can be found under: http://dx.doi.org/10.1002/anie.201603938.

The straightforward reaction of **PN^HP^{iPr}** with AuCl(SMe₂) in a 1:2 ratio provides white solid **1** (³¹P NMR: δ 40.9 ppm) with a ¹H NMR spectrum suggestive of a C₂ symmetric species (Scheme 2). Single crystal X-ray diffraction analysis



Scheme 2. Synthesis of well-defined Au_2 -complexes 2–5 from Au^1-Au^1 precursor 1, generated by reaction of ligand L^H with $AuCl(SMe_2)$ (2 equiv).



Figure 1. Displacement ellipsoid plots (50% probability level) of 1 (left) and **2** (right). For clarity, the counterion and hydrogen atoms are not shown. Selected bond lengths (Å) and angles (°), for **1**: Au1– P1 2.2495(8), Au1–Cl1 2.3010(7), Au1–Au2 3.23791(17), P1-Au1-Cl1 173.45(3), C6-N1-Cl2 126.3(3). For **2**: Au1–P1 2.2655(21), Au1–N1 2.041(7), Au1–Cl1 2.354(2), Au1–Cl2 2.3039(22), Au2–P2 2.2401(23), Au1–Au2 4.641, P1-Au1-Cl1 178.39(8), P1-Au1-N1 84.51(19), N1-Au1-Cl2 172.66(20), C1-N1-C7 116.4(7), torsion \pm C6-C1-C7-C8 76.4(8).

intramolecular Au1-Au2 reveals an distance of 3.23791(17) Å, which suggests an aurophilic $d^{10}-d^{10}$ interaction^[13] in the solid state (Figure 1). Reaction of $\mathbf{1}$ with one equivalent of dichloro- λ^3 -(iodanyl)benzene (PhICl₂) in CH₂Cl₂ instantaneously generates a dark purple solution that shows two signals in the ³¹P NMR spectrum at δ 44.3 (P2) and 105.3 ppm (P1). The strongly downfield shifted signal for P1 supports coordination to a Au^{III} center by site-selective two-electron Au^I oxidation with formation of Au^I-Au^{III} derivative 2. Mass spectrometry supports an overall Au₂Cl₃(L) configuration. The ¹H NMR spectrum contains one strongly upfield shifted aromatic signal at δ 5.90 ppm but no -NH signal could be identified. X-ray structure determination confirms the sole formation of species 2, with the central deprotonated amido nitrogen coordinating to the square-planar Au^{III} center (Figure 1).

The **PNP**-ligand backbone in **2** is severely twisted, with a C6-C1-C7-C8 torsion angle of 76.4(8)°. This distortion relates to the upfield shifted aromatic signal in the ¹H NMR spectrum, which corresponds to the shielded C(6)-H hydrogen that is positioned on top of the second phenyl ring. The intramolecular Au–Au distance of 4.641 Å excludes an aurophilic d^8 – d^{10} interaction. Cyclic voltammetry does not indicate reversible oxidation of the potentially redox-active ligand scaffold in **2**.

A prerequisite for catalysis with a AuX(L) precursor (X is a halide) is the generation of a vacant coordination site by halide abstraction with, for example, Ag⁺ salts or a suitable Lewis acid. Addition of one equivalent of AgNTf₂ (or related silver salts or GaCl₃; see the Supporting Information) to 2, which bears multiple chlorido fragments, leads to a mixture of species according to ³¹P NMR spectroscopy. However, addition of two equivalents results in rapid decoloration of the reaction solution and generation of a single symmetric product (³¹P NMR: δ 40.4 ppm), suggestive of the presence of only Au^I-phosphine fragments. The ¹H NMR spectrum shows only one methyl signal for the ditolylamine backbone and chemically identical isopropyl groups at phosphorus. Most notably, only two instead of the anticipated three aromatic hydrogen signals are observed, together with a downfield signal at δ 10.53 ppm. This signal integrates for one hydrogen and is attributed to an -NH fragment. ESI-MS data suggest that the dinuclear complex, bearing only one chlorido ligand, remains intact during this transformation. The structure of complex 3 could be elucidated by X-ray structure determination of single crystals grown from CH2Cl2-pentane (Figure 2).

Clearly, halide abstraction by $AgNTf_2$, related silver salts, or $GaCl_3$, induces a highly unusual metal-based reduction of mixed-valent species **2** to yield Au^I - Au^I species **3**



Figure 2. Displacement ellipsoid plot (50% probability level) for the cationic part of **3**. The NTf₂ counterion, lattice solvent molecules, and hydrogen atoms are omitted for clarity. Selected bond lengths (Å) and angles (°): Au1–P1 2.2500(12), Au2–P2 2.2500(12), Au1–Cl1 2.3484(12), Au1–Au2 3.0758(3), C2–C8 1.449(7), P1-Au1-Cl1 178.53(4), Au1-Cl1-Au2 81.74(4).

www.angewandte.org

© 2016 Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim

(concomitant with oxidative C-C coupling of the two C-H groups ortho to the nitrogen in the ligand backbone), to generate a carbazole framework wherein the nitrogen is reprotonated. The gold centers are bridged by a single chlorido ligand, leading to an acute 4Au1-Cl1-Au2 of 81.74(4)°. Relative to "open" complex 1, the intramolecular Au-Au distance decreased by approximately 0.16 Å in this "closed" derivative, as a result of increased directional positioning of the phosphine lone pairs, despite the larger natural bite angle.^[14] To the best of our knowledge, this is the first report of redox-chemistry occurring at the ortho C-H positions of the PNP framework within the coordination sphere of any transition metal. The existence of a single chlorido bridgehead between two Au^I centers is relatively rare.^[15] Notably, no single example exists of an intramolecular Au-Cl-Au bridge stabilized by a single dinucleating ligand. Reaction of 1 (featuring the "open" ligand) with one equivalent of AgNTf₂ generates tetranuclear bis(chlorido)bridged complex 4 as a crystalline solid (XAu1-Cl1-Au2 96.78(4)°). This species is likely in equilibrium with the dinuclear monomer in solution (see the Supporting Information for details about complex 4).

Substitution of the chlorido bridge in these dinuclear gold species might release a masked Au^I-cation. Reaction of **3** with excess phenylacetylene leads to broadening of the ³¹P NMR signal and appearance of an additional singlet at δ 43.3 ppm, which converts to a single product 5 (with complete conversion) upon addition of one equivalent of AgNTf₂ or K_2CO_3 . No signal corresponding to the terminal CH of the alkyne was observed for this species by ¹H NMR spectroscopy, suggesting formation of a gold(acetylide) fragment. The overall symmetry of the complex appears to be retained during this transformation, which points to rapid exchange of the phenylacetylide between the two gold centers.^[16] Single crystal X-ray structure determination corroborates the dual interaction of the -C=CPh ligand with the Au^I-Au^I complex, that is, σ -coordination of the terminal phenylacetylide carbon C(15) to Au(1) and π -coordination of the triple bond system



Figure 3. Displacement ellipsoid plot (50% probability level) for the cationic part of 5. The NTf₂ counterion, lattice solvent molecules, and hydrogen atoms are omitted for clarity. Selected bond lengths (Å) and angles (°): Au1–P1 2.2899(10), Au2–P2 2.2676(11), Au1–C15 2.019(5), Au2–Ct(C15-C16) 2.201, C15–C16 1.227(6), Au1–Au2 3.1110(2), P1-Au1-C15 176.81(13), Au1-C15-C16 174.2(4), P2-Au2-Ct(C15-C16) 174.99, C15-C16-C17 170.0(5).

to Au(2) (Figure 3). This is the first crystallographically characterized intramolecular Au₂(σ,π -acetylide) complex with a diphosphine ligand, and serves to illustrate that preorganization of two gold centers may lead to selective substrate coordination. Complex **4** reacts in a similar fashion, leading to the well-defined species **5'** that was characterized by NMR spectroscopy and mass spectrometry (see the Supporting Information).

Having established that dinuclear complex **3** engages in well-defined σ,π -activation of C–C triple bond systems, the heterocyclization (intramolecular hydroamination) of ureafunctionalized alkyne **6** was selected as a proof-of-concept reaction to probe the catalytic competence of this welldefined Au₂ species. Markovnikov addition to generate a six-membered ring (**8**) involves π -activation by a single gold species, whereas dinuclear σ,π -activation results in anti-Markovnikov addition to give a five-membered ring (**7**; Scheme 3).^[17] Using 2.5 mol% of **3** in DMF at 60 °C for five



Scheme 3. Heterocyclization of 1-(ethynylphenyl)urea **(6)** into either *anti*-Markovnikov product **7** or Markovnikov addition product **8**.

hours leads to full conversion and high regioselectivity for the five-membered indole **7** (85%), in accordance with a selective σ,π -acetylide mechanism. External base (such as K₂CO₃) inhibits the reactivity, while addition of one equivalent of AgNTf₂ with respect to the catalyst provides slightly higher regioselectivity for **7** (90%); presumably this is due to faster generation of the σ,π -acetylide species. In the absence of additives (Ag⁺ or base), bis(chlorido)-bridged tetranuclear complex **4** displays a similar preference for formation of **7**. The high regioselectivity achieved with dinuclear catalysts **3** and **4** is attributed to the ligand-enforced proximity of both Au^I centers. Stoichiometric reaction of **3** and **6** shows sole formation of the σ,π -activated substrate by ESI-MS (Supporting Information, Figure S27).

Dilution studies were performed to investigate the effect of decreased catalyst loading on the level of regiocontrol for the conversion of 6 to 7 and 8. The findings clearly validate our hypothesis, as the high regioselectivity for 7 obtained with catalyst 3 is independent of the catalyst concentration (Figure 4). In contrast, dilution experiments with mononuclear AuCl(P'Bu₃)—reported as the best catalyst to form 7 by σ,π -activation^[17]—result in a sharp drop in selectivity.^[18] These results demonstrate the benefits of well-defined preorganization of two gold centers to enforce selective σ,π -activation and to mediate regioselective dual gold catalysis with functionalized alkynes, even at low catalyst loadings.

In summary, we have demonstrated that the redox-active $\mathbf{PN^{H}P^{iPr}}$ ligand is a suitable framework to preorganize two gold centers for selective σ,π -activation of functionalized alkynes. Halide abstraction from mixed-valent Au^I-Au^{III} complex 2 using AgNTf₂ results in highly unusual reactivity

Angew. Chem. Int. Ed. 2016, 55, 1-6

© 2016 Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim

www.angewandte.org



Figure 4. Comparison of regioselectivity toward indole 7 with dinuclear catalyst 3 vs. mononuclear benchmark catalyst $AuCl(P^tBu_3)$ under dilution conditions.

involving two-electron ligand oxidation, generating chlorido-bridged Au^I-Au^I species **3** bearing a highly rigidified carbazolyl backbone. Reaction with phenylacetylene enabled the first crystallographically characterized intramolecular dual gold σ,π -acetylide complex (5) supported by a single diphosphine ligand. The well-defined dinuclear Au^I complexes are excellent precatalysts for dual gold catalysis involving selective σ , π -activation, inducing high regioselectivity in the gold-catalyzed heterocyclization of urea 6, without the need to add base or silver salts. Dilution experiments show that dinuclear catalyst 3 retains high selectivity at decreased catalyst loadings, unlike mononuclear Au^I catalysts typically employed for this reaction. These results illustrate the benefits of preorganization of gold centers to invoke selective substrate activation in dual gold catalysis. The detailed mechanism behind the surprising conversion of 2 into 3 is currently under investigation. This ligand-to-gold two-electron transformation appears to involve several digold intermediates, including the dinuclear "monomer" of complex 4, as well as a unique Au^{III}-Au^{III} species supported by one PNP ligand.^[19]

Acknowledgements

This research was funded by the European Research Council (ERC) through Starting Grant 279097 (*EuReCat*) to J.I.v.d.V.; A.C.H.J. and J.N.H.R. thank the ERC for Advanced Grant 339786. We thank Dr. Marc Devillard for valuable tips and tricks and Ed Zuidinga for ESI-MS measurements.

Keywords: σ,π -activation \cdot bridging acetylide \cdot dinuclear gold complexes \cdot dual gold catalysis \cdot redox-active ligand

a) A. S. K. Hashmi, Acc. Chem. Res. 2014, 47, 864–876; b) M.
 Chiarucci, M. Bandini, Beilstein J. Org. Chem. 2013, 9, 2586–2614; c) B. Ranieri, I. Escofet, A. M. Echavarren, Org. Biomol.

Chem. **2015**, *13*, 7103–7118; d) R. Dorel, A. M. Echavarren, *Chem. Rev.* **2015**, *115*, 9028–9072; e) Y. Wei, M. Shi, *ACS Catal.* **2016**, *6*, 2515–2524; f) D. Pflästerer, A. S. K. Hashmi, *Chem. Soc. Rev.* **2016**, *45*, 1331–1367.

- [2] For selected examples, see ref. [1a] and: a) P. H.-Y. Cheong, P. Morganelli, M. R. Luzung, K. N. Houk, F. D. Toste, J. Am. Chem. Soc. 2008, 130, 4517-4526; b) A. Grirrane, H. Garcia, A. Corma, E. Álvarez, ACS Catal. 2011, 1, 1647-1653; c) A. S. K. Hashmi, I. Braun, P. Nösel, J. Schädlich, M. Wieteck, M. Rudolph, F. Rominger, Angew. Chem. Int. Ed. 2012, 51, 4456-4460; Angew. Chem. 2012, 124, 4532-4536; d) L. Ye, Y. Wang, D. H. Aue, L. Zhang, J. Am. Chem. Soc. 2012, 134, 31-34; e) Y. Oonishi, A. Gómez-Suárez, A. R. Martin, S. P. Nolan, Angew. Chem. Int. Ed. 2013, 52, 9767-9771; Angew. Chem. 2013, 125, 9949-9953; f) M. Højer Larsen, K. N. Houk, A. S. K. Hashmi, J. Am. Chem. Soc. 2015, 137, 10668-10676; g) Y. Jiang, Y. Wei, X. Y. Tang, M. Shi, Chem. Eur. J. 2015, 21, 7675-7681.
- [3] A. S. K. Hashmi, T. Lauterbach, P. Nösel, M. H. Vilhelmsen, M. Rudolph, F. Rominger, *Chem. Eur. J.* 2013, 19, 1058–1065.
- [4] For dppm as a dinucleating ligand, see: a) T. S. Teets, D. G. Nocera, J. Am. Chem. Soc. 2009, 131, 7411-7420; Amino-diphosphine: b) M. D. Levin, F. D. Toste, Angew. Chem. Int. Ed. 2014, 53, 6211-6215; Angew. Chem. 2014, 126, 6325-6329; Bis(carbene) ligands: c) M. Baron, C. Tubaro, M. Basato, A. Biffis, M. M. Natile, C. Graiff, Organometallics 2011, 30, 4607-4615; Bis(methylene)-phosphonium ligands: d) R. G. Raptis, L. C. Porter, R. J. Emrich, H. H. Murray, J. P. Fackler, Jr., Inorg. Chem. 1990, 29, 4408-4412; e) L. A. Méndez, J. Jiménez, E. Cerrada, F. Mohr, M. Laguna, J. Am. Chem. Soc. 2005, 127, 852-853.
- [5] For applications in asymmetric catalysis, see: a) W. Zi, F. D. Toste, *Chem. Soc. Rev.* 2016, 45, DOI: 10.1039/C5CS00929D;
 b) Y. M. Wang, A. D. Lackner, F. D. Toste, *Acc. Chem. Res.* 2014, 47, 889–901; c) R. A. Widenhoefer, *Chem. Eur. J.* 2008, 14, 5382–5391.
- [6] See ref. [4b] and: a) E. Tkatchouk, N. P. Mankad, D. Benitez, W. A. Goddard, F. D. Toste, J. Am. Chem. Soc. 2011, 133, 14293–14300; b) M.-A. Abadie, X. Trivelli, F. Medina, F. Capet, P. Roussel, F. Agbossou-Niedercorn, C. Michon, ChemCatChem 2014, 6, 2235–2239; c) J. M. Serrano-Becerra, A. F. G. Maier, S. González-Gallardo, E. Moos, C. Kaub, M. Gaffga, G. Niedner-Schatteburg, P. W. Roesky, F. Breher, J. Paradies, Eur. J. Org. Chem. 2014, 4515–4522; See also: R. Gramage-Doria, J. Hessels, S. H. A. M. Leenders, O. Tröppner, M. Dürr, I. Ivanović-Burmazović, J. N. H. Reek, Angew. Chem. Int. Ed. 2014, 53, 13380–13384; Angew. Chem. 2014, 126, 13598–13602.
- [7] a) N. Morita, A. Yasuda, M. Shibata, S. Ban, Y. Hashimoto, I. Okamoto, O. Tamura, *Org. Lett.* **2015**, *17*, 2668–2671; b) A. W. Sromek, M. Rubina, V. Gevorgyan, *J. Am. Chem. Soc.* **2005**, *127*, 10500–10501.
- [8] For recent examples, see: a) D. E. Herbert, A. D. Miller, O. V. Ozerov, *Chem. Eur. J.* 2012, *18*, 7696-7704; b) Y.-T. Hung, M.-T. Chen, M.-H. Huang, T.-Y. Kao, Y.-S. Liu, L.-C. Liang, *Inorg. Chem. Front.* 2014, *1*, 405-413; c) C. Yoo, S. Oh, J. Kim, Y. Lee, *Chem. Sci.* 2014, *5*, 3853-3858; d) C.-I. Lee, J. C. DeMott, C. J. Pell, A. Christopher, J. Zhou, N. Bhuvanesh, N. O. V. Ozerov, *Chem. Sci.* 2015, *6*, 6572-6582; e) C. Yoo, M. J. Ajitha, Y. Jung, Y. Lee, *Organometallics* 2015, *34*, 4305-4311; f) V. Vreeken, M. A. Siegler, B. de Bruin, J. N. H. Reek, M. Lutz, J. I. van der Vlugt, *Angew. Chem. Int. Ed.* 2015, *54*, 7055-7058; *Angew. Chem.* 2015, *127*, 7161-7165; g) T. Kurogi, P. J. Carroll, D. J. Mindiola, *J. Am. Chem. Soc.* 2016, *138*, 4306-4309.
- [9] a) S. B. Harkins, N. P. Mankad, A. J. M. Miller, R. K. Szilagyi, J. C. Peters, J. Am. Chem. Soc. 2008, 130, 3478–3485; b) S. B. Harkins, J. C. Peters, J. Am. Chem. Soc. 2005, 127, 2030–2031.

www.angewandte.org

© 2016 Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim

- [10] J. C. DeMott, F. Basuli, U. J. Kilgore, B. M. Foxman, J. C. Huffman, O. V. Ozerov, D. J. Mindiola, *Inorg. Chem.* 2007, 46, 6271–6276.
- [11] Reviews: a) D. L. J. Broere, R. Plessius, J. I. van der Vlugt, *Chem. Soc. Rev.* 2015, 44, 6886–6915; b) V. Lyaskovskyy, B. de Bruin, ACS Catal. 2012, 2, 270–279; c) K. G. Caulton, *Eur. J. Inorg. Chem.* 2012, 435–443.
- [12] a) D. Adhikari, S. Mossin, F. Basuli, J. C. Huffman, R. K. Szilagyi, K. Meyer, D. J. Mindiola, *J. Am. Chem. Soc.* 2008, 130, 3676-3682; b) A. T. Radosevich, J. G. Melnick, S. A. Stoian, D. Bacciu, C. H. Chen, B. M. Foxman, O. V. Ozerov, D. G. Nocera, *Inorg. Chem.* 2009, 48, 9214-9221.
- [13] H. Schmidbaur, A. Schier, Chem. Soc. Rev. 2008, 37, 1931-1951.
- [14] For the bite angle of the oxygen analogue of L^H and benzofuran derivative, see: M. Kranenburg, Y. E. M. van der Burgt, P. C. J. Kamer, P. W. N. M. van Leeuwen, K. Goubitz, J. Fraanje, *Organometallics* **1995**, *14*, 3081–3089.
- [15] a) A. Grirrane, E. Álvarez, H. García, A. Corma, Angew. Chem. Int. Ed. 2014, 53, 7253-7258; Angew. Chem. 2014, 126, 7381-7386; b) N. Phillips, T. Dodson, R. Tirfoin, J. I. Bates, S. Aldridge, Chem. Eur. J. 2014, 20, 16721-16731; c) Y. Zhu, C. S. Day, L. Zhang, K. J. Hauser, A. C. Jones, Chem. Eur. J. 2013, 19, 12264-12271; d) A. Homs, I. Escofet, A. M. Echavarren, Org. Lett. 2013, 15, 5782-5785; e) V. W. W. Yam, C.-L.

Chan, K.-K. Cheung, J. Chem. Soc. Dalton Trans. **1996**, 4019–4022; f) P. G. Jones, G. M. Sheldrick, R. Uson, A. Lapuna, Acta Crystallogr. Sect. B **1980**, 36, 1486; Dimeric structures: g) A. Hamel, N. W. Mitzel, H. Schmidbaur, J. Am. Chem. Soc. **2001**, 123, 5106–5107.

- [16] See ref. [2g] and: a) T. J. Brown, R. A. Widenhoefer, *Organometallics* 2011, *30*, 6003–6009; b) T. N. Hooper, M. Green, C. A. Russell, *Chem. Commun.* 2010, *46*, 2313–2315.
- [17] A. Gimeno, A. B. Cuenca, S. Suárez-Pantiga, C. Ramírez de Arellano, M. Medio-Simón, G. Asensio, *Chem. Eur. J.* 2014, 20, 683–688.
- [18] See also: D. Ye, J. Wang, X. Zhang, Y. Zhou, X. Ding, E. Feng, H. Sun, G. Liu, H. Jianga, H. Liu, *Green Chem.* 2009, 11, 1201–1208.
- [19] Because of the anticipated complexity associated with this multistep "cascade" process, which converts 2 into 3, a detailed mechanistic investigation of this transformation is deemed beyond the scope of this study and will be reported elsewhere.

Received: April 26, 2016 Revised: May 27, 2016 Published online:

© 2016 Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim

einheim www.angewandte.org These are not the final page numbers!





Communications

Gold Catalysis

V. Vreeken, D. L. J. Broere, A. C. H. Jans, M. Lankelma, J. N. H. Reek, M. A. Siegler, J. I. van der Vlugt* _____ IIII - IIII

Well-Defined Dinuclear Gold Complexes for Preorganization-Induced Selective Dual Gold Catalysis



Worth its weight in gold: Coordination of gold(I) to a rigid bis(*o*-tolylamino)diphosphine ligand allows for versatile chemistry, including the isolation of a mixed-valent Au^I–Au^{III} species. Halide abstraction induces ligand-to-gold two-electron transfer, producing a carbazolyl-diphosphine analogue with a bridging chlorido ligand. This preorganized Au^I–Au^{II} species enables σ,π -activation of alkynes and is an isolable precatalyst for highly selective heterocyclizations.

6 www.angewandte.org

C 2016 Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim

Angew. Chem. Int. Ed. 2016, 55, 1-6