




available at www.sciencedirect.com



journal homepage: www.elsevier.com/locate/steroids



Synthesis of steroidal lactone by *penicillium citreo-viride*

Hong-Min Liu^{a,*}, Heping Li^a, Lihong Shan^a, Jian Wu^b

^a New Drug Research & Development Center, Zhengzhou University, Zhengzhou 450052, China

^b Bioengineering Department Zhengzhou University, Zhengzhou 450052, China

ARTICLE INFO

Article history:

Received 29 April 2006

Received in revised form
20 June 2006

Accepted 21 June 2006

Published on line 12 September 2006

Keywords:

Biotransformation

Lactone

Penicillium citreo-viride

Incubation

Testolactone

ABSTRACT

The biotransformations of a series of steroids by the fungus *penicillium citreo-viride* A.C.C.C. 0402 have been investigated, and the conversion to the same product testolactone (1) was observed from progesterone (2), dehydroepiandrosterone (3), 4-androstene-3, 17-dione (4), 5-androstene-3, 17-diol (5) with the exception of pregnenolone (6) and 3 β -hydroxy-5, 16-pregnadien-20-one (7). The possible metabolic pathways of the biotransformations were also discussed in the paper and the fungus *penicillium citreo-viride* A.C.C.C. 0402 was isolated during screening stains from samples collected from Zhengzhou, Henan province of China.

© 2006 Elsevier Inc. All rights reserved.

1. Introduction

Steroidal lactones possess useful biological activities such as anticancer, antibacterial, anticarcinogen and antiandrogenic activities [1–4], and 16-oxa analogues of estrane series express antihypercholesterolemic activities [5]. Plants of genus *Withania* are known to exhibit a variety of pharmacological activities mainly due to the presence of withanolides, which are steroidal lactones. Testolactones are moderately specific first generation inhibitors of human aromatase activity and thereby may contribute to the prevention of hormone dependent tumors such as breast cancer [6], prostatic hyperplasia and prostate cancer [7]. Testolactones are used as therapeutic agents in disorders caused by imbalance between estrogen and androgen action, e.g. gynecomastia [8] or precocious puberty [9]. Aromatase inhibitors are also essential tools for studying the role of estrogens in adults, or during development [10].

Because of the important bioactivities of the testolactones, scientists were prompted to synthesize these steroidal lactones. Although chemical ways of synthesis of steroidal lactones such as Baeyer-Villiger oxidation are possible, the biotransformation is thought to be the more environmental method, especially nowadays, green chemistry is advocated so frequently. In fact a variety of fungi [11–17] have been identified which can stereospecifically degrade the 17 β -acetyl side chain of progesterone 2 in high yield to give the ring D testolactone 1 and the metabolic pathway was also discussed [18]. Recently testosterone (8) was transformed into 1 by *Penicillium notatum* has been reported [19]. With the aim of synthesizing steroidal lactones and doing some further work, we have endeavored to screen microorganisms capable of producing steroidal lactones, and a common strain of *penicillium citreo-virid* A.C.C.C. 0402 which could transform 2–5 into 1 was isolated during screening stains from samples collected from Zhengzhou, Henan province, while 6 and 7 have not any

* Corresponding author. Tel.: +86 371 67767200; fax: +86 371 67767200.

E-mail addresses: liuhm@public2.zz.ha.cn, liuhm@zzu.edu.cn (H.-M. Liu).

0039-128X/\$ – see front matter © 2006 Elsevier Inc. All rights reserved.

doi:10.1016/j.steroids.2006.06.005

changes. *Penicillium* genus have been reported to be used in the biotransformations using steroids as the substrates, but the attention was focused mainly on 15 α -hydroxylation catalyzed by *Penicillium raistrickii* [20,21] and 5 α -reductase of *Penicillium decumbens* [22,23] and *Penicillium crustosum* [24]. Here we first reported on the microbial transformation using the strain of *penicillium citreo-viride*.

2. Experimental

2.1. Instrumental methods

Sterilization was carried out in an HVE-50 Hirayama autoclave. Aseptic operation was carried out in ClassIIA/B3 Biological safety cabinet from Forma Scientific. Incubation was carried out on an HZQ-Q orbital shaker. Melting points (mp) were determined on a XT5 melting point apparatus, which was uncorrected. Infrared (IR) spectra was recorded using KBr discs on a Bruker Vector-22 spectrometer. Mass spectra (MS) was obtained on an Esquire3000 mass spectrometer by electrospray ionization (ESI). The ^1H and ^{13}C nuclear magnetic resonance (NMR) spectra were obtained using a Bruker Avance DPX-400 spectrometer at 400 and 100 MHz, respectively, with tetramethylsilane (TMS) as internal standard in DMSO- d_6 . Chemical shifts (δ) were given in parts per million (ppm) relative to TMS. Coupling constants (J) were given in hertz (Hz). Chromatography was performed with ether acetate/chloroform (1:1) and visualized by spraying the plates with 50% sulfuric acid solution and heating in an oven at 100 °C until the colors developed. Thin layer chromatography (TLC) was performed on 0.25 mm thick layer of silica gel G (Qingdao Marine Chemical Factory, China).

2.2. Microorganism

The strain of *penicillium citreo-viride* A.C.C.C. 0402 was isolated from the samples collected from the suburb of Zhengzhou, Henan province, PRC, and identified by Bioengineering Department Zhengzhou University. Stock cultures were maintained at 4 °C on agar slopes composed of peptone (1.2%), dextrose (3.0%), yeast extract (0.1%), KH_2PO_4 (0.13%), agar (2.0%) (pH 4.5).

2.3. Conditions of cultivation and transformation

Spores freshly obtained from peptone dextrose agar slopes were transferred aseptically into 10, 500 ml Erlenmeyer flasks containing 100 ml of sterile peptone dextrose broth in a biological safety cabinet and were incubated for 48 h at 27.5 °C. The cultures were shaken at 210 rpm on an orbital shaker. Then 1.0 g substrate dissolved in 20 ml acetone was evenly distributed among the flasks under sterile conditions and incubated for a further 5 days at the same conditions after which the metabolites were extracted from the broth.

2.4. Extraction and separation of metabolites

The mycelium was separated from the broth by filtration under vacuum, and rinsed in ultrasonic with ethyl acetate (100 ml \times 5) to ensure that all of the available products were

Table 1 – The biotransformations of the substrates of 2–7

Substrates	Yield of 1 (%)	Yield of 4 (%)
2	60.9	8.4
3	65.4	<0.5
4	73.6	–
5	32.5	45.2
6	–	–
7	–	–

(–) Means that no product was detected on TLC.

isolated from the mycelium. The mycelial broth was then extracted for five times with ethyl acetate. The organic extract altogether was concentrated into about 300 ml, which was then washed with saturated aqueous NaHCO_3 , brine and water, respectively, for three times, then dried with sodium sulfate and the solvent was evaporated under reduced pressure to give the mixed products. The mixture was separated on silica gel column chromatography by using ether acetate/chloroform (1:1) as eluant, and the solvent was collected in aliquots (10 ml) and analysed by thin layer chromatography to identify the separated metabolite fractions. The solvent systems used for running the TLC plates was also ether acetate/chloroform (1:1), and the TLC was visualized by spraying the plates with 50% sulfuric acid solution and heating in an oven at 100 for 3 min until the colors developed.

2.5. Incubation of 2–7

The biotransformations of the substrates of 2–7 were, respectively, shown in Table 1 according to Sections 2.3 and 2.4, and the results were shown in Fig. 1. All of the substrates of 2, 3, 4 and 5 could be converted into testolactone 1, mp 201.2–202.1 °C; IR(KBr): 3420.1, 2954.6, 1719.8, 1667.2 cm^{-1} ; ^1H NMR (CDCl_3 , 400 MHz) δ : 5.76 (s, H4), 1.18 (s, H18), 1.36 (s, H19) ppm; ^{13}C NMR (CDCl_3 , 400 MHz) δ : 38.9 (C1), 32.3(C2), 199.1(C3), 171.1(C4), 124.1(C5), 35.5 (C6), 33.8(C7), 37.9 (C8), 52.5 (C9), 38.4 (C10), 19.8 (C11), 30.4 (C12), 82.7 (C13), 45.7 (C14), 21.8 (C15), 28.5 (C16), 169.3 (C17); 17.4 (C18), 20.0 (C19); HRMS(ESI) m/z : $\text{C}_{19}\text{H}_{26}\text{O}_3[\text{M} + \text{H}^+]$, calcd. 303.1960, found 303.1957, $[\text{M} + \text{Na}^+]$, calcd. 325.1780, found 325.1771, $[\text{M} + \text{K}^+]$ calcd. 341.1579, found 343.1827.

Compound 4, mp 172.6–173.5 °C; IR(KBr): 2952.1, 2919.2, 1735.3, 1664.3, 1616.3, 1378.2 cm^{-1} ; ^1H NMR (CDCl_3 , 400 MHz) δ : 5.75 (s, H4), 0.92 (s, H18), 1.22 (s, H19) ppm; ^{13}C NMR (CDCl_3 , 400 MHz) δ : 35.5 (C1), 33.8 (C2), 199.1 (C3), 124.0 (C4), 170.2 (C5), 32.4 (C6), 31.2 (C7), 35.0 (C8), 53.7 (C9), 38.5 (C10), 20.2 (C11), 35.6 (C12), 47.4 (C13), 50.7 (C14), 21.6 (C15), 31.7 (C16), 220.2 (C17); 13.6 (C18), 17.6 (C19); HRMS(ESI) m/z : $\text{C}_{19}\text{H}_{26}\text{O}_2[\text{M} + \text{H}^+]$, calcd. 287.2006, found 287.2018.

3. Results and discussion

A series of different structural steroids were found to yield the same product 1 in the biotransformation by *penicillium citreo-virid* which has not been reported, so we have investigated the scope of the use of *penicillium citreo-virid* for steroid transformation and discussed the metabolic pathways.




Fig. 1 - The biotransformations of 2-5 by *penicillium citreo-virid*.




Fig. 2 - Metabolic pathways of the biotransformations of the substrates by *penicillium citreo-virid*.

The possible metabolic pathways of the biotransformations of the substrates in *penicillium citreo-virid* were shown in Fig. 2. Conversion of 4 by *penicillium citreo-virid* led to a single product 1 in a considerable yield, that was to say *penicillium citreo-virid* could add an oxygen atom into D-ring of the steroidal substrates; 3 and 5 could be transformed into 4 beside 1, that was to say *penicillium citreo-virid* could oxidate 3 β -hydroxy-5-ene derivatives to 4-en-3-one derivatives of the steroidal substrates, respectively, and 2 could be also changed into 4 and 1, that was to say *penicillium citreo-virid* could degrade the 17 β -acetyl side chain of the steroidal substrates, so we concluded that 4 is a necessary intermediate for the biotransformations, in that way the biotransformation of 4 could carry out quickly and get the highest yield compare to other substrates. The 3 was changed 4 first, then 4 was transformed into 1, in fact very small quantity of 4 was detected by TLC, so we could convince that the first step was taken place fastly. The 5 and 2 also produced 4 first, respectively, through different two steps, and their final product was the same product 1. But we found that there was no any changes in the transformation of 6 and 7, we concluded that the 3-hydroxy of 6 and 7 could not be oxidated into corresponding 3-carbonyl derivatives because of the hindrance of the 17-acetyl of the substrates which was different from the 3 and 5, while 3-carbonyl derivative was thought to be the key intermediate, so the biotransformation of 6 and 7 could not be finished. The 3-hydroxy of 3 and 5 could be easily oxidated into 3-carbonyl corresponding derivatives and then the next step could carry out quickly.

In conclusion, this work investigated the biotransformations of a series of steroids by the fungus *penicillium citreo-viride* A.C.C.C. 0402 and gave their metabolic pathways, respectively, and we believed it would prompt further investigation about the biotransformation and metabolic pathway of the steroidal lactones.

Acknowledgement

We are grateful to NNSF of PRC for financially supporting of this research.

REFERENCES

- [1] Fried J, Thoma RW. 1-Dehydrotestolactone. U.S. Patent 2744. 120: Chem Abstr 50:14812b, 1956.
- [2] Cella JA, Kagawa CM. Steroidal lactone. J Am Chem Soc 1957;79:4808-9.
- [3] Wolf ME, Zanati G, Shanmugasundrum G, Gupte S, Aadahl G, Thia III Steroids. Derivatives of 2-thia-Nor-5 α -androstan-17 β -ol, as probes of steroids-receptor interactions. J Med Chem 1970;13:521-35.
- [4] Seralini GE, Moslemi S. Aromatase inhibitors: past, present and future. Mol Cell Endocrinol 2001;178:117-31.
- [5] Baran JC. The synthesis, stereochemistry, and biology of 16-hetero and 17-oxa-D-homo steroids. J Med Chem 1967;10:1039-47.
- [6] Brodie AMH, Njar VCO. Aromatase inhibitors in advanced breast cancer: mechanism of action and clinical implications. J Steroid Biochem Mol Biol 1998;66:1-10.
- [7] Li S, Parish EJ. Design and action of steroidal aromatase inhibitors. JAOCS 1996;73:1435-51.
- [8] Braunstein GD. Aromatase and gynecomastia. Endocr-Relat Cancer 1999;6:315-24.
- [9] Feuillan P, Merke D, Leschek EW, Cutler GB. Use of aromatase inhibitors in precocious puberty. Endocr-Relat Cancer 1999;6:303-6.
- [10] Seralini GE, Moslemi S. Aromatase inhibitors: past, present and future. Mol Cell Endocrinol 2001;178:117-31.
- [11] Brannon DR, Martin J, Oehlschlager AC, Durham NN, Zalkow LH. Transformation of progesterone and related steroids by *Aspergillus tamarii*. J Org Chem 1965;30:760-2.
- [12] Fonken GS, Murray HC, Reineke LM. Pathway of progesteroneoxidation by *Cladosporium resinae*. J Am Chem Soc 1960;82:5507-8.
- [13] Peterson DH, Eppstein SH, Meister PD, Murray HC, Leigh HM, Weintraub A, et al. Microbiological transformations of steroids. IX. Degradation of C21 steroids to C19 ketones and totestolactone. J Am Chem Soc 1953;75:5768-9.
- [14] Fried J, Thoma RW, Klingsberg A. Oxydation of steroids by microorganisms. III Side chain degradation, ring D-cleavage and dehydrogenation in ring A. J Am Chem Soc 1953;75:5764-5.
- [15] Mihovilovic MD, Müller B, Stanley P. Monooxygenase-mediated Baeyer-Villiger oxidations. Eur J Org Chem 2002;22:3711-30.
- [16] El-Tayeb O, Knight SG, Sih CJ. Substrate specificity in steroid side-chain degradation by microorganisms. Biochim Biophys Acta 1964;93:411-7.
- [17] El-Tayeb O, Knight SG, Sih CJ. Steroid epoxide cleavage by *Cylindrocarpus radicola*. Biochim Biophys Acta 1964;93:402-10.
- [18] Christy Hunter A, Carragher Natasha E. Flexibility of the endogenous progesterone lactonisation pathway in *Aspergillus tamarii* KITA: transformation of a series of cortical steroid analogues. J Steroid Biochem Mol Biol 2003;87:301-8.
- [19] Bartmanska A, Dmochowska-Gładysz J, Huszeza E. Steroids' transformation in *Penicillium notatum* culture. Steroids 2005;70:193-8.
- [20] Irrgang S, Schlosser D, Fritsche W. Involvement of cytochrome P-450 in the 15-hydroxylation of 13-ethyl-gon-4-ene-3, 17-dione by *Penicillium raistrickii*. J Steroid Biochem Mol Biol 1997;60:339-46.
- [21] Irrgang S, Schlosser D, Schmauder H-P. The steroid 15-hydroxylase of *Penicillium raistrickii* I 477 is inducible. Biotechnol Lett 1992;14:33-8.
- [22] Holland HL, Nguyen DH, Pearson NM. Biotransformation of corticosteroids by *Penicillium decumbens* ATCC 10436. Steroids 1995;60:646-9.
- [23] Holland HL, Dore S, Xu W, Brown FM. Formation of 5-steroids by biotransformation involving the 5-reductase activity of *Penicilliumdecumbens*. Steroids 1994;59:642-7.
- [24] Flores E, Cabeza M, Quiroz A, Bratoeff E, Garcia G, Ramirez E. Effect of a novel steroid (PM-9) on the inhibition of 5-reductase present in *Penicillium crustosum* broths. Steroids 2003;68:71-5.