This article was downloaded by: [The Aga Khan University] On: 09 October 2014, At: 05:29 Publisher: Taylor & Francis Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information: http://www.tandfonline.com/loi/gpss20

Synthetic Utility of Enaminonitrile Moiety in Heterocyclic Synthesis: Synthesis of Some New Thienopyrimidines

H. M. F. Madkour $^{\rm a}$, A. A. E. Afify $^{\rm a}$, A. A. Abdalha $^{\rm a}$, G. A. Elsayed $^{\rm a}$ & M. S. Salem $^{\rm a}$

^a Synthetic Organic Chemistry Laboratory, Chemistry Department, Faculty of Science, Ain Shams University, Abbasiya, Cairo, Egypt Published online: 18 Feb 2009.

To cite this article: H. M. F. Madkour , A. A. E. Afify , A. A. Abdalha , G. A. Elsayed & M. S. Salem (2009) Synthetic Utility of Enaminonitrile Moiety in Heterocyclic Synthesis: Synthesis of Some New Thienopyrimidines, Phosphorus, Sulfur, and Silicon and the Related Elements, 184:3, 719-732, DOI: <u>10.1080/10426500802266407</u>

To link to this article: http://dx.doi.org/10.1080/10426500802266407

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with

primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <u>http://www.tandfonline.com/page/terms-and-conditions</u>



Synthetic Utility of Enaminonitrile Moiety in Heterocyclic Synthesis: Synthesis of Some New Thienopyrimidines

H. M. F. Madkour, A. A. E. Afify, A. A. Abdalha, G. A. Elsayed, and M. S. Salem

Synthetic Organic Chemistry Laboratory, Chemistry Department, Faculty of Science, Ain Shams University, Abbasiya, Cairo, Egypt

The hitherto unknown 3-amino-5-bromo-4, 6-dimethylthieno [2, 3-b] pyridine-2carbonitrile (4) was condensed with p-anisaldehyde affording the Schiff base (5). Acylation of the thienopyridine derivative (4) using freshly distilled acetic anhydride gave a mixture of mono and diacetyl derivatives (6) and (7). Condensation of (4) with triethylorthoformate yielded the ethoxymethyleneamino derivative (8), which was treated with hydrazine hydrate to give the hydrazide derivative (9), which in turn was converted to a triazolopyrimidine derivative (10) upon treatment with freshly distilled acetic anhydride. Thiation of (4) with carbon disulfide afforded the pyrimidine dithione derivative (11), which was alkylated with ethyl iodide to give the di-s-ethylpyrimidine derivative (12). On the other hand, treatment of (4) with formamide yielded the aminopyrimidine derivative (13), whereas its treatment by formic acid produced the thienopyrimidinone derivative (14). Chlorination of (14) with a mixture of phosphorus pentachloride and phosphorus oxychloride gave the chloropyrimidine derivative (15), which in turn afforded the hydrazide derivative (9) upon treatment with hydrazine hydrate. Hydrazinolysis of ethyl-3-amino-5-bromo-4,6-dimethylthieno[2,3-b]pyridine-2-carboxylate (17) gave the hydrazino derivative (18), which in turn was converted to 8-bromo-7,9-dimethyl-3formylaminopyrido[3',2':4,5]thieno[3,2-d]pyrimidin-4(3H)-one (19) and 8-bromo-3-diacetylamino-2,7,9-trimethylpyrido[3',2':4,5]thieno[3,2-d]pyrimidin-4(3H)-one (20) upon treatment with formic acid and freshly distilled acetic anhydride, respectively.

Keywords Antimicrobial activity; enaminonitrile; thienopyridine; thienopyrimidine

INTRODUCTION

In continuation of our present program in the utility of enaminonitrile and enaminoester moieties in the synthesis of different

Received 25 February 2008; accepted 11 June 2008.

Address correspondence to M. S. Salem, Synthetic Organic Chemistry Laboratory, Chemistry Department, Faculty of Science, Ain Shams University, Abbasiya, Cairo 11566, Egypt. E-mail: s_s_marwa@yahoo.com

heterocyclic systems,¹⁻⁴ and due to the pronounced biological and pharmacological activities of thienopyrimidine derivatives,⁵⁻⁷ such as anticancer,⁸ antiviral,⁹ antitumor,¹⁰ anti-inflammatory,¹¹ antimicrobial,¹² antifungal,¹³ antihistaminic,¹⁴ analgesic,¹⁵ antibiotic,^{16,17} antidepressant,^{18,19} antihypertensive,^{19,20} antiulcer,²¹ bactericidal,^{22,23} blood platelet aggregation inhibitory,^{24–26} fungicidal,^{27–29} herbicidal,³⁰ hypersensitivity inhibitory,³¹ insecticidal,^{30,32} immunomodulatory, and oncostatic.³³ The therapeutic importance of thienopyrimidines prompted us to construct several analogs by exotic combinations of groups and active moieties. The present work aimed to synthesize some new derivatives of this class of compounds utilizing 3-amino-5-bromo-4, 6-dimethylthieno[2, 3-b]pyridine-2-carbonitrile.

RESULTS AND DISCUSSION

The key starting material, 3-amino-5-bromo-4,6-dimethylthieno[2,3b]pyridine-2-carbonitrile (4), has been obtained in fair yield via the cyclization of 5-bromo-2-[(cyanomethyl)thio]-4,6-dimethylnicotinonitrile (3) with sodium ethoxide in ethanolic solution.³⁴ Pyridine derivative (3) was obtained by treatment the thione derivative (2) with chloroacetonitrile.³⁴ It was found that (2) was obtained upon thiation of (1)by phosphorous pentasulfide (Scheme 1). 3-Cyano-4,6-dimethyl-2-(1*H*)-pyridone was obtained according to Bardhan's method.³⁵ Bromination of the latter compound using N-bromo succinimide in carbon tetrachloride containing a few drops of hydrogen peroxide (30%) afforded 5-bromo-4,6-dimethyl-2-oxo-1,2-dihydropyridine-3-carbonitrile (1).

The structure of compound (1) was confirmed by its mp and mixed mp and by comparison of these with the literature value,³⁶ although it was prepared by an alternative method (Br_2 /glacial AcOH).

The structure of (4) was established by elemental analysis as well as spectral data (IR,¹H-NMR, MS) and chemically via condensation with p-anisaldehyde to give the corresponding Schiff base (5) (Scheme 2).

The effect of acetic anhydride on enaminonitrile has been investigated in many publications,^{37,38} and it has been found that the nature of the product depends mainly upon the time of the reaction. In our present work, acylation of the starting material (4) with acetic anhydride afforded a mixture of both mono and diacetyl derivatives (6) and (7), respectively (Scheme 2).

Alternatively, treatment of the thienopyridine derivative (4) with triethyl orthoformate³⁹ afforded 5-bromo-2-cyano-4, 6-dimethyl-3-ethoxy-methyleneaminothieno[2,3-b]pyridine (8), which upon



SCHEME 1

hydrazinolysis^{40,41} gave 8-bromo-4-hydrazino-7,9-dimethylpyrido[3', 2':4,5]thieno[3,2-*d*]pyrimidine (**9**), whose structure was confirmed by elemental analysis as well as spectral data. Furthermore, chemical proof is achieved via acetylation with acetic anhydride to yield the triazolopyrimidine derivative (**10**) in fair yield (Scheme 2).

Thiation of the thienopyridine derivative (**4**) using carbon disulfide yielded 8-bromo-7,9-dimethylpyrido[3',2':4,5]thieno[3,2-d]pyrimidine-2,4-(1*H*,3*H*)-dithione (**11**), which was alkylated by ethyl iodide in dry acetone to afford 8-bromo-2,4-di(ethylthio)-7,9-dimethylpyrido[3',2':4,5]thieno[3,2-d]pyrimidine (**12**) (Scheme 2).

Treatment of the thienopyridine derivative (4) with formamide^{7,42} gave the expected aminopyrimidine derivative (13) in quantitative yield (Scheme 2). Furthermore, the reaction of (4) with formic acid^{7,43} gave 8-bromo-7,9-dimethylpyrido[3',2':4,5]thieno[3,2-d]pyrimidin-4-(3H)-one (14), which underwent chlorination⁴⁴ using a mixture of phosphorus pentachloride and phosphorus oxychloride as a chlorinating reagent to give the chloropyrimidine derivative (15). The structure of (15) was elucidated via elemental analysis and also spectral data, and chemically through the reaction with hydrazine hydrate to afford 8-bromo-4-hydrazino-7,9-dimethylpyrido[3',2':4,5]thieno[3,2-d]pyrimidine (9) (Scheme 2).



(i) P-anisaldehyde /EtOH, Piperidine, 78°C(ii) $Ac_2O.138°C$; (iii) $CH(OEt)_3.146°C$; (iv) $N_2H_4.H_2O.EtOH.78°C$; (v) $Ac_2O.138°C$; (vi) CS_2 /dry pyridine ,114 °C; (vii) Etl /EtOH.78°C; (viii) HCONH₂, 210 °C; (ix) HCOOH.100°C; (x) PCI₅ / POCI₃, 105 °C.

SCHEME 2

On the other hand, alkylation of the thione derivative (2) using ethyl chloroacetate³⁴ afforded pyridine derivative (16), which in turn was cyclized³⁴ to give ethyl-3-amino-5-bromo-4,6-dimethylthieno[2,3-b]pyridine-2-carboxylate (17) whose structure was confirmed by elemental analysis as well as spectral data (Scheme 3).



SCHEME 3

Br

Furthermore, reaction of (17) with hydrazine hydrate⁴⁵⁻⁴⁷ yielded 3-amino-5-bromo-4,6-dimethylthieno[2,3-b]pyridine-2-carbohydrazide (18), which in turn was subjected to react with formic acid and freshly distilled acetic anhydride to give 8-bromo7,9-dimethyl-3formylaminopyrido[3',2':4,5]thieno[3,2-d]pyrimidin-4(3H)-one (19) and 8-bromo-3-diacetylamino-2,7,9-trimethylpyrido[3',2':4,5]thieno[3,2-d] pyrimidin-4-(3H)-one (20), respectively (Scheme 3).

EXPERIMENTAL (Tables I and II)

All melting points were measured on a Gallenkamp melting point apparatus and are uncorrected. The infrared spectra were recorded using potassium bromide disks on a Pye Unicam SP-3-300 infrared spectrophotometer. ¹H-NMR experiments were run at 300 MHz on a Varian

			Analysis (F	Required/I	Found)	
Compound No.	MP °C (color)	Yield % (solvent)	Mol. Formula (M. wt)	С	н	N
(2)	279–280	71	$C_8H_7BrN_2S$	39.52	2.90	11.52
	(yellow)	Т	(243.12)	39.20	2.62	11.21
(3)	146-148	89	$C_{10}H_8BrN_3S$	42.57	2.86	14.89
	(yellow)	\mathbf{P}_{a}	(282.16)	42.43	2.61	14.58
(4)	258-260	80	$C_{10}H_8BrN_3S$	42.57	2.86	14.89
	(yellow)	\mathbf{E}	(282.16)	42.36	2.54	14.60
(5)	190-192	56	C ₁₈ H ₁₄ BrN ₃ OS	54.01	3.53	10.50
	(yellow)	\mathbf{P}_{a}	(400.29)	53.89	3.30	10.28
(6)	274 - 276	50	$C_{12}H_{10}BrN_3OS$	44.46	3.11	12.96
	(white)	В	(324.20)	44.19	2.92	12.65
(7)	130 - 132	45	$C_{14}H_{12}BrN_3O_2S$	45.91	3.30	11.47
	(yellow)	\mathbf{P}_b	(366.23)	45.59	2.99	11.25
(8)	118 - 120	51	$C_{13}H_{12}BrN_3OS$	46.16	3.58	12.42
	(yellow)	\mathbf{P}_a	(338.22)	45.89	3.27	12.13
(9)	260 - 262	84	$C_{11}H_{10}BrN_5S$	40.75	3.11	21.60
	(white)	Α	(324.20)	40.47	2.87	21.25
(10)	238 - 240	45	$C_{13}H_{10}BrN_5S$	44.84	2.89	20.11
	(white)	\mathbf{P}_{a}	(348.22)	44.57	2.62	19.99
(11)	302 - 305	50	$C_{11}H_8BrN_3S_3$	36.87	2.25	11.73
	(orange)	P_y	(358.30)	36.55	1.97	11.45
(12)	120 - 122	60	$C_{15}H_{16}BrN_3S_3$	43.47	3.89	10.14
	(white)	\mathbf{P}_a	(414.41)	43.15	3.57	9.89
(13)	> 360	46	$C_{11}H_9BrN_4S$	42.73	2.93	18.12
	(white)	D	(309.18)	42.46	2.65	17.87
(14)	> 360	64	$C_{11}H_8BrN_3OS$	42.60	2.60	13.55
	(white)	Α	(310.17)	42.28	2.27	13.28
(15)	176 - 178	54	$C_{11}H_7BrClN_3S$	40.20	2.15	12.79
	(white)	\mathbf{P}_b	(328.62)	39.96	1.98	12.50
(16)	78 - 80	58	$\mathrm{C}_{12}\mathrm{H}_{13}\mathrm{BrN}_{2}\mathrm{O}_{2}\mathrm{S}$	43.78	3.98	8.51
	(yellow)	\mathbf{P}_{a}	(329.21)	43.50	3.70	8.22
(17)	202 - 204	66	$C_{12}H_{13}BrN_2O_2S$	43.78	3.98	8.51
	(yellow)	\mathbf{P}_b	(329.21)	43.50	3.72	8.26
(18)	240 - 242	63	$C_{10}H_{11}BrN_4OS$	38.11	3.52	17.78
	(yellow)	D	(315.18)	37.89	3.25	17.53
(19)	258 - 260	63	$\mathrm{C_{12}H_9BrN_4O_2S}$	40.81	2.57	15.86
	(pale yellow)	\mathbf{E}	(353.19)	40.49	2.30	15.62
(20)	218 - 220	55	$\mathrm{C_{16}H_{15}BrN_4O_3S}$	45.40	3.57	13.24
	(white)	\mathbf{E}	(423.28)	45.16	3.28	12.98

TABLE I Characterization Data of Prepared Compounds

A: Acetic acid,B: Benzene, D: Dioxane, E: Ethanol, P_a: Petroleum ether (60–80°C), P_b: Petroleum ether (80–100°C), P_y: Pyridine

Downloaded by [The Aga Khan University] at 05:29 09 October 2014

Compound			IR	CM^{-1}	
No.	¹ H-NMR (δ in ppm)	MS (m/z %)	VNH,OH	$\nu_{C\equiv N}$	$\nu_{\rm C=0}$
(2)	3.50 (s, 1H), 2.57 (s, 3H), 2.44 (s, 3H)	[M+2] $(+244, 57.1%), [M]$ $(+(242, 52.4%), 107 (A3.6%), 118 (A0.0%), 81 (A1.4%)$	3174	2221	I
(3)	4.05 (s, 2H), 2.77 (s, 3H), 2.58 (s, 3H)	$[M+2]^{+}(283, 690\%), [M]^{+}(281, 85.4\%), 256 (100.0\%), 255 (30.8\%), 131 (74.9\%), 256 (100.0\%), 255 (30.8\%), 131 (74.9\%), 256 (30.8\%$	I	2222	
(4)	$6.51 (s, NH_2), 2.84 (s, 3H), 2.51 (s, 3H)$	$\begin{array}{l} 81 \; (19.2\%) \\ [M+2]^{+}(283, 74.8\%), \; [M]^{+}(281, 62.2\%), \\ 202 \; (23.1\%), \; 134 \; (6.3\%), \; 131 \; (15.9\%), \; 70 \end{array}$	3333, 3238	2203	Ι
(5)	8.62 (s,1H), 7.95—7.92 (d, 2H), 7.06—7.03 (d, 2H), 3.92 (s, 3H), 2.91	$\begin{array}{c} (100.0\%) \\ [M+2]^{.+}(401, \ 75.1\%), \ [M]^{.+}(399, \ 100.0\%), \\ 398 \ (77.3\%), \ 370 \ (25.9\%), \ 292(33.5\%) \end{array}$	I	2204	Ι
(9)	(s, 3 H), 2.79 (s, 3 H) 7.38 (s, 1H), 2.82 (s, 3 H), 2.65 (s, 3 H), 2.39 (s, 3 H)	1	3223	2217	1679
(2)	2.84 (s, 3H), 2.59 (s, 3H), 2.43 (s, 6H)	$\begin{array}{l} [\mathrm{M}{+}2]^{+}(367,00.0\%),[\mathrm{M}]^{-1}(365,00.0\%),\\ 323(38.2\%),283(56.6\%),281(100.0\%),\\ 244(10.4\%)\end{array}$		2222	1723
(8)	8.03 (s, 1H), 4.49—4.47 (q, 2H), 2.84 (s, 3H) 2.81 (s, 3H) 2.81 (s, 3H) 1.47 (t, 3H)	$[M+2]^{+1}(339, 84.7\%), [M]^{+}(337, 75.5\%), 292.(79.9\%), 281.(95.8\%), 215.(13.6\%)$	Ι	2206	I
(6)	8.20 (s, 1H), 7.32 (s, 1H), 5.82 (s, 2H), 3.00 (S. 3H), 2.72 (s, 3H)	[M+2] + (325, 100.0%), [M] + (323, 65.8%), 308 (26.8%), 281 (29.2%), 186 (15.8%)	3305, 3115	I	
(10)	9.33 (s, 11H), 3.15 (s, 3H), 2.88 (s, 3H), 2.72 (s, 3H)	$[M+2]^{+}(349, 81.3\%), [M]^{+}(347, 100.0\%), 350 (14.3\%), 268 (.35.5\%)$		I	
(11)	4.33 (s, 1H), 2.83 (s, 3H), 2.62 (s, 3H), 2.02 (s, 1H)	$[M+2]^{+}(359, 00.0\%), [M]^{+}(357, 00.0\%), 325 (31.1\%), 291 (33.6\%), 210 (31.1\%)$	3412		

TABLE II ¹H-NMR, MS, and IR Data of Prepared Compounds

(Continued on next page)

r 2014
Octobe
60
tt 05:29
] a
rsity
Unive
Khan
Aga
[The
l by
adec
omlo;
Ă

TABLE II ¹H-NMR, MS, and IR Data of Prepared Compounds (Continued)

Compound				$\rm IR~CM^{-1}$	
No.	¹ H-NMR (δ in ppm)	MS (m/z %)	VNH,OH	V⊆N	$\nu_{C=0}$
(12)	3.46—3.44 (q, 2H), 3.38—3.27 (q, 2H),	$[M+2]^{+}(416, 22.0\%), [M]^{+}(414, 31.3\%),$	I		I
(13)	3.20 (s, 3H), 2.86 (s, 3H), 1.47 (t, 6H) 8 54 (s, 1H) 7 58 (s, 2H) 3 09 (s, 3H)	382 (56.9%), 324 (25.9%) [M+2] + (310, 96.3%) [M] + (308, 70, 1%)	3457	I	I
	2.74 (s, 3H)	293 (27.1%), 229 (30.8%), 200 (74.8%)	3292		
(14)	11.89 (s, 1H), 7.87 (s, 1H), 2.81 (s, 3H),	[M+2]· ⁺ (311, 100.0%), $[M]$ · ⁺ (309, 80.1%),	3446	Ι	1659
	2.64 (s, 3H)	281(38.3%),202(30.1%)			
(15)	9.11 (s, 1H), 3.23 (s, 3H), 2.90 (s, 3H)	$[M+2]^{+}(329, 100.0\%), [M]^{+}(327, 66.2\%),$		I	I
		248(45.1%),184(11.3%)			
(16)	4.25 - 4.22 (q, 2H), 3.96 (s, 2H), 2.65 (s, -4.22)	[M+2] + (330, 11.4%), [M] + (328, 12.1%),		2218	1735
	3H), 2.59 (s, $3H$), 1.28 (t, $3H$)	256(49.2%),255(100.0%),209(13.8%)			
(17)	6.17 (br, 2H), 4.39-4.32 (q, 2H), 2.92	$[M+2]^{+}(330, 76.7\%), [M]^{+}(328, 38.9\%),$	3452,		1668
	(s, 3H), 2.79 (s, 3H), 1.42 (t, 3H)	285(23.0%),282(100.0%),234(22.0%)	3338		
(18)	8.99 (br, 1H), 6.75 (br, 2H), 4.42 (br,	$[M+2]^{+}(316, 24.6\%), [M]^{+}(314, 21.9\%),$	3491,	I	1651
	2H), 2.71 (s, 3H), 2.65 (s, 3H)	285(100.0%),283(94.0%),255(25.3%)	3304,		
			3255,		
			3202		
(19)	8.46 (s, 1H), 8.14 (s, 1H), 5.04 (s, 1H),	$[M+2]^{+}(354, 53.7\%), [M]^{+}(352, 51.7\%),$	3210	I	1679
	3.14 (s, 3H), 2.87 (s, 3H)	297 (100.0%), 295 (89.6%), 269 (24.8%)			
(20)	3.14 (s, $3H$), 2.86 (s, $3H$), 2.52 (s, $3H$),	$[M+2]$ $^{+}(424, 20.2\%), [M]$ $^{+}(422, 30.5\%),$		I	1752, 1696
	2.52 (s, 6H)	380(46.2%),365(35.9%),340(100.0%)			

Mercury VX-300 NMR spectrometer using TMS as internal standard in deuterated chloroform or deuterated dimethylsulphoxide. Chemical shifts are quoted δ . The mass spectra were recorded on a Shimadzu GCMS-QP-1000EX mass spectrometers at 70 eV. Elemental analyses were carried out at the Microanalytical Center of Cairo University. All the newly synthesized compounds gave satisfactory elemental analyses.

5-Bromo-4,6-dimethyl-2-thioxo-1, 2-dihydropyridine-3-carbonitrile (2)

A mixture of (1) (10 mmol, 2.26 g) and phosphorus pentasulfide (5 mmol, 1.11 g) in dry toluene (25 mL) was heated at reflux for 4 h. The solid product that was separated out while hot was collected, dried, and recrystallized to give (2).

5-Bromo-2-[(cyanomethyl)thio]-4,6-dimethylnicotinonitrile (3)

A mixture of (2) (10 mmol, 2.42 g) and chloroacetonitrile (11 mmol, 0.75 mL) in ethanol (20 mL) in the presence of fused sodium acetate (15 mmol, 1.23 g) was heated at reflux for 1.5 h and left to cool. The solid product that formed was filtered and washed with water, dried, and recrystallized to afford (3).

3-Amino-5-bromo-4,6-dimethylthieno[2, 3-*b*] pyridine-2-carbonitrile (4)

To a hot solution of (3) in ethanol (20 mL), a few drops of sodium ethoxide solution (prepared by dissolving 0.5 g sodium metal in 20 mL ethanol) was added dropwise. The solution was heated at reflux for 15 min and allowed to cool. The solid that formed was collected, dried, and recrystallized to give (4).

5-Bromo-3-{[(4-methoxyphenyl)methylene]amino}-4, 6-dimethylthieno[2,3-*b*]pyridine-2-carbonitrile (5)

To a hot solution of (4) (1.7 mmol, 0.5 g) in ethanol (20 mL) containing a few drops of piperidine, p-anisaldehyde (1.7 mmol, 0.24 mL) was added, and the reaction mixture was heated under reflux for 2 h. The precipitated solid that separated on cooling was collected by filtration, dried, and recrystallized to furnish (5).

N-(5-Bromo-2-cyano-4,6-dimethylthieno[2,3-*b*]pyridin-3yl)acetamide (6) and *N*-Acetyl-*N*-(5-bromo-2-cyano-4, 6-dimethylthieno[2,3-*b*]pyridin-3-yl)acetamide (7)

A mixture of (4) (1.7 mmol, 0.5 g) and freshly distilled acetic anhydride (10 mL) was heated at reflux for 3 h. The solid that deposited after distilling the excess solvent was collected and recrystallized to afford (6) and (7).

N.B.: The diacetylamino derivative (7) was authentically prepared by acetylating the monoacetylamino derivative (6) using freshly distilled acetic anhydride.

5-Bromo-2-cyano-4,6-dimethyl-3ethoxymethyleneaminothieno[2,3-b]pyridine(8)

A mixture of (4) (1.7 mmol, 0.5 g) and triethyl orthoformate (10 mL) was heated at reflux for 2 h. An excess of triethyl orthoformate was removed by distillation under reduced pressure. The solid obtained was filtered, dried, and recrystallized to afford (8).

8-Bromo-4-hydrazino-7,9-dimethylpyrido[3',2':4,5] thieno[3,2-*d*]pyrimidine (9)

Method A

To a hot solution of (8) (1.48 mmol, 0.5 g) in ethanol (20 mL), hydrazine hydrate (0.5 mL) was added, and the reaction mixture was heated at reflux for 2 h. The solid product that was separated out while hot was collected, dried, and recrystallized to afford (9).

Method B

To a hot solution of (15) (1.5 mmol, 0.5 g) in ethanol (20 mL), hydrazine hydrate (0.5 mL) was added dropwise while the solution was heated at reflux for 2 h. The solid product that separated out while hot was collected, washed with ethanol, dried, and recrystallized to afford (9).

4-Bromo-3,5,9-trimethylpyrido[3',2':4,5]thieno[3,2-*d*] 1,3,4-triazolo[1,2-*f*]pyrimidine (10)

A mixture of (9) (0.92 mmol, 0.3 g) and freshly distilled acetic anhydride (10 mL) was heated at reflux for 5 h. The solid that deposited after distilling off the excess solvent was collected and recrystallized to afford (10).

8-Bromo-7,9-dimethylpyrido[3',2':4,5]thieno [3,2-*d*]pyrimidine-2,4(1*H*,3*H*)-dithione (11)

A mixture of (4) (1.7mmol, 0.5 g) and carbon disulfide (10 mL) in dry pyridine (20 mL) was heated at reflux for 6 h. Upon cooling, the precipitated solid was collected, dried, and recrystallized to afford (11).

8-Bromo-2,4-di(ethylthio)-7,9-dimethylpyrido[3',2':4,5] thieno[3,2-*d*]pyrimidine (12)

A hot solution of (11) (1.54 mmol, 0.55 g) in dry acetone (20 mL) and ethyl iodide (1.54 mmol, 0.47 mL) was heated at reflux for 10 h. The solid product that separated out while hot was collected, dried, and recrystallized to afford (12).

4-Amino-8-bromo-7,9-dimethylpyrido[3',2':4,5]thieno[3,2d]pyrimidine(13)

A mixture of (4) (1.7 mmol, 0.5 g) and formamide (10 mL) was heated at reflux for 4 h. The precipitated solid while hot was filtered, washed with water, dried, and recrystallized to give (13).

8-Bromo-7,9-dimethylpyrido[3',2':4,5]thieno[3,2-*d*] pyrimidin-4(3*H*)-one (14)

A mixture of (4) (1.7 mmol, 0.5 g) and formic acid (10 mL) was heated at reflux for 1.5 h. The solid product that was separated out while hot was collected, washed with water, dried, and recrystallized to afford (14).

8-Bromo-4-chloro-7,9-dimethylpyrido[3',2':4,5] thieno[3,2-*d*]pyrimidine (15)

A mixture of (14) (0.16 mmol, 0.5 g) and phosphorus pentachloride (0.16 mmol, 0.5 g) in phosphorus oxychloride (10 mL) was heated in a water bath at 100°C for 10 h. The reaction mixture was poured into water, and the precipitated solid was filtered, washed with water, dried, and recrystallized to afford (15).

Ethyl[(5-bromo-3-cyano-4,6-dimethylpyridin-2-yl)thio] acetate (16)

A mixture of (2) (10 mmol, 2.42 g) and ethyl chloroacetate (11 mmol, 0.75 mL) in ethanol (20 mL) in the presence of fused sodium acetate (15 mmol, 1.23 g) was heated at reflux for 1.5 h and left to cool. The solid product that deposited was filtered, washed with water, dried, and recrystallized to afford (16).

Ethyl 3-amino-5-bromo-4,6-dimethylthieno[2,3-*b*] pyridine-2-carboxylate (17)

To a hot solution of (16) in ethanol (20 mL), a few drops of sodium ethoxide solution (prepared by dissolving 0.5 g sodium metal in 20 mL ethanol) was added .The solution was heated at reflux for 15 min and allowed to cool. The solid product that formed was collected, dried, and recrystallized to give (17).

3-Amino-5-bromo-4,6-dimethylthieno[2,3-*b*] pyridine-2-carbohydrazide (18)

Method A

A mixture of (17) (15 mmol, 4.92 g) and hydrazine hydrate (15 mL) was heated at reflux for 2 h. The separated solid was filtered, triturated with ethanol, dried, and recrystallized to afford (18).

Method B

A mixture of (16) (15 mmol, 4.92 g) and hydrazine hydrate (15 mL) was heated at reflux in a water bath at 100° C for 2 h. The precipitated solid product was washed several times with ethanol and recrystallized to afford (18).

8-Bromo-7,9-dimethyl-3-formylaminopyrido[3',2':4,5] thieno[3,2-*d*]pyrimidin-4(3*H*)-one (19)

A mixture of (18) (1.59 mmol, 0.5 g) and formic acid (20 mL) was heated at reflux for 4 h. The cooled reaction mixture was diluted with water, and the precipitated solid was recrystallized to afford (19).

8-Bromo-3-diacetylamino2,7,9-trimethylpyrido[3',2':4,5] thieno[3,2-*d*]pyrimidin-4(3*H*)-one (20)

A mixture of (18) (1.59 mmol, 0.5 g) and freshly distilled acetic anhydride (20 mL) was heated at reflux for 3 h. The solid that precipitated upon cooling was recrystallized to afford (20).

REFERENCES

- H. M. F. Madkour, A. A. E. Afify, G. A. El-Sayed, and M. S. Salem, Bull. Chem. Commun., 40(2), 147 (2008).
- [2] H. M. F. Madkour, M. R. Mahmoud, A. M. Sakr, and M. M. Habashy, Sci. Pharm., 69, 33 (2001).
- [3] M. A. I. Salem, H. M. F. Madkour, E. A. Soliman, and N. F. H. Mahmoud, *Heterocycles*, 53(5), 1124 (2000).
- [4] M. E. Azab, M. A. E. Ibraheem, and H. M. F. Madkour, Phosphorus, Sulfur, and Silicon, 181, 1299 (2006).
- [5] M. H. Bhuiyan, M. Rahman, K. Hossain, A. Rahim, M. I. Hossain, and M. Abu Naser, Acta Pharm., 56, 441 (2006).
- [6] A. H. Bedair, N. A. El-Hady, M. S. AbdEl-Latif, A. H. Fakery, and A. M. El-Agrody, *Il Farmaco*, 55, 708 (2000).
- [7] M. M. Ghorab, S. M. Abdel-Gawad, and M. S. A. El-Gaby, Il Farmaco, 55, 249 (2000).
- [8] M. S. A. E.-A. El-Gaby, S. G. Abdel-Hamide, M. M. Ghorab, and S. M. El-Sayed, Acta Pharm., 49, 149 (1999).
- [9] M. N. Nasr and M. M. Gineinah, Arch. Pharm., 335, 289 (2002).
- [10] P. G. Baraldi, M. G. Pavani, M. Nunez, P. Brigidi, B. Vitali, R. Gambari, and R. Romagnoli, *Bioorg. Med. Chem.*, 10, 449 (2002).
- [11] S. M. Sondhi, M. Johar, S. Rajvanshi, S. G. Dastidar, R. Shukla, R. Raghubir, and J. W. Lown, Australian J. Chem., 54, 69 (2001).
- [12] A. Z. M. S. Chowdhury, M. M. Matin, and M. N. Anwar, *Chittagong Univ. Stud. Part II: Sci.*, **21**, 79 (1997); *Chem. Abstr.*, **130**, 237530p (1999).
- [13] G. Mangalagiu, M. Ungureanu, G. Grosu, I. Mangalagiu, and M. Petrovanu, Ann. Pharm. Fr., 59, 139 (2001).
- [14] C. J. Shishoo, V. S. Shirsath, I. S. Rathod, M. J. Patil, and S. S. Bhargava, Arzneim. Forsch, 51, 221 (2001).
- [15] O. Bruno, C. Brullo, S. Schenone, A. Ranise, F. Bondavalli, E. Barocelli, M. Tognolini, F. Magnanini, and V. Bollabeni, *Il Farmaco*, 57, 753 (2002).
- [16] G. E. Robinson and C. J. Squire, Eur. Pat., 253, 507 (1988); Chem. Abstr., 109, 6318 (1988).
- [17] V. I. Shvedov, V. M. Aryuzina, M. D. Mashkovskii, A. J. Polezhaeva, and L. F. Roshchina, USSR Pat. 745, 160 (1983); Chem. Abstr., 99, 82509 (1983).
- [18] K. Ninomiya, K. Nitsuta, A. Tobe, M. Egawa, and R. Kikumoto, Jpn. Kokai, 16, 557 (1994); Chem. Abstr., 120, 290120 (1994).
- [19] J. B. Press and R. K. Russell, U.S. Pat., 835, 157 (1989); Chem. Abstr., 111, 174121 (1989).
- [20] S. E. Delaszlo, A. A. Patchett, E. E. Allen, and W. J. Greenle, *Eur. Pat.*, **502**, 725 (1992); *Chem. Abstr.*, **118**, 22249 (1993).
- [21] F. E. Janssens, I. G. Torremans, J. F. Hens, and T. T. J. M. Offenwert, Eur. Pat., 144, 101 (1985); Chem. Abstr., 104, 68856 (1986).

- [22] R. Bradbury, F. H. Jung, J. J. Lohmann, P. R. Marsham, and G. Pasquet, Aust. Pat., 584, 898 (1989); Chem. Abstr., 112, 20851 (1990).
- [23] K. Fujii, T. Tanaka, and Y. Fukuda, Eur. Pat., 370, 704 (1990); Chem. Abstr., 113, 212005 (1990).
- [24] R. Boehm, G. Hanbold, R. Pech, and G. Laban, Ger. Pat., 226, 893 (1985); Chem. Abstr., 104, 186437 (1986).
- [25] Daiichi Seiyaku Co. Ltd., Jpn. Kokai, 77687 (1982); Chem. Abstr., 97, 163007 (1982).
- [26] H. Fukumi, F. Saitoh, H. Hovikoshi, and S. Kobayashi, Eur. Pat., 82, 023 (1983); Chem. Abstr., 99, 158453 (1983).
- [27] S. El-Bahaie, A. El-Deeb, and M. G. Assy, Pharmazie, 46, 26 (1991).
- [28] D. K. K. Showa, Jpn. Kokai, 53, 681 (1981); Chem. Abstr., 95, 115522 (1981).
- [29] D. K. K. Showa, Jpn. Kokai, 56, 8389 (1981); Chem. Abstr., 95, 81010 (1981).
- [30] M. Wiesenfeldt, K. Etzbach, P. Hofmeister, C. Kueyast, and K. O. Westpalen, Eur. Pat., 447, 891 (1991); Chem. Abstr., 115, 256224 (1991).
- [31] A. A. Larsen and D. A. Owens, Ger. Pat., 231, 103 (1983); Chem. Abstr., 98, 204410 (1983).
- [32] T. Obata, K. Fujii, and I. Narita, Eur. Pat., 356, 158 (1990); Chem. Abstr., 113, 59210 (1990).
- [33] M. Arita, Y. Hukumasu, M. Sano, Y. Hoshino, and H. Komatsu, PCT Int. 08, 113 (1989); Chem. Abstr., 112, 98533 (1990).
- [34] A. Krauze, Z. Bomika, A. M. Shestopalov, L. A. Rodinovskaya, J. Pelcers, G. Duburs, Yu, A. Sharanin, and V. K. Promonenkov, *Khim. Geterotsikl. Soedin.*, **3**, 377 (1981); *Chem. Abstr.*, **95**, 80662 (1981).
- [35] J. C. Bardhan, J. Chem. Soc., 2223 (1929).
- [36] W. J. Thompson and J. Gaudino, J. Org. Chem., 49(26), 5237 (1984).
- [37] T. I. El-Emary, J. Chin. Chem. Soc., 46, 585 (1999).
- [38] M. M. Khafagy, A. H. F. Abd-Elwahab, F. A. Eid, and A. M. El-Agrody, *Il Farmaco*, 57, 715 (2002).
- [39] C. G. Dave and R. D. Shah, J. Heterocycl. Chem., 37, 757 (2000).
- [40] A. Sh. Oganisyan, A. S. Noravyan, and M. Zh. Grigoryan, Chem. Heterocycl. Compd., 40(1), 75 (2004).
- [41] S. G. Pilosyan, Authors Abstract, Dissertations of candidates in chemical sciences (in Russian), Yerevan (1986).
- [42] A. M. M. El-Saghier, Molecules, 7, 756 (2002).
- [43] A.M. Abdel-fattah, S.M. Sherif, A.M. El-Reedy, and S.A. Gad-Alla, *Phosphorus*, Sulfur, and Silicon, 70, 67 (1992).
- [44] C.G. Dave and R.D. Shah, J. Heterocycl. Chem., 35, 1295 (1998).
- [45] P. K. Bridson, R. A. Davis and I. S. Renner, J. Heterocycl. Chem., 22, 753 (1985).
- [46] N. P. Peet and S. Sunder, J. Heterocycl. Chem., 21, 1807 (1984).
- [47] I. Lalerzari and C. A. Stein, J. Heterocycl. Chem., 21, 5 (1984).