

Communication

Enantioselective Si–H Insertion Reactions of Diarylcarbenes for the Synthesis of Silicon-Stereogenic Silanes

Jake R. Jagannathan, James C. Fettinger, Jared T. Shaw, and Annaliese K. Franz

J. Am. Chem. Soc., Just Accepted Manuscript • DOI: 10.1021/jacs.0c04533 • Publication Date (Web): 16 Jun 2020

Downloaded from pubs.acs.org on June 16, 2020

Just Accepted

"Just Accepted" manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides "Just Accepted" as a service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. "Just Accepted" manuscripts appear in full in PDF format accompanied by an HTML abstract. "Just Accepted" manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are citable by the Digital Object Identifier (DOI®). "Just Accepted" is an optional service offered to authors. Therefore, the "Just Accepted" Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the "Just Accepted" Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these "Just Accepted" manuscripts.

is published by the American Chemical Society. 1155 Sixteenth Street N.W., Washington, DC 20036

Published by American Chemical Society. Copyright © American Chemical Society. However, no copyright claim is made to original U.S. Government works, or works produced by employees of any Commonwealth realm Crown government in the course of their duties.

7

8 9 10

11 12

13

14 15

16

17

18

19

20

21

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58 59

60

Enantioselective Si–H Insertion Reactions of Diarylcarbenes for the Synthesis of Silicon-Stereogenic Silanes

Jake R. Jagannathan, James C. Fettinger, Jared T. Shaw*, and Annaliese K. Franz*

Department of Chemistry, University of California, One Shields Avenue, Davis, California 95616, United States Supporting Information Placeholder

ABSTRACT: We report the first example of enantioselective, intermolecular diarylcarbene insertion into Si–H bonds for synthesis of silicon-stereogenic silanes. Dirhodium(II) carboxylates catalyze an Si–H insertion using carbenes derived from diazo compounds where selective formation of an enantioenriched silicon center is achieved using prochiral silanes. Fourteen prochiral silanes were evaluated with symmetrical and prochiral diazo reactants to produce a total of 25 novel silanes. Adding an ortho substituent on one phenyl ring of a prochiral diazo enhances enantioselectivity up to 95:5 er with yields up to 98 %. Using *in situ* IR spectroscopy, the impact of the off-cycle azine formation is supported based on the structural dependence for relative rates of diazo decomposition. A catalytic cycle is proposed with Si–H insertion as the rate-determining step, supported by kinetic isotope experiments. Transformations of an enantioenriched silane derived from this method, including selective synthesis of a novel sila-indane, are demonstrated.

The potential utility of chiral-at-silicon compounds incorporated into more complex structures has not been fully understood due to a shortage of synthetic methods. Silicon-stereogenic molecules are rare in number and diversity of structures as compared to carbon. Selected examples to generate siliconstereogenic silanes include dehydrocouplings,^{1–3} arylation,^{4,5} hydrosilylation,^{6–9} Si–C activation,^{10,11} and reactions controlled by chiral auxillaries.^{12–14} Brief explorations of the effect of silicon chirality on reaction outcome to produce more complex molecules have occurred,^{15–17} yet remain limited.

The catalytic insertion of carbenes into Si–H bonds to generate organosilicon compounds has been intermittently explored since Doyle's original work in 1988.^{18,19} Methods to date have focused on generation of stereogenic carbon centers using donor/acceptor carbenes (Figure 1A).^{20–23} Si–H insertion to generate stereogenic silicon centers has been demonstrated by Katsuki²⁴ and Iwasa²⁵ using donor/acceptor carbenes (Figure 1A). Diarylcarbenes are commonly referred to as donor/donor carbenes because they are typically less reactive, with few reports of intermolecular Si–H insertion, and one report of an enantioselective variant using functionalized alkynes as precursors (Figure 1B). ^{26–29}

Donor/donor carbenes have recently emerged as useful substrates for highly selective C–H insertion reactions.^{30–34} Rhodium carbene complexes demonstrate sufficient reactivity at the insertion carbon despite the presence of two aryl rings for potential stabilization.^{35,36} The Franz group has a long-standing interest in organosilicon chemistry and expertise synthesizing prochiral dihydridosilanes with variation of steric and electronic factors.^{37–39} We envisioned that the additional aryl ring could accomplish an enantioselective intermolecular Si-H insertion process with prochiral silanes (Figure 1C). Herein, we communicate the first enantioselective diarylcarbene Si–H insertion to produce silicon-stereogenic organosilanes. A. Enantioselective Si-H insertion of donor/acceptor carbenes:



Figure 1. Insertion of carbenes into Si-H bonds.

We began our studies screening metal catalysts [Ru(II), Ir(I), Fe(II), Rh(II) and Cu(II)] with diphenyldiazomethane (**2a**) and prochiral methylphenylsilane (**1a**). Inverse addition of **2a** using a syringe pump (over 1 hour) increased yield of **4a** by preventing azine formation, as seen in previous studies with Si–H insertion methodologies.^{21,40,41} Insertion product **4a** was only observed using dirhodium tetraacetate (Table 1, entry 1).⁴² Based on this lead result, we proceeded to screen chiral dirhodium(II)-based catalysts to identify an enantioselective variant.

A screen of well-studied chiral dirhodium compounds highlighted the reactivity of dirhodium tetracarboxylates (Table 1). Carboxylate ligands afforded higher yields compared to amidocontaining ligands due to the increased electrophilicity of the metal center and resulting carbene (entry 2 vs. entries 4-9).⁴³ Of the catalysts studied, $Rh_2(S$ -TCPTTL)₄ provided the highest levels of enantioselectivity when compared to others (entries 5-8 vs 9), which improved further using toluene (entry 10). When an insertion was tested using prochiral diazo carbene **3a**, the enantioselectivity of silane product **5a** increased *from 82:18 to 93:7 er*, with a notable increase in yield (76% to 91% yield, entry 10 vs. 14).

Table 1. Optimization of donor/donor Si-H insertion

1

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57 58 59

60



^a NMR yield using Ph-TMS as an internal standard. ^b Determined using ¹H NMR Spectroscopy. ^c Determined using CSP-HPLC analysis of silanol obtained from Pd/C hydrolysis; major diastereomer if relevant. ^d Toluene used as a solvent. ^e Diazo added via syringe over five minutes (without syringe pump).

Under optimized conditions, slow addition of **3a** over 5 minutes without a syringe pump forms **5a** in comparable yield and selectivity (entry 14 vs. 15). Reducing the reaction temperature below 23 °C did not increase selectivity and no insertion was observed below -30 °C. With optimized conditions in hand, we investigated the effect of substituents with both symmetrical and prochiral diazo compounds.

A series of sterically and electronically varied silanes and symmetrical diazo compounds were evaluated to study the effects on enantioselectivity (Scheme 1, 4a-o). Electron-rich diazo 2b was less reactive than 2a and provides lower yield for 4b (45%, 81:19 er). Yield improved using diazo 2c (91%) and lower enantioselectivity was observed for silane 4c (76:24 er). Electron-withdrawing groups do not strongly affect selectivity (4d, 80:20 er) while electron-donating groups on the silane proved deleterious to enantioselectivity (4e and 4f, 50:50 er and 74:26 er respectively). Additional steric bulk on the aryl ring of the silane generally eroded enantioselectivity (4g-i) but maintained fair to good yields (55-69%). Selectivity similar to 4a (82:18) was also observed using 2-naphthyl silane 1g, with the yield also higher compared to 4h (69 vs 60%). A slight recovery of enantioselectivity was also observed with 4k (52%, 82:18 er) compared to 4h (79:21 er), and comparable to 4a. Studies with varied alkyl substitution on the silicon center were conducted with diazo 2a.

Scheme 1. Scope of enantioselective Si–H insertion with symmetrical diazo compounds^{*a*}



^{*a*} isolated yields; er determined using CSP-HPLC analysis of silanol obtained from Pd/C hydrolysis. ^{*b*} Reaction performed using 1 mmol of **2a**.

Isobutyl-containing **4l** provided the highest enantioselectivity observed using **2a** (86:14 er). However, neopentyl substitution led to loss of enantioselectivity (**4m**, 50:50 er), and cyclohexyl substitution reduced enantioselectivity as well (**4n**, 70:30 er). Lastly, switching to a siloxane also deleteriously affected enantioselectivity while maintaining fair yield (**4o**, 60%, 61:39 er). We next turned our focus to insertion of prochiral diazo reactants. 1

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35 36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57 58 59

60

The ability of the ortho substituent on one phenyl ring of the diazo compounds to control enantioselectivity was explored (Scheme 2). Electron-donating substituents lower diastereose-lectivity (**5b**, 90:10 dr vs 93:7 dr), but slightly improve enantioselectivity (95:5 vs 93:7 er). With an electron-withdrawing group (**5c**), excellent yield and enantioselectivity is observed (93%, 93:7 er) and diastereoselectivity increased (98:2 vs 93:7 dr). Recent work has noted potential synergistic effects of electronics and ortho substitution on the selectivity of donor/donor **Scheme 2.** Scope of enantioselective Si–H insertion with prochiral diazo reagents^{*a*}

inverse addition CHa N₂ 1 mol % Si Rh₂(S-TCPTTL) H Ή ·Δı PhMe (0.05 M), 4Å MS 1a-c: e-g 3a-3e 23 °C, 1.5 h 5a-5i 5 equiv Ph Ph Ph Ph ,H _Η H₃C-H₃C-Si -Śi H₃C--Si H₂C= Si Ph' Ph' Ph' Ar H₃CO H₃C F₃C CI Ar = PMP 5a, 89%^b 5b, 91% 5c, 93% 5d, 98% 98:2 dr 93:7 dr 90:10 dr 90:10 dr 93:7 er 95:5 er 93:7 er 89:11 er H₃CO Ph .Н Si H₃C= н н H₃C Ph Ph Ph Ph H₃C H₃C H₃C 5e. 78% **5g**, 90% 5f. 85% 5h 95% 85:15 dr 94:6 dr 91:9 dr 91:9 dr 93:7 er 61:39 er 94:6 er 93:7 er H₃C= S H₂C= Si Ph Ph H₃C H₂C 5i^d, 55% **5j**, 66% (±) 5i hydrolysis 84:16 dr 82:18 dr product 92:8 ei 93.7 er

^{*a*} Isolated yields; dr determined using ¹H NMR spectroscopy; er Determined using CSP-HPLC analysis of silanol obtained from Pd/C hydrolysis. ^{*b*} Reaction performed using 1.00 g of **3a** and 0.05 mol % catalyst, at 0.1 M in toluene. ^{*c*} dr was determined using ¹⁹F NMR spectroscopy. ^{*d*} Relative configuration assigned by X-ray analysis.

carbene chemistry.⁴⁴ Substitution on both phenyl rings was able to achieve excellent yield and good selectivity in **5d** (98% yield, 90:10 dr . 89:11 er), although slightly lower compared to other substitution patterns. The combined steric and push-pull electronic effects improve enantioselectivity compared to symmetrical diazo compounds. These substrates demonstrate that the presence of ortho-substitution iso-steric to a methyl may induce enantioselectivity. Replacing phenyl with a 1-naphthyl group led to decreased diastereoselectivity (**5e**, 85:15 dr) and low enantioselectivity (61:39 er), suggesting other competitive steric effects are present. We sought to explore varied substitution of silanes with prochiral diazo 3a, given the increase in yield and enantioselectivity compared to using 2a. Prochiral silanes were tested with diazo 3a and all demonstrated above 90:10 er for the major diastereomer (Scheme 2, 5f-5j). Additionally, the reaction performed with 1 gram of 3a using <1 mol% catalyst affords excellent yield, diastereoselectivity and enantioselectivity is substrate controlled, while enantioselectivity is controlled by the rhodium catalyst. Notably, using a diastereoselective reaction with silane 1c promotes enantioselectivity with 5f (94:6 er) compared to 4e (50:50 er). This result highlights the benefit of using prochiral 3a to improve enantioselectivity.

A catalytic cycle for the enantioselective Si–H insertion of diarylcarbenes is proposed (Figure 2A).³⁶ The Rh(II) carboxylate catalyst (I) reacts with the diazo compound (**2a** or **3a**) to form complex II, which is approached by prochiral silane **1a** to produce the silicon-stereogenic silane and regenerate catalyst. Kinetic isotope experiments support the rate-determining insertion step ($k_H/k_D = 1.6$), fitting closely with previous experiments



Figure 2. A. Proposed catalytic cycle with kinetic isotope effect; B. Diagram of proposed selectivity rationale of donor/donor insertions.

of Si–H insertion with donor-acceptor^{20,41,45} and donor/donor carbenes.²⁹ Off-cycle formation of azine (6 or 7) can occur when metal carbene **II** reacts with another diazo reactant. Using

in situ IR spectroscopy, we determined that the ortho-substituted prochiral diazo **3a** has a significantly reduced rate of azine formation (vs 2a), which accounts for higher yields of the Si-H insertion products. Relative rates of azine formation ($k_{rel} > 120$) was observed for decomposition of diazo 2a vs 3a with Rh₂(S-TCPTTL)₄ (in toluene) in the absence of silane (Figure 2A).⁴⁶ Increased yields of Si-H insertion products with ortho-substitution (5a-i) are attributed to steric interactions blocking the approach of diazo 3a to II, which reduces off-cycle azine formation. The addition of 4Å mol sieves reduces off-cycle processes leading to formation of siloxane 8.42 The increase in enantioselectivity observed with prochiral donor/donor diazo 3 is attributed to a twisting of the ortho-substituted aryl ring, which blocks one face of the carbene in II to promote selective approach of the silane (Figure 2B).44,47,48 Davies recently reported that the twisting effect has electronic contributions similar to that of a donor/acceptor carbene;⁴⁴ however, our data supports that the steric effect of an out-of-plane phenyl twist is significant.

1

2

3 4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31 32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58 59

60

To demonstrate the utility of enantioenriched silanes, silane **5a** was transformed to silanol, dehydrocoupling, and intramolecular C–H silylation products. Silanes are useful intermediates in stereoselective synthesis, and have versatile reactivity with the remaining Si–H bond.⁴⁹ It is well know that transition metals are capable of oxidative insertion into Si–H bonds with retention of configuration. ^{50,51} Pd/C-catalyzed silane hydrolysis affords silanol **9** in 90% yield with 90:10 dr and 93:7 er.^{38,51,52} Under attempted hydrosilylation conditions, an unexpected dehydrocoupling product **10** was isolated in good yield (62%) and 93:7 dr.^{53–55} Exploiting the presence of the ortho-methyl group, diasteroenriched sila-indane **11** was accessed in 90% yield with 90:10 dr using Ir-catalyzed C–H silylation methodology developed by the Hartwig group.^{56,57}



^{*a*} Isolated yields; dr determined using ¹H NMR spectroscopy; er determined using CSP-HPLC. ^{*b*} Isolated as a 85:15 (major) mixture with the hydrosilylation product. See SI for more information. ^{*c*} dr determined using ¹⁹F NMR spectroscopy.

In conclusion, the first example of enantioselective diarylcarbene insertion into Si-H bonds has been accomplished with Rh₂(*S*-TCPTTL)₄, yielding silicon-stereogenic benzhydryl silanes. While symmetrical diazo compounds demonstrated initial enantioselectivity, using a prochiral diazo reactant dramatically improved the reaction, providing yields up to 98% with 98:2 dr and 95:5 er. A catalytic cycle is proposed and the impact of the off-cycle azine formation is supported based on the structural dependence for relative rates of diazo decomposition. Transformation of the enantioenriched silane affords access to silicon-stereogenic silanol, dehydrocoupling and intramolecular C–H silylation products.

ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website.

Experimental procedures, spectra for all new compounds, HPLC data, procedures for *in situ* IR experiments, and X-ray structure information for hydrolysis product of **5i** (CCDC 2009546) (PDF).

AUTHOR INFORMATION

Corresponding Authors

* email: akfranz@ucdavis.edu or jtshaw@ucdavis.edu

Funding Sources

We acknowledge the National Science Foundation for support of this research (CHE-1900300, CHE-1363375; AKF) and the dual source diffractometer (CHE-1531193) used in this study. The National Institutes of Health (R01 GM124234; JTS) is also acknowledged for support of this research.

Author Contributions

All authors have given approval to the final version of the manuscript.

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENT

Will Jewell is acknowledged for assistance with acquiring mass spectrometry data. Benjamin Bergstrom, Sarah Dishman and Lucas Souza are acknowledged for helpful discussion.

REFERENCES

- (1) Schmidt, D. R.; O'Malle, S. J.; Leighton, J. L. Catalytic Asymmetric Silane Alcoholysis: Practical Access to Chiral Silanes. J. Am. Chem. Soc. 2003, 125 (5), 1190–1191. See ref. 2 and 3 for more examples of dehydrocoupling methodology to access stereogenic silanes.
- (2) Li, N.; Guan, B. Yttrium–Benzyl Complexes Bearing Chiral Iminophosphonamide Ligands: Synthesis and Application in Catalytic Asymmetric Amine-Silane Dehydrocoupling Reactions. *Adv. Synth. Catal.* 2017, 359 (20), 3526–3531.
- (3) Corriu, R. J. P.; Moreau, J. J. E. Asymmetric Synthesis at Silicon. J. Organomet. Chem. **1976**, 120 (3), 337–346.
- (4) Koga, S.; Ueki, S.; Shimada, M.; Ishii, R.; Kurihara, Y.; Yamanoi, Y.; Yuasa, J.; Kawai, T.; Uchida, T. A.; Iwamura, M.; Nozaki, K.; Nishihara, H. Access to Chiral Silicon Centers for Application to Circularly Polarized Luminescence Materials. *J. Org. Chem.* 2017, 82 (12), 6108–6117. See ref. 5 for more examples of arylation methodology to access stereogenic silanes.
- (5) Kurihara, Y.; Nishikawa, M.; Yamanoi, Y.; Nishihara, H. Synthesis of Optically Active Tertiary Silanes via Pd-Catalyzed Enantioselective Arylation of Secondary Silanes. *Chem. Commun.* **2012**, *48* (94), 11564–11566.
- (6) Igawa, K.; Yoshihiro, D.; Ichikawa, N.; Kokan, N.; Tomooka, K. Catalytic Enantioselective Synthesis of Alkenylhydrosilanes. *Angew*.

1

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

Chemie Int. Ed. **2012**, *51* (51), 12745–12748. See ref. 7-9 for more examples of hydrosilylation methodology to access stereogenic silanes.

- (7) Hayashi, T.; Yamamoto, K.; Kumada, M. Asymmetric Synthesis of Bifunctional Organosilicon Compounds via Hydrosilylation. *Tetrahedron Lett.* **1974**, *15* (4), 331–334.
- (8) Ohta, T.; Ito, M.; Tsuneto, A.; Takaya, H. Asymmetric Synthesis of Silanes with a Stereogenic Centre at Silicon via Hydrosilylation of Symmetric Ketones with Prochiral Diaryl Silanes Catalysed by Binap-Rh Complexes. J. Chem. Soc. Chem. Commun. 1994, No. 21, 2525– 2526.
- (9) Zhao, Z.-Y.; Nie, Y.-X.; Tang, R.-H.; Yin, G.-W.; Cao, J.; Xu, Z.; Cui, Y.-M.; Zheng, Z.-J.; Xu, L.-W. Enantioselective Rhodium-Catalyzed Desymmetric Hydrosilylation of Cyclopropenes. ACS Catal. 2019, 9 (10), 9110–9116.
- (10) Shintani, R. Recent Advances in the Transition-Metal-Catalyzed Enantioselective Synthesis of Silicon-Stereogenic Organosilanes. *Asian J. Org. Chem.* 2015, 4, 510–514. See ref. 11 for more examples of Si–C activation to access stereogenic silanes.
- (11) Shintani, R. Recent Progress in Catalytic Enantioselective Desymmetrization of Prochiral Organosilanes for the Synthesis of Silicon-Stereogenic Compounds. *Synlett* **2018**, *29* (4), 388–396.
- (12) Bauer, J. O.; Strohmann, C. Stereocontrol in Nucleophilic Substitution Reactions at Silicon: The Role of Permutation in Generating Silicon-Centered Chirality. J. Am. Chem. Soc. 2015, 137 (13), 4304–4307. See ref. 13 and 14 for more examples of using chiral auxillaries to access stereogenic silanes.
- (13) Bauer, J. O.; Strohmann, C. Stereoselective Synthesis of Silicon-Stereogenic Aminomethoxysilanes: Easy Access to Highly Enantiomerically Enriched Siloxanes. *Angew. Chemie Int. Ed.* 2014, 53 (3), 720–724.
- (14) Kimiko, K.; Takayuki, K.; Masafumi, U.; Shinji, M. Asymmetric Synthesis of Organosilicon Compounds Using a C2 Chiral Auxiliary. *Bull. Chem. Soc. Jpn.* **1997**, *70* (6), 1393–1401.
- (15) Xu, L.-W.; Li, L.; Lai, G.-Q.; Jiang, J.-X. The Recent Synthesis and Application of Silicon-Stereogenic Silanes: A Renewed and Significant Challenge in Asymmetric Synthesis. *Chem. Soc. Rev.* **2011**, 40 (3), 1777–1790. See ref. 16 and 17 for more examples of transformations using silicon-stereogenic silanes.
- (16) Bauer, J. O.; Strohmann, C. Recent Progress in Asymmetric Synthesis and Application of Difunctionalized Silicon-Stereogenic Silanes. *Eur. J. Inorg. Chem.* **2016**, 2016 (18), 2868–2881.
- (17) Shintani, R. Catalytic Asymmetric Synthesis of Siliconstereogenic Compounds by Enantioselective Desymmetrization of Prochiral Tetraorganosilanes. *Yuki Gosei Kagaku Kyokaishi/Journal Synth.* Org. Chem. 2018, 76 (11), 1163–1169.
- (18) Keipour, H.; Carreras, V.; Ollevier, T. Recent Progress in the Catalytic Carbene Insertion Reactions into the Silicon-Hydrogen Bond. Org. Biomol. Chem. 2017, 15 (26), 5441–5456. See ref 19-25 for work on enantioselective donor/acceptor carbene insertions into Si–H bonds.
- (19) Bagheri, V.; Doyle, M. P.; Taunton, J.; Claxton, E. E. A New and General Synthesis of .Alpha.-Silyl Carbonyl Compounds by Silicon-Hydrogen Insertion from Transition Metal-Catalyzed Reactions of Diazo Esters and Diazo Ketones. J. Org. Chem. 1988, 53 (26), 6158– 6160.
- (20) Chen, D.; Zhu, D.-X.; Xu, M.-H. Rhodium(I)-Catalyzed Highly Enantioselective Insertion of Carbenoid into Si-H: Efficient Access to Functional Chiral Silanes. J. Am. Chem. Soc. 2016, 138 (5), 1498– 1501.
- (21) Gu, H.; Han, Z.; Xie, H.; Lin, X. Iron-Catalyzed Enantioselective Si-H Bond Insertions. Org. Lett. 2018, 20 (20), 6544–6549.
- (22) Dakin, L. A.; Ong, P. C.; Panek, J. S.; Staples, R. J.; Stavropoulos, P. Speciation and Mechanistic Studies of Chiral Copper(I) Schiff Base Precursors Mediating Asymmetric Carbenoid Insertion Reactions of Diazoacetates into the Si–H Bond of Silanes. *Organometallics* 2000, 19 (15), 2896–2908.
- (23) Kan, S. B. J.; Lewis, R. D.; Chen, K.; Arnold, F. H. Directed Evolution of Cytochrome c for Carbon-Silicon Bond Formation: Bringing Silicon to Life. *Science* 2016, *354* (6315), 1048–1051.
- (24) Yasutomi, Y.; Suematsu, H.; Katsuki, T. Iridium(III)-Catalyzed Enantioselective Si-H Bond Insertion and Formation of an Enantioenriched Silicon Center. J. Am. Chem. Soc. 2010, 132 (13), 4510–4511.
- (25) Nakagawa, Y.; Chanthamath, S.; Fujisawa, I.; Shibatomi, K.; Iwasa,

S. Ru(II)-Pheox-Catalyzed Si–H Insertion Reaction: Construction of Enantioenriched Carbon and Silicon Centers. *Chem. Commun.* **2017**, *53* (26), 3753–3756.

- (26) Liu, Z.; Li, Q.; Yang, Y.; Bi, X. Silver(i)-Promoted Insertion into X– H (X = Si, Sn, and Ge) Bonds with N-Nosylhydrazones. *Chem. Commun.* 2017, 53 (16), 2503–2506. See ref. 27-29 for previous work on donor/donor carbene insertion into Si–H bonds.
- (27) Liu, Z.; Huo, J.; Fu, T.; Tan, H.; Ye, F.; Hossain, M. L.; Wang, J. Palladium(0)-Catalyzed C(Sp3)–Si Bond Formation via Formal Carbene Insertion into a Si–H Bond. *Chem. Commun.* **2018**, *54* (81), 11419–11422.
- (28) Wang, E. H.; Ping, Y. J.; Li, Z. R.; Qin, H.; Xu, Z. J.; Che, C. M. Iron Porphyrin Catalyzed Insertion Reaction of N-Tosylhydrazone-Derived Carbenes into X-H (X = Si, Sn, Ge) Bonds. Org. Lett. 2018, 20 (15), 4641–4644.
- (29) Huang, M.-Y.; Yang, J.-M.; Zhao, Y.-T.; Zhu, S.-F. Rhodium-Catalyzed Si–H Bond Insertion Reactions Using Functionalized Alkynes as Carbene Precursors. ACS Catal. 2019, 9 (6), 5353–5357.
- (30) Soldi, C.; Lamb, K. N.; Squitieri, R. A.; González-López, M.; Di Maso, M. J.; Shaw, J. T. Enantioselective Intramolecular C–H Insertion Reactions of Donor–Donor Metal Carbenoids. J. Am. Chem. Soc. 2014, 136 (43), 15142–15145. See ref. 31-34 on recent work with donor/donor carbenes.
- (31) Souza, L. W.; Squitieri, R. A.; Dimirjian, C. A.; Hodur, B. M.; Nickerson, L. A.; Penrod, C. N.; Cordova, J.; Fettinger, J. C.; Shaw, J. T. Enantioselective Synthesis of Indolines, Benzodihydrothiophenes, and Indanes by C-H Insertion of Donor/Donor Carbenes. *Angew. Chemie* **2018**, *130* (46), 15433– 15436.
- (32) Lamb, K. N.; Squitieri, R. A.; Chintala, S. R.; Kwong, A. J.; Balmond, E. I.; Soldi, C.; Dmitrenko, O.; Castiñeira Reis, M.; Chung, R.; Addison, J. B.; Fettinger, J. C.; Hein, J. E.; Tantillo, D. J.; Fox, J. M.; Shaw, J. T. Synthesis of Benzodihydrofurans by Asymmetric C-H Insertion Reactions of Donor/Donor Rhodium Carbenes. *Chem.* - A Eur. J. 2017, 23 (49), 11843–11855.
- (33) Zhu, D.; Chen, L.; Fan, H.; Yao, Q.; Zhu, S. Recent Progress on Donor and Donor–Donor Carbenes. *Chem. Soc. Rev.* **2020**, *49* (3), 908–950.
- (34) Nickerson, L. A.; Bergstrom, B. D.; Gao, M.; Shiue, Y. S.; Laconsay, C. J.; Culberson, M. R.; Knauss, W. A.; Fettinger, J. C.; Tantillo, D. J.; Shaw, J. T. Enantioselective Synthesis of Isochromans and Tetrahydroisoquinolines by C-H Insertion of Donor/Donor Carbenes. *Chem. Sci.* **2020**, *11* (2), 494–498.
- (35) Werlé, C.; Goddard, R.; Philipps, P.; Farès, C.; Fürstner, A. Structures of Reactive Donor/Acceptor and Donor/Donor Rhodium Carbenes in the Solid State and Their Implications for Catalysis. *J. Am. Chem. Soc.* 2016, *138* (11), 3797–3805.
- (36) Davis, P. J.; Harris, L.; Karim, A.; Thompson, A. L.; Gilpin, M.; Moloney, M. G.; Pound, M. J.; Thompson, C. Substituted Diaryldiazomethanes and Diazofluorenes: Structure, Reactivity and Stability. *Tetrahedron Lett.* **2011**, *52* (14), 1553–1556.
- (37) Tran, N. T.; Wilson, S. O.; Franz, A. K. Cooperative Hydrogen-Bonding Effects in Silanediol Catalysis. *Org. Lett.* 2012, *14* (1), 186–189. See ref. 38 amd 39 for our previous work on the synthesis of silanes.
- (38) Diemoz, K. M.; Wilson, S. O.; Franz, A. K. Synthesis of Structurally Varied 1,3-Disiloxanediols and Their Activity as Anion-Binding Catalysts. *Chemistry (Easton).* 2016, *22* (51), 18349–18353.
- (39) Diemoz, K. M.; Hein, J. E.; Wilson, S. O.; Fettinger, J. C.; Franz, A. K. Reaction Progress Kinetics Analysis of 1,3-Disiloxanediols as Hydrogen-Bonding Catalysts. *J. Org. Chem.* **2017**, *82* (13), 6738–6747.
- (40) Keipour, H.; Jalba, A.; Delage-Laurin, L.; Ollevier, T. Copper-Catalyzed Carbenoid Insertion Reactions of α-Diazoesters and α-Diazoketones into Si–H and S–H Bonds. J. Org. Chem. 2017, 82 (6), 3000–3010.
- (41) Landais, Y.; Parra-Rapado, L.; Planchenault, D.; Weber, V. Mechanism of Metal-Carbenoid Insertion into the Si-H Bond. *Tetrahedron Lett.* **1997**, *38* (2), 229–232.
- (42) Hydrolysis of Si-H bonds using dirhodium(II) compounds has been previous observed: Doyle, M. P.; High, K. G.; Bagheri, V.; Pieters, R.; Lewis, P. J.; Pearson, M. M. Rhodium(II) Perfluorobutyrate Catalyzed Silane Alcoholysis. A Highly Selective Route to Silyl Ethers. J. Org. Chem. 1990, 55 (25), 6082–6086.
- (43) Lloret, J.; Carbó, J. J.; Bo, C.; Lledós, A.; Pérez-Prieto, J. Influence

of the Nature of the Ligand on Dirhodium(II) Carbene Species: A Theoretical Analysis. *Organometallics* **2008**, *27* (12), 2873–2876.

- (44) Recent work supports the twisting of the phenyl ring for carbene II caused by ortho substitution, see: Lee, M.; Ren, Z.; Musaev, D. G.; Davies, H. M. L. Rhodium-Stabilized Diarylcarbenes Behaving as Donor/Acceptor Carbenes. ACS Catal. 2020, 6240–6247.
- (45) Yasutomi, Y.; Suematsu, H.; Katsuki, T. Iridium(III)-Catalyzed Enantioselective Si-H Bond Insertion and Formation of an Enantioenriched Silicon Center. J. Am. Chem. Soc. 2010, 132 (13), 4510–4511.
- (46) Azines 6 and 7 Were Formed >90 % yield during kinetic exeperiments as determined using ¹H NMR spectroscopy with Ph-TMS as an internal standard; thereore the rates of azine formation were assumed to correlate directly with the consumption of diazo compounds 2a and 3a.
- (47) The rationale for the "all up" conformation of ligands is based on: Lindsay, V. N. G.; Charette, A. B. Design and Synthesis of Chiral Heteroleptic Rhodium(II) Carboxylate Catalysts: Experimental Investigation of Halogen Bond Rigidification Effects in Asymmetric Cyclopropanation. ACS Catal. 2012, 2 (6), 1221–1225.
- (48) The rationale for a twisted phenyl ring in the structure of carbene II is based of X-ray data from: Werlé, C.; Goddard, R.; Fürstner, A. The First Crystal Structure of a Reactive Dirhodium Carbene Complex and a Versatile Method for the Preparation of Gold Carbenes by Rhodium-to-Gold Transmetalation. *Angew. Chemie Int. Ed.* 2015, *54* (51), 15452–15456. Refer to ref. 44 and 32 for computational support for this structure.
- (49) Fleming, I.; Barbero, A.; Walter, D. Stereochemical Control in Organic Synthesis Using Silicon-Containing Compounds. *Chem. Rev.* 1997, 97 (6), 2063–2192.

- (50) The insertion of Rh(I) and Ir(I) is known to occur with retention of stereochemistry: Oestreich, M. Chirality Transfer from Silicon to Carbon. *Chem. – A Eur. J.* 2006, *12* (1), 30–37. See ref. 51 for more information.
- (51) Corriu, R. J. P.; Guérin, C.; Moreau, J. J. E. Stereochemistry at Silicon. *Topics in Stereochemistry*. January 1, 1984, pp 43–198.
- (52) Hydrolysis of the silane to the silanol using Pd/C/H₂O conditions occurs with inversion of stereochemistry, see ref. 15 and 51 for more information. Hydrolysis of the silane to the silanol under these conditons was performed on all substrates to ensure separation of enantiomers for CSP-HPLC analysis.
- (53) Onopchenko, A.; Sabourin, E. T.; Beach, D. L. Vinyl- and Allylsilanes from the Rhodium(I)-Catalyzed Hydrosilylation of 1-Alkenes with Trialkylsilanes. J. Org. Chem. 1984, 49 (18), 3389– 3392.
- (54) Kakiuchi, F.; Nogami, K.; Chatani, N.; Seki, Y.; Murai, S. Dehydrogenative Silylation of 1,5-Dienes with Hydrosilanes Catalyzed by RhCl(PPh₃)₃. Organometallics **1993**, *12* (12), 4748– 4750.
- (55) Adams, C.; Riviere, P.; Riviere-Baudet, M.; Morales-Verdejo, C.; Dahrouch, M.; Morales, V.; Castel, A.; Delpech, F.; Manríquez, J. M.; Chávez, I. Catalytic Study of Heterobimetallic Rhodium Complexes Derived from Partially Alkylated S-Indacene in Dehydrogenative Silylation of Olefins. J. Organomet. Chem. 2014, 749, 266–274.
- (56) Cheng, C.; Hartwig, J. F. Catalytic Silylation of Unactivated C-H Bonds. Chem. Rev. 2015, 115 (17), 8946–8975.
- (57) Karmel, C.; Chen, Z.; Hartwig, J. F. Iridium-Catalyzed Silylation of C-H Bonds in Unactivated Arenes: A Sterically Encumbered Phenanthroline Ligand Accelerates Catalysis. J. Am. Chem. Soc. 2019, 141 (17), 7063–7072.

59

60

