

Gold-Catalyzed Ammonium Acetate-Assisted Cascade Cyclization of 2-Alkynylarylketones

Maciej E. Domaradzki, Yuhua Long, Zhigang She, Xiaochen Liu, Gan Zhang, and Yu Chen

J. Org. Chem., **Just Accepted Manuscript** • DOI: 10.1021/acs.joc.5b01939 • Publication Date (Web): 23 Oct 2015

Downloaded from <http://pubs.acs.org> on October 25, 2015

Just Accepted

"Just Accepted" manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides "Just Accepted" as a free service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. "Just Accepted" manuscripts appear in full in PDF format accompanied by an HTML abstract. "Just Accepted" manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are accessible to all readers and citable by the Digital Object Identifier (DOI®). "Just Accepted" is an optional service offered to authors. Therefore, the "Just Accepted" Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the "Just Accepted" Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these "Just Accepted" manuscripts.




Gold-Catalyzed Ammonium Acetate-Assisted Cascade Cyclization of 2-Alkynylarylketones

Maciej E. Domaradzki,^a Yuhua Long,^a Zhigang She, Xiaochen Liu, Gan Zhang, and Yu Chen*

Department of Chemistry and Biochemistry, Queens College and the Graduate Center of the City

University of New York, 65-30 Kissena Blvd., Flushing, New York 11367

yu.chen1@qc.cuny.edu



ABSTRACT

An ammonium acetate-assisted gold-catalyzed cascade cyclization reaction of 2-alkynylarylketones is described. Under the reported conditions, a gold-catalyzed intramolecular cyclization of 2-alkynylarylketones takes place through two competing reaction mechanisms—a 5-*exo-dig* or a 6-*endo-dig* cyclization—leading to two regioisomeric intermediates: isobenzofuranium or isobenzopyrylium. In the presence of ammonium acetate, the two intermediate compounds undergo further rearrangement to 2,3-disubstituted indenones and 1,3-disubstituted isoquinolines, respectively. While both reaction pathways proceed via a cyclization-rearrangement cascade,

^a These authors have contributed equally to all the work described herein.

the gold-mediated 5-*exo-dig* process is especially notable as it provides a novel cyclization protocol of 2-alkynylarylketones.


INTRODUCTION

The Lewis acid induced intramolecular electrophilic cyclization of *o*-alkynylarylcarbonyl compounds has proven to be a versatile synthetic approach to a variety of biologically interesting heterocycles and carbocycles.^{1,2} Among the *o*-alkynylarylcarbonyl compounds, *o*-alkynylarylaldehydes have shown great versatility, displaying high reactivity as substrates for cyclization reactions.³ In comparison to the many electrophilic cyclization reactions explored for *o*-alkynylarylaldehydes, examples of cyclization reactions involving *o*-alkynylarylketones are relatively rare,⁴ possibly due to their lower chemical reactivity than the aldehydes. The cyclizations of *o*-alkynylarylketones are usually explored as an extension of the reaction scope of analogous *o*-alkynylarylaldehydes. In general, the known cyclization processes of *o*-alkynylarylaldehydes and -ketones share the same isobenzopyrylium intermediate, which reacts with either a nucleophile to furnish isochromene derivatives^{3a-c,5} (Scheme 1, path a) or undergoes a cycloaddition with an alkene/alkyne moiety to generate polycyclic compounds (Scheme 1, path b).⁶ Besides the two major reaction pathways, a few examples have demonstrated that in the presence of a nitrogen nucleophile, such as ammonia,⁷ ammonium acetate,⁸ or primary amines,⁹ intramolecular cyclizations of *o*-alkynylarylaldehydes and -ketones lead to the formation of isoquinoline derivatives (Scheme 1, path c). Although the two *o*-alkynylarylcarbonyl substrates often show similar chemical reactivity, and lead to the same derivatives, it is known that *o*-alkynylarylaldehydes preferentially undergo 6-*endo-dig* cyclizations to form isobenzopyrylium intermediates, whereas *o*-alkynylarylketones often undergo 5-*exo-dig* cyclizations to form isobenzofuraniums, under similar reaction conditions. While the reactivity of isobenzopyrylium intermediates generated from the cyclization of *o*-alkynylarylcarbonyl compounds has been explored in numerous processes, such as nucleophilic additions⁵ and cycloadditions,⁶ the reactivity of isobenzofuranium intermediates generated in these electrophilic cyclization reactions have not yet been widely elucidated.¹⁰ Most of the known

reactions of isobenzofuranium intermediates are limited to simple nucleophilic additions, which exclusively lead to isobenzofuran derivatives (Scheme 1, path d).^{3h,11}

We previously developed a one-pot synthesis of isoquinolines by a palladium-catalyzed sequential coupling–imination–annulation reaction,^{8a} involving a key mechanistic step—the electrophilic cyclization of *o*-alkynylarylaldehydes in the presence of *in situ* generated ammonia. When we attempted to expand the substrate scope of this synthetic protocol to *o*-alkynylarylketones, we encountered limited success, and the desired isoquinolines were only obtained in low yields (< 20%). The low chemical reactivity observed for *o*-alkynylarylketones in our previous work prompted us to further explore their reactivity in the electrophilic cyclization reactions. We hereby report our recent progress on this topic: an ammonium acetate-assisted gold-catalyzed electrophilic cyclization of *o*-alkynylarylketones, through a cascade cyclization-rearrangement process, leading to the formation of indenones and isoquinolines (Scheme 1, path e).

Scheme 1. Intramolecular Cyclization Reactions of *o*-Alkynylarylaldehydes and –ketones.




RESULTS AND DISCUSSION

Our initial study focused on the cyclization reaction of the model substrate 1-(2-(phenylethynyl)phenyl)ethan-1-one (**1a**). In the presence of 6 mol% of both NaAuCl₄·2H₂O and AgSbF₆, reaction of **1a** led to partial decomposition and no useful product was obtained (Table 1, entry 1). During the course of studying this reaction, we found significant additive effects in this cyclization.

In the presence of 1 equiv of ammonium acetate (NH_4OAc), indenone **2a** was obtained in 48% yield together with 10% of isoquinoline **3a** after 4 hours in a 70% conversion (Table 1, entry 2). Longer reaction time did not improve the chemical yields but caused partial decomposition of the products. Further investigation showed that in the presence of 2 equiv of pyridine *N*-oxide the reaction was completed in 2 hours and the chemical yields of indenone **2a** and isoquinoline **3a** were enhanced to 53% and 15%, respectively (Table 1, entry 3). Other ammonium salts such as ammonium carbonate ($(\text{NH}_4)_2\text{CO}_3$) and ammonium formate (HCO_2NH_4) were also tested (Table 1, entries 4 and 5); however, both producing slightly inferior results to NH_4OAc . When *L*-proline was employed in this reaction to replace NH_4OAc , indenone **2a** was isolated as the sole product, although only in moderate yield (51%; Table 1, entry 6). Further investigation of additives, including several pyridine *N*-oxide derivatives and diphenyl sulfoxide (Table 1, entries 7-11), showed that the highest yields of the cyclization products were produced in the presence of 4-methoxypyridine *N*-oxide (Table 1, entry 7). No enhancement of chemical yields were observed, when the loading of NH_4OAc was increased to 1.5 equiv (Table 1, entry 12). When the loading of 4-methoxypyridine *N*-oxide was reduced to 1.2 equiv, the chemical yields of **2a** and **3a** were further raised to 69% and 19% (Table 1, entry 13). When $\text{NaAuCl}_4 \cdot 2\text{H}_2\text{O}$ was used as the sole catalyst, the yield of **2a** decreased to 46% while **3a** was obtained in the same yield (Table 1, entry 14). On the other hand, **3a** was obtained as the only product in 74% yield when AgSbF_6 was used as the sole catalyst (Table 1, entry 15). Considering the formation of **2a** involving an unprecedented cyclization-rearrangement cascade reaction of *o*-alkynylarylketones, the conditions listed in entry 13 in Table 1 were chosen to be tested with other substrates.

Table 1. Optimization of Reaction Conditions for the Cyclization of *o*-Alkynylarylketones.^a



entry	catalyst (mol %)	additive (equiv)	yield (%) ^b
-------	------------------	------------------	------------------------

			2a, 3a
1 ^c	NaAuCl ₄ ·2H ₂ O /AgSbF ₆ (6/6)	-	-, -
2 ^d	NaAuCl ₄ ·2H ₂ O /AgSbF ₆ (6/6)	NH ₄ OAc (1 equiv)	48, 10
3	NaAuCl ₄ ·2H ₂ O /AgSbF ₆ (6/6)	NH ₄ OAc (1 equiv), Pyridine <i>N</i> -Oxide (2 equiv)	53, 15
4	NaAuCl ₄ ·2H ₂ O/AgSbF ₆ (6/6)	(NH ₄) ₂ CO ₃ (1 equiv), Pyridine <i>N</i> -Oxide (2 equiv)	50, 14
5	NaAuCl ₄ ·2H ₂ O/AgSbF ₆ (6/6)	HCO ₂ NH ₄ (1 equiv), Pyridine <i>N</i> -Oxide (2 equiv)	43, 13
6	NaAuCl ₄ ·2H ₂ O/AgSbF ₆ (6/6)	<i>L</i> -Proline (1 equiv), Pyridine <i>N</i> -Oxide (2 equiv)	51, -
7	NaAuCl ₄ ·2H ₂ O/AgSbF ₆ (6/6)	NH ₄ OAc (1 equiv), 4- Methoxypyridine <i>N</i> -Oxide (2 equiv)	59, 18
8	NaAuCl ₄ ·2H ₂ O/AgSbF ₆ (6/6)	NH ₄ OAc (1 equiv), 4- Nitropyridine <i>N</i> -Oxide (2 equiv)	41, 12
9	NaAuCl ₄ ·2H ₂ O/AgSbF ₆ (6/6)	NH ₄ OAc (1 equiv), 2,6- Dichloropyridine <i>N</i> -Oxide (2 equiv)	37, 12
10	NaAuCl ₄ ·2H ₂ O/AgSbF ₆ (6/6)	NH ₄ OAc (1 equiv), Quinoline <i>N</i> -Oxide (2 equiv)	56, 15
11	NaAuCl ₄ ·2H ₂ O/AgSbF ₆ (6/6)	NH ₄ OAc (1 equiv), Diphenylsulfoxide (2	47, 20

			equiv)	
12	NaAuCl ₄ ·2H ₂ O/AgSbF ₆ (6/6)	NH ₄ OAc (1.5 equiv), 4-	55, 20	
		Methoxypyridine <i>N</i> -Oxide		
		(2 equiv)		
13	NaAuCl ₄ ·2H ₂ O/AgSbF ₆ (6/6)	NH ₄ OAc (1 equiv), 4-	69, 19	
		Methoxypyridine <i>N</i> -Oxide		
		(1.2 equiv)		
14	NaAuCl ₄ ·2H ₂ O (12)	NH ₄ OAc (1 equiv), 4-	46, 19	
		Methoxypyridine <i>N</i> -Oxide		
		(1.2 equiv)		
15	AgSbF ₆ (12)	NH ₄ OAc (1 equiv), 4-	-, 74	
		Methoxypyridine <i>N</i> -Oxide		
		(1.2 equiv)		







^a General procedure: The catalyst, additives, substrate **1a** (110.2 mg, 0.5 mmol), and CH₃CN (4 mL) were added to a 4-dram vial. The reaction mixture was purged with argon, sealed, and stirred at 85 °C for 2 h. ^b Isolated yields after column chromatography. ^c Reaction was stopped after 8 h. ^d Reaction was stopped after 4 h and 30% of **1a** was recovered.


A variety of 2-alkynylarylketones, including those with aryl, alkenyl, and alkyl groups at the distal position of the alkyne triple bond, were then subjected to the optimized reaction conditions (Table 2, entries 1-10). Based on these results, electron-donating and electron-withdrawing groups are found to be compatible on both the distal and proximal phenyl rings, relative to the alkyne triple bond (Table 2, entries 2 to 6), although lower yields for indenones are obtained when an electron-withdrawing substituent such as an ester group is present at the *para*-position of the distal phenyl ring (Table 2, entry 4) or when an electron-donating substituent such as a methoxy group is present at the *para*-position of the proximal phenyl ring (Table 2, entry 5). Additionally, the introduction of a sterically hindered


α -naphthyl group at the distal position of the triple bond leads to a lower chemical yield for indenone (35%) but an increased yield of isoquinoline (42%; Table 2, entry 7). Alkenyl and alkyl moieties at the distal position of the triple bond are also compatible, but likewise lower the yield of the indenone products (Table 2, entries 8-10). When a sterically hindered *tert*-butyl group is present, a low conversion of the starting material (40%) is observed during the cyclization, even after an elongated reaction time (Table 2, entry 10).

Aside from acetophenone substrates, several other species were also investigated to further elucidate the regioselectivity of the cyclization process. As with acetophenone, benzophenone moieties lead to both the indenone and isoquinoline products (Table 2, entries 11 and 12). However, the cyclization of the benzaldehyde substrate occurs only via a 6-*endo-dig* route, leading regiospecifically to isoquinoline (Table 2, entry 13). Likewise, the cyclization of 1-(3-(arylethynyl)thiophen-2-yl)ethan-1-ones, which has the ketone and alkynyl moieties at the *ortho*-positions of the 5-membered heterocyclic ring-system, proceeds exclusively via a 6-*endo-dig* cyclization, affording only thieno[2,3-*c*]pyridines (Table 2, entries 14 and 15).

Table 2. Gold-Catalyzed Cyclization of *o*-Alkynylarylketones and -aldehydes.^a

entry	substrate 1	product 2 & 3 / yield ^b	
1	 1a	 2a / 69%	 3a / 19%
2 ^c	 1b	 2b / 70%	 3b / 14%






^a See the Experimental Section for the general procedure. ^b Isolated yields after column chromatography. ^c When *L*-proline was employed to replace NH₄OAc, indenone **2b** was isolated as the sole product in 29% yield. ^d When *L*-proline was employed to replace NH₄OAc, indenone **2i** was isolated as the sole product in 30% yield. ^e Reaction was stopped after 8 h, and 60% of starting material **1j** was recovered.

The reaction presumably begins with a gold-catalyzed intramolecular electrophilic cyclization, proceeding via one of two possible reaction pathways—a 6-*endo-dig* or 5-*exo-dig* cyclization (Scheme 2). When a 6-*endo-dig* cyclization takes place, the isobenzopyrylium intermediate **4** forms. The nucleophilic addition of ammonia to **4** leads to the isochromene intermediate **5**. After protonation of the oxygen atom of isochromene **5**, the carbon-oxygen bond dissociates to form enol **6**, which subsequently tautomerizes to ketone **7**. An intramolecular nucleophilic addition of the imino group to the ketone carbonyl group forms cyclized intermediate **8**. Following dehydration, the final isoquinoline product is

produced, together with the regenerated catalyst. Alternatively, when a 5-*exo-dig* cyclization takes place, the isobenzofuranium intermediate **9** forms. The nucleophilic addition of ammonia to **9** leads to the isobenzofuran intermediate **10**. After protonation of the oxygen atom of isobenzofuran **10**, the carbon-oxygen bond is cleaved to form enol **11**, which subsequently cyclizes via an intramolecular nucleophilic addition to the imino group to form indanone intermediate **12**. After elimination of ammonia, the indenone is produced together with the regenerated catalyst. In both cases the role of ammonium acetate is to trap the oxonium intermediates (**4** and **9**) and drive the reaction in the forward direction. Our results showed that both NaAuCl₄ and AgSbF₆ were effective catalysts for the cyclization of 2-alkynylarylketones though they displayed different regioselectivity. AgSbF₆ induces a regioselective 6-*endo-dig* cyclization (Table 1, entry 15), while NaAuCl₄ leads to the formation of a mixture of 6-*endo-dig* and 5-*exo-dig* cyclization products (Table 1, entry 14). In the current reaction protocol, AgSbF₆ presumably activates the gold catalyst by abstracting the chloride ions,¹² and the combination of NaAuCl₄ and AgSbF₆ provides superior result to the sole NaAuCl₄ catalyst (Table 1, entries 13 and 14). While the specific role of 4-methoxypyridine *N*-oxide is unclear, it is hypothesized to function as an accelerant that stabilizes the gold(III) catalyst, thus increasing the turnover number of both cyclization processes.¹³ We also cannot rule out the possibility that 4-methoxypyridine *N*-oxide plays the role as a buffer to maintain an optimal pK_a value of the reaction medium for the cascade cyclization.¹⁴

Scheme 2. Proposed Mechanism for Gold-Catalyzed Cyclization of *o*-Alkynylarylketones.



CONCLUSION

An ammonium acetate-assisted gold-catalyzed cascade cyclization of 2-alkynylarylketones is described. The cyclization process is initiated by one of two competing reactions—a 5-*exo-dig* or a 6-*endo-dig* cyclization—of the carbonyl group with the alkyne triple bond, leading to isobenzofuranium or isobenzopyrylium intermediates, respectively. The two intermediates tandemly undergo an ammonium acetate-assisted rearrangement and a subsequent cyclization to generate two different species: indenone and isoquinoline. The methodology reported herein is notable as it describes the first documented example of an isobenzofuranium intermediate undergoing a novel cascade reaction. Further exploration into the processes controlling the regioselectivity of this cascade and into the chemical reactivity of the isobenzofuranium intermediates are currently underway in our laboratory and will be reported in due course.


EXPERIMENTAL SECTION

General information. All reactions were carried out in sealed 4-/6-dram vials, unless otherwise indicated. All commercially available chemicals were used as received without further purification, unless otherwise noted. All ^1H and ^{13}C NMR spectra were recorded at 500 or 400 MHz and 125 or 100 MHz, respectively, using CDCl_3 as the solvent. The chemical shifts of all ^1H and ^{13}C NMR spectra are

referenced to the residual signal of CDCl_3 (δ 7.26 ppm for the ^1H NMR spectra and δ 77.23 ppm for the ^{13}C NMR spectra). High resolution mass analysis was carried out on a high resolution mass spectrometer using electrospray ionization (ESI) mode with a time-of-flight (TOF) mass analyzer. Samples were dissolved in methylene chloride and methanol and analyzed via flow injection into the mass spectrometer at a flow rate of 200 $\mu\text{L}/\text{min}$. The mobile phase was 90:10 methanol: water, with 0.1% formic acid. The mass spectrometer was operated in positive ion mode and the melting points are uncorrected.

The 2-alkynylarylketone/2-alkynylarylaldehyde materials (**1**) were prepared by Sonogashira coupling¹⁵ of 2-bromoarylketone/2-bromoarylaldehyde with terminal alkynes, except compounds **1c**, **1d**, **1g** and **1q**. 1-(2-Ethynylphenyl)ethanone (**1q**) was prepared from 1-(2-((trimethylsilyl)ethynyl)phenyl)ethan-1-one (**1p**) *via* a cesium fluoride mediated desilylation reaction. Compounds **1c**, **1d** and **1g** were prepared by Sonogashira coupling of **1q** with aryl iodides.

General procedure for the preparation of 2-alkynylarylketones and aldehydes (**1**).



A 6-dram vial was charged with 2-bromoarylketone (2.0 mmol), a terminal alkyne (2.2 mmol), $\text{Pd}(\text{PPh}_3)_2\text{Cl}_2$ (28.1 mg, 0.04 mmol), CuI (7.6 mg, 0.04 mmol), and Et_3N (14 mL). The vial was then purged with argon and sealed. The reaction mixture was stirred at 80°C for 18 h, until the disappearance of starting material was observed, as monitored by thin layer chromatography. The reaction mixture was diluted with diethyl ether (40 mL) and washed with brine (40 mL), and the aqueous phase was then extracted with diethyl ether (2×20 mL). The combined organic layers were dried over anhydrous MgSO_4 and concentrated using a rotary evaporator under reduced pressure (20 mmHg); the resulting residue was purified by flash column chromatography on silica gel (eluent: hexane/ethyl acetate).

1-(2-(Phenylethynyl)phenyl)ethanone (1a)

This product was obtained as a light brown oil (286.4 mg, 65% yield); ^1H NMR (500 MHz, CDCl_3) δ 7.76 (d, $J = 7.9$ Hz, 1H), 7.63 (d, $J = 7.6$ Hz, 1H), 7.54–7.57 (m, 2H), 7.47 (t, $J = 7.5$ Hz, 1H), 7.36–7.41 (m, 4H), 2.80 (s, 3H). The ^1H NMR spectral data are in good agreement with the literature data.¹⁶

1-(2-((4-Methoxyphenyl)ethynyl)phenyl)ethanone (1b)

This product was obtained as a light brown oil (160.2 mg, 32% yield); ^1H NMR (500 MHz, CDCl_3) δ 7.75 (dd, $J = 7.8, 1.0$ Hz, 1H), 7.61 (dd, $J = 7.7, 0.6$ Hz, 1H), 7.45–7.51 (m, 3H), 7.38 (dt, $J = 7.7, 1.0$ Hz, 1H), 6.90 (d, $J = 8.9$ Hz, 2H), 3.84 (s, 3H), 2.80 (s, 3H). The ^1H NMR spectral data are in good agreement with the literature data.¹⁷

1-(5-Methoxy-2-((4-methoxyphenyl)ethynyl)phenyl)ethanone (1e)

This product was obtained as a light brown oil (229.9 mg, 41% yield); ^1H NMR (500 MHz, CDCl_3) δ 7.52 (d, $J = 8.6$ Hz, 1H), 7.44–7.47 (m, 2H), 7.26 (d, $J = 2.8$ Hz, 1H), 7.01 (dd, $J = 8.6, 2.7$ Hz, 1H), 6.87–6.90 (m, 2H), 3.86 (s, 3H), 3.83 (s, 3H), 2.82 (s, 3H); ^{13}C NMR (125 MHz, CDCl_3) δ 200.8, 160.0, 159.4, 142.1, 135.3, 133.0, 118.3, 115.4, 114.6, 114.3, 113.1, 94.1, 87.4, 55.7, 55.5, 30.5; IR (neat, cm^{-1}) ν 2922, 1715, 1697, 1595, 1509, 1490, 1455, 1379, 1208, 1083, 1025, 853, 829, 755, 712, 696; HRMS (ESI) calcd for $\text{C}_{18}\text{H}_{17}\text{O}_3$ ($\text{M} + \text{H}$)⁺ 281.1172, Found: 281.1175.

1-(4-Fluoro-2-(phenylethynyl)phenyl)ethanone (1f)

This product was obtained as a light yellow oil (333.6 mg, 70% yield); ^1H NMR (500 MHz, CDCl_3) δ 7.82 (dd, $J = 8.8, 5.9$ Hz, 1H), 7.53–7.57 (m, 2H), 7.37–7.39 (m, 3H), 7.31 (dd, $J = 9.1, 2.4$ Hz, 1H), 7.07–7.11 (m, 1H), 2.78 (s, 3H); ^{13}C NMR (125 MHz, CDCl_3) δ 198.7, 164.1 (d, $J_{\text{C-F}} = 250.9$ Hz), 136.9 (d, $J_{\text{C-F}} = 3.0$ Hz), 131.83, 131.75, 129.3, 128.7, 124.6 (d, $J_{\text{C-F}} = 10.4$ Hz), 122.5, 120.6 (d, $J_{\text{C-F}} = 23.2$ Hz), 115.9 (d, $J_{\text{C-F}} = 21.5$ Hz), 96.4, 87.7 (d, $J_{\text{C-F}} = 2.6$ Hz), 30.1; IR (neat, cm^{-1}) ν 2970, 1735, 1683, 1595, 1558, 1490, 1447, 1418, 1364, 1266, 1228, 1216, 1098, 1066, 1024, 925, 877, 826, 756, 688; HRMS (ESI) calcd for $\text{C}_{16}\text{H}_{12}\text{FO}$ ($\text{M} + \text{H}$)⁺ 239.0867. Found: 239.0868.

1-(2-(Cyclohex-1-en-1-ylethynyl)phenyl)ethanone (1h)

This product was obtained as a light brown oil (152.5 mg, 34% yield); ^1H NMR (500 MHz, CDCl_3) δ 7.70 (dd, $J = 7.9, 1.1$ Hz, 1H), 7.50 (d, $J = 7.9, 1.2$ Hz, 1H), 7.41 (dt, $J = 7.3, 1.2$ Hz, 1H), 7.33 (dt, $J = 7.7, 1.3$ Hz, 1H), 6.24–6.26 (m, 1H), 2.74 (s, 3H), 2.22–2.25 (m, 2H), 2.14–2.18 (m, 2H), 1.66–1.71 (m, 2H), 1.60–1.65 (m, 2H); ^{13}C NMR (125 MHz, CDCl_3) δ 201.0, 140.8, 136.4, 133.9, 131.4, 128.8, 127.9, 122.5, 120.8, 97.4, 86.2, 30.3, 29.0, 26.0, 22.4, 21.6; IR (neat, cm^{-1}) ν 2925, 1696, 1587, 1464, 1426, 1355, 1281, 1239, 1091, 1025, 958, 756; HRMS (ESI) calcd for $\text{C}_{16}\text{H}_{17}\text{O}$ ($\text{M} + \text{H}$) $^+$ 225.1274, Found: 225.1275.

6-(2-Acetylphenyl)hex-5-ynenitrile (1i)

This product was obtained as a yellow oil (169.0 mg, 40% yield); ^1H NMR: (400 MHz, CDCl_3) δ 7.66 (dd, $J = 7.7, 1.0$ Hz, 1H), 7.48 (dd, $J = 7.6, 1.0$ Hz, 1H), 7.40 (td, $J = 7.6, 1.5$ Hz, 1H), 7.34 (td, $J = 7.6, 1.4$ Hz, 1H), 2.60–2.65 (m, 5H), 2.41–2.50 (m, 2H), 1.93–1.99 (m, 2H). The ^1H NMR spectral data are in good agreement with the literature data.¹⁸

1-(2-(3,3-Dimethylbut-1-yn-1-yl)phenyl)ethan-1-one (1j)

This product was obtained as yellow oil (280.4 mg, 70% yield); ^1H NMR: (400 MHz, CDCl_3) δ 7.64 (dd, $J = 7.8, 1.0$ Hz, 1H), 7.44 (dd, $J = 7.7, 0.9$ Hz, 1H), 7.35 (td, $J = 7.6, 1.5$ Hz, 1H), 7.28 (td, $J = 7.6, 1.4$ Hz, 1H), 2.71 (s, 3H), 1.32 (s, 9H). The ^1H NMR spectral data are in good agreement with the literature data.¹⁹

Phenyl(2-(phenylethynyl)phenyl)methanone (1k)

This product was obtained as a light yellow oil (254.1 mg, 45% yield); ^1H NMR (500 MHz, CDCl_3) δ 7.88–7.90 (m, 2H), 7.62 (dd, $J = 7.3, 1.1$ Hz, 1H), 7.57–7.60 (m, 1H), 7.51–7.55 (m, 2H), 7.44–7.50 (m, 3H), 7.18–7.25 (m, 3H), 7.03–7.05 (m, 2H). The ^1H NMR spectral data are in good agreement with the literature data.²⁰

(2-((4-Methoxyphenyl)ethynyl)phenyl)(phenyl)methanone (1l)

This product was obtained as a light brown oil (324.9 mg, 52% yield); ^1H NMR (500 MHz, CDCl_3) δ 7.89 (d, $J = 7.3$ Hz, 2H), 7.57–7.60 (m, 2H), 7.41–7.53 (m, 5H), 6.97 (d, $J = 8.8$ Hz, 2H), 6.73 (d, $J = 8.7$ Hz, 2H), 3.76 (s, 3H); ^{13}C NMR (125 MHz, CDCl_3) δ 197.4, 159.9, 141.4, 137.5, 133.3, 133.1,

132.5, 130.5, 130.4, 128.9, 128.5, 128.0, 122.4, 114.8, 113.9, 95.5, 86.5, 55.4; IR (neat, cm^{-1}) ν 2932, 2834, 2216, 1735, 1665, 1605, 1594, 1510, 1465, 1449, 1436, 1365, 1316, 1289, 1250, 1178, 1149, 1091, 1026, 960, 936, 880, 830, 805, 788, 770; HRMS (ESI) calcd for $\text{C}_{22}\text{H}_{17}\text{O}_2$ ($\text{M} + \text{H}$)⁺ 313.1223, Found: 313.1226.

2-(Phenylethynyl)benzaldehyde (1m)

This product was obtained as a light brown oil (325.9 mg, 79% yield); ^1H NMR (500 MHz, CDCl_3) δ 10.66 (s, 1H), 7.96 (d, $J = 7.6$ Hz, 1H), 7.66 (d, $J = 7.6$ Hz, 1H), 7.61 (d, $J = 7.2$ Hz, 1H), 7.57–7.59 (m, 2H), 7.47 (t, $J = 7.5$ Hz, 1H), 7.39–7.41 (m, 3H). The ^1H NMR spectral data are in good agreement with the literature data.²¹

1-(3-(Phenylethynyl)thiophen-2-yl)ethanone (1n)

This product was obtained as a yellow solid (353.0 mg, 78% yield): mp = 45.5–46.8 °C; ^1H NMR (500 MHz, CDCl_3) δ 7.53–7.75 (m, 3H), 7.38–7.39 (m, 3H), 7.23 (d, $J = 5.0$ Hz, 1H), 2.83 (s, 3H); ^{13}C NMR (125 MHz, CDCl_3) δ 191.3, 145.8, 133.0, 132.1, 131.6, 129.3, 128.8, 125.6, 122.5, 96.9, 84.8, 29.2; IR (neat, cm^{-1}) ν 3079, 2205, 1738, 1648, 1595, 1517, 1485, 1442, 1405, 1379, 1293, 1243, 1216, 1098, 1070, 1043, 1018, 992, 920, 846, 755; HRMS (ESI) calcd for $\text{C}_{14}\text{H}_{11}\text{OS}$ ($\text{M} + \text{H}$)⁺ 227.0525, Found: 227.0529.


1-(3-((4-Methoxyphenyl)ethynyl)thiophen-2-yl)ethanone (1o)

This product was obtained as a light yellow solid (256.3 mg, 50% yield): mp = 97.2–98.8 °C; ^1H NMR (500 MHz, CDCl_3) δ 7.54 (d, $J = 4.9$ Hz, 1H), 7.47–7.49 (m, 2H), 7.21 (d, $J = 5.1$ Hz, 1H), 6.90–6.92 (m, 2H), 3.85 (s, 3H), 2.83 (s, 3H); ^{13}C NMR (125 MHz, CDCl_3) δ 191.4, 160.5, 145.2, 133.2, 132.9, 132.1, 126.1, 114.6, 114.4, 97.2, 83.8, 55.6, 29.2; IR (neat, cm^{-1}) ν 2915, 2839, 2200, 1638, 1601, 1519, 1497, 1405, 1384, 1291, 1242, 1176, 1161, 1111, 1027, 991, 921, 850, 835, 811, 747; HRMS (ESI) calcd for $\text{C}_{15}\text{H}_{13}\text{O}_2\text{S}$ ($\text{M} + \text{H}$)⁺ 257.0631, Found: 257.0633.

1-(2-((Trimethylsilyl)ethynyl)phenyl)ethan-1-one (1p)


This product was obtained as a light yellow oil (341.8 mg, 79% yield); ^1H NMR (500 MHz, CDCl_3) δ 7.68 (dd, $J = 7.6, 1.4$ Hz, 1H), 7.55 (dd, $J = 7.6, 1.3$ Hz, 1H), 7.42 (dt, $J = 7.6, 1.6$ Hz, 1H), 7.37 (dt, $J = 7.6, 1.4$ Hz, 1H), 2.74 (s, 3H), 0.26 (s, 9H). The ^1H NMR spectral data are in good agreement with the literature data.²²

Procedure for the preparation of 1-(2-ethynylphenyl)ethanone (1q).



A 6-dram vial was charged with 1-(2-((trimethylsilyl)ethynyl)phenyl)ethanone (**1p**, 432.8 mg, 2.0 mmol), CsF (455.7 mg, 3.0 mmol) and methanol (12 mL). The reaction mixture was stirred at 35 °C for 18 h, until the disappearance of the starting material as monitored by thin layer chromatography. Methanol was removed using a rotary evaporator, under reduced pressure (20 mmHg). The residue was dissolved in diethyl ether (30 mL) and washed with brine (30 mL). The aqueous phase was extracted with diethyl ether (20 mL). The combined organic layers were dried over anhydrous MgSO_4 , and concentrated using a rotary evaporator, under reduced pressure (20 mmHg). The residue was purified by flash column chromatography on silica gel (eluent: hexanes/ethyl acetate), to afford a light yellow oil (181.7 mg, 63% yield); ^1H NMR (500 MHz, CDCl_3) δ 7.71 (d, $J = 7.8$ Hz, 1H), 7.60 (d, $J = 7.5$ Hz, 1H), 7.41–7.47 (m, 2H), 3.39 (s, 1H), 2.72 (s, 3H). The ^1H NMR spectral data are in good agreement with the literature data.²²

General procedure for the preparation of 2-alkynylarylketones (1c, 1d, 1g).



A 6-dram vial was charged with 1-(2-ethynylphenyl)ethanone (**1q**, 144.2 mg, 1.0 mmol), an aryl iodide (1.1 mmol), $\text{Pd}(\text{PPh}_3)_2\text{Cl}_2$ (14.0 mg, 0.02 mmol), CuI (3.8 mg, 0.02 mmol), and Et_3N (7 mL). The vial was then purged with argon and sealed. The reaction mixture was stirred at room temperature for 18 h, until the disappearance of starting material was observed, as monitored by thin layer chromatography. The reaction mixture was diluted with diethyl ether (40 mL) and washed with brine (40 mL), and the aqueous phase was then extracted with diethyl ether (2×20 mL). The combined organic layers were dried over anhydrous MgSO_4 , concentrated using a rotary evaporator, under reduced pressure (20 mmHg). The resulting residue was purified by flash column chromatography on silica gel (eluent: hexanes/ethyl acetate).

1-(2-(*p*-Tolylethynyl)phenyl)ethanone (**1c**)

This product was obtained as a light brown oil (142.9 mg, 61% yield); ^1H NMR (500 MHz, CDCl_3) δ 7.75 (dd, $J = 7.8, 1.3$ Hz, 1H), 7.62 (dd, $J = 7.7, 1.1$ Hz, 1H), 7.44–7.49 (m, 3H), 7.39 (dt, $J = 7.5, 1.2$ Hz, 1H), 7.18 (d, $J = 7.9$ Hz, 2H), 2.80 (s, 3H), 2.38 (s, 3H). The ^1H NMR spectral data are in good agreement with the literature data.¹⁷

Methyl 4-((2-acetylphenyl)ethynyl)benzoate (**1d**)

This product was obtained as a yellow solid (200.4 mg, 72% yield): mp = 118.3–119.1 °C; ^1H NMR (500 MHz, CDCl_3) δ 8.03–8.05 (m, 2H), 7.78 (d, $J = 7.8$ Hz, 1H), 7.66 (d, $J = 7.6$ Hz, 1H), 7.62 (d, $J = 8.0$ Hz, 2H), 7.51 (t, $J = 7.5$ Hz, 1H), 7.44 (t, $J = 7.6$ Hz, 1H), 3.94 (s, 3H), 2.78 (s, 3H); ^{13}C NMR (125 MHz, CDCl_3) δ 200.2, 166.7, 140.9, 134.3, 131.69, 131.67, 130.1, 129.8, 129.1, 129.0, 127.8, 121.3,

94.0, 91.5, 52.5, 30.0; IR (neat, cm^{-1}) ν 2924, 1719, 1684, 1431, 1407, 1365, 1275, 1217, 1104, 1017, 931, 876, 823, 796, 762, 729, 697; HRMS (ESI) calcd for $\text{C}_{18}\text{H}_{15}\text{O}_3$ ($\text{M} + \text{H}$)⁺ 279.1016, Found: 279.1019.

1-(2-(Naphthalen-1-ylethynyl)phenyl)ethanone (1g)

This product was obtained as a yellow solid (183.8 mg, 68% yield): mp = 92.1–93.2 °C; ^1H NMR (500 MHz, CDCl_3) δ 8.52 (d, J = 8.3 Hz, 1H), 7.88 (d, J = 8.2 Hz, 2H), 7.77–7.82 (m, 3H), 7.64 (dt, J = 8.2, 1.2 Hz, 1H), 7.52–7.57 (m, 2H), 7.43–7.50 (m, 2H), 2.83 (s, 3H); ^{13}C NMR (125 MHz, CDCl_3) δ 200.4, 140.7, 134.4, 133.5, 133.4, 131.6, 130.9, 129.5, 129.1, 128.5, 127.3, 126.8, 126.4, 125.5, 122.1, 120.8, 93.4, 30.1 (fewer ^{13}C signals were observed due to signal overlapping); IR (neat, cm^{-1}) ν 3056, 2201, 1734, 1682, 1589, 1540, 1506, 1478, 1436, 1354, 1290, 1268, 1243, 1162, 1014, 956, 879, 802, 768; HRMS (ESI) calcd for $\text{C}_{20}\text{H}_{15}\text{O}$; ($\text{M} + \text{H}$)⁺ 271.1117, Found: 271.1120.

General procedure for the gold-catalyzed intramolecular cyclization of 2-alkynylarylketones and -aldehydes.

A 4-dram vial was charged with $\text{NaAuCl}_4 \cdot 2\text{H}_2\text{O}$ (11.9 mg, 0.03 mmol), AgSbF_6 (10.3 mg, 0.03 mmol), NH_4OAc (38.5 mg, 0.5 mmol), 4-methoxypyridine *N*-oxide (75.1 mg, 0.6 mmol), 2-alkynylarylketone (**1**, 0.5 mmol) and acetonitrile (4 mL). The vial was purged with argon and sealed. The reaction mixture was stirred at 85 °C for 2 h. After cooling to room temperature, the resulting mixture was diluted with diethyl ether (15 mL) and washed with brine (15 mL). The aqueous phase was extracted with diethyl ether (2 \times 10 mL). The combined organic layers were dried over anhydrous MgSO_4 , and concentrated using a rotary evaporator, under reduced pressure (20 mmHg). The residue was purified by flash column chromatography on silica gel (eluent: hexanes/ethyl acetate).

3-Methyl-2-phenyl-1*H*-inden-1-one (2a)

This compound was obtained as a light brown oil (76.0 mg, 69% yield): ^1H NMR (500 MHz, CDCl_3) δ 7.49 (d, J = 6.9 Hz, 1H), 7.41–7.46 (m, 5H), 7.34–7.36 (m, 1H), 7.25–7.28 (m, 1H), 7.17 (d, J = 7.2 Hz, 1H), 2.32 (s, 3H); ^{13}C NMR (125 MHz, CDCl_3) δ 196.7, 154.9, 146.1, 133.8, 133.6, 131.3, 130.6, 129.7,

129.1, 128.5, 127.9, 122.3, 119.6, 12.8; HRMS (ESI) calcd for $C_{16}H_{13}O$ ($M + H$)⁺ 221.0961, Found: 221.0961. The ¹H and ¹³C NMR spectral data are in good agreement with the literature data.²³

1-Methyl-3-phenylisoquinoline (3a)

This compound was obtained as a light yellow oil (20.8 mg, 19% yield): ¹H NMR (500 MHz, CDCl₃) δ 8.14 (dd, $J = 8.2, 1.3$ Hz, 3H), 7.93 (s, 1H), 7.86 (d, $J = 8.2$ Hz, 1H), 7.68 (dt, $J = 6.8, 1.1$ Hz, 1H), 7.56–7.59 (m, 1H), 7.49–7.52 (m, 2H), 7.40 (tt, $J = 7.4, 1.8$ Hz, 1H), 3.05 (s, 3H); ¹³C NMR (125 MHz, CDCl₃) δ 158.8, 150.2, 140.0, 136.9, 130.3, 128.9, 128.5, 127.8, 127.2, 127.0, 126.8, 125.9, 115.5, 22.9; HRMS (ESI) calcd for $C_{16}H_{14}N$ ($M + H$)⁺ 220.1121, Found: 220.1119. The ¹H and ¹³C NMR spectral data are in good agreement with the literature data.²⁴

2-(4-Methoxyphenyl)-3-methyl-1H-inden-1-one (2b)

This compound was obtained as an orange solid (87.6 mg, 70% yield): mp = 106.6–107.4 °C; ¹H NMR (500 MHz, CDCl₃) δ 7.47 (d, $J = 7.0$ Hz, 1H), 7.41 (td, $J = 7.7, 1.0$ Hz, 1H), 7.37–7.39 (m, 2H), 7.25 (t, $J = 7.3$ Hz, 1H), 7.15 (d, $J = 7.2$ Hz, 1H), 6.98 (dd, $J = 6.9, 1.9$ Hz, 2H), 3.85 (s, 3H), 2.31 (s, 3H); ¹³C NMR (125 MHz, CDCl₃) δ 197.1, 159.3, 153.6, 146.3, 133.8, 133.1, 131.0, 130.6, 128.8, 123.7, 122.2, 119.4, 114.0, 55.5, 12.8; IR (neat, cm⁻¹) ν 2919, 2849, 1707, 1611, 1591, 1508, 1547, 1378, 1247, 1176, 1113, 1080, 1024, 1004, 861, 822, 795, 758; HRMS (ESI) calcd for $C_{17}H_{15}O_2$ ($M + H$)⁺ 251.1067, Found: 251.1068.

3-(4-Methoxyphenyl)-1-methylisoquinoline (3b)

This compound was obtained as a beige solid (17.5 mg, 14% yield): mp = 56.1–57.3 °C; ¹H NMR (500 MHz, CDCl₃) δ 8.08–8.11 (m, 3H), 7.84 (s, 1H), 7.82 (d, $J = 8.2$ Hz, 1H), 7.63–7.66 (m, 1H), 7.52–7.55 (m, 1H), 7.02–7.05 (m, 2H), 3.88 (s, 3H), 3.03 (s, 3H); ¹³C NMR (125 MHz, CDCl₃) δ 160.1, 158.6, 149.9, 137.0, 132.7, 130.1, 128.4, 127.6, 126.6, 126.4, 125.8, 114.30, 114.28, 55.6, 22.9; HRMS (ESI) calcd for $C_{17}H_{16}NO$ ($M + H$)⁺ 250.1226, Found: 250.1227. The ¹H and ¹³C NMR spectral data are in good agreement with the literature data.²⁵

3-Methyl-2-(p-tolyl)-1H-inden-1-one (2c)

The product was obtained as a beige solid (59.8 mg, 51% yield): mp = 80.3–82.3 °C; ^1H NMR (500 MHz, CDCl_3) δ 7.47 (d, J = 7.2 Hz, 1H), 7.40 (dt, J = 7.6, 1.0 Hz, 1H), 7.31–7.33 (m, 2H), 7.23–7.26 (m, 3H), 7.14 (d, J = 7.2 Hz, 1H), 2.39 (s, 3H), 2.30 (s, 3H); ^{13}C NMR (125 MHz, CDCl_3) δ 196.9, 154.3, 146.2, 137.8, 133.8, 133.6, 130.6, 129.6, 129.2, 129.0, 128.4, 122.3, 119.5, 21.6, 12.8; IR (neat, cm^{-1}) ν 2922, 2852, 1716, 1635, 1540, 1418, 1375, 1324, 1265, 1216, 1204, 1165, 1114, 1066, 1015, 926, 861, 806, 760; HRMS (ESI) calcd for $\text{C}_{17}\text{H}_{15}\text{O}$ ($\text{M} + \text{H}$) $^+$ 235.1117, Found: 235.1118.

1-Methyl-3-(*p*-tolyl)isoquinoline (3c)

This compound was obtained as a yellow solid (36.2 mg, 31% yield): mp = 83.2–84.8 °C; ^1H NMR (500 MHz, CDCl_3) δ 8.12 (dd, J = 8.5, 0.8 Hz, 1H), 8.04 (dd, J = 6.5, 1.6 Hz, 2H), 7.89 (s, 1H), 7.84 (d, J = 8.2 Hz, 1H), 7.64–7.67 (m, 1H), 7.54–7.57 (m, 1H), 7.31 (d, J = 8.0 Hz, 2H), 3.04 (s, 3H), 2.42 (s, 3H); ^{13}C NMR (125 MHz, CDCl_3) δ 158.7, 150.2, 138.4, 137.2, 137.0, 130.2, 129.7, 127.7, 127.0, 126.8, 126.6, 125.8, 114.9, 22.9, 21.5; HRMS (ESI) calcd for $\text{C}_{17}\text{H}_{16}\text{N}$ ($\text{M} + \text{H}$) $^+$ 234.1277, Found: 234.1276. The ^1H and ^{13}C NMR spectral data are in good agreement with the literature data.²⁶

Methyl 4-(3-methyl-1-oxo-1*H*-inden-2-yl)benzoate (2d)

This compound was obtained as a yellow solid (54.3 mg, 39% yield): mp = 158.9–159.5 °C; ^1H NMR (500 MHz, CDCl_3) δ 8.10 (d, J = 8.4 Hz, 2H), 7.49–7.51 (m, 3H), 7.44 (dt, J = 7.6, 0.9 Hz, 1H), 7.29 (t, J = 7.4 Hz, 1H), 7.20 (d, J = 7.2 Hz, 1H), 3.93 (s, 3H), 2.35 (s, 3H); ^{13}C NMR (125 MHz, CDCl_3) δ 196.0, 167.1, 156.4, 145.7, 136.1, 134.0, 132.6, 130.5, 129.7, 129.5, 129.2, 122.5, 120.0, 52.4, 13.0 (fewer ^{13}C signals were observed due to signal overlapping); IR (neat, cm^{-1}) ν 2923, 2851, 1716, 1705, 1609, 1591, 1507, 1456, 1428, 1408, 1279, 1190, 1106, 1081, 1028, 1004, 969, 946, 884, 867, 847, 820, 761, 754; HRMS (ESI) calcd for $\text{C}_{18}\text{H}_{15}\text{O}_3$ ($\text{M} + \text{H}$) $^+$ 279.1016, Found: 279.1018.

Methyl 4-(1-methylisoquinolin-3-yl)benzoate (3d)

This compound was obtained as a white solid (37.4 mg, 27% yield): mp = 125.1–126.9 °C; ^1H NMR (500 MHz, CDCl_3) δ 8.23 (dt, J = 8.7, 1.9 Hz, 2H), 8.17 (t, J = 1.6 Hz, 1H), 8.14–8.16 (m, 2H), 8.01 (s, 1H), 7.89 (d, J = 8.5 Hz, 1H), 7.69–7.72 (m, 1H), 7.60–7.63 (m, 1H), 3.96 (s, 3H), 3.05 (s, 3H); ^{13}C NMR (125 MHz, CDCl_3) δ 167.3, 159.1, 148.8, 144.3, 136.7, 130.5, 130.3, 129.8, 128.0, 127.6, 127.1,

127.0, 125.9, 116.4, 52.4, 22.9; IR (neat, cm^{-1}) ν 2922, 2852, 1738, 1711, 1617, 1565, 1456, 1443, 1414, 1383, 1364, 1332, 1317, 1272, 1230, 1217, 1103, 1015, 960, 902, 889, 863, 852, 824, 785; HRMS (ESI) calcd for $\text{C}_{18}\text{H}_{16}\text{NO}_2$ ($\text{M} + \text{H}$)⁺ 278.1176, Found: 278.1178.

5-Methoxy-2-(4-methoxyphenyl)-3-methyl-1*H*-inden-1-one (2e)

This compound was obtained as a pink solid (25.2 mg, 18% yield): mp = 119.3–120.9 °C; ¹H NMR (500 MHz, CDCl_3) δ 7.43 (d, J = 8.1 Hz, 1H), 7.37–7.39 (m, 2H), 6.97–6.98 (m, 2H), 6.70 (d, J = 2.2 Hz, 1H), 6.62 (dd, J = 7.7, 2.2 Hz, 1H), 3.88 (s, 3H), 3.84 (s, 3H), 2.27 (s, 3H); ¹³C NMR (125 MHz, CDCl_3) δ 195.9, 164.9, 159.3, 151.1, 149.1, 134.5, 131.0, 124.1, 123.9, 123.4, 114.0, 109.9, 108.6, 55.9, 55.5, 12.6; IR (neat, cm^{-1}) ν 2915, 2848, 1700, 1662, 1653, 1594, 1576, 1507, 1473, 1457, 1437, 1306, 1289, 1251, 1180, 1115, 1070, 1023, 1001, 829, 799, 780; HRMS (ESI) calcd for $\text{C}_{18}\text{H}_{16}\text{NaO}_3$ ($\text{M} + \text{Na}$)⁺ 303.0992, Found: 303.0991.

7-Methoxy-3-(4-methoxyphenyl)-1-methylisoquinoline (3e)

This compound was obtained as an orange solid (41.9 mg, 30% yield): mp = 130.5–132.2 °C; ¹H NMR (500 MHz, CDCl_3) δ 8.05–8.06 (m, 2H), 7.78 (s, 1H), 7.74 (d, J = 8.8 Hz, 1H), 7.29–7.33 (m, 2H), 7.01–7.03 (m, 2H), 3.96 (s, 3H), 3.87 (s, 3H), 2.98 (s, 3H); ¹³C NMR (125 MHz, CDCl_3) δ 159.9, 158.0, 156.9, 148.3, 132.9, 132.5, 129.2, 128.1, 127.3, 122.9, 114.3, 114.2, 103.7, 55.63, 55.55, 23.0; IR (neat, cm^{-1}) ν 2914, 2849, 1593, 1569, 1511, 1308, 1283, 1247, 1233, 1218, 1175, 1069, 1030, 876, 824, 817, 779, 752; HRMS (ESI) calcd for $\text{C}_{18}\text{H}_{18}\text{NO}_2$ ($\text{M} + \text{H}$)⁺ 280.1332, Found: 280.1335.

6-Fluoro-3-methyl-2-phenyl-1*H*-inden-1-one (2f)

This compound was obtained as a beige solid (47.7 mg, 40% yield): mp = 105.7–107.8 °C; ¹H NMR (500 MHz, CDCl_3) δ 7.43–7.46 (m, 2H), 7.39–7.41 (m, 2H), 7.34–7.37 (m, 1H), 7.21 (dd, J = 7.1, 2.4 Hz, 1H), 7.10–7.13 (m, 1H), 7.07 (dt, J = 8.8, 2.4 Hz, 1H), 2.32 (s, 3H); ¹³C NMR (125 MHz, CDCl_3) δ 195.0, 163.9 (d, $J_{\text{C-F}}$ = 250.5 Hz), 155.1, 141.4 (d, $J_{\text{C-F}}$ = 3.6 Hz), 134.0 (d, $J_{\text{C-F}}$ = 4.6 Hz), 133.0 (d, $J_{\text{C-F}}$ = 7.4 Hz), 131.1, 129.6, 128.6, 128.0, 120.7 (d, $J_{\text{C-F}}$ = 7.8 Hz), 119.0 (d, $J_{\text{C-F}}$ = 22.2 Hz), 111.1 (d, $J_{\text{C-F}}$ = 24.7 Hz), 13.0; IR (neat, cm^{-1}) ν 3061, 2927, 1703, 1618, 1604, 1473, 1428, 1373, 1355, 1330, 1309,

1271, 1220, 1203, 1098, 1076, 1027, 1013, 887, 839, 813, 793, 769; HRMS (ESI) calcd for C₁₆H₁₂FO (M + H)⁺ 239.0867, Found: 239.0871.

6-Fluoro-1-methyl-3-phenylisoquinoline (3f)

The product was obtained as a brown oil (46.3 mg, 39% yield): ¹H NMR (500 MHz, CDCl₃) δ 8.11–8.14 (m, 3H), 7.85 (s, 1H), 7.51 (t, *J* = 7.8 Hz, 2H), 7.40–7.45 (m, 2H), 7.31 (dt, *J* = 8.8, 2.5 Hz, 1H), 3.02 (s, 3H); ¹³C NMR (125 MHz, CDCl₃) δ 163.3 (d, *J*_{C-F} = 251.9 Hz), 158.7, 151.2, 139.6, 138.6 (d, *J*_{C-F} = 10.4 Hz), 129.0, 128.9 (d, *J*_{C-F} = 10.6 Hz), 128.8, 127.2, 124.0, 117.1 (d, *J*_{C-F} = 25.0 Hz), 115.0 (d, *J*_{C-F} = 5.0 Hz), 110.9 (d, *J*_{C-F} = 20.4 Hz), 23.0; IR (neat, cm⁻¹) ν 3062, 1716, 1625, 1601, 1570, 1443, 1362, 1268, 1225, 1215, 1135, 1027, 964, 882, 815, 775, 758; HRMS (ESI) calcd for C₁₆H₁₃FN (M + H)⁺ 238.1027, Found: 238.1031.

3-Methyl-2-(naphthalen-1-yl)-1*H*-inden-1-one (2g)

This product was obtained as a red solid (47.3 mg, 35% yield): mp = 100.4–102.2 °C; ¹H NMR (500 MHz, CDCl₃) δ 7.89 (t, *J* = 6.9 Hz, 2H), 7.70 (d, *J* = 8.2 Hz, 1H), 7.53–7.56 (m, 2H), 7.43–7.51 (m, 3H), 7.31–7.35 (m, 2H), 7.23 (d, *J* = 7.2 Hz, 1H), 2.11 (s, 3H); ¹³C NMR (125 MHz, CDCl₃) δ 196.7, 157.6, 145.9, 134.2, 133.9, 133.8, 132.3, 130.9, 129.3, 129.2, 128.8, 128.7, 128.3, 126.2, 126.1, 125.5, 122.5, 119.7, 13.2 (fewer ¹³C signals were observed due to signal overlapping); IR (neat, cm⁻¹) ν 3055, 2922, 285, 1701, 1596, 1459, 1378, 1314, 1171, 1085, 1026, 879, 807, 798; HRMS (ESI) calcd for C₂₀H₁₅O (M + H)⁺ 271.1117, Found: 271.1114.

1-Methyl-3-(naphthalen-1-yl)isoquinoline (3g)

This product was obtained as a beige solid (56.6 mg, 42% yield): mp = 86.5–88.0 °C; ¹H NMR (500 MHz, CDCl₃) δ 8.22 (d, *J* = 8.4 Hz, 1H), 8.13 (d, *J* = 8.5 Hz, 1H), 7.92 (d, *J* = 8.4 Hz, 2H), 7.88 (d, *J* = 8.1 Hz, 1H), 7.79 (s, 1H), 7.69–7.75 (m, 2H), 7.64–7.67 (m, 1H), 7.57–7.60 (m, 1H), 7.51 (dt, *J* = 6.8, 1.0 Hz, 1H), 7.45–7.48 (m, 1H), 3.09 (s, 3H); ¹³C NMR (125 MHz, CDCl₃) δ 158.6, 151.8, 139.1, 136.6, 134.2, 131.8, 130.4, 128.7, 128.5, 127.9, 127.7, 127.3, 126.5, 126.4, 126.2, 126.0, 125.9, 125.6, 120.2, 22.8; IR (neat, cm⁻¹) ν 3056, 2221, 1763, 1696, 1618, 1564, 1390, 1358, 1325, 1260, 1213, 1160,

1147, 1018, 964, 903, 890, 873, 855, 804, 776; HRMS (ESI) calcd for $C_{20}H_{16}N$ ($M + H$)⁺ 270.1277, Found: 270.1278.

2-(Cyclohex-1-en-1-yl)-3-methyl-1*H*-inden-1-one (2h)

This product was obtained as a light brown oil (20.2 mg, 18% yield): ¹H NMR (500 MHz, CDCl₃) δ 7.33–7.38 (m, 2H), 7.18 (t, *J* = 7.4 Hz, 1H), 7.05 (d, *J* = 6.7 Hz, 1H), 5.74–5.76 (m, 1H), 2.25–2.26 (m, 2H), 2.18–2.20 (m, 2H), 2.18 (s, 3H), 1.70–1.73 (m, 2H), 1.65–1.68 (m, 2H); ¹³C NMR (125 MHz, CDCl₃) δ 197.6, 153.2, 146.4, 136.7, 133.5, 130.8, 130.0, 129.8, 128.5, 121.9, 119.2, 28.3, 25.7, 22.9, 22.2, 12.7; IR (neat, cm⁻¹) ν 2930, 2855, 2430, 2360, 2017, 1704, 1612, 1590, 1170, 759; HRMS (ESI) calcd for $C_{16}H_{17}O$ ($M + H$)⁺ 225.1274, Found: 225.1275.

3-(Cyclohex-1-en-1-yl)-1-methylisoquinoline (3h)

This product was obtained as a brown oil (43.6 mg, 39% yield): ¹H NMR (500 MHz, CDCl₃) δ 8.06 (dd, *J* = 8.2, 0.7 Hz, 1H), 7.76 (d, *J* = 8.1 Hz, 1H), 7.60–7.63 (m, 1H), 7.48–7.52 (m, 1H), 7.44 (s, 1H), 7.06–7.07 (m, 1H), 2.96 (s, 3H), 2.55–2.58 (m, 2H), 2.31–2.34 (m, 2H), 1.83–1.87 (m, 2H), 1.69–1.74 (m, 2H); ¹³C NMR (125 MHz, CDCl₃) δ 157.8, 151.2, 136.9, 135.8, 130.0, 128.3, 127.7, 126.6, 126.3, 125.8, 113.1, 26.23, 26.19, 23.2, 22.9, 22.5; HRMS (ESI) calcd for $C_{16}H_{18}N$ ($M + H$)⁺ 224.1434, Found: 224.1430. The ¹H and ¹³C NMR spectral data are in good agreement with the literature data.²⁷

4-(3-Methyl-1-oxo-1*H*-inden-2-yl)butanenitrile (2i)

This product was obtained as an orange oil (37.0 mg, 35% yield): ¹H NMR (400 MHz, CDCl₃) δ 7.32–7.35 (m, 2H), 7.17 (td, *J* = 7.1, 0.7 Hz, 1H), 7.04 (dd, *J* = 6.9, 1.0 Hz, 1H), 2.42 (t, *J* = 7.4 Hz, 2H), 2.33 (t, *J* = 7.1 Hz, 2H), 2.15 (s, 3H), 1.85 (p, *J* = 7.2 Hz, 2H); ¹³C NMR (100 MHz, CDCl₃) δ 197.8, 155.8, 145.8, 133.6, 132.3, 130.6, 128.7, 121.8, 119.6, 119.2, 24.6, 21.8, 16.8, 11.7; IR (neat, cm⁻¹) ν 2937, 2245, 1701, 1701, 1628, 1608, 1593, 1455, 1427, 1384, 1148, 1084, 987, 918, 755; HRMS (ESI) calcd for $C_{14}H_{14}NO$ ($M + H$)⁺ 212.1070, Found: 212.1074.

4-(1-Methylisoquinolin-3-yl)butanenitrile (3i)

This product was obtained as a brown oil (16.8 mg, 16% yield): ¹H NMR (400 MHz, CDCl₃) δ 8.08 (d, *J* = 8.5 Hz, 1H), 7.74 (d, *J* = 8.3 Hz, 1H), 7.65 (t, *J* = 7.2 Hz, 1H), 7.55 (t, *J* = 7.9 Hz, 1H), 7.36 (s, 1H),

3.02 (t, $J = 7.2$ Hz, 2H), 2.93 (s, 3H), 2.39 (t, $J = 7.0$ Hz, 2H), 2.14–2.22 (m, 2H); ^{13}C NMR (100 MHz, CDCl_3) δ 158.9, 151.5, 136.6, 130.4, 127.0, 126.8, 126.3, 125.8, 119.9, 117.7, 36.5, 25.5, 22.5, 16.7; IR (neat, cm^{-1}) ν 2939, 2245, 1717, 1617, 1592, 1568, 1496, 1432, 1390, 1365, 1331, 1216, 1023, 954, 899, 828, 785; HRMS (ESI) calcd for $\text{C}_{14}\text{H}_{15}\text{N}_2$ ($\text{M} + \text{H}$) $^+$ 211.1230, Found: 211.1240.

2-(*tert*-Butyl)-3-methyl-1H-inden-1-one (2j)

This product was obtained as a yellow solid (12.0 mg, 12% yield): mp = 57.0–58.0 °C; ^1H NMR (500 MHz, CDCl_3) δ 7.32–7.35 (m, 2H), 7.16 (td, $J = 7.4$, 0.6 Hz, 1H), 7.04–7.06 (m, 1H), 2.29 (s, 3H), 1.36 (s, 9H); ^{13}C NMR (125 MHz; CDCl_3) δ 198.8, 151.2, 147.1, 140.3, 133.4, 130.7, 128.3, 121.4, 118.6, 34.1, 30.7, 13.4; HRMS (ESI) calcd for $\text{C}_{14}\text{H}_{17}\text{O}$ ($\text{M} + \text{H}$) $^+$ 201.1274, Found: 201.1276. The ^1H and ^{13}C NMR spectral data are in good agreement with the literature data.²⁸

3-(*tert*-Butyl)-1-methylisoquinoline (3j)

This product was obtained as a yellow oil (20.0 mg, 20% yield); ^1H NMR (400 MHz, CDCl_3) δ 8.06 (d, $J = 8.4$ Hz, 1H), 7.75 (d, $J = 8.2$ Hz, 1H), 7.60 (td, $J = 7.6$, 0.8 Hz, 1H), 7.50 (td, $J = 7.6$, 1.2 Hz, 1H), 7.44 (s, 1H), 2.95 (s, 3H), 1.44 (s, 9H); ^{13}C NMR (100 MHz, CDCl_3) δ 162.0, 157.4, 136.7, 129.6, 127.4, 126.1, 125.8, 125.6, 112.8, 37.1, 30.4, 22.8; IR (neat, cm^{-1}) ν 2953, 2863, 1738, 1624, 1571, 1480, 1442, 1388, 1355, 1329, 1216, 1203, 1155, 928, 877, 847, 785; HRMS (ESI) calcd for $\text{C}_{14}\text{H}_{18}\text{N}$ ($\text{M} + \text{H}$) $^+$ 200.1434, Found: 200.1434.

2,3-Diphenyl-1H-inden-1-one (2k)

This product was obtained as a red solid (55.1 mg, 39% yield): mp = 150.0–152.0 °C; ^1H NMR (500 MHz, CDCl_3) δ 7.59 (d, $J = 6.7$ Hz, 1H), 7.36–7.42 (m, 6H), 7.25–7.30 (m, 6H), 7.15 (d, $J = 7.4$ Hz, 1H); ^{13}C NMR (125 MHz, CDCl_3) δ 196.8, 155.5, 145.4, 133.7, 132.9, 132.6, 130.93, 130.91, 130.2, 129.5, 129.2, 129.0, 128.7, 128.3, 127.9, 123.2, 121.5; HRMS (ESI) calcd for $\text{C}_{21}\text{H}_{15}\text{O}$ ($\text{M} + \text{H}$) $^+$ 283.1117, Found: 283.1120. The ^1H and ^{13}C NMR spectral data are in good agreement with the literature data.²⁹

1,3-Diphenylisoquinoline (3k)

This product was obtained as an orange solid (35.2 mg, 25% yield): mp = 76.0–77.0 °C; ¹H NMR (400 MHz, CDCl₃) δ 8.21–8.23 (m, 2H), 8.13 (dd, *J* = 8.5, 0.7 Hz, 1H), 8.08 (s, 1H), 7.941 (d, *J* = 8.2 Hz, 1H), 7.81–7.83 (m, 2H), 7.69 (ddd, *J* = 8.1, 6.9, 1.1 Hz, 1H), 7.48–7.59 (m, 6H), 7.42 (dm, *J* = 7.3 Hz, 1H); ¹³C NMR (100 MHz, CDCl₃) δ 160.6, 150.4, 140.1, 139.8, 138.0, 130.4, 130.3, 128.9, 128.8, 128.7, 128.5, 127.8, 127.7, 127.3, 127.1, 126.0, 115.9; HRMS (ESI) calcd for C₂₁H₁₆N (M + H)⁺ 282.1277, Found: 282.1274. The ¹H and ¹³C NMR spectral data are in good agreement with the literature data.³⁰

2-(4-Methoxyphenyl)-3-phenyl-1*H*-inden-1-one (2l)

This compound was obtained as a red solid (56.2 mg, 36% yield): mp = 117.0–117.5 °C; ¹H NMR (500 MHz, CDCl₃) δ 7.55 (d, *J* = 7.0 Hz, 1H), 7.37–7.42 (m, 5H), 7.34 (dt, *J* = 7.4, 1.0 Hz, 1H), 7.21–7.24 (m, 3H), 7.10 (d, *J* = 7.1 Hz, 1H), 6.78–6.80 (m, 2H), 3.77 (s, 3H); ¹³C NMR (125 MHz, CDCl₃) δ 197.2, 159.4, 154.0, 145.7, 133.6, 133.2, 132.0, 131.5, 130.9, 129.3, 129.0, 128.9, 128.7, 123.2, 123.1, 121.2, 113.8, 55.4; HRMS (ESI) calcd for C₂₂H₁₇O₂ (M + H)⁺ 313.1223, Found: 313.1228. The ¹H and ¹³C NMR spectral data are in good agreement with the literature data.³¹

3-(4-Methoxyphenyl)-1-phenylisoquinoline (3l)

This compound was obtained as a yellow solid (42.0 mg, 27% yield): mp = 113.0–115.0 °C; ¹H NMR (500 MHz, CDCl₃) δ 8.17–8.19 (m, 2H), 8.11 (d, *J* = 8.6 Hz, 1H), 8.00 (s, 1H), 7.90 (d, *J* = 8.4 Hz, 1H), 7.81–7.83 (m, 2H), 7.65–7.68 (m, 1H), 7.55–7.58 (m, 2H), 7.46–7.53 (m, 2H), 7.02–7.04 (m, 2H), 3.88 (s, 3H); ¹³C NMR (125 MHz, CDCl₃) δ 160.4, 160.3, 150.1, 140.1, 138.1, 132.4, 130.4, 130.2, 128.7, 128.48, 128.46, 127.7, 127.5, 126.7, 125.6, 114.8, 114.3, 55.6; IR (neat, cm⁻¹) ν 3057, 2921, 2221, 1763, 1712, 1604, 1561, 1512, 1437, 1387, 1336, 1285, 1247, 1170, 1159, 1142, 1023, 976, 828, 806, 769; HRMS (ESI) calcd for C₂₂H₁₈NO (M + H)⁺ 312.1383, Found: 312.1384.

3-Phenylisoquinoline (3m)

This product was obtained as a light red solid (100.6 mg, 98% yield): mp = 103.1–104.2 °C; ¹H NMR (500 MHz, CDCl₃) δ 9.35 (s, 1H), 8.13–8.15 (m, 2H), 8.07 (s, 1H), 7.98 (d, *J* = 8.3 Hz, 1H), 7.86 (d, *J* = 8.1 Hz, 1H), 7.69 (t, *J* = 8.3 Hz, 1H), 7.58 (t, *J* = 7.7 Hz, 1H), 7.51–7.54 (m, 2H), 7.41–7.45 (m, 1H);

¹³C NMR (125 MHz, CDCl₃) δ 152.6, 151.4, 139.8, 136.8, 130.7, 129.0, 128.7, 127.9, 127.7, 127.23, 127.17, 127.07, 116.7; HRMS (ESI) calcd for C₁₅H₁₂N (M + H)⁺ 206.0964, Found: 206.0964. The ¹H and ¹³C NMR spectral data are in good agreement with the literature data.³⁰

7-Methyl-5-phenylthieno[2,3-*c*]pyridine (3n)

This product was obtained as a yellow oil (72.1 mg, 64% yield): ¹H NMR (500 MHz, CDCl₃) δ 8.06 (d, *J* = 7.4 Hz, 2H), 7.95 (s, 1H), 7.65 (d, *J* = 5.4 Hz, 1H), 7.49 (t, *J* = 7.3 Hz, 2H), 7.39–7.42 (m, 2H), 2.89 (s, 3H); ¹³C NMR (125 MHz, CDCl₃) δ 152.7, 152.0, 146.1, 140.2, 134.4, 131.4, 128.9, 128.4, 127.3, 124.2, 112.7, 23.9; IR (neat, cm⁻¹) ν 3058, 1575, 1545, 1480, 1439, 1392, 1377, 1348, 1117, 1086, 1065, 1032, 862, 820, 772; HRMS (ESI) calcd for C₁₄H₁₂NS (M + H)⁺ 226.0685, Found: 226.0687.

5-(4-Methoxyphenyl)-7-methylthieno[2,3-*c*]pyridine (3o)

This product was obtained as a yellow solid (108.5 mg, 85% yield): mp = 82.1–83.3 °C; ¹H NMR (500 MHz, CDCl₃) δ 8.00 (d, *J* = 9.0 Hz, 2H), 7.89 (s, 1H), 7.65 (d, *J* = 5.5 Hz, 1H), 7.39 (d, *J* = 5.1 Hz, 1H), 7.01 (d, *J* = 8.7 Hz, 2H), 3.87 (s, 3H), 2.86 (s, 3H); ¹³C NMR (125 MHz, CDCl₃) δ 160.1, 152.6, 151.8, 146.2, 133.8, 132.9, 131.4, 128.5, 124.2, 114.3, 111.9, 55.6, 24.0; IR (neat, cm⁻¹) ν 1650, 1605, 1575, 1545, 1515, 1387, 1300, 1280, 1250, 1180, 1035, 830; HRMS (ESI) calcd for C₁₅H₁₄NOS (M + H)⁺ 256.0791, Found: 256.0795.

Acknowledgments This paper is dedicated to the memory of Dr. Cliff Soll at Hunter College. We thank Queens College and the City University of New York for financial support. YL thanks South China Normal University for partial financial support.

Supporting Information Available Copies of ¹H and ¹³C NMR spectra for the new compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.

References:

-
- (1) For reviews, see: (a) Yamamoto, Y.; Gridnev, I. D.; Patil, N. T.; Jin, T. *Chem. Commun.* **2009**, 5075–5087. (b) Larock, R. C. Edited by Diederich, F.; Stang, P. J.; Tykwinski, R. R. *Acetylene Chemistry* **2005**, 51–99. (c) Bhunia, S.; Liu, R.-S. *Pure Appl. Chem.* **2012**, 84, 1749–1757.

(2) For recent examples, see: (a) Mariaule, G.; Newsome, G.; Toullec, P. Y.; Belmont, P.; Michelet, V. *Org. Lett.* **2014**, *16*, 4570-4573. (b) Yu, S.-Y.; Zhang, H.; Gao, Y.; Mo, L.; Wang, S.; Yao, Z.-J. *J. Am. Chem. Soc.* **2013**, *135*, 11402-11407. (c) Long, Y.; She, Z.; Liu, X.; Chen, Y. *J. Org. Chem.* **2013**, *78*, 2579-2588. (d) Terada, M.; Li, F.; Toda, Y. *Angew. Chem., Int. Ed.* **2014**, *53*, 235-239.

(3) For representative examples, see: (a) Barluenga, J.; Vazquez-Villa, H.; Ballesteros, A.; Gonzalez, J. M. *J. Am. Chem. Soc.* **2003**, *125*, 9028-9029. (b) Asao, N.; Nogami, T.; Takahashi, K.; Yamamoto, Y. *J. Am. Chem. Soc.* **2002**, *124*, 764-765. (c) Dyker, G.; Hildebrandt, D.; Liu, J.; Merz, K. *Angew. Chem., Int. Ed.* **2003**, *42*, 4399-4402.

(4) (a) Asao, N.; Sato, K. *Org. Lett.* **2006**, *8*, 5361-5363. (b) Patil, N. T.; Pahadi, N. K.; Yamamoto, Y. *J. Org. Chem.* **2005**, *70*, 10096-10098. (c) Tang, J.-M.; Liu, T.-A.; Liu, R.-S. *J. Org. Chem.* **2008**, *73*, 8479-8483. (d) Terada, M.; Li, F.; Toda, Y. *Angew. Chem., Int. Ed.* **2014**, *53*, 235-239. (e) Wang, J.; Zhu, H.-T.; Li, Y.-X.; Wang, L.-J.; Qiu, Y.-F.; Qiu, Z.-H.; Zhong, M.-J.; Liu, X.-Y.; Liang, Y.-M. *Org. Lett.* **2014**, *16*, 2236-2239.

(5) (a) Yue, D.; Della Ca, N.; Larock, R. C. *J. Org. Chem.* **2006**, *71*, 3381-3388. (b) Asao, N.; Chan, C. S.; Takahashi, K.; Yamamoto, Y. *Tetrahedron* **2005**, *61*, 11322-11326. (c) Patil, N. T.; Yamamoto, Y. *J. Org. Chem.* **2004**, *69*, 5139-5142.

(6) (a) Shu, X.-Z.; Zhao, S.-C.; Ji, K.-G.; Zheng, Z.-J.; Liu, X.-Y.; Liang, Y.-M. *Eur. J. Org. Chem.* **2009**, 117-122. (b) Kloeckner, U.; Finkbeiner, P.; Nachtsheim, B. J. *J. Org. Chem.* **2013**, *78*, 2751-2756. (c) Asao, N.; Kasahara, T.; Yamamoto, Y. *Angew. Chem., Int. Ed.* **2003**, *42*, 3504-3506. (d) Zhu, S.; Zhang, Z.; Huang, X.; Jiang, H.; Guo, Z. *Chem. Eur. J.* **2013**, *19*, 4695-4700. (e) Bhunia, S.; Wang, K.-C.; Liu, R.-S. *Angew. Chem., Int. Ed.* **2008**, *47*, 5063-5066. (f) Asao, N.; Takahashi, K.; Lee, S.; Kasahara, T.; Yamamoto, Y. *J. Am. Chem. Soc.* **2002**, *124*, 12650-12651.

(7) (a) Tovar, J. D.; Swager, T. M. *J. Org. Chem.* **1999**, *64*, 6499-6504. (b) Dell'Acqua, M.; Abbiati, G.; Rossi, E. *Synlett* **2010**, 2672-2676. (c) Alfonsi, M.; Dell'Acqua, M.; Facchetti, D.; Arcadi, A.; Abbiati, G.; Rossi, E. *Eur. J. Org. Chem.* **2009**, 2852-2862.

(8) (a) Yang, D.; Burugupalli, S.; Daniel, D.; Chen, Y. *J. Org. Chem.* **2012**, *77*, 4466-4472. (b) Reddy, V.; Jadhav, A. S.; Anand, R. V. *Org. Biomol. Chem.* **2015**, *13*, 3732-3741.

(9) (a) Asao, N.; Iso, K.; Salprima, Y. S. *Org. Lett.* **2006**, *8*, 4149-4151. (b) Ding, Q.; Wu, J. *Org. Lett.* **2007**, *9*, 4959-4962.

(10) (a) Jiang, D.; Herndon, J. W. *Org. Lett.* **2000**, *2*, 1267-1269. (b) Luo, Y.; Herndon, J. W.; Cervantes-Lee, F. *J. Am. Chem. Soc.* **2003**, *125*, 12720-12721.

(11) Godet, T.; Vaxelaire, C.; Michel, C.; Milet, A.; Belmont, P. *Chem. Eur. J.* **2007**, *13*, 5632-5641.

(12) For review on gold catalysis, see: (a) Fürstner, A.; Davies, P. W. *Angew. Chem., Int. Ed.* **2007**, *46*, 3410-3449. (b) Hashmi, A. S. K. *Chem. Rev.* **2007**, *107*, 3180-3211.

(13) The control experiments showed that in the absence of 4-methoxypyridine *N*-oxide cyclization of substrate **1a** led to the formation of a mixture of **2a** and **3a** in 48% and 10% yield respectively after 4 hours. About 30% of **1a** was recovered from the reaction mixture. No enhancement of the reaction conversion was observed even when the reaction time was extended to 8 hours.

(14) For discussion on the role of pyridine *N*-oxides in gold catalysis, see review: Zhang, L. *Acc. Chem. Res.* **2014**, *47*, 877-888.

- (15) (a) Sonogashira, K.; Tohda, Y.; Hagihara, N. *Tetrahedron Lett.* **1975**, *16*, 4467-4470. (b) Sonogashira, K. *Metal-Catalyzed Cross-Coupling Reactions*, F. Diederich; P. J. Stang, Eds.; Wiley-VCH: Weinheim, 1998; pp 203–229.
- (16) Dong, C.; Liao, Z.; Xu, X.; Zhou, H. *J. Heterocyclic Chem.* **2014**, *51*, 1282-1286.
- (17) Gorelsky, S. I.; Gevorgyan, V. *Angew. Chem. Int. Ed.* **2011**, *50*, 2342-2345.
- (18) Mukherjee, A.; Liu, R.-S. *Org. Lett.* **2011**, *13*, 660-663.
- (19) Chen, X.; Jin, J.; Wang, N.; Lu, P.; Wang, Y. *Eur. J. Org. Chem.* **2012**, 824-830.
- (20) Zhang, H.; Karasawa, T.; Yamada, H.; Wakamiya, A.; Yamaguchi, S. *Org. Lett.* **2009**, *11*, 3076-3079.
- (21) Mariaule, G.; Newsome, G.; Toullec, P. Y.; Belmont, P.; Michelet, V. *Org. Lett.* **2014**, *16*, 4570-4573.
- (22) Shen, H.; Fu, J.; Gong, J.; Yang, Z. *Org. Lett.* **2014**, *16*, 5588-5591.
- (23) Morimoto, T.; Yamasaki, K.; Hirano, A.; Tsutsumi, K.; Kagawa, N.; Kakiuchi, K.; Harada, Y.; Fukumoto, Y.; Chatani, N.; Nishioka, T. *Org. Lett.* **2009**, *11*, 1777-1780.
- (24) Zhang, Z.-W.; Lin, A.; Yang, J. *J. Org. Chem.* **2014**, *79*, 7041-7050.
- (25) Arambasic, M.; Hooper, J. F.; Willis, M. C. *Org. Lett.* **2013**, *15*, 5162-5165.
- (26) Zhao, M.-N.; Ren, Z.-H.; Wang, Y.-Y.; Guan, Z.-H. *Chem. Commun.* **2012**, 48, 8105-8107.
- (27) Hwang, S.; Lee, Y.; Lee, P. H.; Shin, S. *Tetrahedron Lett.* **2009**, *50*, 2305-2308.
- (28) Liebeskind, L. S.; South, M. S. *J. Org. Chem.* **1980**, *45*, 5426-5429.
- (29) Kuninobu, Y.; Matsuki, T.; Takai, K. *Org. Lett.* **2010**, *12*, 2948-2950.
- (30) Niu, Y.-N.; Yan, Z.-Y.; Gao, G.-L.; Wang, H.-L.; Shu, X.-Z.; Ji, K.-G.; Liang, Y.-M. *J. Org. Chem.* **2009**, *74*, 2893-2896.
- (31) Gao, K.; Yoshikai, N. *Chem. Commun.* **2012**, 48, 4305-4307.