

Cite this: *Org. Biomol. Chem.*, 2012, **10**, 5137

www.rsc.org/obc

PAPER

The effects of solvent on switchable stereoselectivity: copper-catalyzed asymmetric conjugate additions using D_2 -symmetric biphenyl phosphoramidite ligands†

Han Yu, Fang Xie, Zhenni Ma, Yangang Liu and Wanbin Zhang*

Received 14th April 2012, Accepted 6th May 2012

DOI: 10.1039/c2ob25730k

A highly enantioselective copper-catalyzed conjugate addition of diethylzinc to acyclic aromatic enones was developed with phosphoramidite ligands bearing a D_2 -symmetric biphenyl backbone. This type of reaction demonstrated that toluene and THF solvents can completely reverse the absolute configuration of the products, thus simplifying the process of accessing either enantiomer (*S*: 92% ee, 94% yield; *R*: 99% ee, 96% yield).

Introduction

Catalytic asymmetric syntheses of both enantiomers of a chiral compound are of significant interest to the chemical community.¹ Commonly, in a catalytic asymmetric protocol, both isomers of a chiral ligand are required to access each enantiomer of chiral compounds. Alternatives include using different central metals in the presence of an identical ligand² or changing reaction parameters (temperature, additives, and solvents *etc.*).^{3,4}

A number of effective chiral ligand systems have been developed for Cu-catalyzed enantioselective addition reactions of diethylzinc to structurally diverse enones.^{5–8} Among the successful chiral ligands, BINOL-,⁵ TADDOL-,⁶ and BIPOL-derived phosphoramidites⁷ have shown remarkable enantioselectivities in the reactions of diethylzinc to enones. Very recently, our group reported new types of biphenyl phosphoramidite ligands **L1–5** (Fig. 1) containing a D_2 -symmetric backbone. These ligands demonstrated excellent activity and selectivity in Cu-catalyzed asymmetric 1,4-addition reactions of diethylzinc to α,β -unsaturated ketones and nitroalkenes. Their excellent chiral environment and their ability to be easily modified at the 3,3',5,5'-position of the biphenyl backbone, makes them excellent ligands.⁹ Interestingly, we found that the substituents at the 3,3',5,5'-position of the biphenyl backbone of the ligands have a dramatic effect on the stereocontrol in Cu-catalyzed asymmetric 1,4-addition reactions.^{9c,d}

In addition, Cu-catalyzed asymmetric additions to acyclic aromatic α,β -unsaturated enones such as chalcone, can provide




Fig. 1 Phosphoramidite ligands with a D_2 -symmetric biphenyl backbone.

valuable compounds for use as intermediates or building blocks in organic chemistry.¹⁰ To explore the behaviour of the phosphoramidite ligands with a D_2 -symmetric biphenyl backbone, the ligands were used in asymmetric conjugate additions of diethylzinc to α,β -unsaturated acyclic aromatic enones. Both enantiomeric products were obtained by changing reaction parameters, a method of which has not been significantly reported.³

Results and discussion

Trans-L2 provided excellent enantioselectivities for the asymmetric conjugate addition of diethylzinc to α,β -unsaturated carbonyl substrates and nitroalkenes.^{9a–c} The conjugate addition of diethylzinc to chalcone **1a** was performed in the presence of 2 mol% of *trans-L2* and 1 mol% of copper salt in toluene, at a temperature of -40 °C (Table 1). Cu(OAc)₂·H₂O was found to be the most suitable catalyst precursor because of its high reactivity and enantioselectivity (entries 1–10). With Cu(OAc)₂·H₂O as the catalyst precursor and *trans-L2* as the ligand, we investigated the solvent effect (entries 10–16) and discovered that toluene provided the most promising results. When Et₂O, methyl *t*-butyl ether (MTBE) or diisopropyl ether (DIPE) were used as

School of Chemistry and Chemical Engineering, Shanghai Jiao Tong University, 800 Dongchuan Road, Shanghai 200240, P. R. China.
E-mail: wanbin@sjtu.edu.cn; Fax: +86-21-5474-3265;
Tel: +86-21-5474-3265

† Electronic supplementary information (ESI) available. See DOI: 10.1039/c2ob25730k

Table 1 Screening of reaction conditions^a

| Entry | Cu salt | Solvent | T (°C) | Yield ^b (%) | ee ^c (%) |
|-----------------|--|-------------------|--------|------------------------|---------------------|
| 1 | Cu(I)(MeCN) ₄ PF ₆ | Toluene | -40 | 96 | 68 (R) |
| 2 | Cu(II)(OTf) ₂ | Toluene | -40 | 95 | 78 (R) |
| 3 | Cu(II)(acac) ₂ | Toluene | -40 | 98 | 65 (R) |
| 4 | Cu(II)Br ₂ | Toluene | -40 | 95 | 53 (R) |
| 5 | Cu(II)ClO ₄ ·6H ₂ O | Toluene | -40 | 93 | 79 (R) |
| 6 | Cu(I)(MeCN) ₄ BF ₄ | Toluene | -40 | 90 | 76 (R) |
| 7 | Cu(I)(MeCN) ₄ PF ₆ | Toluene | -40 | 91 | 73 (R) |
| 8 | Cu(I)TC | Toluene | -40 | 99 | 67 (R) |
| 9 | Cu(I)(C ₆ H ₆)OTf | Toluene | -40 | 94 | 66 (R) |
| 10 | Cu(II)(OAc) ₂ ·H ₂ O | Toluene | -40 | 96 | 92 (R) |
| 11 | Cu(II)(OAc) ₂ ·H ₂ O | PhCF ₃ | -40 | 95 | 73 (R) |
| 12 | Cu(II)(OAc) ₂ ·H ₂ O | Et ₂ O | -40 | 99 | 39 (R) |
| 13 | Cu(II)(OAc) ₂ ·H ₂ O | MTBE | -40 | 97 | 42 (R) |
| 14 | Cu(II)(OAc) ₂ ·H ₂ O | DIPE | -40 | 97 | 43 (R) |
| 15 | Cu(II)(OAc) ₂ ·H ₂ O | DCM | -40 | 93 | 70 (R) |
| 16 | Cu(II)(OAc) ₂ ·H ₂ O | THF | -40 | 96 | 81 (S) |
| 17 | Cu(II)(OAc) ₂ ·H ₂ O | THF | -30 | 97 | 75 (S) |
| 18 | Cu(II)(OAc) ₂ ·H ₂ O | THF | -50 | 95 | 92 (S) |
| 19 | Cu(II)(OAc) ₂ ·H ₂ O | THF | -60 | 93 | 91 (S) |
| 20 ^d | Cu(II)(OAc) ₂ ·H ₂ O | THF | -78 | 93 | 82 (S) |
| 21 ^e | Cu(II)(OAc) ₂ ·H ₂ O | Toluene | -30 | 98 | 81 (R) |
| 22 | Cu(II)(OAc) ₂ ·H ₂ O | Toluene | -50 | 96 | 99 (R) |
| 23 ^d | Cu(II)(OAc) ₂ ·H ₂ O | Toluene | -78 | 92 | 83 (R) |

^a 1 mol% Cu salt, 2 mol% *trans*-L2, 1.5 eq. ZnEt₂. ^b Yield of the isolated product. ^c Determined by HPLC, Chiralcel AD-H column. The absolute configuration was determined by comparison with literature data. ^d The reaction was stirred for 20 h. ^e The reaction was stirred for 12 h.

the reaction solvent, excellent reactivity but low enantioselectivities for the conjugate addition reactions were obtained (Table 1, entries 12–14). To our surprise, when THF was employed as the reaction solvent, the other enantiomer was obtained in excellent yield and enantioselectivity (entry 16), most probably due to THF's strong coordinating ability. These results indicated that the ether solvent possessed appropriate coordinating ability to be involved in the catalytic cycle, acting as a hemilabile ligand and altering the structure of the catalyst.

Since both enantiomers could be obtained by using toluene and THF as solvents, different temperatures were screened to optimize the reaction conditions. Reducing the reaction temperature from -30 °C to -50 °C provided excellent yields and increased enantioselectivity (entries 18 and 22). The highest enantioselectivity was observed at -50 °C in both toluene and THF. Furthermore, decreasing the temperature to -78 °C resulted in a slight decrease in yield, and an obvious drop in enantioselectivity in toluene or THF at this temperature (entries 20 and 23).

Encouraged by the results described above, reactions involving several phosphoramidite ligands containing a D₂-symmetric biphenyl backbone were investigated to identify the most efficient ligand for use in toluene and THF (Table 2).

Ligand screening revealed that the phosphoramidite ligands L1–2 in Fig. 1 were effective for this transformation (91–96% yield, *R*-2a: 52–99% ee, *S*-2a: 86–92%). Both *cis*- and *trans*-L1

Table 2 Screening of ligands^a

| Entry | L | Solvent | Yield ^b (%) | ee ^c (%) |
|-------|------------------|---------|------------------------|---------------------|
| 1 | <i>cis</i> -L1 | Toluene | 92 | 77 (R) |
| 2 | <i>cis</i> -L1 | THF | 91 | 52 (R) |
| 3 | <i>trans</i> -L1 | Toluene | 93 | 79 (R) |
| 4 | <i>trans</i> -L1 | THF | 94 | 57 (R) |
| 5 | <i>cis</i> -L2 | Toluene | 96 | 93 (R) |
| 6 | <i>cis</i> -L2 | THF | 93 | 86 (S) |
| 7 | <i>trans</i> -L2 | Toluene | 96 | 99 (R) |
| 8 | <i>trans</i> -L2 | THF | 92 | 92 (S) |

^a 1 mol% Cu salt, 2 mol% L, 1.5 eq. ZnEt₂. ^b Yield of the isolated product. ^c Determined by HPLC, Chiralcel AD-H column. The absolute configuration was determined by comparison with literature data.

lacking substituents at the 3,3',5,5'-position, gave *R* configuration products (entries 1–4). Substituents at the 3,3',5,5'-position of the biphenyl backbone of the ligands showed significant influence on catalysis. No matter whether the 3,3',5,5'-position contained substituents or not, both *cis*- and *trans*-L provided *R* configuration product when using toluene as the solvent. *Trans*-L gave higher enantioselectivity compared to its *cis*-L counterpart (entries 1, 3, 5 and 7), and up to 99% ee was obtained with methyl-substituted *trans*-L2 (entries 1, 3, 5 and 7). When THF was used as the reaction solvent, ligands *cis*-L2 and *trans*-L2, containing substituents on the biphenyl backbone, all provided *S* configuration product (entries 6 and 8). Up to 92% ee was obtained with methyl-substituted *trans*-L2 (entry 8). *Trans*-L3–5 were also examined, however no improvement was observed with these ligands (ESI†).

With toluene and THF used as a reversal solvent pair, we expected to induce the formation of both absolute configurations in the products. A wide range of aromatic enones were examined to investigate substrate scope and the reversal in stereoselectivity (Table 3).

To our delight, a complete switch in stereoselectivity was observed for all of the substrates. The introduction of a methyl group at the *ortho*-, *meta*- and *para*-position on the phenyl ring R¹ or R² of substrate 1, led to a drop in enantioselectivity when using both solvents, which may be a result of steric hindrance (entries 1–4 and 8–10). When a methoxy group was introduced at the *para*-position of the phenyl ring R¹ or R², decreased enantioselectivity was observed with the exception of substrate 1e, whose enantioselectivity was similar to that of substrate 1a (entries 1, 5 and 11). Addition of an electron-withdrawing substituent (such as *para*-bromo group) to the phenyl ring R¹ or addition of a *para*-chloro group to the phenyl ring R², was detrimental to enantioselectivity when compared to substrates lacking substituted groups on the phenyl ring (entries 1, 6 and 12). Neither the electron-donating nor the electron-withdrawing groups at the *para*-position of the phenyl ring of R¹ or R² improved the enantioselectivity of this reaction. Furthermore, replacement of R¹ or R² with 2-naphthyl groups resulted in a decrease in enantioselectivity when the reaction was performed

Table 3 Substrate scope^a

| Entry | Substrate | R ¹ | R ² | Solvent | Yield ^b (%) | ee ^c (%) |
|-----------------|-----------|--|--|---------|------------------------|---------------------|
| 1 | 1a | Ph | Ph | THF | 94 | 92 (<i>S</i>) |
| | | | | Toluene | 96 | 99 (<i>R</i>) |
| 2 | 1b | <i>o</i> -MeC ₆ H ₄ | Ph | THF | 91 | 80 (<i>S</i>) |
| | | | | Toluene | 92 | 71 (<i>R</i>) |
| 3 | 1c | <i>m</i> -MeC ₆ H ₄ | Ph | THF | 92 | 78 (<i>S</i>) |
| | | | | Toluene | 94 | 85 (<i>R</i>) |
| 4 | 1d | <i>p</i> -MeC ₆ H ₄ | Ph | THF | 93 | 86 (<i>S</i>) |
| | | | | Toluene | 95 | 86 (<i>R</i>) |
| 5 | 1e | <i>p</i> -MeOC ₆ H ₄ | Ph | THF | 93 | 92 (<i>S</i>) |
| | | | | Toluene | 96 | 90 (<i>R</i>) |
| 6 | 1f | <i>p</i> -BrC ₆ H ₄ | Ph | THF | 95 | 86 (<i>S</i>) |
| | | | | Toluene | 96 | 96 (<i>R</i>) |
| 7 | 1g | 2-naphthyl | Ph | THF | 90 | 87 (<i>S</i>) |
| | | | | Toluene | 92 | 91 (<i>R</i>) |
| 8 | 1h | Ph | <i>o</i> -MeC ₆ H ₄ | THF | 90 | 60 (<i>S</i>) |
| | | | | Toluene | 91 | 79 (<i>R</i>) |
| 9 | 1i | Ph | <i>m</i> -MeC ₆ H ₄ | THF | 92 | 90 (<i>S</i>) |
| | | | | Toluene | 93 | 75 (<i>R</i>) |
| 10 | 1j | Ph | <i>p</i> -MeC ₆ H ₄ | THF | 91 | 83 (<i>S</i>) |
| | | | | Toluene | 95 | 91 (<i>R</i>) |
| 11 | 1k | Ph | <i>p</i> -MeOC ₆ H ₄ | THF | 93 | 65 (<i>S</i>) |
| | | | | Toluene | 97 | 92 (<i>R</i>) |
| 12 | 1l | Ph | <i>p</i> -ClC ₆ H ₄ | THF | 92 | 81 (<i>S</i>) |
| | | | | Toluene | 93 | 92 (<i>R</i>) |
| 13 | 1m | Ph | 2-Naphthyl | THF | 93 | 87 (<i>S</i>) |
| | | | | Toluene | 94 | 92 (<i>R</i>) |
| 14 ^d | 1n | Ph | Me | THF | 93 | 86 (<i>S</i>) |
| | | | | Toluene | 90 | 97 (<i>R</i>) |

^a 1 mol% Cu salt, 2 mol% *trans*-**L2**, 1.5 eq. ZnEt₂. ^b Yield of the isolated product. ^c Determined by HPLC, Chiralcel AD-H column. The absolute configuration was determined by comparison with literature data. ^d The ee value (toluene as solvent) is cited from our previous report for comparison.^{9a}

in THF or toluene (entries 1, 7 and 13). The above results suggested that the effect of steric hindrance and the electron-donating/withdrawing properties of the *para*-position substituent on the phenyl ring R¹ or R², decreased enantioselectivity. We also performed the reaction using a typical acyclic enone, benzalacetone (**1n**). Using toluene and THF as a solvent pair, products were obtained with high enantioselectivities and reversed absolute configurations (entry 14).

In order to explain the reversal phenomenon, a stereochemical pathway has been proposed according to our experimental results (Fig. 2). As a copper ion coordinates to one phosphorus on the ligand,¹¹ the other phosphorus remains relatively far away from the reaction center. Both the *cis* and *trans* configurations of the ligands therefore have little effect on the absolute configuration of the products. The steric repulsion between the substituents at the 3,3'- or 5,5'-position of the biphenyl backbone with the aryl groups of the substrate or THF, may determine the stereochemical outcome of the 1,4-addition. For the transition states involving THF as a solvent in this reaction, the oxygen atom of THF coordinates with the zinc ion. Steric repulsion between the substituent at the 3,3',5,5'-position of the biphenyl backbone of ligand **L2** and THF in **L2-THF-TS2**, is larger than that in **L2-THF-TS1** between the substituent at the 3,3',5,5'-




Fig. 2 Proposed stereochemical pathway showing the facial coordination of the copper atom with the enone substrate.

position of the biphenyl backbone of ligand **L2** and the aryl group of the substrate. This leads to the 1,4-adduct with a *S* isomer configuration. However, when toluene is used as a solvent, there are no coordination effects. The steric repulsion between the aryl group of ligand **L2** and the aryl group of the substrate in transition state **L2-TS2**, is smaller than that between

the substituent at the 3,3',5,5'-position of the biphenyl backbone, and the aryl group of the substrate in **L2-TS1**. This leads to the 1,4-adduct with the *R* isomer configuration.

Conclusions

In summary, we have developed a highly enantioselective copper-catalyzed conjugate addition of diethylzinc to acyclic aromatic enones with phosphoramidite ligands bearing a *D*₂-symmetric biphenyl backbone. These types of reactions demonstrated that toluene and THF solvents can be used to completely reverse the absolute configuration of the products, thus simplifying the process of accessing either enantiomer (*S*: 92% ee, 94% yield; *R*: 99% ee, 96% yield).

Experimental section

General details

Commercially available reagents were used without further purification other than those described below. Substrates **1** were prepared according to modified methods of reported procedures.¹⁰ All air- and moisture-sensitive manipulations were carried out with standard Schlenk techniques under nitrogen. Toluene, PhCF₃, DMF, THF, Et₂O, DIPE, MTBE, and dichloromethane were dried according to published procedures. Column chromatography was run on silica gel (100–200 mesh). ¹H NMR (400 MHz) spectra and ¹³C NMR (100 MHz) spectra were obtained on a Varian MERCURY plus-400 spectrometer. HRMS was performed on a Micromass LCT TM at the Instrumental Analysis Center of Shanghai Jiao Tong University.

General procedures for conjugate addition

A flame dried Schlenk tube was charged with Cu(OAc)₂·H₂O 1.0 mg (0.005 mmol) and two equivalents of ligand (0.010 mmol) under nitrogen, and the mixture was dissolved in dry toluene (1.5 mL), resulting in a colorless solution. The solution was stirred at 25 °C for 2 h and then cooled to –50 °C. The substrate **1** (0.50 mmol dissolved in 1.0 mL dry toluene) was then added dropwise over 3 min. The solution was stirred for 5 min at –50 °C and gradually turned to light yellow. Diethylzinc (0.75 mmol, 0.75 mL of 1 M sol. in hexane) was added dropwise over 3 min. The reaction mixture was stirred at –50 °C for 16 h and monitored by TLC until full conversion of product was observed. The reaction mixture was quenched with aqueous saturated NH₄Cl and extracted with ethyl acetate (5 mL × 2). The organic extracts were combined, concentrated and the residue was purified by silica gel column chromatography to afford the product **2**. Enantiomeric excess was determined by chiral HPLC. Melting points of racemic product **2** was measured with SGW X-4 micro melting point apparatus.

1,3-Diphenylpentan-1-one (**2a**)¹⁰

A white solid, mp: 55–56 °C; using THF as solvent: *S*-**2a**, 92% ee, 94% yield; using toluene as solvent: *R*-**2a**, 99% ee, 96% yield; ¹H NMR (400 MHz, CDCl₃) δ 7.92–7.87 (m, 2H),

7.55–7.50 (m, 1H), 7.45–7.40 (m, 2H), 7.31–7.15 (m, 5H), 3.31–3.22 (m, 3H), 1.85–1.74 (m, 1H), 1.70–1.57 (m, 1H), 0.80 (t, *J* = 7.4 Hz, 3H). HPLC conditions: Chiralcel AD-H column, *i*-PrOH–hexane 2 : 98, flow rate: 1.0 mL min^{–1}, UV detection at 254 nm, *t*_s = 13.1 min, *t*_R = 15.1 min.

1-Phenyl-3-*o*-tolylpentan-1-one (**2b**)

A white solid, mp: 56–57 °C; using THF as solvent: *S*-**2b**, 80% ee, 91% yield; using toluene as solvent: *R*-**2b**, 71% ee, 92% yield; ¹H NMR (400 MHz, CDCl₃) δ 7.92–7.88 (m, 2H), 7.55–7.50 (m, 1H), 7.45–7.40 (m, 2H), 7.22–7.05 (m, 4H), 3.66–3.55 (m, 1H), 3.30–3.20 (m, 2H), 2.38 (s, 3H), 1.85–1.57 (m, 2H), 0.80 (t, *J* = 7.4 Hz, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 198.2, 163.5, 145.0, 130.6, 128.5, 127.8, 126.4, 113.8, 55.7, 45.5, 43.4, 29.5, 12.3. HRMS calcd for C₁₇H₁₉O [M + Na]⁺ 253.1592, found 253.1599. ESI-MS: 253. HPLC conditions: Chiralcel OJ-H column, *i*-PrOH–hexane 2 : 98, flow rate: 1.0 mL min^{–1}, UV detection at 254 nm, *t*_s = 14.4 min, *t*_R = 16.8 min.

1-Phenyl-3-*m*-tolylpentan-1-one (**2c**)¹⁰

A white solid, mp: 57–58 °C; using THF as solvent: *S*-**2c**, 78% ee, 92% yield; using toluene as solvent: *R*-**2c**, 85% ee, 94% yield; ¹H NMR (400 MHz, CDCl₃) δ 7.93–7.90 (m, 2H), 7.56–7.50 (m, 1H), 7.46–7.40 (m, 2H), 7.21–6.80 (m, 4H), 3.31–3.15 (m, 3H), 2.34 (s, 3H), 1.84–1.60 (m, 2H), 0.80 (t, *J* = 7.4 Hz, 3H). HPLC conditions: Chiralcel AD-H column, *i*-PrOH–hexane 2 : 98, flow rate: 1.0 mL min^{–1}, UV detection at 254 nm, *t*_s = 16.6 min, *t*_R = 18.2 min.

1-Phenyl-3-*p*-tolylpentan-1-one (**2d**)¹⁰

A white solid, mp: 59–60 °C; using THF as solvent: *S*-**2d**, 86% ee, 93% yield; using toluene as solvent: *R*-**2d**, 86% ee, 95% yield; ¹H NMR (400 MHz, CDCl₃) δ 7.93–7.88 (m, 2H), 7.56–7.51 (m, 1H), 7.46–7.40 (m, 2H), 7.16–7.05 (m, 4H), 3.30–3.15 (m, 3H), 2.34 (s, 3H), 1.84–1.60 (m, 2H), 0.80 (t, *J* = 7.4 Hz, 3H). HPLC conditions: Chiralcel AD-H column, *i*-PrOH–hexane 2 : 98, flow rate: 1.0 mL min^{–1}, UV detection at 254 nm, *t*_s = 18.0 min, *t*_R = 23.1 min.

3-(4-Methoxyphenyl)-1-phenylpentan-1-one (**2e**)¹⁰

A white solid, mp: 62–63 °C; using THF as solvent: *S*-**2e**, 92% ee, 93% yield; using toluene as solvent: *R*-**2e**, 90% ee, 96% yield; ¹H NMR (400 MHz, CDCl₃) δ 7.90–7.86 (m, 2H), 7.57–7.51 (m, 1H), 7.46–7.39 (m, 2H), 7.15–7.11 (m, 2H), 6.85–6.80 (m, 2H), 3.77 (s, 3H), 3.30–3.17 (m, 3H), 1.83–1.58 (m, 2H), 0.80 (t, *J* = 7.4 Hz, 3H). HPLC conditions: Chiralcel AD-H column, *i*-PrOH–hexane 2 : 98, flow rate: 1.0 mL min^{–1}, UV detection at 254 nm, *t*_s = 39.6 min, *t*_R = 65.3 min.

3-(4-Bromophenyl)-1-phenylpentan-1-one (**2f**)¹⁰

A white solid, mp: 65–66 °C; using THF as solvent: *S*-**2f**, 86% ee, 95% yield; using toluene as solvent: *R*-**2f**, 96% ee,

96% yield; $^1\text{H NMR}$ (400 MHz, CDCl_3) δ 7.93–7.88 (m, 2H), 7.55–7.50 (m, 1H), 7.45–7.39 (m, 4H), 7.13–7.08 (m, 2H), 3.26–3.18 (s, 3H), 1.82–1.70 (m, 1H), 1.66–1.54 (m, 1H), 0.79 (t, $J = 7.2$ Hz, 3H). HPLC conditions: Chiralcel AD-H column, i-PrOH–hexane 2 : 98, flow rate: 1.0 mL min^{-1} , UV detection at 254 nm, $t_s = 25.3$ min, $t_R = 35.3$ min.

3-(Naphthalen-2-yl)-1-phenylpentan-1-one (2g)¹⁰

A white solid, mp: 55–56 °C; using THF as solvent: *S*-**2g**, 87% ee, 90% yield; using toluene as solvent: *R*-**2g**, 91% ee, 92% yield; $^1\text{H NMR}$ (400 MHz, CDCl_3) δ 7.93–7.89 (m, 2H), 7.80–7.76 (m, 3H), 7.66 (s, 1H), 7.56–7.50 (m, 1H), 7.47–7.38 (m, 5H), 3.45–3.10 (m, 3H), 1.95–1.64 (m, 2H), 0.80 (t, $J = 7.4$ Hz, 3H). HPLC conditions: Chiralcel AD-H column, i-PrOH–hexane 2 : 98, flow rate: 1.0 mL min^{-1} , UV detection at 254 nm, $t_s = 28.1$ min, $t_R = 38.9$ min.

3-Phenyl-1-*o*-tolylpentan-1-one (2h)

A white solid, mp: 57–58 °C; using THF as solvent: *S*-**2h**, 60% ee, 90% yield; using toluene as solvent: *R*-**2h**, 79% ee, 91% yield; $^1\text{H NMR}$ (400 MHz, CDCl_3) δ 7.48 (d, $J = 8.0$ Hz, 1H), 7.35–7.14 (m, 8H), 3.21–3.12 (m, 3H), 2.27 (s, 3H), 1.82–1.56 (m, 2H), 0.81 (t, $J = 7.4$ Hz, 3H); $^{13}\text{C NMR}$ (100 MHz, CDCl_3) δ 204.2, 144.5, 138.8, 137.7, 131.9, 131.1, 128.6, 128.2, 127.9, 126.5, 125.7, 48.8, 43.6, 29.6, 20.9, 12.3. HRMS calcd for $\text{C}_{17}\text{H}_{19}\text{O}$ [$\text{M} + \text{Na}$]⁺ 253.1592, found 253.1586. ESI-MS: 253. HPLC conditions: Chiralcel AD-H column, i-PrOH–hexane 2 : 98, flow rate: 1.0 mL min^{-1} , UV detection at 254 nm, $t_s = 13.0$ min, $t_R = 16.5$ min.

3-Phenyl-1-*m*-tolylpentan-1-one (2i)

A white solid, mp: 60–61 °C; using THF as solvent: *S*-**2i**, 90% ee, 92% yield; using toluene as solvent: *R*-**2i**, 75% ee, 93% yield; $^1\text{H NMR}$ (400 MHz, CDCl_3) δ 7.72–7.68 (m, 2H), 7.36–7.15 (m, 7H), 3.30–3.20 (m, 3H), 2.38 (s, 3H), 1.84–1.56 (m, 2H), 0.80 (t, $J = 7.4$ Hz, 3H); $^{13}\text{C NMR}$ (100 MHz, CDCl_3) δ 199.7, 145.1, 138.4, 137.5, 133.8, 128.8, 128.3, 127.8, 126.4, 125.4, 45.6, 43.2, 29.3, 21.5, 12.3. HRMS calcd for $\text{C}_{17}\text{H}_{19}\text{O}$ [$\text{M} + \text{Na}$]⁺ 253.1592, found 253.1588. ESI-MS: 253. HPLC conditions: Chiralcel AD-H column, i-PrOH–hexane 2 : 98, flow rate: 1.0 mL min^{-1} , UV detection at 254 nm, $t_s = 15.6$ min, $t_R = 18.4$ min.

3-Phenyl-1-*p*-tolylpentan-1-one (2j)¹⁰

A white solid, mp: 59–60 °C; using THF as solvent: *S*-**2j**, 83% ee, 91% yield; using toluene as solvent: *R*-**2j**, 91% ee, 95% yield; $^1\text{H NMR}$ (400 MHz, CDCl_3) δ 7.81 (d, $J = 8.0$ Hz, 2H), 7.31–7.15 (m, 7H), 3.27–3.19 (m, 3H), 2.39 (s, 3H), 1.84–1.55 (m, 2H), 0.80 (t, $J = 7.4$ Hz, 3H). HPLC conditions: Chiralcel AD-H column, i-PrOH–hexane 2 : 98, flow rate: 1.0 mL min^{-1} , UV detection at 254 nm, $t_s = 26.6$ min, $t_R = 35.2$ min.

1-(4-Methoxyphenyl)-3-phenylpentan-1-one (2k)¹⁰

A white solid, mp: 60–62 °C; using THF as solvent: *S*-**2k**, 65% ee, 93% yield; using toluene as solvent: *R*-**2k**, 92% ee, 97% yield; $^1\text{H NMR}$ (400 MHz, CDCl_3) δ 7.90–7.87 (d, $J = 12$ Hz, 2H), 7.31–7.14 (m, 5H), 6.92–6.87 (m, 2H), 3.85 (s, 3H), 3.26–3.17 (m, 3H), 1.82–1.56 (m, 2H), 0.80 (t, $J = 7.4$ Hz, 3H). HPLC conditions: Chiralcel AD-H column, i-PrOH–hexane 2 : 98, flow rate: 1.0 mL min^{-1} , UV detection at 254 nm, $t_s = 57.2$ min, $t_R = 77.6$ min.

1-(4-Chlorophenyl)-3-phenylpentan-1-one (2l)¹⁰

A white solid, mp: 65–67 °C; using THF as solvent: *S*-**2l**, 81% ee, 92% yield; using toluene as solvent: *R*-**2l**, 92% ee, 93% yield; $^1\text{H NMR}$ (400 MHz, CDCl_3) δ 7.85–7.80 (m, 2H), 7.41–7.37 (m, 2H), 7.31–7.25 (m, 2H), 7.23–7.15 (m, 3H), 3.27–3.17 (m, 3H), 1.81–1.60 (m, 2H), 0.80 (t, $J = 7.4$ Hz, 3H). HPLC conditions: Chiralcel AD-H column, i-PrOH–hexane 2 : 98, flow rate: 1.0 mL min^{-1} , UV detection at 254 nm, $t_s = 24.3$ min, $t_R = 29.1$ min.

1-(Naphthalen-2-yl)-3-phenylpentan-1-one (2m)¹⁰

A white solid, mp: 58–60 °C; using THF as solvent: *S*-**2m**, 87% ee, 93% yield; using toluene as solvent: *R*-**2m**, 92% ee, 94% yield; $^1\text{H NMR}$ (400 MHz, CDCl_3) δ 8.42 (s, 1H), 8.00 (d, $J = 8.6$, 1H), 7.95 (d, $J = 8.0$, 1H), 7.87 (d, $J = 8.6$, 2H), 7.60 (t, $J = 7.9$, 1H), 7.55 (t, $J = 7.4$, 1H), 7.34–7.28 (m, 5H), 3.42 (t, $J = 6.9$, 2H), 3.38–3.29 (m, 1H), 1.92–1.82 (m, 1H), 1.77–1.66 (m, 1H), 0.86 (t, $J = 7.4$, 3H). HPLC conditions: Chiralcel AD-H column, i-PrOH–hexane 2 : 98, flow rate: 1.0 mL min^{-1} , UV detection at 254 nm, $t_s = 33.4$ min, $t_R = 41.6$ min.

4-Phenylhexan-2-one (2n)¹⁰

Colorless oil; using THF as solvent: *S*-**2n**, 85% ee, 80% yield; using toluene as solvent: *R*-**2n**, 98% ee, 82% yield; $^1\text{H NMR}$ (400 MHz, CDCl_3) δ 7.33–7.15 (m, 5H), 3.05–3.00 (m, 1H), 2.72 (d, $J = 7.2$ Hz, 2H), 2.03 (s, 3H), 1.69–1.58 (m, 2H), 0.78 (t, $J = 7.4$ Hz, 3H). GC, Supelco γ -DEX-120 column (30 m \times 0.25 mm, ID): 115 °C; Retention time: $t_s = 29$ min, $t_R = 30$ min.

Acknowledgements

This work was supported by the National Nature Science Foundation of China (No. 20872091 and 20972095), Science and Technology Commission of Shanghai Municipality (No. 10dz1910105), Nippon Chemical Industrial Co., Ltd and Shanghai Jiao Tong University. We thank Professor T. Imamoto and Dr M. Sugiya for helpful discussions and the Instrumental Analysis Center of Shanghai Jiao Tong University.

Notes and references

- (a) M. P. Sibi and M. Liu, *Curr. Org. Chem.*, 2001, **5**, 719; (b) G. Zanoni, F. Castronovo, M. Franzini, G. Vidari and E. Giannini,

- Chem. Soc. Rev.*, 2003, **32**, 115; (c) T. Tanaka and M. Hayashi, *Synthesis*, 2008, 3361; (d) M. Bartok, *Chem. Rev.*, 2010, **110**, 1663.
- 2 For selected examples, see: (a) D. Du, M. S. F. Lu, T. Fang and J. X. Xu, *J. Org. Chem.*, 2005, **70**, 3712; (b) A. Frollander and C. Moberg, *Org. Lett.*, 2007, **9**, 1371; (c) K. Y. Spangler and C. Wolf, *Org. Lett.*, 2009, **11**, 4724; (d) H. Y. Kim, H. J. Shih, W. E. Knabe and K. Oh, *Angew. Chem., Int. Ed.*, 2009, **48**, 7420; (e) Z. Wang, Z. G. Yang, D. H. Chen, X. H. Liu, L. L. Lin and X. M. Feng, *Angew. Chem., Int. Ed.*, 2011, **50**, 4928.
- 3 For selected examples, see: (a) S. Kobayashi and H. Ishitani, *J. Am. Chem. Soc.*, 1994, **116**, 4083; (b) J. P. G. Seerden, M. M. Kuypers and M. H. W. Scheeren, *Tetrahedron: Asymmetry*, 1995, **6**, 1441; (c) K. V. Gothelf, R. G. Hazell and K. A. Jorgensen, *J. Org. Chem.*, 1998, **63**, 5483; (d) D. A. Evans, M. C. Kozlowski, J. A. Murry, C. S. Burgey, K. R. Campos, B. T. Connell and R. J. Staples, *J. Am. Chem. Soc.*, 1999, **121**, 669; (e) M. Kawamura and S. Kobayashi, *Tetrahedron Lett.*, 1999, **40**, 3213; (f) R. Kuwano, M. Sawamura and Y. Ito, *Bull. Chem. Soc. Jpn.*, 2000, **73**, 2571; (g) M. C. Perry, X. Cui, M. T. Powell, D. R. Hou, J. H. Reibenspies and K. Burgess, *J. Am. Chem. Soc.*, 2003, **125**, 113; (h) C. P. Casey, S. C. Martins and M. A. Fagan, *J. Am. Chem. Soc.*, 2004, **126**, 5585; (i) J. Zhou, M. C. Ye, Z. Z. Huang and Y. J. Tang, *J. Org. Chem.*, 2004, **69**, 1309; (j) T. Yamamoto, T. Yamada, Y. Nagata and M. Suginome, *J. Am. Chem. Soc.*, 2010, **132**, 7899; (k) T. Yamamoto, Y. Akai, Y. Nagata and M. Suginome, *Angew. Chem., Int. Ed.*, 2011, **50**, 8844.
- 4 Selected examples using tunable ligands, see: (a) S. Kobayashi and M. Horibe, *J. Am. Chem. Soc.*, 1994, **116**, 9805; (b) W. Zeng, G. Y. Chen, Y. G. Zhou and Y. X. Li, *J. Am. Chem. Soc.*, 2007, **129**, 750; (c) W. Q. Wu, Q. Peng, D. X. Dong, X. L. Hou and Y. D. Wu, *J. Am. Chem. Soc.*, 2008, **130**, 9717; (d) X. X. Yan, Q. Peng, Q. Li, K. Zhang, J. Yao, X. L. Hou and Y. D. Wu, *J. Am. Chem. Soc.*, 2008, **130**, 14362; (e) S. Mouri, Z. H. Chen, H. Mitsunuma, M. Furutachi, S. Matsunaga and M. Shibasaki, *J. Am. Chem. Soc.*, 2010, **132**, 1255; (f) Y. L. Liu, D. J. Shang, X. Zhou, Y. Zhu, L. L. Lin, X. H. Liu and X. M. Feng, *Org. Lett.*, 2010, **12**, 180; Selected examples based on other methods in metal-catalyzed reactions, see: (g) A. B. Zaitsev and H. Adolfsson, *Org. Lett.*, 2006, **8**, 5129; (h) J. D. Huber and J. L. Leighton, *J. Am. Chem. Soc.*, 2007, **129**, 14552; (i) M. I. Burguete, M. Collado, J. Escorihuela and S. V. Luis, *Angew. Chem., Int. Ed.*, 2007, **46**, 9002; For selected examples of stereo inversion in organocatalytic reactions, see: (j) S. E. Denmark, X. Su and Y. Nishigaichi, *J. Am. Chem. Soc.*, 1998, **120**, 12990; (k) K. Nakayama and K. Maruoka, *J. Am. Chem. Soc.*, 2008, **130**, 17666; (l) N. Li, X. H. Chen, J. Song, S. W. Luo, W. Fan and L. Z. Gong, *J. Am. Chem. Soc.*, 2009, **131**, 15301; (m) D. G. Blackmond, A. Moran, M. Hughes and A. Armstrong, *J. Am. Chem. Soc.*, 2010, **132**, 7598; (n) M. Messerer and H. Wennemers, *Synlett*, 2011, 499; (o) J. B. Wang and B. L. Feringa, *Science*, 2011, **331**, 1429; (p) K. Endo, D. Hamada, S. Yakeishi, M. Ogawa and T. Shibata, *Org. Lett.*, 2012, **14**, 2342.
- 5 (a) A. H. M. de Vries, A. Meetsma and B. L. Feringa, *Angew. Chem., Int. Ed. Engl.*, 1996, **35**, 2374; (b) B. L. Feringa, M. Pineschi, L. A. Arnold, R. Imbos and A. H. M. de Vries, *Angew. Chem., Int. Ed. Engl.*, 1997, **36**, 2620; (c) F. Zhang and A. S. C. Chan, *Tetrahedron: Asymmetry*, 1998, **9**, 1179; (d) N. Sewald and V. Wendisch, *Tetrahedron: Asymmetry*, 1998, **9**, 1341; (e) A. Alexakis and C. Benhaim, *Org. Lett.*, 2000, **2**, 2579; (f) A. Mandoli, L. A. Arnold, A. H. M. de Vries, P. Salvadori and B. L. Feringa, *Tetrahedron: Asymmetry*, 2001, **12**, 1929; (g) A. Duursma, A. J. Minnaard and B. L. Feringa, *Tetrahedron*, 2002, **58**, 5773; (h) A. Duursma, A. J. Minnaard and B. L. Feringa, *J. Am. Chem. Soc.*, 2003, **125**, 3700; (i) E. Fillion and A. Wilsily, *J. Am. Chem. Soc.*, 2006, **128**, 2774; (j) R. G. Sebesta, M. Pizzuti, A. J. Minnaard and B. L. Feringa, *Adv. Synth. Catal.*, 2007, **349**, 1931; (k) D. Zou, Z. Duan, X. Hu and Z. Zheng, *Tetrahedron: Asymmetry*, 2009, **20**, 235; (l) A. Wilsily and E. Fillion, *J. Org. Chem.*, 2009, **74**, 8583; (m) M. Fañanás-Mastral and B. L. Feringa, *J. Am. Chem. Soc.*, 2010, **132**, 13152.
- 6 (a) E. Keller, J. Maurer, R. Naasz, T. Schader, A. Meetsma and B. L. Feringa, *Tetrahedron: Asymmetry*, 1998, **9**, 2409; (b) A. Alexakis, J. Vastra, J. Burton, C. Benhaim and P. Mangeney, *Tetrahedron Lett.*, 1998, **39**, 7869; (c) A. Alexakis, J. Burton, J. Vastra, C. Benhaim, X. Fournieux, A. van den Heuvel, J. M. Leveque, F. Maze and S. Rosset, *Eur. J. Org. Chem.*, 2000, 4011; (d) A. Mandoli, L. A. Arnold, A. H. M. de Vries, P. Salvadori and B. L. Feringa, *Tetrahedron: Asymmetry*, 2001, **12**, 1929.
- 7 (a) A. Alexakis, S. Rosset, J. Allamand, S. March, F. Guillen and C. Benhaim, *Synlett*, 2001, 1375; (b) A. Alexakis, C. Benhaim, S. Rosset and M. Humam, *J. Am. Chem. Soc.*, 2002, **124**, 5262; (c) Z. Hu, V. C. Vassar, H. Choi and I. Ojima, *Proc. Natl. Acad. Sci. U. S. A.*, 2004, **101**, 5411; (d) H. Choi and I. Ojima, *Org. Lett.*, 2004, **6**, 2689; (e) A. Alexakis, D. Polet, S. Rosset and S. March, *J. Org. Chem.*, 2004, **69**, 5660; (f) M. d'Augustin, L. Palais and A. Alexakis, *Angew. Chem., Int. Ed.*, 2005, **44**, 1376; (g) E. Raluy, O. Pàmies, M. Diéguez, K. Biswas, S. Rosset and A. Alexakis, *Tetrahedron: Asymmetry*, 2009, **20**, 2167; (h) L. Palais, L. Babel, A. Quintard, S. Belot and A. Alexakis, *Org. Lett.*, 2010, **12**, 1988.
- 8 For other selected papers see: (a) Y. Yamanoi and T. Imamoto, *J. Org. Chem.*, 1999, **64**, 2988; (b) I. H. Escher and A. Pfaltz, *Tetrahedron*, 2000, **56**, 2879; (c) P. Müller, P. Nury and G. Bernardinelli, *Helv. Chim. Acta*, 2000, **83**, 843; (d) C. G. Arena, G. Calabrò, G. Franciò and F. Faraone, *Tetrahedron: Asymmetry*, 2000, **11**, 2387; (e) O. Huttenloch, J. Spieler and H. Waldmann, *Chem.-Eur. J.*, 2000, **6**, 671; (f) S. J. Degrado, H. Mizutani and A. H. Hoveyda, *J. Am. Chem. Soc.*, 2001, **123**, 755; (g) O. Huttenloch, E. Laxman and H. Waldmann, *Chem. Commun.*, 2002, 673; (h) R. Shintani and G. C. Fu, *Org. Lett.*, 2002, **4**, 3699; (i) H. Mizutani, S. J. Degrado and A. H. Hoveyda, *J. Am. Chem. Soc.*, 2002, **124**, 779; (j) S. J. Degrado, H. Mizutani and A. H. Hoveyda, *J. Am. Chem. Soc.*, 2002, **124**, 13362; (k) H. Zhou, W. Wang, Y. Fu, J. Xie, W.-J. Shi, L. X. Wang and Q.-L. Zhou, *J. Org. Chem.*, 2003, **68**, 1582; (l) T. Morimoto, N. Mochizuki and M. Suzuki, *Tetrahedron Lett.*, 2004, **45**, 5717; (m) A. P. Duncan and J. L. Leighton, *Org. Lett.*, 2004, **6**, 4117; (n) W. Zhang, C. Wang, W. Gao and X. Zhang, *Tetrahedron Lett.*, 2005, **46**, 6087; (o) L.-T. Liu, M.-C. Wang, W.-X. Zhao, Y.-L. Zhou and X.-D. Wang, *Tetrahedron: Asymmetry*, 2006, **17**, 136; (p) F. Boeda, D. Rix, H. Cravier and C. M. Mauduit, *Tetrahedron: Asymmetry*, 2006, **17**, 2726; (q) K. Wakabayashi, K. Aikawa, S. Kawauchi and K. Mikami, *J. Am. Chem. Soc.*, 2008, **130**, 5012; (r) D. B. Biradar and H.-M. Gau, *Tetrahedron: Asymmetry*, 2008, **19**, 733; (s) M. A. Calter and J. Wang, *Org. Lett.*, 2009, **11**, 2205; (t) X. M. Feng, Z. H. Dong, J. H. Feng, X. A. Fu, X. H. Liu and L. L. Lin, *Chem.-Eur. J.*, 2011, **17**, 1118; (u) S.-F. Zhu, B. Xu, G.-P. Wang and Q.-L. Zhou, *J. Am. Chem. Soc.*, 2012, **134**, 436.
- 9 (a) H. Zhang, F. Fang, F. Xie, H. Yu, G. Yang and W. Zhang, *Tetrahedron Lett.*, 2010, **51**, 3119; (b) F. Fang, H. Zhang, F. Xie, G. Yang and W. Zhang, *Tetrahedron*, 2010, **66**, 3593; (c) B. Yang, F. Xie, H. Yu, K. Shen, Z. Ma and W. Zhang, *Tetrahedron Lett.*, 2011, **67**, 6197; (d) H. Yu, F. Xie, Z. Ma and W. Zhang, *Adv. Synth. Catal.*, 2012, DOI: 10.1002/adsc.201101007.
- 10 (a) X. Hu, H. Chen and X. Zhang, *Angew. Chem., Int. Ed.*, 1999, **38**, 3518; (b) M. Shi, C. Wang and W. Zhang, *Chem.-Eur. J.*, 2004, **10**, 5507; (c) L. Liu and M. Wang, *Tetrahedron: Asymmetry*, 2006, **17**, 136; (d) M. Fañanás-Mastral and B. L. Feringa, *J. Am. Chem. Soc.*, 2010, **132**, 13152; (e) W. Zhang and M. Shi, *Synlett*, 2007, 19; (f) A. Isleyen and Z. Dogan, *Tetrahedron: Asymmetry*, 2007, **18**, 679; (g) Y. Xie, H. Huang and X. Hu, *Tetrahedron: Asymmetry*, 2009, **20**, 1425.
- 11 (a) H. Zhang and R. M. Gschwind, *Angew. Chem., Int. Ed.*, 2006, **45**, 6391; (b) H. Zhang and R. M. Gschwind, *Chem.-Eur. J.*, 2007, **13**, 6691.