

Organozinc Reagents in DMSO Solvent: Remarkable Promotion of S_N2' Reaction for Allene Synthesis

Koji Kobayashi,[†] Hiroshi Naka,[†] Andrew E. H. Wheatley,[‡] and Yoshinori Kondo^{*,†}

Graduate School of Pharmaceutical Sciences, Tohoku University, Aramaki-aza Aoba 6-3, Aoba-ku, Sendai 980-8578, Japan, and Department of Chemistry, University of Cambridge, Lensfield Road, Cambridge, U.K. CB2 1EW

ykondo@mail.pharm.tohoku.ac.jp

Received June 2, 2008

ABSTRACT



The S_N2' reaction of propargyl mesylates with organozinc reagents was dramatically improved in DMSO solvent, and the stereoselective conversion of chiral substrates was successfully achieved using LiCl-free diorganozinc without the loss of optical purity.

Organozinc reagents have been widely used as soft nucleophiles in organic synthesis, where the control of the nucleophilicity is highly important for many selective transformations.¹ Various methods have been utilized for the promotion of reactivity, and conventionally ate complexation or transmetalation has been employed.¹ Meanwhile, allenes have gained attraction as unique functionalities in natural product chemistry² and as important building blocks in transition metal catalyzed cyclization chemistry.³ In connection with our recent studies on the reactivity control of organozinc reagents,⁴ we focused our interest on solvent effects for promoting the reactivity of organozinc

reagents. In this paper, we report allene synthesis using the dramatic effect of DMSO on the reaction of propargyl electrophiles with organozinc reagents.

Solvent is one of the most important parameters in organic reaction. The solvent effects on organometallic reactions have been studied energetically,⁵ not only in terms of the improvement of solubility of metal salt but also in terms of the change of the oligomeric complex structures,^{5a,b} acceleration of reaction,^{5c} and influencing the reaction selectivity.⁵ These effects alter according to the physicochemical nature of the solvents (polarity, donor number, and acceptor number, etc.). Therefore, it is considered to be an attractive research subject to investigate the relationship between solvents and reactivity

[†] Tohoku University.

[‡] University of Cambridge.

(1) Knochel, P.; Jones, P. *Organozinc Reagents*; Oxford University Press: Oxford, 1999.

(2) (a) Hoffmann-Röder, A.; Krause, N. *Angew. Chem., Int. Ed.* **2004**, *43*, 1196–1216. (b) Hoffmann-Röder, A.; Krause, N. *Angew. Chem., Int. Ed.* **2002**, *41*, 2933–2935, and references therein.

(3) (a) Tsukamoto, H.; Matsumoto, T.; Kondo, Y. *J. Am. Chem. Soc.* **2008**, *130*, 388–389. (b) Volz, F.; Krause, N. *Org. Biomol. Chem.* **2007**, *5*, 1519–1521. (c) Song, M.; Montgomery, J. *Tetrahedron* **2005**, *61*, 11440–11448. (d) Bates, R. W.; Satcharoen, V. *Chem. Soc. Rev.* **2002**, *31*, 12–21.

(4) (a) Kondo, Y.; Morey, J. V.; Morgan, J. C.; Naka, H.; Nobuto, D.; Raithby, P. R.; Uchiyama, M.; Wheatley, A. E. H. *J. Am. Chem. Soc.* **2007**, *129*, 12734–12738. (b) Mulvey, R. E.; Mongin, F.; Uchiyama, M.; Kondo, Y. *Angew. Chem., Int. Ed.* **2007**, *46*, 3802–3824. (c) Ueno, M.; Wheatley, A. E. H.; Kondo, Y. *Chem. Commun.* **2006**, 3549–3550. (d) Kobayashi, K.; Ueno, M.; Naka, H.; Kondo, Y. *Chem. Commun.* **2008**, DOI:10.1039/B807169A.

(5) (a) Rutherford, J. L.; Collum, D. B. *J. Am. Chem. Soc.* **2001**, *123*, 199–202. (b) Majewski, M.; Nowak, P. *Tetrahedron Lett.* **1998**, *39*, 1661–1664. (c) Winkle, M. R.; Ronald, R. C. *J. Org. Chem.* **1982**, *47*, 2101–2108. (d) Shibli, A.; Varghese, J. P.; Knochel, P.; Marek, I. *Synlett* **2001**, 818–820.

(6) (a) Buncel, E.; Wilson, H. *Adv. Phys. Org. Chem.* **1977**, *14*, 133–202. (b) Martin, D.; Weise, A.; Niclas, H. *J. Angew. Chem., Int. Ed. Engl.* **1967**, *6*, 318–334.

(7) Some unique behaviors of the organozinc compounds in or with DMSO have been reported. (a) Okhlobystin, Y. O.; Zakharkin, I. L. *J. Organomet. Chem.* **1965**, 257–258. (b) Masao, T.; Hiroyuki, K.; Teruo, O.; Hiroshi, K. *Nippon Kagaku Kaishi* **1976**, 356–358. (c) Berk, S. C.; Yeh, M. C. P.; Jeong, N.; Knochel, P. *Organometallics* **1990**, *9*, 3053–3064. (d) Grondin, J.; Sebban, M.; Vottero, P. *J. Organomet. Chem.* **1989**, *362*, 237–242. (e) Hlavinka, M.; Hagadorn, J. R. *Organometallics* **2007**, *26*, 4105–4108.

for organometallic reagents. DMSO has long been regarded as an excellent polar solvent for a wide variety of organic reactions⁶ but has been hitherto thought of as an unlikely solvent for organozinc reactions.⁷

The synthesis of substituted allenes has been extensively studied,⁸ and the S_N2' reaction of propargyl electrophiles is one of the important methods by which to achieve this. However, the use of organozinc reagent as nucleophiles for this S_N2' reaction⁹ has been limited due to their poor reactivity, and this has led to organozinc reagents being employed for other modes of allene synthesis instead.¹⁰ To clarify solvent effects on the reactivity of organozinc reagents, the S_N2' reaction of propargyl electrophiles with organozinc reagents was investigated using DMSO as a solvent.

In our preliminary experiment, the reaction of diethylzinc and 3-phenylpropargyl mesylate (**1**) in various organic solvents was carried out at room temperature (Table 1). In nonpolar

(E_T)¹¹ of CH_3CN (**46**) is largely similar to those of DMF (43.8) and DMSO (45.0). On the other hand, the donor numbers (DN)¹² of DMF (26.6) and DMSO (29.8) are higher values than that of CH_3CN (14.1). The highest DN solvent gave the most excellent result.

To examine further the scope and limitations of the procedure, the reactions of other substrates were investigated. 3-Phenyl-1-phenethylpropargyl mesylate (**1a**) and 3-butyl-1-phenethylpropargyl mesylate (**1b**) reacted smoothly to give corresponding 1,1,3-trisubstituted allenes **2a** and **2b** (Table 2, entries 1 and 2). Various substituents on the aromatic ring of the 3-arylpropargyl mesylates (**2c–h**) were also compatible with this allene synthesis. Substitution with an electron-donating methoxy group is acceptable to the reaction, and the corresponding allene **2c** was obtained in good yield (Table 2, entry 3). However, yields of allenes **2d,e** were moderate when methyl-substituted substrates **1d,e** were utilized (Table 2, entries 4 and 5). The tolerance of the alkoxycarbonyl group (**1f**), the cyano group (**1g**), and the bromo substituent (**1h**) to the allene synthesis is considered to be synthetically important (Table 2, entries 6–8). The reaction of the substrate with terminal alkyne **1i** was also found to be successful, and 1,3-disubstituted allene **2i** was selectively obtained in high yield (Table 2, entry 9). It is noteworthy that no deprotonation of the terminal alkyne was observed during the reaction.^{10a,b} The allenylsilyl compound **2j** was obtained from the trimethylsilylated substrate **1j** using similar reaction conditions (Table 2, entry 10). The high functional group tolerance of this allene synthesis using organozinc reagents in DMSO is considered to offer significant advantages over the use of conventional organometallic nucleophiles.^{8b} Lastly, di-*n*-butylzinc, prepared from *n*-butyllithium and zinc chloride, can be used for the reaction, allowing the corresponding allene **2k** to be obtained quantitatively (Table 2, entry 11).

A functionalized alkylzinc halide has also been employed for this allene synthesis, with ethoxycarbonylpropylzinc bromide prepared using Rieke's zinc¹³ having been reacted with **1a** in DMSO to give **3** in 57% yield (Scheme 1).

To examine the stereochemistry of allene formation, an enantiomerically pure substrate was employed for the reaction. The chiral propargyl mesylate **5**¹⁴ was prepared from (*R*)-(+)-3-butyn-2-ol and reacted with LiCl-free dibutylzinc¹⁵ to give chiral allene **6**. The absolute configuration of **6** was determined as (*R*) by converting **6** to the known chiral furanone **8**,¹⁶ the

Table 1. Solvent Effect on S_N2' Reaction of **1**^a

entry	solvent	E_T	DN	time (h)	yield (%) ^b
1	CH_2Cl_2	41.1	-	24	n.r.
2	toluene	33.9	-	24	n.r.
3	THF	37.4	20.0	24	n.r.
4	CH_3CN	46.0	14.1	24	n.r.
5	DMF	43.8	26.6	24	43
6	DMSO	45.0	29.8	18	90

^a All reactions were carried out using 3-phenylpropargyl mesylate (**1**) (0.3 mmol) and diethylzinc (0.6 mmol) at rt. ^b Isolated yield.

solvents such as CH_2Cl_2 and toluene, the reaction did not proceed at all (Table 1, entries 1 and 2). In polar solvents such as THF and CH_3CN , failure to react was again observed (Table 1, entries 3 and 4). To our surprise, in polar and strongly donor solvents such as DMF and DMSO, the S_N2' reaction was dramatically promoted, and the target allene **2** was obtained (Table 1, entries 5 and 6). The best result (90%) was observed when DMSO was employed as solvent. The solvent polarity

(8) (a) Krause, N.; Hashmi, A. S. K., *Modern Allene Chemistry*; Wiley-VCH, Weinheim, Germany, 2004; Vols. 1, 2. (b) Krause, N.; Hoffmann-Roder, A. *Tetrahedron* **2004**, *60*, 11671–11694. (c) Brummond, K. M.; DeForrest, *Synthesis* **2007**, 795–818. (d) Nakamura, H.; Kamakura, T.; Onagi, S. *Org. Lett.* **2006**, *8*, 2095–2098. (e) Ogasawara, M.; Nagano, T.; Hayashi, T. *J. Org. Chem.* **2005**, *70*, 5764–5767.

(9) On the use of organozinc reagents for S_N2' reactions of allyl electrophiles in the presence of activators, see: (a) Arai, M.; Kawasuji, T.; Nakamura, E. *Chem. Lett.* **1993**, 357–360. (b) Kacprzyński, M. A.; Hoveyda, A. H. *J. Am. Chem. Soc.* **2004**, *126*, 10676–10681. (c) Merphy, K. E.; Hoveyda, A. H. *Org. Lett.* **2005**, *7*, 1255–1258.

(10) (a) Katsuhira, T.; Harada, T.; Maejima, K.; Osada, A.; Oku, A. *J. Org. Chem.* **1993**, *58*, 6166–6188. (b) Harada, T.; Katsuhira, T.; Osada, A.; Iwazaki, K.; Maejima, K.; Oku, A. *J. Am. Chem. Soc.* **1996**, *118*, 11377–11390. (c) Varghese, J. P.; Knochel, P.; Marek, I. *Org. Lett.* **2000**, *2*, 2849–2852.

(11) Reichardt, C. *Angew. Chem., Int. Ed.* **1979**, *18*, 98–100.

(12) Gutmann, V.; Wychera, E. *Inorg. Nucl. Chem. Lett.* **1966**, 257–260.


(13) Zhu, L.; Wehmeyer, R. M.; Rieke, R. D. *J. Org. Chem.* **1991**, *56*, 1445–1453.












(14) Grée, D.; Grée, R. *Tetrahedron Lett.* **2007**, *48*, 5435–5438.

(15) Côté, A.; Charette, A. B. *J. Am. Chem. Soc.* **2008**, *130*, 2771–2773. When $\text{Zn}(\text{nBu})_2$ containing LiCl was used for the S_N2' reaction, complete racemization was observed.

(16) (a) Marshall, J. A.; Wolf, M. A.; Wallace, E. M. *J. Org. Chem.* **1997**, *62*, 367–371. (b) Corbera, J.; Font, J.; Monsalvatje, M.; Ortuno, R. M. *J. Org. Chem.* **1988**, *53*, 4393–4595. (c) He, Y.-T.; Yang, H.-N.; Yao, Z.-J. *Tetrahedron* **2002**, *58*, 8805–8810.

(17) (a) Elsevier, C. J.; Vermeer, P. J. *J. Org. Chem.* **1989**, *54*, 3726–3730. (b) Marchall, J. A.; Robinson, E. D.; Zapata, A. *J. Org. Chem.* **1989**, *54*, 5854–5855. (c) Myers, A. G.; Condroski, K. R. *J. Am. Chem. Soc.* **1995**, *117*, 3057–3083. (d) Laux, M.; Krause, N.; Koop, U. *Synlett* **1996**, 87–88. (e) Brummond, K. M.; Kerekes, A. D.; Wan, H. *J. Org. Chem.* **2002**, *67*, 5156–5163.


Table 2. Addition to Other Propargyl Mesylates^a


entry	product	time(h)	yield (%) ^b
1		24	90
2		24	69
3		24	74
4		24	44
5		24	59
6		22	88
7		16	71
8		16	74
9		24	86
10		24	59
11		24	100


^a All reactions were carried out using 3-phenylpropargyl mesylate (**1**) (0.3 mmol) and diethylzinc (0.6 mmol) at rt. ^b Isolated yield.

configuration of which was assigned as (*R*) with an optical purity of 98% ee (Scheme 2). This fact indicates that the butyl group attacks the triple bond with anti stereochemistry in this S_N2'-type reaction and that selectivity is the same as with the conventional organocopper reactions.¹⁷

¹H NMR spectra of dimethylzinc in different solvents were measured for the preliminary estimation of the activation

Scheme 1. Addition of Functionalized Alkylzinc Halide

mechanism. The methyl signals of dimethylzinc in DMSO and THF at room temperature were observed as sharp singlets at −0.99 ppm (DMSO) and −0.84 ppm (THF), respectively

Scheme 2. Reaction of Chiral Propargyl Mesylate

(see Supporting Information). This result suggests that dimethylzinc acts more carbanionically in DMSO by virtue of the stronger solvent donation to zinc. It is known that the coordination of donors to alkylzinc compounds causes higher-field shifts of the alkyl protons in the ¹H NMR spectra.¹⁸

In conclusion, the S_N2' reaction of propargyl electrophiles with organozinc reagents was dramatically improved in DMSO solvent, and stereoselective conversion of chiral substrate was successfully achieved using LiCl-free diorganozinc. Further investigations on the scope and limitation of these DMSO-promoted organozinc reactions and associated mechanistic studies are underway.

Acknowledgment. This work was partly supported by the Grant-in Aid for Scientific Research on Priority Areas “Advanced Molecular Transformations of Carbon Resources” (No.19020005) and “Synergistic Effects for Creation of Functional Molecules” (No.20036007) and other Research Grants (No.19390002) from the Ministry of Education, Science, Sports and Culture, Japan, and a grant from Yamada Science Foundation.

Supporting Information Available: Experimental procedures and characterization data for all new compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.

OL801249W

(18) (a) Uchiyama, M.; Koike, M.; Kameda, M.; Kondo, Y.; Sakamoto, T. *J. Am. Chem. Soc.* **1996**, *118*, 8793–8794. (b) Ishimori, M.; Tsuruta, T. *Kogyo Kagaku Zasshi* **1967**, *70*, 152–158.