


On the Synthesis and Reactivity of 2,3-Dihydropyrrolo[1,2-*a*]quinazolin-5(1*H*)-ones

Charlotte L. Sutherland^{a,b,1}Steven V. Ley^a^a Department of Chemistry, University of Cambridge, Lensfield Road, Cambridge, CB2 1EW, U.K.^b Cancer Research UK Cambridge Institute, Li Ka Shing Centre, University of Cambridge, Cambridge, CB2 0RE, U.K.

c.sutherland@imperial.ac.uk

Dedicated with respect and best wishes to our friend Dieter Enders on the occasion of his 70th birthday

Received: 15.06.2016

Accepted after revision: 08.07.2016

Published online: 24.08.2016

DOI: 10.1055/s-0035-1562792; Art ID: ss-2016-z0435-op

Abstract An improved, scalable synthetic route to the quinazolinone natural product 2,3-dihydropyrrolo[1,2-*a*]quinazolin-5(1*H*)-one is reported. The applicability of this method to analogue synthesis and the synthesis of related natural products is explored. Finally, reactivity of the scaffold to a variety of electrophilic reagents, generating products stereoselectively, is reported.

Key words quinazolinone, alkaloid, 2,3-dihydropyrrolo[1,2-*a*]quinazolin-5(1*H*)-one, deoxyvasicinone

Quinazolinone alkaloids are commonly found as the structural basis of natural products and pharmaceuticals that display biological activities in diverse areas, including cancer, CNS systems, inflammation, and hypertension.^{2,3} Many quinazolinone alkaloids contain the quinazolinone ring fused with pyrrole or piperidine rings.⁴ These are often derivatives of deoxyvasicinone (**1**) or vasicinone (**2**), isolated from *Adhatoda vasica*, and mackinazolinone (**3**), isolated from *Mackinalaya* species (Figure 1).^{5,6} These alkaloids have been extensively studied and a wide variety of routes for their synthesis, and of their derivatives, have been developed.⁴

2,3-Dihydropyrrolo[1,2-*a*]quinazolin-5(1*H*)-one (**4**), an isomer of deoxyvasicinone (**1**), was only isolated as a natural product in 2015 from *Castanea crenata* Sieb (Chestnut honey).⁷ It has, however, been obtained by synthetic methods for much longer.⁸ The biological activities of **4** and derivatives based on its alkaloid core have not been extensively investigated compared with related alkaloids **1** and **2**. A brominated derivative of **4** has some antibacterial activity at µg/mL levels against *Bacillus subtilis* and fungicidal activity against *Candida albicans* and *Aspergillus niger*.⁹ Com-





Figure 1 Quinazolinone alkaloids with fused pyrrole or piperidine rings demonstrate a variety of biological activities

pounds containing a core 2,3-dihydropyrrolo[1,2-*a*]quinazolin-5(1*H*)-one motif were found to be inhibitors of proteins in the bromodomain family of epigenetic reader proteins.¹⁰

In order to fully explore the potential of compound **4** as a scaffold for pharmaceutical or agrichemical applications, a robust and readily adaptable synthetic route to **4** and related analogues would be useful. Alkaloid **4** has been synthesised through several methods (Scheme 1). However, some of these routes were low yielding or would be unsuitable for analogue synthesis due to the starting materials or reagents used.^{11–14}

Furthermore, work has not been extensively undertaken to test the reactivity of the scaffold and explore its chemistry in comparison to the related deoxyvasicinone (**1**). In this paper, we report the development of an improved synthetic route to **4**, exploration of the scope of the route, and investigations into the reactivity of **4**. Our synthetic route, based on the existing synthesis by Dunn et al.,¹² enabled the reliable and rapid preparation of **4** at reasonable scale and with



good yields. Our strategy uses inexpensive reagents and was found to be readily adaptable for the synthesis of analogues and some related compounds. With a route that generated **4** in appreciable quantities, we also investigated the reactivity of the scaffold, highlighting the weakly nucleophilic characteristics of the core.

Optimisation of a Synthesis to 2,3-Dihydropyrrolo[1,2-*a*]quinazolin-5(1*H*)-one (**4**)

The route to 2,3-dihydropyrrolo[1,2-*a*]quinazolin-5(1*H*)-one (**4**) reported by Dunn et al. was chosen for exploration, since it used readily accessible, inexpensive reagents and we proposed it would be able to tolerate substitution.¹² In agreement with the literature procedure, reaction of anthranilamide (**5**) with acid chloride **6** gave the precursor secondary amide **7**. Cyclisation using sodium methoxide gave the two isomers deoxyvasicinone (**1**) and the desired compound 2,3-dihydropyrrolo[1,2-*a*]quinazolin-5(1*H*)-one (**4**) in low yields comparable to those reported (Table 1).

We proposed that yields might be improved by use of a stronger base. Reaction of secondary amide **7** with two equivalents of potassium *tert*-butoxide in THF led to rapid cyclisation, with some reactions complete in less than 10 minutes. A significant switch in product distribution was observed, with more of the desired isomer **4** rather than **1** being produced, and the overall yield was also greatly increased. Another strong base, NaHMDS, also demonstrated this effect, but potassium *tert*-butoxide was used subsequently due to its lower cost and ease of use.

The two isomers could be separated readily by flash column chromatography (FCC) rather than using a Soxhlet extractor as reported.¹² Identification of the isomers was con-

Table 1 Optimisation of Reaction Conditions for the Cyclisation of **7** to **4**^a

Entry	Base	Equiv	Time	Yield (%) of 4	Yield (%) of 1
1	25% wt NaOMe	10	4.00	13	19
2	<i>t</i> BuOK	1	4.00	35	22
3	<i>t</i> BuOK	2	0.25	77	15
4	<i>t</i> BuOK	3	0.25	59	29
5	NaHMDS	2	0.25	71	24

^a Reaction conditions: (a) Et₃N (2.0 equiv), THF, 0 °C to r.t., 12 h, 92%; (b) base (equiv), THF, time.


firmed by the presence of a strong NOE signal in **4** between the aromatic ring and pyrrole ring as well as the large difference in their melting points. The reaction was also performed on a gram scale, using **7**, with yields up to 69% being reliably obtained on this scale.

Versatility of the Synthetic Route for Analogue Synthesis

An advantage of this route was the accessibility of substituted 2-aminobenzamides, enabling the robustness of the route for analogue synthesis to be tested. A range of substituted 2-aminobenzamides were obtained through a variety of routes,^{15–17} of which microwave-assisted hydrolysis from 2-aminobenzonitriles proved the most versatile and reliable method (see Supporting Information).¹⁸ These 2-aminobenzamides were then reacted with acid chloride **6** as before (Table 2). Open-chain intermediates **8–18** could be accessed in moderate to excellent yields, with a variety of substitution patterns being tolerated, although substitution *ortho* to the amine was low yielding for steric reasons.

These secondary amides were then reacted with potassium *tert*-butoxide and cyclised successfully with good functional group tolerance. However, the yields and ratio of isomers obtained were sensitive to substitution pattern. Substitution *meta* to the secondary amide generally gave the best yields and the most favourable isomer distribution for the desired 2,3-dihydropyrrolo[1,2-*a*]quinazolin-5(1*H*)-one products. For example, on cyclisation of chlorinated in-

Table 2 Synthesis of Substituted 2-Aminobenzonitriles and Subsequent Reaction with 4-Chlorobutanoyl Chloride and Cyclisation To Give Analogues of **4**



Entry	R	Compound	Yield (%) of 8–18	Compound	Yield (%) of 19–29	Yield (%) of alternate isomer ^b
1	3-Me	8	39	19	53	26
2	4-Me	9	94	20	74	11
3	5-OMe	10	52	21	20	15
4	5-Me	11	79	22	37	8
5	5-F	12	93	23	33	6
6	5-Cl	13	81	24	65	24
7	6-OMe	14	79	25	82	–
8	6-Me	15	60	26	90	<4
9	6-F	16	94	27	73	<5
10	6-Cl	17	93	28	96	<1
11	6-Br	18	78	29	88	<5

^a Reaction conditions: (a) 4-chlorobutanoyl chloride (1.2 equiv), Et₃N (2.0 equiv), THF, 0 °C to r.t., 4 h to 24 h; (b) ^tBuOK (2.0 equiv), THF (0.1 M), 1 h.

^b Yields were determined on the basis of a crude NMR and subsequent purification of the major product.


intermediate **17** the desired product **28** was obtained in 91% yield after flash column chromatography, and none of the alternative isomer was detected by crude NMR analysis.

The methodology was also tested with electron-poor systems, and the nucleophilicity of the primary amine was found to be a key determinant of reactivity. Reaction of 2-aminonicotinamide failed to give the desired product; in-

stead compound **30**, product of the reaction of both the primary amine and the primary amide, was isolated (Scheme 2). By contrast, 3-aminonicotinamide could be successfully reacted to give **31**, as the electron-withdrawing group was not in direct conjugation with the primary amine. This amide was then successfully cyclised to compound **32** in 31% yield.


Generation of Expanded-Ring Analogues

Analogues of the scaffold with an expanded aliphatic ring system, such as mackinazolinone (**3**), have been synthesised previously using the methods of Dunn,¹² Sun,¹³ and Jones¹⁴ as highlighted in Figure 1, amongst many others.⁴ We assessed whether our improved synthetic route would be applicable to these systems. Precursor compounds **33** and **34** were synthesised using the appropriate acid chloride in 79% and 86% yield, respectively. Cyclisation of **33** with potassium *tert*-butoxide led to mackinazolinone (**3**) in 92% yield, rather than the isomer analogous to **4**. Similarly, compound **34** gave **41** in 52% with no isolation of **38** (Table 3). However, substitution of the aromatic ring affected the product distribution. Unlike the unsubstituted system, cyclisation of chlorinated amide **35** bearing a chlorine group gave access to both isomers **39** and **42** in yields of 51% and 48%, respectively. However, cyclisation of **36** gave only compound **43** in 20% yield.



Scheme 2 Use of electron-poor nicotinamides to generate analogues of **4**. Reaction conditions: (a) 4-chlorobutanoyl chloride (1.2 equiv), Et₃N (2.0 equiv), THF, 0 °C to r.t., 21%; (b) 4-chlorobutanoyl chloride (1.5 equiv), Et₃N (1.5 equiv), THF, 0 °C to r.t., 12 h, 58%; (c) ^tBuOK (2.0 equiv), THF (0.1 M), r.t., 2 h, 31%.

Table 3 Synthesis of Analogues with Expanded Ring Systems^a




Entry	R	n	Starting material	Product A	Yield (%)	Product B	Yield (%)
1	H	2	33	37	0	3	92
2	H	3	34	38	0	41	52
3	Cl	2	35	39	51	42	48
4	Cl	3	36	40	0	43	20

^a ^tBuOK (2.0 equiv), THF, r.t.

Proposed Mechanism

All these observations taken in combination agree with the report that the system can be cyclised through two competing mechanisms.¹⁹ We suggest that use of stronger bases, such as potassium *tert*-butoxide, favours deprotonation of the secondary amide in **7** followed by 5-*exo-tet* cyclisation and subsequent condensation of the lactam with the primary amide (route A). However, when the chain length increases the lactamisation is slower. Instead, the condensation of the primary amide onto the secondary amide in **7** may occur first followed by subsequent cyclisation (route B). As these routes would be in competition, their relative rates determine product distribution.

Changing the electronics of the ring will affect both the acidity of the secondary amide and the electrophilicity of the primary amide, altering the ratio of isomers produced. No lactam intermediates from proposed route A were isolated. However, with chlorinated substrates **35** and **36**, with longer chain lengths, reactions were stopped prior to completion, and intermediates **42a** and **43a** could be isolated in 9% and 95% yield, respectively, providing some evidence for the existence of route B (Figure 2).




C-Nucleophilicity of the 2,3-Dihydropyrrolo[1,2-*a*]quinazolin-5(1*H*)-one Scaffold

2,3-Dihydropyrrolo[1,2-*a*]quinazolin-5(1*H*)-one (**4**) is closely related to the alkaloid deoxyvasicinone (**1**) and, like deoxyvasicinone, shows reactivity towards electrophilic reagents at the C3 carbon position.²⁰ In studies of the scaffold as the basis for polymethine dyes, **4** was shown to react with conjugated aldehydes and condensed with benzaldehyde.^{12,21} We conducted initial investigations into the extent of this reactivity using a variety of electrophilic reagents to generate a series of substituted derivatives of compound **4**.


Reaction with Aromatic Aldehydes

The reaction of **4** with aromatic aldehydes was studied first. As reported, heating alkaloid **4** with benzaldehyde with acetic acid and sodium acetate led to rapid condensation and formation of the alkene **44** (Scheme 3).¹² None of the intermediate aldol product was isolated, likely due to the stability of the highly conjugated product. Only one stereoisomer, **44**, was obtained in 94% yield and given the thermodynamic conditions used, the *E*-geometry was proposed to be favoured. This was confirmed by NOESY experiments; a distinctive interaction between the condensed aromatic ring and the aliphatic ring was observed, whilst the potential interaction between the alkene proton and ring in proposed *Z*-isomer were not identified (Scheme 3 B). The methodology was applicable to substituted aldehydes: both electron-withdrawing and electron-donating groups were tolerated to give the final products **45** and **46** in 94% and 66% yield, respectively. The solubility of these products in both organic and aqueous solvent was low.



Reaction with Aliphatic Aldehydes


Unlike aromatic aldehydes, refluxing aliphatic aldehydes with 2,3-dihydropyrrolo[1,2-*a*]quinazolin-5(1*H*)-one (**4**) in acetic acid gave an inseparable mixture of unidentified products. However, treatment of **4** with potassium *tert*-butoxide to form an extended enolate and subsequent addition of 2-ethylbutyraldehyde at room temperature led to aldol reaction and then E1cb elimination of water, to give **47** as the *E*-isomer in 69% yield (Scheme 4). The chemistry tolerated functionalisation of the aromatic ring and varied aldehydes, for example, in synthesis of compound **49** and **50**. In some cases the intermediate aldol products could be isolated as a 1:1 mixture of diastereoisomers.



Scheme 4 Reactions of 2,3-dihydropyrrolo[1,2-*a*]quinazolin-5(1*H*)-ones with aliphatic aldehydes. Reagents and conditions: (a) ^tBuOK, CH₂Cl₂, 10 min, then aldehyde.

Reaction with Formamides

Analogous to reported reactions with deoxyvasicinone (**1**),²² compound **4** was reacted with phosphoryl chloride in dimethylformamide at 60 °C in a Vilsmeier–Haack style




Scheme 5 Reaction of **4** and related analogues with formamides and amines. Reagents and conditions: (a) POCl₃, DMF, 70 °C, 3 h, 74%; (b) formamide, POCl₃, CH₂Cl₂, 50 °C, 3–48 h; (c) amine, DMAP, EtOH, 70 °C, 24–96 h.

formylation, giving the dimethylaminomethylene derivative **51** in 74% yield (Scheme 5). The *E*-isomer alone was observed under the thermodynamic control of these reaction conditions. No aldehyde product from imine hydrolysis was observed, likely due to the stability of the conjugation of the dimethylaminomethylene system once formed.

The chemistry could be extended to other simple formamides to give aminomethylene derivatives such as **52** and **53** in moderate yield. Formamides containing additional nitrogen groups such as 1-methylpiperazine did not give products using these conditions. The dimethylaminomethylene product **51** was weakly reactive to nucleophilic substitution, allowing further elaboration of the dihydropyrrolo[1,2-*a*]quinazolin-5(1*H*)-one core. Upon refluxing with excess secondary amine and catalytic DMAP, the dimethyl group in **51** could be displaced giving compounds **54–56**.

Ring-expanded analogues such as **39** could also be reacted using this methodology to give derivatives **57** and **58** in 49% and 89% yield, respectively (Scheme 6).



Scheme 6 Reaction of ring-expanded analogue **39** with formamide and 1-methylpiperazine. Reagents and conditions: (a) POCl₃, DMF, 70 °C, 6 h, 49%; (b) 1-methylpiperazine, DMAP, EtOH, 70 °C, 6 h, 89%.

These reactions indicate the reactivity of the 2,3 dihydropyrrolo[1,2-*a*]quinazolin-5(1*H*)-one core to a variety of electrophiles is comparable to deoxyvasicinone (**1**). The latter can be brominated and oxidised, and other similar reactivity may also be applicable to **4**. In these reactions, nucleophilicity was only observed at carbon 3. When this position is blocked, for example, in isoindolo[2,1-*a*]quinazolin-5-ones,²³ carbon 1 directly adjacent to the nitrogen becomes the most reactive to electrophiles.

In conclusion, a versatile, scalable synthetic route to the 2,3-dihydropyrrolo[1,2-*a*]quinazolin-5(1*H*)-one **4** has been optimised, and shown to be suitable for the synthesis of a variety of analogues and some related compounds. The nucleophilic reactivity of **4** and analogues has been demonstrated with a variety of electrophiles. Condensation with electron-poor and electron-rich aromatic aldehydes occurs

in acidic medium whilst aliphatic aldehydes can be reacted under basic conditions, in both cases stereoselectively. Additionally, **4** has been derivatised using Vilsmeier condensation chemistry, and the products of this reaction such as **51** can be used to access other compounds through nucleophilic displacement. These results enable the synthesis of a wider variety of structures based on **4** than previously demonstrated, allowing their chemical and biological activity to be explored.

All reactions were performed using oven-dried glassware (200 °C) under a dry argon atmosphere unless otherwise stated or aqueous reagents were employed. All solvents used in reactions were dried and distilled according to standard protocols. All reagents were used as supplied or purified using standard procedures as necessary.

Flash column chromatography (FCC) was performed using Sigma Aldrich silica gel pore size 60 Å, 230–400 mesh under air pressure. The solvents used for FCC were distilled before use. 'Hexanes' refers to petroleum ether distillate (boiling point 40–60 °C). Analytical TLC was performed using silica gel 60 F₂₅₄ precoated glass-backed plates and visualised by ultraviolet radiation (254 nm), KMnO₄ or ninhydrin as appropriate. Preparative TLC was performed using silica gel GF 500 micron UNIPATES (Analtech).

¹H NMR spectra²⁴ were recorded on Bruker Avance DPX-600 (600 MHz), Bruker DPX-400 (400 MHz), or Bruker DCH Cryoprobe (500 MHz). High-resolution mass spectrometry (HRMS) was performed on a Waters Micromass LCT Premier spectrometer using electrospray ionisation and Micromass MS software, or on a ThermoFinnigan Orbitrap Classic using positive ion electrospray. IR spectra²⁴ were recorded as thin films on a PerkinElmer Spectrum One FTIR spectrometer. Melting points were collected using a Stanford Research Systems Optimel automated melting point system using a gradient of 1 °C per min.

2-(4-Chlorobutanamido)benzamides; General Procedure A

A solution of the appropriate substituted 2-aminobenzamide (1.0 equiv) in THF (2.5 mL per mmol substrate) was cooled to 0 °C. Et₃N (2.0 equiv) followed by the appropriate acid chloride (1.2 equiv) in THF (2 mL per mmol substrate) were added to the stirred solution. The reaction mixture was stirred at r.t. until completion as indicated by TLC, when the mixture was diluted with EtOAc and quenched with aq NaHSO₄ (20 mL). The aqueous phase was extracted with EtOAc (3 × 20 mL). The combined organic phases were dried (MgSO₄), excess solvent was removed in vacuo, and the residue purified by FCC.

2,3-Dihydropyrrolo[1,2-*a*]quinazolin-5(1*H*)-ones; General Procedure B

To a solution of the appropriate substituted 2-(4-chlorobutanamido)benzamides substrate (1.0 equiv) in THF (10 mL per mmol substrate) was added ^tBuOK (2.0 equiv). The reaction mixture was stirred at r.t. until TLC indicated completion, then solvent was removed in vacuo, the resulting residue was partitioned between CH₂Cl₂ (20 mL) and aq NaHCO₃ (15 mL), and the aqueous layer extracted with CH₂Cl₂ (5 × 20 mL). The combined organic phases were dried (MgSO₄) and solvent removed in vacuo. The products were obtained after purification of the residue by FCC.

Characterisation data for selected compounds are reported below. For other analogues, see further data in the Supporting Information.

Deoxyvavasicinone (**1**)

To **7** (120 mg, 0.500 mmol, 1.0 equiv) was added NaOMe (25% wt in MeOH, 1.14 mL, 5.00 mmol, 10.0 equiv) and the reaction mixture was heated at reflux for 4 h. The mixture was cooled, diluted with CH₂Cl₂ (10 mL), quenched with aq NaHCO₃ (10 mL), the layers separated, and the aqueous layer extracted with CH₂Cl₂ (3 × 10 mL). The combined organic layers were dried (MgSO₄), and excess solvent removed in vacuo. Purification by FCC (2% MeOH/CH₂Cl₂ to 5% MeOH/CH₂Cl₂) gave deoxyvavasicinone **1** (17.7 mg, 0.0951 mmol, 19%) as an off-white solid; mp 105–106 °C (Lit.²⁵ mp 105–107 °C); *R*_f = 0.31 (2% MeOH/EtOAc).

IR (neat): 2960, 2925, 1671, 1619, 1611, 1465, 1384, 771, 694 cm⁻¹.

¹H NMR (600 MHz, CDCl₃): δ = 8.27 (dd, *J* = 8.0, 1.3 Hz, 1 H), 7.72 (ddd, *J* = 8.2, 7.2, 1.5 Hz, 1 H), 7.64 (d, *J* = 8.3 Hz, 1 H), 7.44 (ddd, *J* = 8.0, 7.2, 1.1 Hz, 1 H), 4.20 (app t, *J* = 7.3 Hz, 2 H), 3.17 (t, *J* = 8.0 Hz, 2 H), 2.28 (app quint, *J* = 7.3 Hz, 2 H).

¹³C NMR (150 MHz, CDCl₃): δ = 161.2, 159.6, 149.3, 134.3, 127.0, 126.6, 126.4, 120.7, 46.7, 32.7, 19.7.

HRMS (ESI+): *m/z* [M + H]⁺ calcd for C₁₁H₁₁N₂O: 187.0871; found: 187.0875.

6,7,8,9-Tetrahydro-11*H*-pyrido[2,1-*b*]quinazolin-11-one (**3**)

Compound **33** (133 mg, 0.523 mmol) was cyclised according to general procedure B. Purification by FCC (1% MeOH/CH₂Cl₂) gave **3** (96.2 mg, 0.481 mmol, 92%) as a white amorphous solid; mp 81–83 °C; *R*_f = 0.47 (5% MeOH/CH₂Cl₂).

IR (neat): 3373, 2949, 1654, 1613, 1588, 1566, 1476, 1398 cm⁻¹.

¹H NMR (600 MHz, CDCl₃): δ = 8.19 (dd, *J* = 8.0, 1.2 Hz, 1 H), 7.65 (ddd, *J* = 8.4, 7.2, 1.2 Hz, 1 H), 7.53 (d, *J* = 8.0 Hz, 1 H), 7.36 (ddd, *J* = 8.0, 7.2, 1.2 Hz, 1 H), 4.02 (t, *J* = 6.2 Hz, 2 H), 2.94 (t, *J* = 6.6 Hz, 2 H), 1.99–1.93 (m, 2 H), 1.92–1.87 (m, 2 H).

¹³C NMR (150 MHz, CDCl₃): δ = 162.2, 154.9, 147.4, 134.2, 126.6, 126.4, 126.1, 120.5, 42.3, 32.0, 22.2, 19.4.

HRMS (ESI+): *m/z* [M + H]⁺ calcd for C₁₂H₁₃N₂O: 201.1022; found: 201.1014.

Data were in agreement with a reported example.¹³

2,3-Dihydropyrrolo[1,2-*a*]quinazolin-5(1*H*)-one (**4**)

Compound **7** (987 mg, 4.10 mmol) was cyclised according to general procedure B, and after FCC (3% MeOH/CH₂Cl₂), the title compound **4** (524 mg, 2.81 mmol, 69%) was obtained as a white amorphous solid; mp 213–216 °C (Lit.¹² mp 217–218 °C); *R*_f = 0.17 (5% MeOH/CH₂Cl₂).

IR (neat): 3059, 2953, 2925, 2897, 1634, 1593, 1533, 1501, 1461, 1420, 778 cm⁻¹.

¹H NMR (600 MHz, CDCl₃): δ = 8.25 (dd, *J* = 8.0, 1.1 Hz, 1 H), 7.66 (ddd, *J* = 8.4, 7.3, 1.5 Hz, 1 H), 7.40 (ddd, *J* = 8.1, 7.3, 0.9 Hz, 1 H), 7.17 (d, *J* = 8.2 Hz, 1 H), 4.22 (app t, *J* = 7.4 Hz, 2 H), 3.15 (app t, *J* = 7.8 Hz, 2 H), 2.41–2.36 (m, 2 H).

¹³C NMR (150 MHz, CDCl₃): δ = 170.3, 166.5, 138.8, 133.7, 128.9, 125.9, 118.9, 114.6, 48.8, 32.9, 18.7.

HRMS (ESI+): *m/z* [M + H]⁺ calcd for C₁₁H₁₁N₂O: 187.0866; found: 187.0870.

2-(4-Chlorobutanamido)benzamide (7)

Anthranilamide (**5**; 4.95 g, 36.4 mmol) and 4-chlorobutanoyl chloride (**6**; 6.17 g, 43.7 mmol) were reacted according to general procedure A. Purification by FCC (gradient 50% EtOAc/hexanes to 5% MeOH/EtOAc) gave **7** (8.08 g, 33.6 mmol, 92%) as an off-white amorphous solid; mp 117–119 °C, (Lit.¹² mp 114–115 °C); R_f = 0.36 (50% EtOAc/hexanes).

IR (neat): 3400, 3270, 3254, 3221, 1669, 1667, 1615, 1591, 1578, 1519, 757 cm⁻¹.

¹H NMR (600 MHz, CDCl₃): δ = 11.24 (s, NH, 1 H), 8.61 (d, J = 8.3 Hz, 1 H), 7.54 (d, J = 7.8 Hz, 1 H), 7.49 (dt, J = 8.1, 0.8 Hz, 1 H), 7.08 (t, J = 7.5 Hz, 1 H), 6.32 (br, NH, 1 H), 5.88 (br, NH, 1 H), 3.65 (t, J = 6.4 Hz, 2 H), 2.60 (t, J = 7.1 Hz, 2 H), 2.20 (quint, J = 6.8 Hz, 2 H).

¹³C NMR (150 MHz, CDCl₃): δ = 171.4, 170.8, 140.2, 133.5, 127.4, 122.9, 121.7, 118.7, 44.4, 35.2, 28.1.

HRMS (ESI⁺): m/z [M + H]⁺ calcd for C₁₁H₁₄³⁵ClN₂O₂: 241.0744; found: 241.0744.

The data were in agreement with a reported example.¹²

2-(4-Chlorobutanamido)-3-methylbenzamide (8)

2-Amino-3-methylbenzamide (90.0 mg, 0.599 mmol) was reacted with 4-chlorobutanoyl chloride according to general procedure A. Purification by FCC (100% EtOAc) gave the title compound **8** (59.8 mg, 0.235 mmol, 39%) as a white amorphous solid; mp 145–146 °C; R_f = 0.50 (50% EtOAc/hexanes).

IR (neat): 3377, 3291, 3190, 1652, 1588, 1513, 1395 cm⁻¹.

¹H NMR (600 MHz, CDCl₃): δ = 8.97 (s, NH, 1 H), 7.34–7.32 (m, 2 H), 7.16 (t, J = 7.6 Hz, 2 H), 6.21 (br, NH, 1 H), 5.63 (br, NH, 1 H), 3.64 (t, J = 6.3 Hz, 2 H), 2.58 (t, J = 7.0 Hz, 2 H), 2.24 (s, 3 H), 2.21–2.16 (m, 2 H).

¹³C NMR (150 MHz, CDCl₃): δ = 171.2, 171.0, 136.7, 134.6, 134.1, 129.7, 126.3, 125.0, 44.5, 33.8, 28.3, 19.1.

HRMS (ESI⁺): m/z [M + Na]⁺ calcd for C₁₂H₁₅³⁵ClN₂O₂²³Na: 277.0714; found: 277.0709.

9-Methyl-2,3-dihydropyrrolo[1,2-*a*]quinazolin-5(1*H*)-one (19)

Compound **8** (42.8 mg, 0.168 mmol) was cyclised according to general procedure B. Purification by FCC (4% MeOH/CH₂Cl₂) gave **19** (17.9 mg, 0.0894 mmol, 53%) as a white amorphous solid; mp 247–250 °C; R_f = 0.14 (4% MeOH/CH₂Cl₂).

IR (neat): 1632, 1586, 1549, 1481, 1411, 785 cm⁻¹.

¹H NMR (600 MHz, CDCl₃): δ = 8.19 (dd, J = 7.9, 1.0 Hz, 1 H), 7.39 (d, J = 7.3 Hz, 1 H), 7.24 (t, J = 7.6 Hz, 1 H), 4.66 (app t, J = 7.2 Hz, 2 H), 3.07 (t, J = 8.0 Hz, 2 H), 2.73 (s, 3 H), 2.32–2.27 (m, 2 H).

¹³C NMR (125 MHz, CDCl₃): δ = 170.3, 167.8, 138.7, 137.7, 127.6, 125.9, 124.8, 120.3, 54.1, 32.7, 22.6, 19.7.

HRMS (ESI⁺): m/z [M + H]⁺ calcd for C₁₂H₁₃N₂O: 201.2028; found: 201.1020.

7-Chloro-1,2,3,4-tetrahydro-6*H*-pyrido[1,2-*a*]quinazolin-6-one (39)

Compound **35** (324 mg, 1.12 mmol) was cyclised according to general procedure B. Purification by FCC (5% MeOH/CH₂Cl₂) gave the desired product **39** (133 mg, 0.567 mmol, 51%) and isomer **42** as white amorphous powders; mp 223–225 °C; R_f = 0.33 (6% MeOH/CH₂Cl₂).

IR (neat): 3064, 2949, 2883, 1660, 1587, 1552, 1458 cm⁻¹.

¹H NMR (600 MHz, CDCl₃): δ = 7.50 (t, J = 7.8 Hz, 1 H), 7.37 (dd, J = 7.8, 0.7 Hz, 1 H), 7.31 (d, J = 8.4 Hz, 1 H), 3.98 (app t, J = 6.3 Hz, 2 H), 2.98 (app t, J = 6.6 Hz, 2 H), 2.17–2.13 (m, 2 H), 1.96–1.92 (m, 2 H).

¹³C NMR (150 MHz, CDCl₃): δ = 166.2, 160.5, 143.5, 135.8, 132.6, 128.8, 117.6, 112.8, 47.4, 32.8, 22.8, 19.1.

HRMS (ESI⁺): m/z [M + H]⁺ calcd for C₁₂H₁₂³⁵ClN₂O: 235.0633; found: 235.0623.

7,8,9,10-Tetrahydroazepino[2,1-*b*]quinazolin-12(6*H*)-one (41)

Compound **34** (96.5 mg, 0.360 mmol) was cyclised according to general procedure B. Purification by FCC (4% MeOH/CH₂Cl₂) gave the product **41** (40.2 mg, 0.188 mmol, 52%) as a white amorphous powder; R_f = 0.38 (4% MeOH/CH₂Cl₂).

IR (neat): 2921, 2855, 1660, 1587, 1568, 1473, 1393 cm⁻¹.

¹H NMR (600 MHz, CDCl₃): δ = 8.25 (dd, J = 8.0, 1.4 Hz, 1 H), 7.70 (ddd, J = 8.4, 7.2, 1.5 Hz, 1 H), 7.60 (dd, J = 8.3, 0.6 Hz, 1 H), 7.43 (ddd, J = 7.9, 7.2, 1.1 Hz, 1 H), 4.40–4.38 (m, 2 H), 3.08–3.06 (m, 2 H), 1.88–1.81 (m, 6 H).

¹³C NMR (150 MHz, CDCl₃): δ = 162.1, 159.9, 147.6, 134.3, 127.2, 126.9, 126.5, 120.4, 43.0, 37.9, 29.7, 28.3, 25.6.

HRMS (ESI⁺): m/z [M + H]⁺ calcd for C₁₃H₁₅N₂O: 215.1184; found: 215.1180.

1-Chloro-6,7,8,9-tetrahydro-11*H*-pyrido[2,1-*b*]quinazolin-11-one (42)

Side-product **42** was obtained from the cyclisation of **35** as a white amorphous solid (113 mg, 0.538 mmol, 48%); mp 124–127 °C; R_f = 0.47 (EtOAc).

IR (neat): 2949, 2883, 1660, 1587, 1552, 1458 cm⁻¹.

¹H NMR (600 MHz, CDCl₃): δ = 7.54 (t, J = 7.8 Hz, 1 H), 7.49 (d, J = 7.8 Hz, 1 H), 7.40 (dd, J = 7.8, 1.2 Hz, 1 H), 4.02 (app t, J = 6.0 Hz, 2 H), 2.97 (app t, J = 6.6 Hz, 2 H), 2.03–1.97 (m, 2 H), 1.95–1.91 (m, 2 H).

¹³C NMR (150 MHz, CDCl₃): δ = 160.3, 155.9, 149.6, 134.0, 133.7, 129.0, 125.7, 117.6, 42.8, 31.9, 22.3, 19.3.

HRMS (ESI⁺): m/z [M + H]⁺ calcd for C₁₂H₁₃³⁵ClN₂O: 235.0638; found: 235.0641.

5-Chloro-2-(4-chlorobutyl)quinazolin-4(3*H*)-one (42a)

Side-product **42a** (7.0 mg, 0.0258 mmol, 9%) was obtained from quenching the cyclisation of **35** with ^tBuOK after 1 h and purification by FCC (50% MeOH/hexanes) as a white amorphous solid; mp 215 °C (dec.); R_f = 0.36 (50% EtOAc/hexanes).

IR (neat): 3174, 3037, 2962, 2901, 1679, 1619, 1594, 1457, 817 cm⁻¹.

¹H NMR (600 MHz, CDCl₃): δ = 11.71 (s, NH), 7.63–7.59 (m, 2 H), 7.46 (dd, J = 6.8, 2.1 Hz, 1 H), 3.61 (t, J = 6.6 Hz, 2 H), 2.81 (t, J = 7.7 Hz, 2 H), 2.09–2.03 (m, 2 H), 1.97–1.92 (m, 2 H).

¹³C NMR (150 MHz, CDCl₃): δ = 161.7, 155.9, 151.0, 133.4, 133.2, 128.4, 125.8, 117.0, 43.5, 33.8, 31.1, 23.6.

HRMS (ESI⁺): m/z [M + H]⁺ calcd for C₁₂H₁₃³⁵Cl₂N₂O: 271.0399; found: 271.0387.

1-Chloro-7,8,9,10-tetrahydroazepino[2,1-*b*]quinazolin-12(6*H*)-one (43)

Compound **36** (13.0 mg, 0.0456 mmol, 1.0 equiv) was cyclised according to general procedure B. Purification by preparative TLC (4% MeOH/CH₂Cl₂) gave the product **43** (2.3 mg, 0.00925 mmol, 20%) as a white amorphous powder; R_f = 0.27 (4% MeOH/CH₂Cl₂).

IR (neat): 2919, 2851, 1664, 1595, 1548, 1457 cm⁻¹.

^1H NMR (600 MHz, CDCl_3): δ = 7.55 (t, J = 7.9 Hz, 1 H), 7.50 (dd, J = 8.2, 1.3 Hz, 1 H), 7.42 (dd, J = 7.7, 1.3 Hz, 1 H), 4.36–4.35 (m, 2 H), 3.04–3.02 (m, 2 H), 1.90–1.79 (m, 6 H).

^{13}C NMR (150 MHz, CDCl_3): δ = 160.6, 160.2, 150.0, 134.4, 133.7, 129.3, 126.3, 117.6, 43.1, 37.8, 29.7, 28.1, 25.5.

HRMS (ESI+): m/z [$M + H$] $^+$ calcd for $\text{C}_{13}\text{H}_{14}^{35}\text{ClN}_2\text{O}$: 249.0795; found: 249.0797.

5-Chloro-2-(5-chloropentyl)quinazolin-4(3H)-one (43a)

Compound **43a** (13.0 mg, 0.0456 mmol, 95%) was obtained from quenching the cyclisation of **36** (14.5 mg, 0.0478 mmol) with $^t\text{BuOK}$ after 24 h and purification by FCC (2% MeOH/ CH_2Cl_2) as a white amorphous solid; mp 160–161 $^\circ\text{C}$; R_f = 0.37 (2% MeOH/ CH_2Cl_2).

IR (neat): 2956, 2932, 2902, 2869, 1674, 1617, 1600, 1462, cm^{-1} .

^1H NMR (600 MHz, CDCl_3): δ = 11.91 (s, NH), 7.64–7.60 (m, 2 H), 7.46 (dd, J = 6.8, 2.1 Hz, 1 H), 3.56 (t, J = 6.5 Hz, 2 H), 2.80 (app t, J = 7.9 Hz, 2 H), 1.97–1.92 (m, 2 H), 1.90–1.85 (m, 2 H), 1.65–1.60 (m, 2 H).

^{13}C NMR (150 MHz, CDCl_3): δ = 162.8, 157.4, 152.1, 134.3, 134.1, 129.3, 126.7, 118.0, 44.9, 35.6, 32.3, 26.69, 26.67.

HRMS (ESI+): m/z [$M + \text{Na}$] $^+$ calcd for $\text{C}_{13}\text{H}_{14}^{35}\text{Cl}_2\text{N}_2\text{ONa}$: 307.0375; found: 307.0369.

(E)-3-Benzylidene-2,3-dihydropyrrolo[1,2-a]quinazolin-5(1H)-one (44)

The procedure was adapted from a literature protocol.²⁶ To a solution of **4** (59.2 mg, 0.318 mmol, 1.0 equiv) in AcOH (1.00 mL) was added benzaldehyde (65.7 μL , 0.350 mmol, 1.1 equiv), followed by NaOAc (21.0 mg, 0.260 mmol, 0.8 equiv). The reaction mixture was heated at reflux for 4 h, cooled to r.t., then the solids were filtered, and washed with H_2O , CH_2Cl_2 , and EtOH successively, to give **52** (82.9 mg, 0.302 mmol, 95%) as a white amorphous solid; mp 290–293 $^\circ\text{C}$; R_f = 0.25 (4% MeOH/ CH_2Cl_2).

IR (neat): 1634, 1598, 1538, 1493, 1465, 1426, 1139, 757, 688 cm^{-1} .

^1H NMR (600 MHz, CDCl_3): δ = 8.27 (d, J = 7.8 Hz, 1 H), 7.93 (s, 1 H), 7.65 (t, J = 7.6 Hz, 1 H), 7.49 (d, J = 7.4 Hz, 2 H), 7.41–7.32 (m, 3 H), 7.20 (d, J = 8.3 Hz, 1 H), 4.30 (t, J = 7.0 Hz, 2 H), 3.38–3.37 (m, 2 H).

^{13}C NMR (150 MHz, CDCl_3): δ = 170.5, 161.4, 138.9, 135.3, 133.8, 133.5, 130.2, 129.5, 129.03, 128.97, 126.3, 119.9, 114.5, 46.3, 25.4.

HRMS (ESI+): m/z [$M + H$] $^+$ calcd for $\text{C}_{18}\text{H}_{15}\text{N}_2\text{O}$: 275.1184; found: 275.1176.

(E)-3-(2-Ethylbutylidene)-2,3-dihydropyrrolo[1,2-a]quinazolin-5(1H)-one (47)

To a solution of **4** (50.0 mg, 0.269 mmol) in CH_2Cl_2 (920 μL) was added $^t\text{BuOK}$ (36.1 mg, 0.322 mmol, 1.2 equiv). The mixture was stirred vigorously for 5 min, then 2-ethylbutyraldehyde (33.1 μL , 0.269 mmol, 1.0 equiv) was added. After 1 h, aq NaHCO_3 (10 mL) and CH_2Cl_2 (10 mL) were added, the layers separated, and the aqueous layer was extracted with CH_2Cl_2 (3 $\times$ 10 mL). The combined organic layers were dried (MgSO_4) and excess solvent removed in vacuo. Purification by preparative TLC (5% MeOH/ CH_2Cl_2) gave the product **47** (49.8 mg, 0.186 mmol, 69%) as an off-white amorphous solid; mp 113–115 $^\circ\text{C}$ (dec.); R_f = 0.30 (4% MeOH/ CH_2Cl_2).

IR (neat): 2958, 2929, 2874, 2857, 1637, 1626, 1598, 1532, 1496, 1461, 1423, 741 cm^{-1} .

^1H NMR (600 MHz, CDCl_3): δ = 8.31 (dd, J = 7.9, 1.3 Hz, 1 H), 7.66 (dt, J = 7.4, 1.6 Hz, 1 H), 7.41 (dt, J = 7.4, 0.8 Hz, 1 H), 7.19 (d, J = 8.2 Hz, 1 H), 6.92 (dt, J = 10.8, 2.7 Hz, 1 H), 4.20 (t, J = 7.4 Hz, 2 H), 3.02 (m, 2 H), 2.19–2.14 (m, 1 H), 1.61–1.50 (m, 2 H), 1.42–1.34 (m, 2 H), 0.87 (t, J = 7.8 Hz, 6 H).

^{13}C NMR (150 MHz, CDCl_3): δ = 170.6, 160.3, 142.4, 139.0, 133.7, 131.4, 129.0, 126.1, 119.9, 114.3, 45.8, 44.3, 27.7, 23.2, 12.1.

HRMS (ESI+): m/z calcd for $\text{C}_{17}\text{H}_{21}\text{N}_2\text{O}$: 269.164; found: 269.1658.

(E)-6-Chloro-3-[(tetrahydro-2H-pyran-4-yl)methylene]-2,3-dihydropyrrolo[1,2-a]quinazolin-5(1H)-one (50)

To a solution of **28** (50.0 mg, 0.227 mmol, 1.0 equiv) in CH_2Cl_2 (920 μL) was added $^t\text{BuOK}$ (30.5 mg, 0.272 mmol, 1.2 equiv). After stirring the mixture vigorously for 5 min, tetrahydro-2H-pyran-4-carbaldehyde (0.250 mmol, 1.1 equiv) was added. After 1 h, aq NaHCO_3 (10 mL) and CH_2Cl_2 (10 mL) were added, the layers separated, and the aqueous layer was extracted with CH_2Cl_2 (3 $\times$ 10 mL). The combined organic layers were dried (MgSO_4) and excess solvent removed in vacuo. Purification by preparative TLC (5% MeOH/ CH_2Cl_2) gave the product **50** (48.4 mg, 0.153 mmol, 67%) as an off-white amorphous solid; mp 211–214 $^\circ\text{C}$ (dec.); R_f = 0.33 (5% MeOH/ CH_2Cl_2).

IR (neat): 2931, 2832, 1645, 1602, 1585, 1543, 1488 cm^{-1} .

^1H NMR (600 MHz, CDCl_3): δ = 7.52 (t, J = 8.1 Hz, 1 H), 7.41 (dd, J = 7.8, 0.7 Hz, 1 H), 7.08 (dd, J = 8.3, 0.7 Hz, 1 H), 6.98 (dt, J = 9.7, 2.8 Hz, 1 H), 4.19 (t, J = 7.3 Hz, 2 H), 4.01 (dt, J = 11.5, 3.4 Hz, 2 H), 3.50–3.45 (m, 2 H), 3.06 (dt, J = 7.3, 2.8 Hz, 2 H), 2.59–2.53 (m, 1 H), 1.66–1.62 (m, 4 H).

^{13}C NMR (150 MHz, CDCl_3): δ = 168.1, 159.5, 141.2, 140.5, 136.6, 133.0, 130.0, 129.1, 116.9, 113.2, 67.3, 46.5, 36.6, 31.3, 22.6.

HRMS (ESI+): m/z calcd for $\text{C}_{17}\text{H}_{18}^{35}\text{ClN}_2\text{O}_2$: 317.1052; found: 317.1056.

(E)-3-[(Dimethylamino)methylene]-2,3-dihydropyrrolo[1,2-a]quinazolin-5(1H)-one (51)

To a solution of **4** (372 mg, 2.00 mmol, 1.0 equiv) in DMF (6.7 mL) was added POCl_3 (373 μL , 4.00 mmol, 2.0 equiv). The crude product was purified by FCC (4% MeOH/ CH_2Cl_2) to give **51** (357 mg, 1.48 mmol, 74%) as a tan amorphous solid; mp 251–252 $^\circ\text{C}$ (dec.); R_f = 0.21 (5% MeOH/ CH_2Cl_2).

IR (neat): 1646, 1622, 1598, 1514, 1495, 1397, 1373, 1314, 1109, 776 cm^{-1} .

^1H NMR (600 MHz, $\text{DMSO}-d_6$): δ = 7.92 (dd, J = 7.8, 1.2 Hz, 1 H), 7.64 (ddd, J = 7.8, 7.2, 1.2 Hz, 1 H), 7.39 (t, J = 1.2 Hz, 1 H), 7.27 (dt, J = 7.5, 0.6 Hz, 1 H), 7.23 (d, J = 8.4 Hz, 1 H), 4.10–4.08 (m, 2 H), 3.21 (t, J = 7.8 Hz, 2 H), 3.09 (s, 6 H).

^{13}C NMR (150 MHz, CDCl_3): δ = 168.2, 163.9, 144.0, 139.5, 132.9, 127.1, 123.4, 118.8, 114.2, 94.5, 45.4, 40.0, 22.7.

HRMS (ESI+): m/z [$M + H$] $^+$ calcd for $\text{C}_{14}\text{H}_{16}\text{N}_3\text{O}$: 242.1293; found: 242.1295.

(E)-3-(Piperidin-1-ylmethylene)-2,3-dihydropyrrolo[1,2-a]quinazolin-5(1H)-one (52)

To a solution of **4** (59.0 mg, 0.317 mmol, 1.0 equiv) in CH_2Cl_2 (1.00 mL) was added 1-formylpiperazine (107 μL , 0.951 mmol, 3.0 equiv) and POCl_3 (59.1 μL , 0.634 mmol, 2.0 equiv). The reaction mixture was refluxed for 28 h, cooled to r.t., diluted with CH_2Cl_2 (10 mL), and quenched with aq NaHCO_3 (10 mL). The aqueous layer was extracted with CH_2Cl_2 (5 $\times$ 10 mL), the combined organic layers dried (MgSO_4),

and solvent removed in vacuo. Purification by FCC (2% MeOH/CH₂Cl₂ to 5% MeOH/CH₂Cl₂) gave the title compound **52** (28.9 mg, 0.103 mmol, 32%) as a tan amorphous solid; *R*_f = 0.30 (5% MeOH/CH₂Cl₂).

IR (neat): 2934, 2853, 1643, 1600, 1587, 1510, 1486, 1439, 1402 cm⁻¹.

¹H NMR (600 MHz, DMSO-*d*₆): δ = 7.93 (dd, *J* = 7.8, 1.2 Hz, 1 H), 7.64 (ddd, *J* = 8.4, 7.8, 1.2 Hz, 1 H), 7.40 (s, 1 H), 7.27 (dt, *J* = 7.8, 0.6 Hz, 1 H), 7.24 (d, *J* = 8.4 Hz, 1 H), 4.11 (app t, *J* = 7.8 Hz, 2 H), 3.47–3.45 (m, 4 H), 3.11 (app t, *J* = 7.8 Hz, 2 H), 1.63–1.57 (m, 6 H).

¹³C NMR (100 MHz, CDCl₃): δ = 168.1, 164.1, 142.4, 139.5, 132.9, 127.2, 123.4, 118.8, 114.2, 93.9, 50.9, 45.5, 26.1, 23.6, 23.2.

HRMS (ESI+): *m/z* [M + H]⁺ calcd for C₁₇H₂₀N₃O: 282.1606; found: 282.1615.

(E)-3-(Morpholinomethylene)-2,3-dihydropyrrolo[1,2-*a*]quinazolin-5(1H)-one (53)

To a solution of **4** (50.0 mg, 0.269 mmol, 1.0 equiv) in CH₂Cl₂ (1.35 mL) was added morpholine-4-carbaldehyde (310 mg, 2.69 mmol, 10.0 equiv) and POCl₃ (50.1 μL, 0.538 mmol, 2.0 equiv). The reaction mixture was refluxed for 3 h, diluted with CH₂Cl₂ (10 mL), cooled to r.t., and quenched with aq NaHCO₃ (15 mL). The aqueous layer was extracted with CH₂Cl₂ (5 × 15 mL), the combined organic layers dried (Na₂SO₄), and solvent removed in vacuo. Purification by FCC (5% MeOH/CH₂Cl₂ to 10% MeOH/CH₂Cl₂) gave the title compound **53** (26.7 mg, 0.0942 mmol, 35%) as a tan amorphous solid; mp 275 °C (dec.); *R*_f = 0.25 (5% MeOH/CH₂Cl₂).

IR (neat): 2961, 2925, 2887, 1634, 1600, 1594, 1525, 1510, 1110, 760 cm⁻¹.

¹H NMR (600 MHz, CDCl₃): δ = 8.29 (dd, *J* = 7.9, 1.3 Hz, 1 H), 7.65 (s, 1 H), 7.59 (ddd, *J* = 8.3, 7.4, 1.6 Hz, 1 H), 7.32 (ddd, *J* = 8.0, 7.5, 0.9 Hz, 1 H), 7.03 (d, *J* = 8.2 Hz, 1 H), 4.14–4.11 (m, 2 H), 3.76–3.75 (m, 4 H), 3.51–3.48 (m, 4 H), 3.50 (d, *J* = 4.8 Hz, 2 H), 3.18 (ddd, *J* = 8.3, 6.7, 1.7 Hz, 2 H).

¹³C NMR (100 MHz, CDCl₃): δ = 170.3, 164.4, 143.0, 139.5, 133.2, 128.9, 124.5, 119.4, 113.3, 95.6, 66.7, 50.4, 45.0, 24.0.

HRMS (ESI+): *m/z* [M + H]⁺ calcd for C₁₆H₁₈N₃O₂: 284.1399; found: 284.1396.

(E)-3-[(4-Ethylpiperazin-1-yl)methylene]-2,3-dihydropyrrolo[1,2-*a*]quinazolin-5(1H)-one (54)

To a solution of **51** (50.0 mg, 0.207 mmol, 1.0 equiv) in EtOH (1 mL) was added 1-ethylpiperazine (80.0 μL, 0.630 mmol, 0.33 equiv) and *N,N*-dimethylaminopyridine (2.5 mg, 0.020 mmol, 0.1 equiv), and heated at reflux for 32 h. The solvent was removed in vacuo and purification by FCC (5% MeOH/CH₂Cl₂) gave the product **54** (39.6 mg, 0.128 mmol, 62%) as a pale brown amorphous solid; mp 165–168 °C; *R*_f = 0.30 (5% MeOH/CH₂Cl₂).

IR (neat): 2970, 2901, 2819, 1645, 1613, 1598, 1588, 1509, 1491 cm⁻¹.

¹H NMR (600 MHz, CDCl₃): δ = 8.27 (dd, *J* = 7.9, 1.3 Hz, 1 H), 7.65 (s, 1 H), 7.57 (dd, *J* = 8.4, 7.4, 1.5 Hz, 1 H), 7.29 (dt, *J* = 7.8, 0.6 Hz, 1 H), 7.00 (d, *J* = 7.8 Hz, 1 H), 4.10 (app t, *J* = 7.8 Hz, 2 H), 3.52 (app t, *J* = 5.0 Hz, 4 H), 3.16 (dt, *J* = 7.8, 1.2 Hz, 2 H), 2.51 (app t, *J* = 4.8 Hz, 4 H), 2.46 (q, *J* = 7.2 Hz, 2 H), 1.10 (t, *J* = 7.2 Hz, 3 H).

¹³C NMR (150 MHz, CDCl₃): δ = 170.3, 164.6, 142.9, 139.6, 133.1, 128.9, 124.3, 119.5, 113.2, 94.7, 52.8, 52.4, 50.3, 45.9, 24.0, 12.0.

HRMS (ESI+): *m/z* [M + H]⁺ calcd for C₁₈H₂₃N₄O: 311.1861; found: 311.1866.

(E)-3-[(4-Methyl-1,4-diazepan-1-yl)methylene]-2,3-dihydropyrrolo[1,2-*a*]quinazolin-5(1H)-one (55)

To a solution of **51** (21.0 mg, 0.0870 mmol) in EtOH (0.5 mL) was added 1-methylhomopiperazine (32 μL, 0.257 mmol, 3.4 equiv), and *N,N*-dimethylaminopyridine (1.1 mg, 0.0087 mmol, 0.1 equiv), and heated at reflux for 96 h. The solvent was removed in vacuo and purification by FCC (5% MeOH/CH₂Cl₂) gave the product **55** (13.2 mg, 0.0425 mmol, 49%) as a pale brown amorphous solid; mp 92–94 °C; *R*_f = 0.32 (10% MeOH/CH₂Cl₂ + 1% NH₄OH).

IR (neat): 2939, 2913, 2857, 2810, 1641, 1602, 1587, 1506, 1496 cm⁻¹.

¹H NMR (500 MHz, CDCl₃): δ = 8.27 (dd, *J* = 7.9, 1.4 Hz, 1 H), 7.74 (t, *J* = 1.5 Hz, 1 H), 7.57 (ddd, *J* = 8.3, 7.3, 1.5 Hz, 1 H), 7.29 (ddd, *J* = 8.1, 7.5, 1.0 Hz, 1 H), 7.00 (d, *J* = 8.0 Hz, 1 H), 4.11–4.07 (m, 2 H), 3.62 (t, *J* = 6.3 Hz, 2 H), 3.59–3.57 (m, 2 H), 3.21 (m, 2 H), 2.67–2.65 (m, 2 H), 2.61–2.59 (m, 2 H), 2.39 (s, 3 H), 1.95–1.91 (m, 2 H).

¹³C NMR (125 MHz, CDCl₃): δ = 170.4, 164.6, 144.5, 139.7, 133.0, 128.8, 124.1, 119.5, 113.1, 94.1, 59.1, 57.0, 54.1, 51.7, 46.9, 45.8, 28.5, 23.9.

HRMS (ESI+): *m/z* [M + H]⁺ calcd for C₁₈H₂₃N₄O: 311.1872; found: 311.1884.

tert-Butyl (E)-4-[(5-Oxo-1,2-dihydropyrrolo[1,2-*a*]quinazolin-3(5H)-ylidene)methyl]piperazine-1-carboxylate (56)

To a solution of **51** (21.0 mg, 0.0870 mmol) in EtOH (0.5 mL) was added with 1-Boc-piperazine (49.0 mg, 0.263 mmol, 3.0 equiv) and *N,N*-dimethylaminopyridine (1.1 mg, 0.0087 mmol, 0.1 equiv), and heated at reflux for 96 h. The solvent was removed in vacuo and purification by FCC (5% MeOH/CH₂Cl₂) gave the product **56** (20.0 mg, 0.0522 mmol, 60%) as a pale brown amorphous solid; mp 140 °C (dec.); *R*_f = 0.25 (5% MeOH/CH₂Cl₂).

IR (neat): 2974, 2918, 1688, 1644, 1605, 1591, 1518, 1494, 1405 cm⁻¹.

¹H NMR (600 MHz, CDCl₃): δ = 8.24 (dd, *J* = 7.4, 1.2 Hz, 1 H), 7.57 (m, 1 H), 7.54 (ddd, *J* = 8.4, 7.4, 1.4 Hz, 1 H), 7.27 (dt, *J* = 7.9, 0.6 Hz, 1 H), 6.97 (d, *J* = 8.1 Hz, 1 H), 4.07 (app t, *J* = 7.9 Hz, 2 H), 3.48–3.46 (m, 4 H), 3.39 (br, 4 H), 3.10 (dt, *J* = 7.8, 1.4 Hz, 2 H), 1.46 (s, 9 H).

¹³C NMR (150 MHz, CDCl₃): δ = 170.2, 164.3, 154.5, 142.5, 139.5, 133.0, 128.6, 124.2, 119.4, 113.4, 95.8, 80.8, 46.0, 28.6, 24.0. *C13 and C14 were weak and very broad in ¹³C NMR and could not be reliably assigned.

HRMS (ESI+): *m/z* [M + H]⁺ calcd for C₂₁H₂₇N₄O₃: 383.2078; found: 383.2070.

(E)-7-Chloro-4-[(dimethylamino)methylene]-1,2,3,4-tetrahydro-6H-pyrido[1,2-*a*]quinazolin-6-one (57)

To a stirred solution of **40** (88.0 mg, 0.37 mmol, 1.0 equiv) in DMF (740 μL) was added POCl₃ (69.9 μL, 0.74 mmol, 2.0 equiv) and the reaction mixture was heated to 60 °C for 6 h. The mixture was cooled to r.t., diluted with CH₂Cl₂ (15 mL), and quenched with aq NaHCO₃ (15 mL). The aqueous layer was extracted with CH₂Cl₂ (5 × 15 mL), the combined organic layers dried (Na₂SO₄) and solvent removed in vacuo. The residue was co-evaporated with toluene (3 × 5 mL) and purification by FCC (6% MeOH/CH₂Cl₂) gave the product **57** (52.0 mg, 0.18 mmol, 49%) as a yellow amorphous solid; mp 197–200 °C; *R*_f = 0.43 (6% MeOH/CH₂Cl₂).

IR (neat): 3111, 2944, 2885, 2813, 1607, 1587, 1499, 1481, 1464 cm⁻¹.

¹H NMR (600 MHz, CDCl₃): δ = 8.27 (s, 1 H), 7.39 (t, *J* = 8.2 Hz, 1 H), 7.29 (dd, *J* = 7.9, 0.9 Hz, 1 H), 7.14 (dd, *J* = 8.5, 0.6 Hz, 1 H), 3.85 (app t, *J* = 6.0 Hz, 2 H), 3.12 (s, 6 H), 2.73–2.71 (m, 2 H), 2.09–2.05 (m, 2 H).

^{13}C NMR (150 MHz, CDCl_3): δ = 168.9, 158.6, 151.1, 144.4, 135.2, 131.2, 127.0, 117.7, 111.7, 94.2, 46.9, 43.8, 23.4, 22.5.

HRMS (ESI⁺): m/z [$M + H$]⁺ calcd for $\text{C}_{15}\text{H}_{17}^{35}\text{ClN}_3\text{O}$: 290.1055; found: 290.1048.

(E)-7-Chloro-4-[(4-methylpiperazin-1-yl)methylene]-1,2,3,4-tetrahydro-6H-pyrido[1,2-a]quinazolin-6-one (58)

To a solution of **57** (26.0 mg, 0.09 mmol, 1.0 equiv) in EtOH (450 μL) in a microwave vial was added 1-methylpiperazine (49.7 μL , 0.45 mmol, 5.0 equiv) and DMAP (1.1 mg, 0.009 mmol, 0.1 equiv). The vial was sealed and the reaction mixture heated at 70 °C for 6 h, then cooled to r.t. and diluted with CH_2Cl_2 (10 mL) and NaHCO_3 (10 mL). The aqueous layer was extracted with CH_2Cl_2 (5 $\times$ 10 mL), the combined organic layers were dried (Na_2SO_4), and excess solvent was removed in vacuo. Purification by FCC (10% MeOH/ CH_2Cl_2 + 0.5% NH_4OH) gave the product **58** (27.6 mg, 0.080 mmol, 89%) as a yellow amorphous solid; mp 174–177 °C; R_f = 0.22 (10% MeOH/ CH_2Cl_2 + 0.5% NH_4OH).

IR (neat): 2934, 2845, 2790, 1615, 1586, 1494, 1474, 1419 cm^{-1} .

^1H NMR (600 MHz, CDCl_3): δ = 8.24 (s, 1 H), 7.39 (t, J = 8.2 Hz, 1 H), 7.28 (dd, J = 7.8, 0.7 Hz, 1 H), 7.14 (d, J = 8.4 Hz, 1 H), 3.85 (app t, J = 6.0 Hz, 2 H), 3.53 (m, 4 H), 2.60 (app t, J = 6.1 Hz, 2 H), 2.43 (m, 4 H), 2.30 (s, 3 H), 2.08 (app quint, J = 6.0 Hz, 2 H).

^{13}C NMR (150 MHz, CDCl_3): δ = 165.9, 158.3, 149.3, 144.2, 135.1, 131.7, 127.0, 117.7, 111.7, 94.8, 55.2, 51.2, 46.7, 46.2, 24.2, 22.3.

HRMS (ESI⁺): m/z [$M + H$]⁺ calcd for $\text{C}_{18}\text{H}_{22}^{35}\text{ClN}_4\text{O}$: 345.1482; found: 345.1494.

Acknowledgment

Charlotte Sutherell was funded by the Cambridge Ph.D. Training Programme in Chemical Biology and Molecular Medicine. We gratefully acknowledge the EPSRC (SVL, Grant Nos. EP/K099494/1 and EP/K039520/1).

Supporting Information

Supporting information for this article is available online at <http://dx.doi.org/10.1055/s-0035-1562792>.

References

- (1) Department of Chemistry, Imperial College, London, SW7 2AZ.
- (2) Horton, D. A.; Bourne, G. T.; Smythe, M. L. *Chem. Rev.* **2003**, *103*, 893.

- (3) Connolly, D. J.; Cusack, D.; Sullivan, T. P. O.; Guiry, P. J. *Tetrahedron* **2005**, *61*, 10153.
- (4) Mhaske, S. B.; Argade, N. P. *Tetrahedron* **2006**, *62*, 9787.
- (5) Johns, S.; Lamberton, J. A. J. *Chem. Soc., Chem. Commun.* **1965**, 267.
- (6) Fitzgerald, J. S.; Johns, S. R.; Lamberton, J. A.; Redcliffe, A. H. *Aust. J. Chem.* **1966**, *19*, 151.
- (7) Cho, J.; Bae, S.; Kim, H.; Lee, M.; Choi, Y.; Jin, B.; Lee, H. J.; Jeong, H. Y.; Lee, Y. G.; Moon, J. J. *Agric. Food Chem.* **2015**, *63*, 3587.
- (8) Moehrle, H.; Hemmerling, H. J. *Arch. Pharm. (Weinheim)* **1978**, *311*, 586.
- (9) Hassanzadeh, F.; Jafari, E.; Hakimelahi, G. H.; Rahmani Khajouei, M.; Jalali, M.; Khodarahmi, G. A. *Res Pharm Sci* **2012**, *7*, 87.
- (10) Sutherell, C. L.; Tallant, C.; Monteiro, O. P.; Yapp, C.; Fuchs, J. E.; Fedorov, O.; Siejka, P.; Müller, S.; Knapp, S.; Brenton, J. D.; Brennan, P. E.; Ley, S. V. *J. Med. Chem.* **2016**, *57*, 5095.
- (11) Nazarenko, K. G.; Shyroka, T. I.; Tolmachev, A. A. *Synth. Commun.* **2008**, *33*, 303.
- (12) Dunn, A. D.; Kinnear, K. I. J. *Heterocycl. Chem.* **1986**, *23*, 53.
- (13) Sun, X.; Hu, Y.; Nie, S.; Yan, Y.; Zhang, X. *Adv. Synth. Catal.* **2013**, *355*, 2179.
- (14) Jones, A. M.; Lebl, T.; Patterson, S.; van Mourik, T.; Fruchtl, H. A.; Philp, D.; Slawin, A. M. Z.; Westwood, N. J. *Tetrahedron* **2009**, *65*, 563.
- (15) Zhu, Y.; Fei, Z.; Liu, M.; Jia, F.; Wu, A. *Org. Lett.* **2013**, *15*, 378.
- (16) Moorthy, J. N.; Senapati, K.; Parida, K. N.; Jhulki, S.; Sooraj, K.; Nair, N. N. *J. Org. Chem.* **2011**, *76*, 9593.
- (17) Cheng, X.; Vellalath, S.; Goddard, R.; List, B. *J. Am. Chem. Soc.* **2008**, *130*, 15786.
- (18) Tu, T.; Wang, Z.; Liu, Z.; Feng, X.; Wang, Q. *Green Chem.* **2012**, *14*, 921.
- (19) Möhrle, H.; Seidel, C.-M. *Arch. Pharm. (Weinheim)* **1976**, *309*, 542.
- (20) Elmuradov, B. Z.; Abdurazakov, A. S.; Shakhidoyatov, K. M. *Chem. Nat. Compd.* **2008**, *44*, 475.
- (21) Il'chenko, A. Y.; Nazarenko, K. G.; Shirokaya, T. I.; Tolmachev, A. A.; Tolmachev, A. I. *Ukr. Khim. Zh.* **2003**, *69*, 48.
- (22) Bubenyák, M.; Pálfi, M.; Takács, M.; Béni, S.; Szökó, E.; Noszál, B.; Kökösi, J. *Tetrahedron Lett.* **2008**, *49*, 4937.
- (23) Voitenko, Z. V.; Halaev, O. I.; Samoylenko, V. P.; Kolotilov, S. V.; Lepetit, C.; Donnadieu, B.; Chauvin, R. *Tetrahedron* **2010**, *66*, 8214.
- (24) Primary data related to this publication (NMR files, processed NMR spectra, IR files, IR pdf files) are available at <http://dx.doi.org/10.17863/CAM.523>.
- (25) Darras, F. H.; Kling, B.; Heilmann, J.; Decker, M. *ACS Med. Chem. Lett.* **2012**, *3*, 914.
- (26) Yan, J. W.; Li, Y. P.; Ye, W. J.; Chen, S. B.; Hou, J. Q.; Tan, J. H.; Ou, T. M.; Li, D.; Gu, L. Q.; Huang, Z. S. *Bioorg. Med. Chem.* **2012**, *20*, 2527.