

Intramolecular Heck Cyclization of α -Sulfenyl Enol Triflates. Asymmetric Synthesis of a Pentacyclic Cardenolide Precursor Having Functionality at C-11.

John Hynes, Jr., Larry E. Overman*, Talal Nasser and Paul V. Rucker

Department of Chemistry, 516 Physical Sciences I, University of California, Irvine, CA 92697-2025, USA

Received 5 March 1998; accepted 13 April 1998

Abstract. A concise route to the core of complex cardenolides is described. The sequence features use of a sulfone to join enantioenriched A and D ring fragments and also control intramolecular aldolization to generate ring C and an intramolecular Heck cyclization of an α -sulfenyl enol triflate to form the steroid skeleton. © 1998 Elsevier Science Ltd. All rights reserved.

Keywords: aldol reactions; Heck reactions; enantiospecificity; sterols; sulfones

In the preceding communication, we described use of an intramolecular Heck reaction to construct an enantioenriched steroid having much of the functional signature of complex cardenolides: *cis* A/B and C/D ring fusions, oxygen functionality at carbons 5, 14 and 19, and a β -oriented substituent at C-17.¹ This intermediate also contained a 9,11 double bond, which potentially could be elaborated to install the 11-hydroxyl group of complex cardenolides such as ouabain (1). However, in light of the difficulty often encountered in elaborating the 9,11-alkene of *cis* A,B-fused steroids,² we wished to develop a companion strategy that directly delivered a cardenolide precursor having C-11 at the carbonyl oxidation state. Such a strategy is outlined in Figure 1. In this communication we report a streamlined method for assembling cyclization precursors and successful intramolecular Heck cyclization to form enantiopure cardenolide precursors containing vinylsulfide functionality at C-11.






Figure 1

An improved preparation of tetracyclic ketone intermediate **11**, which exploits methyl phenyl sulfone to link enantiopure A and C ring units and an intramolecular aldol reaction to form ring C,³ is outlined in Scheme 1. Hydrindanone nitrile **4**, which is available in three steps and high overall yield from (*S*)-Hajos-Parrish ketone,^{4,5} was cleaved with ozone and esterified to give **5**.⁶ Addition of 1 equiv of this ester to the lithium salt of sulfone **7** (2.5 equiv, available in quantitative yield from enantioenriched iodide **6**⁷), provided **8**, a mixture of sulfone epimers, in high yield. Treatment of **8** with 2 equiv of SmI_2 ⁸ yielded a 1:1 mixture of hydrindanone epimers **9**.⁹ Without separation, this mixture was converted to the desired stereoisomer **11** by initial treatment with HMDS at 80 °C in DMF to generate **10** and subsequent exposure of this intermediate to excess TBAF in CHCl_3 containing silica gel.¹⁵ This intramolecular aldol approach provided **11** in 60% overall yield from **8** and is four steps shorter than the first route we developed.¹



Scheme 1

Bis-silylation of **11** with excess KHMDS and TMSCl followed by selective desilylation yielded **12** (Scheme 2).¹⁶ Sulfonylation of the potassium enolate of **12** at -78°C ,¹⁷ followed by a second enolization with KHMDS and trapping with PhNTf_2 provided enol triflates **13** in high yield. Attempted Heck closure of **13a** was complicated by loss of the TMS group, so this group was removed. Heck cyclization of **14a** using conditions optimized in our previous study¹ produced pentacycle **15a**¹⁸ in 70% yield. This crystalline product (mp $125\text{--}127^\circ\text{C}$) provided single crystals which allowed the structure of this product to be confirmed by X-ray diffraction analysis.¹⁹ In contrast, intramolecular Heck cyclization of **14b** ($\text{R} = \text{Me}$) proceeded in low yield due to competing reduction of the enol triflate.



Scheme 2

The sequence reported here represents a practical way to prepare highly functionalized steroids that contain much of the functionality found in complex cardenolides. The overall sequence from commercially available 1,3-cyclohexanedione and (*S*)-Hajos-Parrish ketone involves 17 total steps and 14 isolated and purified intermediates. The synthesis strategy disclosed here has several notable aspects of potential general utility in synthesis: (a) use of a sulfone group to join fragments and then control regiochemistry of a subsequent reductive enolization and aldol cyclization, (b) the first demonstration of intramolecular Heck reaction of an α -sulfenyl enol triflate,²⁰ and (c) an additional illustration that highly substituted alkenes, in this case tetra- and trisubstituted, can be joined by intramolecular Heck reactions. We anticipate that intermediates such as **14** will prove useful in the total synthesis of complex cardenolides such as ouabain (**1**).

Acknowledgments. Support of this investigation by NIH NIGMS grant GM-30859 is gratefully acknowledged. NMR and mass spectra were determined at UCI using instruments acquired with the assistance of the NSF and NIH shared instrumentation programs. We also thank NIH for a NRSA postdoctoral fellowship for J.H. (CA-72229-02) and UCI for a Presidential Undergraduate Research Fellowship for T. N. We are grateful to Dr. Mark Jensen for his early development of the aldol strategy in another system and to Drs. Joseph Ziller and John Greaves for obtaining X-ray and mass spectrometric analysis.

References

- Overman, L. E.; Rucker, P. V. *Tetrahedron Lett.* **1998**, 39, 4643.
- Nussim, M.; Mazur, Y.; Sondheimer, F. *J. Org. Chem.* **1964**, 29, 1120.
- Intramolecular aldolization has been employed previously to form 14-hydroxy steroids [see references 10–14]. However, to the best of our knowledge, the nucleophilic component has not previously been a 9-keto group.
- Inubushi, Y.; Kikuchi, T.; Ibuka, T.; Tanaka, K.; Saji, I.; Tokane, K. *Chem. Pharm. Bull.* **1974**, 22, 349.
- Caine, D.; Kotian, P. L. *J. Org. Chem.* **1992**, 57, 6587.
- Yang, Y.; Haino, T.; Usui, S.; Fukazawa, Y. *Tetrahedron* **1996**, 52, 2325.

7. Deng, W.; Jensen, M. S.; Overman, L. E.; Rucker, P. V.; Vionnet, J.-P. *J. Org. Chem.* **1996**, *61*, 6760.
8. Molander, G. A.; Hahn, G. *J. Org. Chem.* **1986**, *51*, 1135.
9. This mixture undoubtedly arises by indiscriminate formation of a mixture of *E*- and *Z*- samarium enolates. This scenario has been rigorously established in a related system. Unpublished studies of M. Jensen of this laboratory.
10. Yates, P.; Douglas, S. P.; Datta, S. K.; Sawyer, J. F. *Can. J. Chem.* **1988**, *66*, 2268.
11. Daniewski, A. R.; Valenta, Z.; White, P. S. *Bull. Polish Acad. Sci.* **1986**, *32*, 29.
12. Ruel, R.; Deslongchamps, P. *Can. J. Chem.* **1990**, *68*, 1917.
13. Ruel, R.; Deslongchamps, P. *Can. J. Chem.* **1992**, *70*, 1939.
14. Barlaam, B.; Boivin, J.; Elkaim, L.; Elton-Farr, S.; Zard, S. Z. *Tetrahedron* **1995**, *51*, 1675.
15. We have independently demonstrated that the β epimer of **9** is converted to **11** under these conditions. It is notable that **10** can be prepared and cleaved without unraveling to an enone.
16. $[\alpha]_D^{25} +81.9^\circ$ (*c* 1.0, CHCl₃); FTIR (thin film) 2237 and 1713 cm⁻¹; ¹H NMR (500 MHz, CDCl₃) δ 5.52 (br s, 1H), 4.98 (d, *J* = 6.5 Hz, 1H), 4.88 (d, *J* = 6.5 Hz, 1H), 4.38 (br d, *J* = 12.4 Hz, 1H), 4.05 (d, *J* = 12.4 Hz, 1H), 2.65 (app t, *J* = 6.6 Hz, 1H), 2.57-2.49 (m, 2H), 2.26-1.90 (m, 6H), 1.81-1.52 (m, 9H), 1.45-1.40 (m, 2H), 1.48 (s, 3H), 0.26 (s, 9H); ¹³C NMR (125 MHz, CDCl₃) δ 209.1, 134.9, 124.6, 121.7, 91.3, 88.1, 75.7, 68.5, 56.5, 49.9, 38.4, 38.0, 35.1, 33.9, 32.1, 30.0, 26.0, 24.6, 18.8, 18.1, 17.7, 2.7; HRMS (FAB, mNBA) *m/z* 432.2578 (432.2570 calcd for C₂₄H₃₈O₄NSi, MH).
17. Trost, B. M.; Salzmann, T. N.; Hiroi, K. *J. Am. Chem. Soc.* **1976**, *98*, 4887.
18. $[\alpha]_D^{25} +184.2^\circ$ (*c* 0.45, CHCl₃); FTIR (thin film) 3474, 2237 and 1731 cm⁻¹; ¹H NMR (500 MHz, CDCl₃) δ 7.35-7.32 (m, 2H), 7.24-7.21 (m, 3H), 5.95 (br d, *J* = 10.1 Hz, 1H), 5.63 (dt, *J* = 10.1, 3.2 Hz, 1H), 5.11 (br s, 1H), 5.10 (d, *J* = 6.2 Hz, 1H), 4.95 (d, *J* = 6.2 Hz, 1H), 3.73 (d, *J* = 11.7 Hz, 1H), 3.17 (br d, *J* = 11.1 Hz, 1H), 2.61 (br s, 1H), 2.51 (dd, *J* = 9.1, 4.5 Hz, 1H), 2.29-2.00 (m, 8H), 1.80-1.75 (m, 3H), 1.60 (br m, 1H), 1.46-1.30 (m, 2H), 1.28 (s, 3H); ¹³C NMR (125 MHz, CDCl₃) δ 135.6, 129.2, 128.8, 128.3, 127.5, 126.3, 122.3, 87.2, 83.8, 74.3, 72.1, 47.3, 45.8, 45.2, 43.4, 37.4, 32.6, 29.8, 29.7, 25.2, 23.7, 20.9, 15.7; HRMS (FAB, mNBA) *m/z* 449.2023 (449.2024 calcd for C₂₇H₃₁O₃NS, M).
19. The authors have deposited atomic coordinates for this compound with the Cambridge Crystallographic Data Centre. The coordinates can be obtained, on request, from the Director, Cambridge Crystallographic Data Centre, 12 Union Road, Cambridge, CB2 1EZ, UK.
20. Use of α -sulphenyl enol phosphates²¹ and ortho-iodoaryl sulfides²² in organopalladium reactions has been described.
21. Sato, M.; Takai, K.; Oshima, K.; Nozaki, H. *Tetrahedron Lett.* **1981**, *22*, 1609.
22. Arnau, N.; Moreno-Mañas, M.; Pleixats, R. *Tetrahedron* **1994**, *49*, 11019.