This article was downloaded by: [Tufts University] On: 09 October 2014, At: 12:46 Publisher: Taylor & Francis Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



# Nucleosides, Nucleotides and Nucleic Acids

Publication details, including instructions for authors and subscription information: <u>http://www.tandfonline.com/loi/lncn20</u>

# Synthesis of Some Novel 6-Benzyl(or Substituted Benzyl)-2-β-d-Glucopyranosyl-1,2,4-Triazolo[4,3-b] [1,2,4]Triazines as Potential Antimicrobial Chemotherapeutics

Nasser S. A. M. Khalil<sup>a</sup>, Abdel Kader Mansour<sup>b</sup> & Mohga M. Eid<sup>b</sup> <sup>a</sup> Central Laboratory for Food and Feed, Agricultural Research Center, Giza, Egypt <sup>b</sup> Department of Chemistry, Faculty of Science, Cairo University, Giza, Egypt Published online: 17 Aug 2006.

To cite this article: Nasser S. A. M. Khalil , Abdel Kader Mansour & Mohga M. Eid (2004) Synthesis of Some Novel 6-Benzyl(or Substituted Benzyl)-2-β-d-Glucopyranosyl-1,2,4-Triazolo[4,3-b][1,2,4]Triazines as Potential Antimicrobial Chemotherapeutics, Nucleosides, Nucleotides and Nucleic Acids, 23:12, 1889-1910, DOI: <u>10.1081/NCN-200040663</u>

To link to this article: http://dx.doi.org/10.1081/NCN-200040663

### PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <a href="http://www.tandfonline.com/page/terms-and-conditions">http://www.tandfonline.com/page/terms-and-conditions</a>

## Synthesis of Some Novel 6-Benzyl(or Substituted Benzyl)-2-β-D-Glucopyranosyl-1,2,4-Triazolo[4,3-*b*][1,2,4]Triazines as Potential Antimicrobial Chemotherapeutics

Nasser S. A. M. Khalil,<sup>1,\*</sup> Abdel Kader Mansour,<sup>2</sup> and Mohga M. Eid<sup>2</sup>

<sup>1</sup>Central Laboratory for Food and Feed, Agricultural Research Center, Giza, Egypt <sup>2</sup>Department of Chemistry, Faculty of Science, Cairo University, Giza, Egypt

#### ABSTRACT

Glucosidation of the new 8-amino-6-benzyl(or substituted benzyl)-2,8-dihydro-1,2,4-triazolo[4,3-b][1,2,4]triazin-7(3H)-ones (**3a-d**) with 2,3,4,6-tetra-O-acetyl- $\alpha$ -D-glucosyl bromide **4** gave the corresponding N-glucosides **5a-d**. Chemical transformations leading to new functionalities have also been achieved to give compounds **7–12**. Antimicrobial activity of compounds **5a-c** against Aspergillus fumigatus, Penicillium italicum, Syncephalastrum racemosum, Candida albicans, Staphylococcus aureus, Pseudomonas aeruginosa, Bacillus subtilis, Escherichia coli is described.

*Key Words:* Synthesis; 1,2,4-Triazolo[4,3-*b*][1,2,4]triazines;  $\beta$ -D-Glucopyranosyls; Antimicrobial chemotherapeutics.

1889

DOI: 10.1081/NCN-200040663 Copyright © 2004 by Marcel Dekker, Inc. 1525-7770 (Print); 1532-2335 (Online) www.dekker.com

<sup>\*</sup>Correspondence: Nasser S. A. M. Khalil, Central Laboratory for Food and Feed, Agricultural Research Center, Giza, Egypt; E-mail: Nasserkhalil\_23@hotmail.com.

#### **INTRODUCTION**

1,2,4-Triazoles,<sup>[1-5]</sup> 1,2,4-triazines<sup>[6]</sup> and their glycosides<sup>[7-20]</sup> are reported to present a plethora of biological activities. The 1,2,4-triazole nucleus is associated with diverse pharmacological activities such as antibacterial, antifungal, hypoglycemic, antihypertensive and analgesic properties.<sup>[1-5]</sup> The synthesis and investigation of biological activity of 1,2,4-triazole glycosides have been stimulated by the finding that Ribavirin ( $\beta$ -D-ribofuranosyl-1,2,4-triazole-3-carboxamide),<sup>[9]</sup> is remarkable in its broad spectrum activity against DNA and RNA viruses.<sup>[10,11]</sup> Ribavirin has been developed clinically<sup>[12]</sup> and approved for human use in many pharmaceutical preparations. Many *N*-glycosides of 1,2,4-triazines possess cytotoxic, antiviral, enzyme inhibiting, immunosuppressive, antiphlogistic, antipsoriatic, bacteriostatic, and antitumor activity.<sup>[7,8,13-20]</sup> Recently, we have synthesized some 6-substituted-2-glycosyl-1,2,4-triazines that showed in vitro activity against MCF7 (Breast) and SF-268 cell lines in a primary human anticancer assay.<sup>[21,22]</sup> All these facts animated us to synthesize some new 6-benzyl(or substituted benzyl)-2- $\beta$ -D-glucopyranosyl-1,2,4-triazolo[4,3-*b*][1,2,4]triazin-7(3*H*)-ones and report the antimicrobial screening of three selected compounds.

#### **RESULTS AND DISCUSSION**

Hydrazinolysis of 4-amino-6-(3,4-dimethoxybenzyl)-3-methylthio-1,2,4-triazin-5(4*H*)one (**1e**) followed by its cyclocondensation with carbon disulfide was previously reported to offer a covenient synthesis of the 8-amino-6-(3,4-dimethoxybenzyl)-2,8dihydro-3-thioxo-1,2,4-triazolo[4,3-*b*][1,2,4]-triazin-7(3*H*)-one (**3e**).<sup>[23]</sup> In the present

Table 1. Antimicrobial activity of compounds 5a-c compared to standard antimicrobial agents.

Test organisms	Compounds											
	5a			5b			5c			St.		
	5	2.5	1	5	2.5	1	5	2.5	1	5	2.5	1
Aspergillus fumigatus	0	0	0	0	0	0	+	0	0	+++	+++	++
Penicillium italicum	0	0	0	0	0	0	0	0	0	+++	+++	++
Syncephalastrum racemosum	0	0	0	0	0	0	0	0	0	+++	+++	+++
Candida albicans	0	0	0	0	0	0	+	+	+	++	++	++
Staphylococcus aureus	0	0	0	0	0	0	0	0	0	++	++	++
Pseudomonas aeruginosa	0	0	0	+	+	+	0	0	0	+++	++	++
Bacillus subtilis	0	0	0	0	0	0	0	0	0	++	++	++
Escherichia coli	0	0	0	0	0	0	0	0	0	+++	++	++

St. = Reference standard; Chloramphenicol was used as a standard antibacterial agent and Terbinafin was used as a standard antifungal agent.

100  $\mu l$  of each conc. (5, 2.5, 1 mg/mL) was tested.

The test was done using the diffusion agar technique.

Inhibition values = 0.1-0.5 cm beyond contol = +; Inhibition values = 0.6-1.0 cm beyond contol = ++; Inhibition values = 1.1-1.5 cm beyond contol = +++; 0 = Not detected.



investigation, the new 8-amino-6-benzyl (or substituted benzyl)-2,8-dihydro-3-thioxo-1,2,4-triazolo[4,3-*b*][1,2,4]-triazin-7(3*H*)-ones (**3a–d**) were analogously synthesized and allowed to react with 2,3,4,6-tetra-*O*-acetyl- $\alpha$ -D-glucopuranosyl bromide (**4**) in pyridine containing triethylamine to afford the corresponding 8-amino-6-benzyl (or substituted benzyl)-2-(2,3,4,6-tetra-*O*-acetyl- $\beta$ -D-glucopyranosyl)-2,8-dihydro-3-thioxo-1,2,4-triazolo[4,3-*b*][1,2,4]-triazin-7(3*H*)-one pyridinium salts (**5a–d**). The structure of compounds **5a–d** was inferred from their chemical and spectral evidences (a chemical





evidence comes from oxidation of compounds **9a–d** to compounds **11a–d** as will be seen later). Thus, the position of the anomeric proton at  $\delta$  7.2–6.9 (J = 9.0-9.4 Hz) consistent with similar reported data<sup>[21,22,24–28]</sup> confirms the  $\beta$ -*N*- structure of compounds **5a–d** and excludes the possible isomeric  $\beta$ -*S*- structure of compounds **6a–d**. The anomeric proton of similar  $\beta$ -*N*- glucosides having an adjacent C = S was reported to appear more downfield (7.0–6.0) than that for the  $\beta$ -*S* glucoside ( $\delta$  5.7–5.5) due to the anisotropic deshielding effect of the C = S.<sup>[29]</sup> The <sup>1</sup>H NMR of Compounds **5a–d** showed the pyridinyl protons at  $\delta$  8.9–7.8. Treatment of compounds **5a–d** with acetic acid gave the corresponding 8-amino-6-benzyl (or substituted benzyl)-2-(2,3,4,6tetra-*O*-acetyl- $\beta$ -D-glucopyranosyl)-2,8-dihydro-3-thioxo-1,2,4-triazolo[4,3-*b*][1,2,4]-triazin-7(3*H*)-ones (**7a–d**) whose <sup>1</sup>H NMR spectra showed the absence of the pyridinyl protons at  $\delta$  8.9–7.8.



Condensation of compounds **7a-d** with aromatic aldehydes gave the corresponding 8-arylideneamino derivatives **8a-z**. The <sup>1</sup>H NMR spectra of these compounds not only showed the absence of the NH<sub>2</sub> proton signal at  $\delta$  6.50–6.38, but also revealed the N = CH proton signal at  $\delta$  8.43–8.29 consistent with similar reported data.<sup>[21,22,24–30]</sup>

Deamination of compounds 7a-d was achieved by the action of nitrous acid to give the 6-benzyl (or substituted benzyl)-2-(2,3,4,6-tetra-*O*-acetyl- $\beta$ -D-glucopyranosyl)-2,8-dihydro-3-thioxo-1,2,4-triazolo[4,3-*b*][1,2,4]-triazin-7(3*H*)-ones (**9a-d**). Assignment of the structure of compounds **9a-d** was based on spectroscopic and chemical evidences. Thus, the <sup>1</sup>H NMR spectra of these compounds showed the absence of the NH<sub>2</sub> proton signal at  $\delta$  6.50–6.38 and the appearance of the NH proton signal at  $\delta$  9.57–9.41. While thiation of compounds **9a-d** gave the corresponding dithiones **10a-d**, their oxidation gave the corresponding diones **11a-d**. The structure of compounds **10a-d**, **11a-d** was confirmed from their correct analytical and spectral data.

Deacetylation of compounds 9a-d via methanolic ammonia treatment afforded the corresponding free  $\beta$ -D-glucopyranosides 12a-d.

#### **BIOLOGICAL EVALUATION**

Compounds 5a-c were tested for antimicrobial activity (in vitro) against Aspergillus fumigatus, Penicillium italicum, Syncephalastrum racemosum, Candida albicans, Staphylococcus aureus, Pseudomonas aeruginosa, Bacillus subtilis, Escherichia coli. The inhibitory effects of compounds 5a-c against these organisms are given in Table 1. The screening results of compound 5b showed moderate activity against Pseudomonas aeruginosa at all tested concentrations (5, 2.5, 1 mg/mL). Compound 5cnot only showed moderate activity against Candida albicans at all tested concentrations (5, 2.5, 1 mg/mL), but also showed moderate activity against Aspergillus fumigatus at a concentration of 5 mg/mL (Schemes 1–3: Synthesis of some 2-â-D-glucopyranosyl-1,2,4-triazolo[4,3-b][1,2,4]triazines possessing different functionalities.).

#### **EXPERIMENTAL**

All melting points are uncorrected. IR spectra were recorded on a Perkin Elmer 1430 spectrometer. <sup>1</sup>H NMR spectra were recorded at 200 MHz with a Varian GEMINI 200 spectrometer. Elemental analyses were carried out at the Micro Analytical Center, Cairo University, Giza, Egypt. Antimicrobial screening of compounds **5a–c** was carried out at the Medical Mycology Lab., The regional center for Mycology and Biotechnology, Al Azhar University, Cairo, Egypt.

**4-Amino-6-benzyl(or substituted benzyl)-3-methylthio-4,5-dihydro-1,2,4-triazin-5-ones (1a–d).** General procedure: The appropriate 4-amino-6-benzyl(or substituted benzyl)-3-thioxo-2,3-dihydro-1,2,4-triazin-5(4H)-one<sup>[21,31–33]</sup> (10 mmol) was dissolved in cold methanolic sodium methoxide solution [prepared from sodium metal (10 mmol) and 10 mL methanol], then methyl iodide (10 mmol) was added. The reaction mixture was stirred for 30 minutes and left overnight at room temperature. The precipitate obtained was collected by filtration, washed with water, dried at room temperature and recrystallised from ethanol to give colorless crystals of compounds 1a-d.

**4-Amino-6-benzyl-3-methylthio-4,5-dihydro-1,2,4-triazin-5-one (1a).** Using the general procedure, 4-amino-6-benzy-3-thioxo-2,3-dihydro-1,2,4-triazin-5(4*H*)-one<sup>[21,32,33]</sup> gave **1a** (87%); mp 196–198°C; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>)  $\delta$  7.72–6.99 (m, 5H, ArH), 6.35 (s, 2H, NH<sub>2</sub>-exchangeable), 4.11 (s, 2H, C<sub>6</sub>H<sub>5</sub>CH<sub>2</sub>), 2.62 (s, 3H, SCH<sub>3</sub>). Anal. Calcd. for C<sub>11</sub>H<sub>12</sub>N<sub>4</sub>OS: C, 53.21; H, 4.87; N, 22.56. Found: C, 53.25; H, 5.00; N, 22.49.

4-Amino-6-(4-methylbenzyl)-3-methylthio-4,5-dihydro-1,2,4-triazin-5-one (1b). Using the general procedure, 4-amino-6-(4-methylbenzyl)-3-thioxo-2,3-dihydro-1,2,4-triazin-5(4*H*)-one<sup>[21,33]</sup> gave 1b (92%); mp 182–184°C; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>)  $\delta$  7.42 (d, 2H, J = 7.3 Hz, ArH), 6.73 (d, 2H, J = 7.6 Hz, ArH), 6.41 (s, 2H, NH<sub>2</sub>-exchangeable),

#### 1,2,4-Triazolo[4,3-b][1,2,4]Triazines

4.07 (s, 2H, 4-CH<sub>3</sub>C<sub>6</sub>H<sub>4</sub>CH<sub>2</sub>), 2.69 (s, 3H, SCH<sub>3</sub>), 2.24 (s, 3H, 4-CH<sub>3</sub>C<sub>6</sub>H<sub>4</sub> CH<sub>2</sub>). Anal. Calcd. for  $C_{12}H_{14}N_4OS$ : C, 54.94; H, 5.38; N, 21.36. Found: C, 55.02; H, 5.42; N, 21.58.

**4-Amino-6-(4-methoxybenzyl)-3-methylthio-4,5-dihydro-1,2,4-triazin-5-one** (**1c**). Using the general procedure, 4-amino-6-(4-methoxybenzyl)-3-thioxo-2,3-dihydro-1,2,4-triazin-5(4*H*)-one<sup>[21,33]</sup> gave **1c** (85%); mp 140°C; (lit. mp,<sup>[33]</sup> 140°C); <sup>1</sup>H NMR (DMSO-d<sub>6</sub>)  $\delta$  7.48 (d, 2H, *J* = 7.8 Hz, ArH), 6.78 (d, 2H, *J* = 7.6 Hz, ArH), 6.37 (s, 2H, NH<sub>2</sub>-exchangeable), 4.08 (s, 2H, 4-CH<sub>3</sub>OC<sub>6</sub>H<sub>4</sub>CH<sub>2</sub>), 3.77 (s, 3H, 4-CH<sub>3</sub>OC<sub>6</sub>H<sub>4</sub>CH<sub>2</sub>). Anal. Calcd. for C<sub>12</sub>H<sub>14</sub>N<sub>4</sub>O<sub>2</sub>S: C, 51.78; H, 5.07; N, 20.13. Found: C, 51.81; H, 5.09; N, 20.26.

**4-Amino-6-(4-chlorobenzyl)-3-methylthio-4,5-dihydro-1,2,4-triazin-5-one (1d).** Using the general procedure, 4-amino-6-(4-chlorobenzyl)-3-thioxo-2,3-dihydro-1,2,4-triazin-5(4*H*)-one<sup>[21,32,33]</sup> gave **1d** (89%); mp 190°C; (lit. mp,<sup>[33]</sup> 188°C); <sup>1</sup>H NMR (DMSO-d<sub>6</sub>)  $\delta$  7.33 (d, 2H, *J* = 7.5Hz, ArH), 6.75 (d, 2H, *J* = 7.8 Hz, ArH), 6.45 (s, 2H, NH<sub>2</sub>-exchangeable), 4.11 (s, 2H, 4-ClC<sub>6</sub>H<sub>4</sub>CH<sub>2</sub>). Anal. Calcd. for C<sub>11</sub>H<sub>11</sub>ClN<sub>4</sub>OS: C, 46.73; H, 3.92; N, 19.81. Found: C, 46.81; H, 3.87; N, 19.77.

4-Amino-6-benzyl(or substituted benzyl)-3-hydrazino-4,5-dihydro -1,2,4-triazin-5-ones (2a-d). General procedure: A solution of each of compounds 1a-d (10 mmol) and hydrazine hydrate (80%; 10 mL) in 2-propanol (50 mL) was heated at reflux temperature for 3 hours. The precipitate formed on cooling was collected by filtration and recrystallised from DMF/AcOH to give colorless crystals of compounds 2a-d.

**4-Amino-6-benzyl-3-hydrazino-4,5-dihydro-1,2,4-triazin-5-one (2a).** Using the general procedure, **1a** gave **2a** (65%); mp 254°C; IR (KBr) 3394, 3284 (NH, NH<sub>2</sub>), 1672 (C = O amide); <sup>1</sup>H NMR (DMSO-d<sub>6</sub>)  $\delta$  9–8 (brs, 1H, NH-exchangeable), 7.52–7.15 (m, 5H, ArH), 6.41, 4.53 (2s, 2H each, NH<sub>2</sub>-exchangeable and NHNH<sub>2</sub>-exchangeable), 4.08 (s, 2H, C<sub>6</sub>H<sub>5</sub>CH<sub>2</sub>). Anal. Calcd. for C<sub>10</sub>H<sub>12</sub>N<sub>6</sub>O: C, 51.72; H, 5.21; N, 36.19. Found: C, 51.84; H, 5.43; N, 36.38.

**4-Amino-6-(4-methylbenzyl)-3-hydrazino-4,5-dihydro-1,2,4-triazin-5-one** (2b). Using the general procedure, **1b** gave **2b** (74%); mp 264°C; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>)  $\delta$  8.32 (s, 1H, NH-exchangeable), 7.35 (d, 2H, J = 7.4 Hz, ArH), 6.89 (d, 2H, J = 7.8 Hz, ArH), 6.35, 4.50 (2s, 2H each, NH<sub>2</sub>-exchangeable and NHNH<sub>2</sub>-exchangeable), 4.12 (s, 2H, 4-CH<sub>3</sub>C<sub>6</sub>H<sub>4</sub>CH<sub>2</sub>), 2.26 (s, 3H, 4-CH<sub>3</sub>C<sub>6</sub>H<sub>4</sub> CH<sub>2</sub>). Anal. Calcd. for C<sub>11</sub>H<sub>14</sub>N<sub>6</sub>O: C, 53.65; H, 5.73; N, 34.12. Found: C, 53.82; H, 5.84; N, 34.29.

**4-Amino-6-(4-methoxybenzyl)-3-hydrazino-4,5-dihydro-1,2,4-triazin-5-one (2c).** Using the general procedure, **1c** gave **2c** (70%); mp 243–244°C ; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 8.89 (s, 1H, NH-exchangeable), 7.29 (d, 2H, J = 7.7 Hz, ArH), 6.99 (d, 2H, J = 7.5 Hz, ArH), 6.29, 4.54 (2s, 2H each, NH<sub>2</sub>–exchangeable and NHNH<sub>2</sub>-exchangeable), 4.07 (s, 2H, 4-CH<sub>3</sub>OC<sub>6</sub>H<sub>4</sub>CH<sub>2</sub>), 3.78 (s, 3H, 4-CH<sub>3</sub>OC<sub>6</sub>H<sub>4</sub>CH<sub>2</sub>). Anal. Calcd. for C<sub>11</sub>H<sub>14</sub>N<sub>6</sub>O<sub>2</sub>: C, 50.38; H, 5.38; N, 32.04. Found: C, 50.49; H, 5.29; N, 32.09.

4-Amino-6-(4-chlorobenzyl)-3-hydrazino-4,5-dihydro-1,2,4-triazin-5-one (2d). Using the general procedure, 1d gave 2d (70%); mp  $262-264^{\circ}C$  (lit. mp,<sup>[33]</sup>  $260^{\circ}C$ );

<sup>1</sup>H NMR (DMSO-d<sub>6</sub>)  $\delta$  9.2 (s, 1H, NH-exchangeable), 7.35 (d, 2H, J = 7.6 Hz, ArH), 6.84 (d, 2H, J = 7.3 Hz, ArH), 6.41, 4.51 (2s, 2H each, NH<sub>2</sub>-exchangeable and NHNH<sub>2</sub>-exchangeable), 4.12 (s, 2H, 4-CH<sub>3</sub>OC<sub>6</sub>H<sub>4</sub>CH<sub>2</sub>). Anal. Calcd. for C<sub>10</sub>H<sub>11</sub>ClN<sub>6</sub>O: C, 45.04; H, 4.16; N, 31.51. Found: C, 45.10; H, 4.02; N, 31.72.

4-Amino-6-benzyl(or substituted benzyl)-2,8-dihydro-3-thioxo-1,2,4-triazolo[4, 3-b][1,2,4]-triazin-7(3H)-ones (3a-d). General procedure: A mixture of each of compounds 2a-d (10 mmol) and carbon disulfide (20 mL) in pyridine (50 mL) was heated at reflux temperature for 3 hours. After cooling, the reaction mixture was diluted with water and the formed precipitate was collected by filtration and recrystallized from DMF/AcOH to give pale crystals of compounds 3a-d.

8-Amino-6-benzyl-2,8-dihydro-3-thioxo-1,2,4-triazolo[4,3-*b*][1,2,4]-triazin-7(3*H*)-one (3a). Using the general procedure, 2a gave 3a (70%); mp 218°C; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 9.29 (s, 1H, NH-exchangeable), 7.42–7.13 (m, 5H, ArH), 6.38 (s, 2H, NH<sub>2</sub>-exchangeable), 4.11 (s, 2H, C<sub>6</sub>H<sub>5</sub>CH<sub>2</sub>). Anal. Calcd. for C<sub>11</sub>H<sub>10</sub>N<sub>6</sub>OS: C, 48.17; H, 3.67; N, 30.64. Found: C, 47.95; H, 3.74; N, 30.79.

8-Amino-6-(4-methylbenzyl)-2,8-dihydro-3-thioxo-1,2,4-triazolo[4,3-*b*][1,2,4]-triazin-7(3*H*)-one (3b). Using the general procedure, 2b gave 3b (60%); mp 223°C; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 9.49 (s, 1H, NH-exchangeable), 7.25 (d, 2H, J = 7.4 Hz, ArH), 6.86 (d, 2H, J = 7.9 Hz, ArH), 6.37 (s, 2H, NH<sub>2</sub>-exchangeable), 4.09 (s, 2H, 4-CH<sub>3</sub>C<sub>6</sub>H<sub>5</sub>CH<sub>2</sub>). 2.24 (s, 3H, 4-CH<sub>3</sub>C<sub>6</sub>H<sub>4</sub>CH<sub>2</sub>). Anal. Calcd. for C<sub>12</sub>H<sub>12</sub>N<sub>6</sub>OS: C, 49.99; H, 4.20; N, 29.15. Found: C, 50.21; H, 4.29; N, 28.98.

8-Amino-6-(4-methoxybenzyl)-2,8-dihydro-3-thioxo-1,2,4-triazolo[4,3-*b*][1,2,4]-triazin-7(3*H*)-one (3c). Using the general procedure, 2c gave 3c (67%); mp 203°C; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 9.36 (s, 1H, NH-exchangeable), 7.33, 7.24 (2d, 2H each, J = 7.2 Hz, ArH), 6.39 (s, 2H, NH<sub>2</sub>-exchangeable), 4.11 (s, 2H, 4-CH<sub>3</sub>OC<sub>6</sub>H<sub>4</sub>CH<sub>2</sub>) 3.84 (s, 3H, 4-CH<sub>3</sub>OC<sub>6</sub>H<sub>4</sub>CH<sub>2</sub>). Anal. Calcd. for C<sub>12</sub>H<sub>12</sub>N<sub>6</sub>O<sub>2</sub>S: C, 47.36; H, 3.97; N, 27.61. Found: C, 47.42; H, 4.04; N, 27.72.

8-Amino-6-(4-chlorobenzyl)-2,8-dihydro-3-thioxo-1,2,4-triazolo[4,3-*b*][1,2,4]triazin-7(3*H*)-one (3d). Using the general procedure, 2d gave 3d (75%); mp 237°C; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 9.26 (s, 1H, NH-exchangeable), 7.39, 7.19 (2d, 2H each, J = 7.6 Hz, ArH), 6.44 (s, 2H, NH<sub>2</sub>-exchangeable), 4.07 (s, 2H, 4-ClC<sub>6</sub>H<sub>4</sub>CH<sub>2</sub>). Anal. Calcd. for C<sub>11</sub>H<sub>9</sub>ClN<sub>6</sub>OS: C, 42.79; H, 2.94; N, 27.22. Found: C, 42.88; H, 3.02; N, 27.14.

8-Amino-6-benzyl(or substituted benzyl)-2-(2,3,4,6-tetra-O-acetyl- $\beta$ -D-glucopyranosyl)-2,8-dihydro-3-thioxo-1,2,4-triazolo[4,3-b][1,2,4]-triazin-7(3H)-one pyridinium salts (5a-d). General procedure: 2,3,4,6-Tetra-O-acetyl- $\alpha$ -D-glucopyranosyl bromide (4) (10 mmol) was added to a solution of each of compounds 3a-d in pyridine (5 mL) and triethylamine (2 mL, 14 mmol). After shaking the reaction mixture for 15 minutes, it was left overnight at room temperature. The next day, the reaction mixture was diluted with water and the precipitate formed was collected by filtration, dried at room temperature and dissolved in dichloromethane (25 mL) to which was

#### 1,2,4-Triazolo[4,3-b][1,2,4]Triazines

added charcoal (0.5 g), celite (0.4 g), and silica gel 60 GF254 (70–230 mesh; 0.1 g). The later mixture was shaken for 1 hour, left at room temperature for another 1 hour, filtered and the solvent was evaporated on a rotavap. The obtained residue was recrystallised from dichloromethane/petroleum ether (bp 40–60°C) to give colorless crystals of compounds **5a–d**.

8-Amino-6-benzyl-2-(2,3,4,6-tetra-*O*-acetyl-β-D-glucopyranosyl)-2,8-dihydro-3thioxo-1,2,4-triazolo[4,-3-*b*][1,2,4]-triazin-7(3*H*)-one pyridinium salt (5a). Using the general procedure, 3a gave 5a (61%); mp 196°C; IR (KBr) 3425, 3256 (NH<sub>2</sub>), 1751 (C = O acetate), 1674 (C = O amide) cm<sup>-1</sup>; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 8.78 (d, 2H, J = 4.9 Hz, C<sub>5</sub>H<sub>5</sub>N), 8.31 (dt, 1H, J = 1.5, 7.8 Hz, C<sub>5</sub>H<sub>5</sub>N), 7.85 (t, 2H, J = 6.83 Hz, C<sub>5</sub>H<sub>5</sub>N), 7.59–7.21 (m, 5H, ArH), 7.22 (d, 1H, J = 9.4 Hz, H-1'), 6.53 (s, 2H, NH<sub>2</sub>exchangeable), 5.48 (t, 1H, J = 9.5 Hz, H-2'), 5.41 (t, 1H, J = 9.5 Hz, H-3'), 5.01 (t, 1H, J = 9.5 Hz, H-4'), 4.23–4.07 (m, 5H, H-5', H-6', C<sub>6</sub>H<sub>5</sub>CH<sub>2</sub>), 2.07–1.93 (4s, 12H, CH<sub>3</sub>CO). Anal. Calcd. for C<sub>25</sub>H<sub>28</sub>N<sub>6</sub>O<sub>10</sub>S.C<sub>5</sub>H<sub>5</sub>N: C, 52.70; H, 4.87; N, 14.34; S, 4.69. Found: C, 52.78; H, 4.73; N, 14.26; S, 4.85.

8-Amino-6-(4-methylbenzyl)-2-(2,3,4,6-tetra-*O*-acetyl-β-D-glucopyranosyl)-2,8dihydro-3-thioxo-1,2,4-triazolo[4,3-*b*][1,2,4]-triazin-7(3*H*)-one pyridinium salt (5b). Using the general procedure, **3b** gave **5b** (69%); mp 176°C; IR (KBr) 3448, 3248 (NH<sub>2</sub>), 1751 (C = O acetate), 1674 (C = O amide) cm<sup>-1</sup>; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 8.84 (d, 2H, J = 4.9 Hz, C<sub>5</sub>H<sub>5</sub>N), 8.37 (dt, 1H, J = 1.5, 7.1 Hz, C<sub>5</sub>H<sub>5</sub>N), 7.89 (t, 2H, J = 6.43 Hz, C<sub>5</sub>H<sub>5</sub>N), 7.39, 7.32 (2d, 2H each, J = 7.3 Hz, ArH), 7.08 (d, 1H, J = 9.0 Hz, H-1′), 6.47 (s, 2H, NH<sub>2</sub>-exchangeable), 5.49 (t, 1H, J = 9.2 Hz, H-2′), 5.41 (t, 1H, J = 9.2 Hz, H-3′), 5.01 (t, 1H, J = 9.2 Hz, H-4′), 4.22–3.98 (m, 5H, H-5′, H-6′, 4-CH<sub>3</sub>C<sub>6</sub>H<sub>4</sub>CH<sub>2</sub>), 2.26 (s, 3H, 4-CH<sub>3</sub>C<sub>6</sub>H<sub>4</sub>CH<sub>2</sub>), 2.07–1.94 (4s, 12H, CH<sub>3</sub>CO). Anal. Calcd. for C<sub>26</sub>H<sub>30</sub>N<sub>6</sub>O<sub>10</sub>S.C<sub>5</sub>H<sub>5</sub>N: C, 53.37; H, 5.06; N, 14.05. Found: C, 53.12; H, 4.97; N, 13.95.

8-Amino-6-(4-methoxybenzyl)-2-(2,3,4,6-tetra-*O*-acetyl-β-D-glucopyranosyl)-2,8-dihydro-3-thioxo-1,2,4-triazolo[4,3-*b*][1,2,4]-triazin-7(3*H*)-one pyridinium salt (5c). Using the general procedure, 3c gave 5c (79%); mp 182°C; IR (KBr) 3448, 3248 (NH<sub>2</sub>), 1751 (C = O acetate), 1674 (C = O amide) cm<sup>-1</sup>; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 8.91 (dt, 2H, J = 0.8, 5.9 Hz, C<sub>5</sub>H<sub>5</sub>N), 8.54 (t, 1H, J = 7.4 Hz, C<sub>5</sub>H<sub>5</sub>N), 8.03 (t, 2H, J = 7.04 Hz, C<sub>5</sub>H<sub>5</sub>N), 7.36, 7.22 (2d, 2H each, J = 7.4 Hz, ArH), 6.91 (d, 1H, J = 9.0 Hz, H-1'), 6.45 (s, 2H, NH<sub>2</sub>-exchangeable), 5.51 (t, 1H, J = 9.3 Hz, H-2'), 5.41 (t, 1H, J = 9.3 Hz, H-3'), 5.01 (t, 1H, J = 9.3 Hz, H-4'), 4.22–3.95 (m, 5H, H-5', H-6', 4-CH<sub>3</sub>OC<sub>6</sub>H<sub>4</sub>CH<sub>2</sub>), 3.77 (s, 3H, 4-CH<sub>3</sub>OC<sub>6</sub>H<sub>4</sub>CH<sub>2</sub>), 2.08–1.99 (4s, 12H, CH<sub>3</sub>CO). Anal. Calcd. for C<sub>26</sub>H<sub>30</sub>N<sub>6</sub>O<sub>11</sub>S.C<sub>5</sub>H<sub>5</sub>N: C, 52.17; H, 4.94; N, 13.74. Found: C, 52.13; H, 4.72; N, 13.66.

8-Amino-6-(4-chlorobenzyl)-2-(2,3,4,6-tetra-*O*-acetyl-β-D-glucopyranosyl)-2,8dihydro-3-thioxo-1,2,4-triazolo[4,3-*b*][1,2,4]-triazin-7(3*H*)-one pyridinium salt (5d). Using the general procedure, 3d gave 5d (72%); mp 198–200°C; IR (KBr) 3447, 3242 (NH<sub>2</sub>), 1751 (C = O acetate), 1674 (C = O amide) cm<sup>-1</sup>; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 8.92 (dt, 2H, J = 0.9, 5.8 Hz, C<sub>5</sub>H<sub>5</sub>N), 8.56 (t, 1H, J = 7.3 Hz, C<sub>5</sub>H<sub>5</sub>N), 8.01 (t, 2H, J = 7.1Hz, C<sub>5</sub>H<sub>5</sub>N), 7.35, 7.21 (2d, 2H each, J = 7.3 Hz, ArH), 6.95 (d, 1H, J = 9.0 Hz, H-1'), 6.44 (s, 2H, NH<sub>2</sub>-exchangeable), 5.50 (t, 1H, J = 9.1 Hz, H-2'), 5.40 (t, 1H, J = 9.1 Hz, H-3'), 5.01 (t, 1H, J = 9.1 Hz, H-4'), 4.21–3.96 (m, 5H, H-5', H-6', 4-ClC<sub>6</sub>H<sub>4</sub>CH<sub>2</sub>), 2.09–1.98 (4s, 12H, CH<sub>3</sub>CO). Anal. Calcd. for C<sub>25</sub>H<sub>27</sub>ClN<sub>6</sub>O<sub>10</sub>S.C<sub>5</sub>H<sub>5</sub>N: C, 50.18; H, 4.49; N, 13.65. Found: C, 49.97 H, 4.70; N, 13.61.

8-Amino-6-benzyl(or substituted benzyl)-2-(2,3,4,6-tetra-*O*-acetyl-β-D-glucopyranosyl)-2,8-dihydro-3-thioxo-1,2,4-triazolo[4,3-*b*][1,2,4]-triazin-7(3*H*)-ones (7a-d). General procedure: Each of compounds 5a-d (10 mmol) was dissolved in ethanol (25 mL) then acetic acid (80%; 25 mL) was added. The previous solution was just boiled and stirred overnight at room temperature. The excess ethanol was evaporated on a rotavap (till the solution came nearly to its half volume), then water (50 mL) was added and the mixture was left overnight in the refrigerator. The formed precipitate was collected by filtration, washed well with water, dried at room temperature and recrystallised from dichloromethane/petroleum ether (bp. 40–60°C) to give colorless crystals of compounds 7a-d.

8-Amino-6-benzyl-2-(2,3,4,6-tetra-*O*-acetyl-β-D-glucopyranosyl)-2,8-dihydro-3thioxo-1,2,4-triazolo[4,3-*b*][1,2,4]-triazin-7(3*H*)-one (7a). Using the general procedure, 5a gave 7a (70%); mp 212°C; IR (KBr) 3420, 3258 (NH<sub>2</sub>), 1751 (C = O acetate), 1674 (C = O amide) cm<sup>-1</sup>; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 7.59–7.21 (m, 5H, ArH), 7.12 (d, 1H, J = 9.4 Hz, H-1'), 6.50 (s, 2H, NH<sub>2</sub>-exchangeable), 5.47 (t, 1H, J = 9.5 Hz, H-2'), 5.42 (t, 1H, J = 9.5 Hz, H-3'), 5.03 (t, 1H, J = 9.5 Hz, H-4'), 4.22–4.08 (m, 5H, H-5', H-6', C<sub>6</sub>H<sub>5</sub>CH<sub>2</sub>), 2.09–1.92 (4s, 12H, CH<sub>3</sub>CO). Anal. Calcd. for C<sub>25</sub>H<sub>28</sub>N<sub>6</sub>O<sub>10</sub>S: C, 49.67; H, 4.67; N, 13.90. Found: C, 49.57; H, 4.70; N, 14.00.

8-Amino-6-(4-methylbenzyl)-2-(2,3,4,6-tetra-*O*-acetyl-β-D-glucopyranosyl)-2,8dihydro-3-thioxo-1,2,4-triazolo[4,3-*b*][1,2,4]-triazin-7(3*H*)-one (7b). Using the general procedure, **5b** gave 7b (72%); mp 202°C; IR (KBr) 3447, 3242 (NH<sub>2</sub>), 1751 (C = O acetate), 1674 (C = O amide) cm<sup>-1</sup>; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 7.41, 7.31 (2d, 2H each, J = 7.2 Hz, ArH), 6.99 (d, 1H, J = 9.1 Hz, H-1'), 6.45 (s, 2H, NH<sub>2</sub>-exchangeable), 5.48 (t, 1H, J = 9.4 Hz, H-2'), 5.42 (t, 1H, J = 9.4 Hz, H-3'), 5.00 (t, 1H, J = 9.4 Hz, H-4'), 4.23–3.97 (m, 5H, H-5', H-6', 4-CH<sub>3</sub>C<sub>6</sub>H<sub>4</sub>CH<sub>2</sub>), 2.26 (s, 3H, 4-CH<sub>3</sub>C<sub>6</sub>H<sub>4</sub>CH<sub>2</sub>), 2.08–1.93 (4s, 12H, CH<sub>3</sub>CO). Anal. Calcd. for C<sub>26</sub>H<sub>30</sub>N<sub>6</sub>O<sub>10</sub>S: C, 50.48; H, 4.89; N, 13.58. Found: C, 50.50; H, 4.90; N, 13.52.

8-Amino-6-(4-methoxybenzyl)-2-(2,3,4,6-tetra-*O*-acetyl-β-D-glucopyranosyl)-2,8-dihydro-3-thioxo-1,2,4-triazolo[4,3-*b*][1,2,4]-triazin-7(3*H*)-one (7c). Using the general procedure, 5c gave 7c (67%); mp 198°C; IR (KBr) 3450, 3250 (NH<sub>2</sub>), 1751 (C = O acetate), 1674 (C = O amide) cm<sup>-1</sup>; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 7.33, 7.20 (2d, 2H each, J = 7.4 Hz, ArH), 6.91 (d, 1H, J = 9.1 Hz, H-1'), 6.44 (s, 2H, NH<sub>2</sub>-exchangeable), 5.51 (t, 1H, J = 9.0 Hz, H-2'), 5.41 (t, 1H, J = 9.0 Hz, H-3'), 5.03 (t, 1H, J = 9.0 Hz, H-4'), 4.22–3.95 (m, 5H, H-5', H-6', 4-CH<sub>3</sub>OC<sub>6</sub>H<sub>4</sub>CH<sub>2</sub>), 3.81 (s, 3H, 4-CH<sub>3</sub>OC<sub>6</sub>H<sub>4</sub>CH<sub>2</sub>), 2.09– 1.97 (4s, 12H, CH<sub>3</sub>CO). Anal. Calcd. for C<sub>26</sub>H<sub>30</sub>N<sub>6</sub>O<sub>11</sub>S: C, 49.21; H, 4.76; N, 13.24. Found: C, 49.30; H, 4.72; N,13.20.

8-Amino-6-(4-chlorobenzyl)-2-(2,3,4,6-tetra-*O*-acetyl-β-D-glucopyranosyl)-2,8dihydro-3-thioxo-1,2,4-triazolo[4,3-*b*][1,2,4]-triazin-7(3*H*)-one (7d). Using the general procedure, 5d gave 7d (75%); mp 179°C; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) 7.39, 7.19 (2d, 2H each, J = 7.3 Hz, ArH), 6.95 (d, 1H, J = 9.0 Hz, H-1'), 6.38 (s, 2H, NH<sub>2</sub>-exchangeable), 5.55 (t, 1H, J = 9.4 Hz, H-2'), 5.39 (t, 1H, J = 9.4 Hz, H-3'), 5.01 (t, 1H, J = 9.4 Hz, H-4'),

#### 1,2,4-Triazolo[4,3-b][1,2,4]Triazines

4.21–3.96 (m, 5H, H-5', H-6', 4-ClC<sub>6</sub>H<sub>4</sub>CH<sub>2</sub>), 2.09–1.98 (4s, 12H, CH<sub>3</sub>CO). Anal. Calcd. for C<sub>25</sub>H<sub>27</sub>ClN<sub>6</sub>O<sub>10</sub>S: C, 46.99; H, 4.26; N,13.15. Found: C, 47.05; H, 4.34; N, 12.99.

8-Arylideneamino-6-benzyl(or substituted benzyl)-2-(2,3,4,6-tetra-*O*-acetyl-β-D-glucopyranosyl)-2,8-dihydro-3-thioxo-1,2,4-triazolo[4,3-*b*][1,2,4]-triazin-7(3*H*)-ones (8a–z). General procedure: A mixture of each of compounds 7a–d (0.5 g) and the appropriate aldehyde (0.5 mL or 0.5 g) was heated at  $150-160^{\circ}$ C in an oil bath for 15 minutes. The reaction mixture was removed from the oil bath, then ethanol (1 mL) was added and the reaction mixture was heated at reflux temperature for 1 hour. After cooling, the formed precipitate was collected by filtration, dried at room temperature and recrystallised from ethanol to give pale colorless crystals of compounds 8a–z.

**8-Benzylideneamino-6-benzyl-2-(2,3,4,6-tetra-***O***-acetyl-β-D-glucopyranosyl)-2,8-dihydro-3-thioxo-1,2,4-triazolo[4,3-***b***][<b>1,2,4**]**-triazin-7(3***H***)-one (8a).** Using the general procedure, **7a** gave **8a** (74%); mp 216°C; IR (KBr) 1743 (C = O acetate), 1674 (C = O amide) cm<sup>-1</sup>; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 8.33 (s, 1H, N = CH), 7.91–7.15 (m, 10H, ArH), 7.18 (d, 1H, J = 9.2 Hz, H-1'), 5.49 (t, 1H, J = 9.3 Hz, H-2'), 5.42 (t, 1H, J = 9.3 Hz, H-3'), 5.02 (t, 1H, J = 9.3 Hz, H-4'), 4.23–4.07 (m, 5H, H-5', H-6', C<sub>6</sub>H<sub>5</sub>CH<sub>2</sub>), 2.09–1.92 (4s, 12H, CH<sub>3</sub>CO). Anal. Calcd. for C<sub>32</sub>H<sub>32</sub>N<sub>6</sub>O<sub>10</sub>S: C, 55.49; H, 4.66; N, 12.13; Found: C, 55.53; H, 4.79; N, 12.17.

8-(4-Methylbenzylideneamino)-6-benzyl-2-(2,3,4,6-tetra-*O*-acetyl-β-D-glucopyranosyl)-2,8-dihydro-3-thioxo-1,2,4-triazolo[4,3-*b*][1,2,4]-triazin-7(3*H*)-one (8b). Using the general procedure,7a gave 8b (75%); mp 214°C; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 8.33 (s, 1H, N = CH), 7.88–7.13 (m, 9H, ArH), 7.15 (d, 1H, J = 9.1Hz, H-1′), 5.51 (t, 1H, J = 9.5 Hz, H-2′), 5.39 (t, 1H, J = 9.5 Hz, H-3′), 5.01 (t, 1H, J = 9.5 Hz, H-4′), 4.26–4.06 (m, 5H, H-5′, H-6′, C<sub>6</sub>H<sub>5</sub>CH<sub>2</sub>), 2.44 (s, 3H, 4-CH<sub>3</sub>C<sub>6</sub>H<sub>4</sub>CH = N), 2.07–1.93 (4s, 12H, CH<sub>3</sub>CO). Anal. Calcd. for C<sub>33</sub>H<sub>34</sub>N<sub>6</sub>O<sub>10</sub>S C, 56.08; H, 4.85; N, 11.89. Found: C, 55.98; H, 4.84; N, 11.97.

8-(2-Methoxyenzylideneamino)-6-benzyl-2-(2,3,4,6-tetra-*O*-acetyl-β-D-glucopyranosyl)-2,8-dihydro-3-thioxo-1,2,4-triazolo[4,3-*b*][1,2,4]-triazin-7(3*H*)-one (8c). Using the general procedure, **7a** gave 8c (76%) mp 210°C; IR (KBr) 1743 (C = O acetate), 1674 (C = O amide) cm<sup>-1</sup>; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 8.39 (s, 1H, N = CH), 7.92-7.13 (m, 9H, ArH), 7.18 (d, 1H, J = 9.4 Hz, H-1'), 5.49 (t, 1H, J = 9.0 Hz, H-2'), 5.42 (t, 1H, J = 9.0 Hz, H-3'), 5.02 (t, 1H, J = 9.0 Hz, H-4'), 4.21-3.99 (m, 5H, H-5', H-6', C<sub>6</sub>H<sub>5</sub>CH<sub>2</sub>), 3.83 (s, 3H, 2-CH<sub>3</sub>OC<sub>6</sub>H<sub>4</sub>CH = N), 2.09-1.92 (4s, 12H, CH<sub>3</sub>CO). Anal. Calcd. for C<sub>33</sub>H<sub>34</sub>N<sub>6</sub>O<sub>11</sub>S: C, 54.84; H, 4.74; N, 11.63. Found: C, 54.78; H, 4.73; N, 11.79.

**8-(4-Methoxybenzylideneamino)-6-benzyl-2-(2,3,4,6-tetra-***O***-acetyl-β-D-glucopyranosyl)-2,8-dihydro-3-thioxo-1,2,4-triazolo[4,3-b][1,2,4]-triazin-7(3H)-one (8d).** Using the general procedure, **7a** gave **8d** (70%); mp 218°C; IR (KBr) 1743 (C = O acetate), 1674 (C = O amide) cm<sup>-1</sup>; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 8.29 (s, 1H, N = CH), 7.87– 7.12 (m, 9H, ArH), 7.14 (d, 1H, J = 9.2 Hz, H-1'), 5.48 (t, 1H, J = 9.5 Hz, H-2'), 5.41 (t, 1H, J = 9.5 Hz, H-3'), 5.01 (t, 1H, J = 9.5 Hz, H-4'), 4.23–4.07 (m, 5H, H-5', H-6', C<sub>6</sub>H<sub>5</sub>CH<sub>2</sub>), 3.79 (s, 3H, 4-CH<sub>3</sub>OC<sub>6</sub>H<sub>4</sub>CH = N), 2.07–1.93 (4s, 12H, CH<sub>3</sub>CO). Anal. Calcd. for C<sub>33</sub>H<sub>34</sub>N<sub>6</sub>O<sub>11</sub>S: C, 54.84; H, 4.74; N, 11.63. Found: C, 54.81; H, 4.65; N, 11.82. 8-(4-Chlorobenzylideneamino)-6-benzyl-2-(2,3,4,6-tetra-*O*-acetyl-β-D-glucopyranosyl)-2,8-dihydro-3-thioxo-1,2,4-triazolo[4,3-*b*][1,2,4]-triazin-7(3*H*)-one (8e). Using the general procedure, **7a** gave **8e** (79%); mp 222°C (decomp.); <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 8.34 (s, 1H, N = CH), 7.90–7.13 (m, 9H, ArH), 7.18 (d, 1H, J = 9.2 Hz, H-1'), 5.49 (t, 1H, J = 9.5 Hz, H-2'), 5.43 (t, 1H, J = 9.5 Hz, H-3'), 5.00 (t, 1H, J = 9.5 Hz, H-4'), 4.22-3.96 (m, 5H, H-5', H-6', C<sub>6</sub>H<sub>5</sub>CH<sub>2</sub>), 2.09–1.92 (4s, 12H, CH<sub>3</sub>CO). Anal. Calcd. for C<sub>32</sub>H<sub>31</sub>ClN<sub>6</sub>O<sub>10</sub>S: C, 52.86; H, 4.30; N, 11.56. Found: C, 52.74; H, 4.26; N, 11.38.

8-(3,4-Methylenedioxybenzylideneamino)-6-benzyl-2-(2,3,4,6-tetra-*O*-acetyl-β- **D**-glucopyranosyl)-2,8-dihydro-3-thioxo-1,2,4-triazolo[4,3-*b*][1,2,4]-triazin-7(3*H*)one (8f). Using the general procedure, 7a gave 8f (73%); mp 204°C; IR (KBr) 1751 (C = O acetate), 1674 (C = O amide) cm<sup>-1</sup>; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 8.41 (s, 1H, N = CH), 7.88-7.12 (m, 8H, ArH), 7.21 (d, 1H, J = 9.4 Hz, H-1'), 5.85 (s, 2H, 3,4-  $CH_2O_2C_6H_3CH = N$ ), 5.48 (t, 1H, J = 9.2 Hz, H-2'), 5.41 (t, 1H, J = 9.2 Hz, H-3'), 5.01 (t, 1H, J = 9.2 Hz, H-4'), 4.23-4.07 (m, 5H, H-5', H-6', C<sub>6</sub>H<sub>5</sub>CH<sub>2</sub>), 2.08-1.92 (4s, 12H, CH<sub>3</sub>CO). Anal. Calcd. for C<sub>33</sub>H<sub>32</sub>N<sub>6</sub>O<sub>12</sub>S: C, 53.80; H, 4.38; N, 11.41. Found: C, 53.76; H, 4.33; N, 11.45.

8-[4-(Dimethylamino)benzylideneamino]-6-benzyl-2-(2,3,4,6-tetra-*O*-acetyl-β-D-glucopyranosyl)-2,8-dihydro-3-thioxo-1,2,4-triazolo[4,3-*b*][1,2,4]-triazin-7(3*H*)-one (8g). Using the general procedure, 7a gave 8g (80%); mp 206°C; IR (KBr) 1743 (C = O acetate), 1674 (C = O amide) cm<sup>-1</sup>; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 8.36 (s, 1H, N = CH), 7.91-7.07 (m, 9H, ArH), 7.17 (d, 1H, J = 9.4 Hz, H-1'), 5.49 (t, 1H, J = 9.5 Hz, H-2'), 5.42 (t, 1H, J = 9.5 Hz, H-3'), 5.03 (t, 1H, J = 9.5 Hz, H-4'), 4.22-4.01 (m, 5H, H-5', H-6', C<sub>6</sub>H<sub>5</sub>CH<sub>2</sub>), 3.03 (s, 6H, 4-(CH<sub>3</sub>)<sub>2</sub>NC<sub>6</sub>H<sub>4</sub>CH = N), 2.07-1.93 (4s, 12H, CH<sub>3</sub>CO). Anal. Calcd. for C<sub>34</sub>H<sub>37</sub>N<sub>7</sub>O<sub>10</sub>S: C, 55.50; H, 5.07; N, 13.33. Found: C, 55.47; H, 4.93; N, 13.44.

**8-(3,4-Dimethoxybenzylideneamino)-6-benzyl-2-(2,3,4,6-tetra-***O*-acetyl-β-D-glucopyranosyl)-2,8-dihydro-3-thioxo-1,2,4-triazolo[4,3-b][1,2,4]-triazin-7(3*H*)-one (**8h**). Using the general procedure, 7a gave 8h (76%); mp 210°C; IR (KBr) 1743 (C = O acetate), 1674 (C = O amide) cm<sup>-1</sup>; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 8.32 (s, 1H, N = CH), 7.90– 7.10 (m, 8H, ArH), 7.14 (d, 1H, J = 9.4 Hz, H-1'), 5.48 (t, 1H, J = 9.0 Hz, H-2'), 5.41 (t, 1H, J = 9.0 Hz, H-3'), 5.01 (t, 1H, J = 9.0 Hz, H-4'), 4.23–4.07 (m, 5H, H-5', H-6', C<sub>6</sub>H<sub>5</sub>CH<sub>2</sub>), 2.41, 2.33 [2s, 6H, 3,4-(CH<sub>3</sub>O)<sub>2</sub>C<sub>6</sub>H<sub>3</sub>CH = N], 2.09–1.92 (4s, 12H, CH<sub>3</sub>CO). Anal. Calcd. for C<sub>34</sub>H<sub>36</sub>N<sub>6</sub>O<sub>12</sub>S: C, 54.25; H, 4.82; N, 11.16;. Found: C, 54.30; H, 4.73; N, 11.26.

8-[(1-Naphthylmethylidene)amino]-6-benzyl-2-(2,3,4,6-tetra-*O*-acetyl-β-D-glucopyranosyl)-2,8-dihydro-3-thioxo-1,2,4-triazolo[4,3-*b*][1,2,4]-triazin-7(3*H*)-one (8i). Using the general procedure, **7a** gave **8i** (79%); mp 221°C; IR (KBr) 1743 (C = O acetate), 1674 (C = O amide) cm<sup>-1</sup>; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 8.43 (s, 1H, N = CH), 8.0–7.11 (m, 12H, ArH), 7.02 (d, 1H, J = 9.4 Hz, H-1'), 5.49 (t, 1H, J = 9.5 Hz, H-2'), 5.42 (t, 1H, J = 9.5 Hz, H-3'), 5.01 (t, 1H, J = 9.5 Hz, H-4'), 4.22–3.99 (m, 5H, H-5', H-6', C<sub>6</sub>H<sub>5</sub>CH<sub>2</sub>), 2.07–1.93 (4s, 12H, CH<sub>3</sub>CO). Anal. Calcd. for C<sub>36</sub>H<sub>34</sub>N<sub>6</sub>O<sub>10</sub>S: C, 58.21; H, 4.61; N, 11.31. Found: C, 58.26; H, 4.74; N, 11.21. 8-Benzylideneamino-6-(4-methylbenzyl)-2-(2,3,4,6-tetra-*O*-acetyl-β-D-glucopyranosyl)-2,8-dihydro-3-thioxo-1,2,4-triazolo[4,3-*b*][1,2,4]-triazin-7(3*H*)-one (8j). Using the general procedure, 7b gave 8j (82%); mp 198–200°C; IR (KBr) 1751 (C = O acetate), 1674 (C = O amide) cm<sup>-1</sup>; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 8.34 (s, 1H, N = CH), 7.92–7.07 (m, 9H, ArH), 7.06 (d, 1H, J = 9.0 Hz, H-1'), 5.49 (t, 1H, J = 9.2 Hz, H-2'), 5.41 (t, 1H, J = 9.2 Hz, H-3'), 5.01 (t, 1H, J = 9.2 Hz, H-4'), 4.23-3.97 (m, 5H, H-5', H-6', 4-CH<sub>3</sub>C<sub>6</sub>H<sub>4</sub>CH<sub>2</sub>), 2.25 (s, 3H, 4-CH<sub>3</sub>C<sub>6</sub>H<sub>4</sub>CH<sub>2</sub>), 2.08–1.92 (4s, 12H, CH<sub>3</sub>CO). Anal. Calcd. for C<sub>33</sub>H<sub>34</sub>N<sub>6</sub>O<sub>10</sub>S: C, 56.08; H, 4.85; N, 11.89. Found: C, 56.01; H, 4.94; N, 12.00.

8-(4-Methylbenzylideneamino)-6-(4-methylbenzyl)-2-(2,3,4,6-tetra-*O*-acetyl-β-D-glucopyranosyl)-2,8-dihydro-3-thioxo-1,2,4-triazolo[4,3-*b*][1,2,4]-triazin-7(3*H*)one (8k). Using the general procedure, 7b gave 8k (71%); mp 208–209°C; IR (KBr) 1751, 1720 (C = O acetate), 1674 (C = O amide) cm<sup>-1</sup>; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 8.34 (s, 1H, N = CH), 7.96–7.04 (m, 8H, ArH), 6.98 (d, 1H, J = 9.1 Hz, H-1'), 5.50 (t, 1H, J = 9.0 Hz, H-2'), 5.42 (t, 1H, J = 9.0 Hz, H-3'), 5.03 (t, 1H, J = 9.0 Hz, H-4'), 4.21– 3.96 (m, 5H, H-5', H-6', 4-CH<sub>3</sub>C<sub>6</sub>H<sub>4</sub>CH<sub>2</sub>), 2.30 (s, 3H, 4-CH<sub>3</sub>C<sub>6</sub>H<sub>4</sub>CH<sub>2</sub>), 2.44 (s, 3H, 4-CH<sub>3</sub>C<sub>6</sub>H<sub>4</sub>CH = N), 2.07–1.94 (4s, 12H, CH<sub>3</sub>CO). Anal. Calcd. for C<sub>34</sub>H<sub>36</sub>N<sub>6</sub>O<sub>10</sub>S: C, 56.66; H, 5.03; N, 11.66. Found: C, 56.79; H, 4.99; N, 11.48.

8-(2-Methoxybenzylideneamino)-6-(4-methylbenzyl)-2-(2,3,4,6-tetra-*O*-acetyl-β-D-glucopyranosyl)-2,8-dihydro-3-thioxo-1,2,4-triazolo[4,3-*b*][1,2,4]-triazin-7(3*H*)one (8l). Using the general procedure, 7b gave 8l (74%); mp 201–202°C; IR (KBr) 1751, 1720 (C = O acetate), 1674 (C = O amide) cm<sup>-1</sup>; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 8.40 (s, 1H, N = CH), 7.98–7.03 (m, 8H, ArH), 7.01 (d, 1H, J = 9.0 Hz, H-1'), 5.49 (t, 1H, J = 9.2 Hz, H-2'), 5.42 (t, 1H, J = 9.2 Hz, H-3'), 5.01 (t, 1H, J = 9.2 Hz, H-4'), 4.22– 3.98 (m, 5H, H-5', H-6', 4-CH<sub>3</sub>C<sub>6</sub>H<sub>4</sub>CH<sub>2</sub>), 3.79 (s, 3H, 2-CH<sub>3</sub>OC<sub>6</sub>H<sub>4</sub>CH = N), 2.26 (s, 3H, 4-CH<sub>3</sub>C<sub>6</sub>H<sub>4</sub>CH<sub>2</sub>), 2.09–1.92 (4s, 12H, CH<sub>3</sub>CO). Anal. Calcd. for C<sub>34</sub>H<sub>36</sub>N<sub>6</sub>O<sub>11</sub>S: C, 55.43; H, 4.93; N, 11.41. Found: C, 55.49; H, 4.96; N, 11.42.

8-(4-Methoxybenzylideneamino)-6-(4-methylbenzyl)-2-(2,3,4,6-tetra-*O*-acetyl-β-D-glucopyranosyl)-2,8-dihydro-3-thioxo-1,2,4-triazolo[4,3-*b*][1,2,4]-triazin-7(3*H*)one (8m). Using the general procedure, 7b gave 8m (72%); mp 198°C; IR (KBr) 1751 (C = O acetate), 1674 (C = O amide) cm<sup>-1</sup>; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 8.38 (s, 1H, N = CH), 7.96-7.04 (m, 8H, ArH), 6.98 (d, 1H, J = 9.0 Hz, H-1'), 5.47 (t, 1H, J = 9.2Hz, H-2'), 5.40 (t, 1H, J = 9.0 Hz, H-3'), 5.01 (t, 1H, J = 9.0 Hz, H-4'), 4.22-3.98 (m, 5H, H-5', H-6', 4-CH<sub>3</sub>C<sub>6</sub>H<sub>4</sub>CH<sub>2</sub>), 3.84 (s, 3H, 4-CH<sub>3</sub>OC<sub>6</sub>H<sub>4</sub>CH = N), 2.25 (s, 3H, 4-CH<sub>3</sub>C<sub>6</sub>H<sub>4</sub> CH<sub>2</sub>), 2.07-1.94 (4s, 12H, CH<sub>3</sub>CO). Anal. Calcd. for C<sub>34</sub>H<sub>36</sub>N<sub>6</sub>O<sub>11</sub>S: C, 55.43; H, 4.93; N, 11.41. Found: C, 55.39; H, 4.87; N, 11.32.

8-(4-Chlorobenzylideneamino)-6-(4-methylbenzyl)-2-(2,3,4,6-tetra-*O*-acetyl-β-D-glucopyranosyl)-2,8-dihydro-3-thioxo-1,2,4-triazolo[4,3-*b*][1,2,4]-triazin-7(3*H*)-one (8n). Using the general procedure, 7b gave 8n (76%); mp 210°C; IR (KBr) 1751, 1720 (C = O acetate), 1674 (C = O amide) cm<sup>-1</sup>; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 8.32 (s, 1H, N = CH), 7.99–7.01 (m, 8H, ArH), 6.95 (d, 1H, J = 9.0 Hz, H-1'), 5.49 (t, 1H, J = 9.2 Hz, H-2'), 5.42 (t, 1H, J = 9.2 Hz, H-3'), 5.01 (t, 1H, J = 9.2 Hz, H-4'), 4.22-3.98 (m, 5H, H-5', H-6', 4-CH<sub>3</sub>C<sub>6</sub>H<sub>4</sub>CH<sub>2</sub>), 2.27 (s, 3H, 4-CH<sub>3</sub>C<sub>6</sub>H<sub>4</sub>CH<sub>2</sub>), 2.08–1.93 (4s, 12H,

CH<sub>3</sub>CO). Anal. Calcd. for C<sub>33</sub>H<sub>33</sub>ClN<sub>6</sub>O<sub>10</sub>S: C, 53.48; H, 4.49; N, 11.34. Found: C, 53.50; H, 4.62; N, 11.36.

8-(3,4-Methylenedioxybenzylideneamino)-6-(4-methylbenzyl)-2-(2,3,4,6-tetra-*O*-acetyl-β-D-glucopyranosyl)-2,8-dihydro-3-thioxo-1,2,4-triazolo[4,3-*b*][1,2,4]-triazin-7(3*H*)-one (80). Using the general procedure, 7b gave 8o (73%); mp 190°C; IR (KBr) 1743 (C = O acetate), 1674 (C = O amide) cm<sup>-1</sup>; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 8.36 (s, 1H, N = CH), 7.94–7.05 (m, 7H, ArH), 7.00 (d, 1H, J = 9.0 Hz, H-1'), 5.87 (s, 2H, 3,4-CH<sub>2</sub>O<sub>2</sub>C<sub>6</sub>H<sub>3</sub>CH = N), 5.49 (t, 1H, J = 9.0 Hz, H-2'), 5.41 (t, 1H, J = 9.0 Hz, H-3'), 5.01 (t, 1H, J = 9.0 Hz, H-4'), 4.24–3.96 (m, 5H, H-5', H-6', 4-CH<sub>3</sub>C<sub>6</sub>H<sub>4</sub>CH<sub>2</sub>), 2.25 (s, 3H, 4-CH<sub>3</sub>C<sub>6</sub>H<sub>4</sub>CH<sub>2</sub>), 2.07–1.94 (4s, 12H, CH<sub>3</sub>CO). Anal. Calcd. for C<sub>34</sub>H<sub>34</sub>N<sub>6</sub>O<sub>12</sub>S: C, 54.40; H, 4.56; N, 11.19. Found: C, 54.42; H, 4.67; N, 11.24.

8-(3,4-Dimethoxybenzylideneamino)-6-(4-methylbenzyl)-2-(2,3,4,6-tetra-*O*-acetyl-β-D-glucopyranosyl)-2,8-dihydro-3-thioxo-1,2,4-triazolo[4,3-*b*][1,2,4]-triazin-7(3*H*)-one (8p). Using the general procedure, 7b gave 8p (70%); mp 198°C; IR (KBr) 1720 (C = O acetate), 1674 (C = O amide) cm<sup>-1</sup>; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 8.31 (s, 1H, N = CH), 7.91-7.02 (m, 7H, ArH),7.04 (d, 1H, J = 9.1 Hz, H-1'), 5.52 (t, 1H, J = 9.0 Hz, H-2'), 5.42 (t, 1H, J = 9.0 Hz, H-3'), 5.01 (t, 1H, J = 9.0 Hz, H-4'), 4.23-3.97 (m, 5H, H-5', H-6', 4-CH<sub>3</sub>C<sub>6</sub>H<sub>4</sub>CH<sub>2</sub>), 2.44, 2.32 [2s, 6H, 3,4-(CH<sub>3</sub>O)<sub>2</sub>C<sub>6</sub>H<sub>3</sub>CH = N], 2.28 (s, 3H, 4-CH<sub>3</sub>C<sub>6</sub>H<sub>4</sub>CH<sub>2</sub>), 2.07-1.92 (4s, 12H, CH<sub>3</sub>CO). Anal. Calcd. for C<sub>35</sub>H<sub>38</sub>N<sub>6</sub>O<sub>12</sub>S: C, 54.82; H, 5.00; N, 10.96. Found: C, 54.96; H, 4.97; N, 11.03.

8-[(1-Naphthylmethylidene)amino]-6-(4-methylbenzyl)-2-(2,3,4,6-tetra-*O*-acetyl-β-D-glucopyranosyl)-2,8-dihydro-3-thioxo-1,2,4-triazolo[4,3-*b*][1,2,4]-triazin-7(*3H*)-one (8q). Using the general procedure, 7b gave 8q (74%); mp 190°C; IR (KBr) 1751 (C = O acetate), 1674 (C = O amide) cm<sup>-1</sup>; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 8.42 (s, 1H, N = CH), 7.99-7.00 (m, 11H, ArH),7.01 (d, 1H, J = 9.0 Hz, H-1'), 5.49 (t, 1H, J = 9.2 Hz, H-2'), 5.41 (t, 1H, J = 9.2 Hz, H-3'), 5.01 (t, 1H, J = 9.2 Hz, H-4'), 4.21-3.98 (m, 5H, H-5', H-6', 4-CH<sub>3</sub>C<sub>6</sub>H<sub>4</sub>CH<sub>2</sub>), 2.24 (s, 3H, 4-CH<sub>3</sub>C<sub>6</sub>H<sub>4</sub>CH<sub>2</sub>), 2.07-1.94 (4s, 12H, CH<sub>3</sub>CO). Anal. Calcd. for C<sub>37</sub>H<sub>36</sub>N<sub>6</sub>O<sub>10</sub>S: C, 58.72; H, 4.79; N, 11.10. Found: C, 58.96; H, 4.94; N, 11.14.

8-Benzylideneamino-6-(4-methoxybenzyl)-2-(2,3,4,6-tetra-*O*-acetyl-β-D-glucopyranosyl)-2,8-dihydro-3-thioxo-1,2,4-triazolo[4,3-*b*][1,2,4]-triazin-7(3*H*)-one (8r). Using the general procedure, 7c gave 8r (77%); mp 217°C; IR (KBr) 1743 (C = O acetate), 1674 (C = O amide) cm<sup>-1</sup>; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 8.34 (s, 1H, N = CH), 7.89–7.00 (m, 9H, ArH), 6.84 (d, 1H, J = 9.0 Hz, H-1'), 5.51 (t, 1H, J = 9.3 Hz, H-2'), 5.41 (t, 1H, J = 9.3 Hz, H-3'), 5.01 (t, 1H, J = 9.3 Hz, H-4'), 4.21–3.96 (m, 5H, H-5', H-6', 4-CH<sub>3</sub>OC<sub>6</sub>H<sub>4</sub>CH<sub>2</sub>), 3.77 (s, 3H, 4-CH<sub>3</sub>OC<sub>6</sub>H<sub>4</sub>CH<sub>2</sub>), 2.08–1.99 (4s, 12H, CH<sub>3</sub>CO). Anal. Calcd. for C<sub>33</sub>H<sub>34</sub>N<sub>6</sub>O<sub>11</sub>S: C, 54.84; H, 4.74; N, 11.63. Found: C, 54.89; H, 4.70; N, 11.46.

8-(4-Methylbenzylideneamino)-6-(4-methoxybenzyl)-2-(2,3,4,6-tetra-*O*-acetyl-β-D-glucopyranosyl)-2,8-dihydro-3-thioxo-1,2,4-triazolo[4,3-*b*][1,2,4]-triazin-7(3*H*)one (8s). Using the general procedure, 7c gave 8s (69%); mp 212°C; IR (KBr) 1743 (C = O acetate), 1666 (C = O amide) cm<sup>-1</sup>; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>)  $\delta$  8.32 (s, 1H, N = CH), 7.84–6.99 (m, 8H, ArH), 6.91 (d, 1H, *J* = 9.2 Hz, H-1'), 5.53 (t, 1H, *J* = 9.0 Hz, H-2'), 5.41 (t, 1H, *J* = 9.0 Hz, H-3'), 5.04 (t, 1H, *J* = 9.0 Hz, H-4'), 4.24–3.95 (m, 5H, H-5', H-6', 4-CH<sub>3</sub>OC<sub>6</sub>H<sub>4</sub>CH<sub>2</sub>), 3.79 (s, 3H, 4-CH<sub>3</sub>OC<sub>6</sub>H<sub>4</sub>CH<sub>2</sub>), 2.45 (s, 3H, 4-CH<sub>3</sub>C<sub>6</sub>H<sub>4</sub>CH = N), 2.08–1.99 (4s, 12H, CH<sub>3</sub>CO). Anal. Calcd. for C<sub>34</sub>H<sub>36</sub>N<sub>6</sub>O<sub>11</sub>S: C, 55.43; H, 4.93; N, 11.41. Found: C, 55.23; H, 4.82; N, 11.49.

8-(2-Methoxybenzylideneamino)-6-(4-methoxybenzyl)-2-(2,3,4,6-tetra-*O*-acetylβ-D-glucopyranosyl)-2,8-dihydro-3-thioxo-1,2,4-triazolo[4,3-*b*][1,2,4]-triazin-7(3*H*)one (8t). Using the general procedure,7c gave 8t (80%); mp 216°C; IR (KBr) 1743 (C = O acetate), 1666 (C = O amide) cm<sup>-1</sup>; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 8.32 (s, 1H, N = CH), 7.88–7.01 (m, 8H, ArH), 6.89 (d, 1H, J = 9.1 Hz, H-1'), 5.51 (t, 1H, J = 9.2Hz, H-2'), 5.42 (t, 1H, J = 9.2 Hz, H-3'), 5.04 (t, 1H, J = 9.2 Hz, H-4'), 4.22–3.93 (m, 5H, H-5', H-6', 4-CH<sub>3</sub>OC<sub>6</sub>H<sub>4</sub>CH<sub>2</sub>), 3.77, 3.80 (2s, 6H, 4-CH<sub>3</sub>OC<sub>6</sub>H<sub>4</sub>CH<sub>2</sub>, 2-CH<sub>3</sub>OC<sub>6</sub>H<sub>4</sub>CH = N), 2.08–1.95 (4s, 12H, CH<sub>3</sub>CO). Anal. Calcd. for C<sub>34</sub>H<sub>36</sub>N<sub>6</sub>O<sub>12</sub>S: C, 54.25; H, 4.82; N, 11.16. Found: C, 54.19; H, 4.80; N, 11.39.

**8-(4-Methoxybenzylideneamino)-6-(4-methoxybenzyl)-2-(2,3,4,6-tetra-***O*-acetylβ-D-glucopyranosyl)-2,8-dihydro-3-thioxo-1,2,4-triazolo[4,3-b][1,2,4]-triazin-7(3H)one (8u). Using the general procedure, 7c gave 8u (76%); mp 219–220°C; IR (KBr) 1743 (C = O acetate), 1674 (C = O amide) cm<sup>-1</sup>; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 8.32 (s, 1H, N = CH), 7.84–7.03 (m, 8H, ArH), 6.84 (d, 1H, J = 9.0 Hz, H-1'), 5.50 (t, 1H, J = 9.1 Hz, H-2'), 5.40 (t, 1H, J = 9.1 Hz, H-3'), 5.02 (t, 1H, J = 9.1 Hz, H-4'), 4.21–3.92 (m, 5H, H-5', H-6', 4-CH<sub>3</sub>OC<sub>6</sub>H<sub>4</sub>CH<sub>2</sub>), 3.79, 3.84 (2s, 6H, 4-CH<sub>3</sub>OC<sub>6</sub>H<sub>4</sub>CH<sub>2</sub>, 4-CH<sub>3</sub>OC<sub>6</sub>H<sub>4</sub>CH = N), 2.07–1.95 (4s, 12H, CH<sub>3</sub>CO). Anal. Calcd. for C<sub>34</sub>H<sub>36</sub>N<sub>6</sub>O<sub>12</sub>S: C, 54.25; H, 4.82; N, 11.16. Found: C, 54.13; H, 4.98; N, 11.27.

8-(4-Chlorobenzylideneamino)-6-(4-methoxybenzyl)-2-(2,3,4,6-tetra-*O*-acetyl-β- **D**-glucopyranosyl)-2,8-dihydro-3-thioxo-1,2,4-triazolo[4,3-*b*][1,2,4]-triazin-7(3*H*)one (8v). Using the general procedure, 7c gave 8v (81%); mp 213°C; IR (KBr) 1743 (C = O acetate), 1666 (C = O amide) cm<sup>-1</sup>; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 8.39 (s, 1H, N = CH), 7.82–7.00 (m, 8H, ArH), 6.85 (d, 1H, J = 9.1 Hz, H-1'), 5.50 (t, 1H, J = 9.2 Hz, H-2'), 5.40 (t, 1H, J = 9.2 Hz, H-3'), 5.02 (t, 1H, J = 9.2 Hz, H-4'), 4.22–3.91 (m, 5H, H-5', H-6', 4-CH<sub>3</sub>OC<sub>6</sub>H<sub>4</sub>CH<sub>2</sub>), 3.79 (s, 3H, 4-CH<sub>3</sub>OC<sub>6</sub>H<sub>4</sub>CH<sub>2</sub>), 2.09–1.94 (4s, 12H, CH<sub>3</sub>CO). Anal. Calcd. for C<sub>33</sub>H<sub>33</sub>ClN<sub>6</sub>O<sub>11</sub>S: C, 52.35; H, 4.39; N, 11.10. Found: C, 52.22; H, 4.54; N, 11.16.

8-(3,4-Methylenedioxybenzylideneamino)-6-(4-methoxybenzyl)-2-(2,3,4,6-tetra-*O*-acetyl-β-D-glucopyranosyl)-2,8-dihydro-3-thioxo-1,2,4-triazolo[4,3-*b*][1,2,4]triazin-7(3*H*)-one (8*w*). Using the general procedure, 7c gave 8*w* (70%); mp 221– 222°C; IR (KBr) 1743 (C = O acetate), 1666 (C = O amide) cm<sup>-1</sup>; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 8.34 (s, 1H, N = CH), 7.81–7.03 (m, 7H, ArH), 6.87 (d, 1H, J = 9.2Hz, H-1'), 5.90 (s, 2H, 3,4-CH<sub>2</sub>O<sub>2</sub>C<sub>6</sub>H<sub>3</sub>CH = N), 5.52 (t, 1H, J = 9.3 Hz, H-2'), 5.41 (t, 1H, J = 9.3 Hz, H-3'), 5.03 (t, 1H, J = 9.3 Hz, H-4'), 4.24–3.92 (m, 5H, H-5', H-6', 4-CH<sub>3</sub>OC<sub>6</sub>H<sub>4</sub>CH<sub>2</sub>), 3.79 (s, 3H, 4-CH<sub>3</sub>OC<sub>6</sub>H<sub>4</sub>CH<sub>2</sub>), 2.08–1.95 (4s, 12H, CH<sub>3</sub>CO). Anal. Calcd. for C<sub>34</sub>H<sub>34</sub>N<sub>6</sub>O<sub>13</sub>S: C, 53.26; H, 4.47; N, 10.96. Found: C, 53.30; H, 4.52; N, 11.18. 8-[4-(Dimethylamino)benzylideneamino]-6-(4-methoxybenzyl)-2-(2,3,4,6-tetra-*O*-acetyl-β-D-glucopyranosyl)-2,8-dihydro-3-thioxo-1,2,4-triazolo[4,3-*b*][1,2,4]triazin-7(3*H*)-one (8x). Using the general procedure, 7a gave 8x (80%); mp 211°C; IR (KBr) 1751 (C = O acetate), 1666 (C = O amide) cm<sup>-1</sup>; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 8.36 (s, 1H, N = CH), 7.90–7.07 (m, 8H, ArH), 6.79 (d, 1H, J = 9.3Hz, H-1'), 5.49 (t, 1H, J = 9.5 Hz, H-2'), 5.44 (t, 1H, J = 9.5 Hz, H-3'), 5.01 (t, 1H, J = 9.5 Hz, H-4'), 4.24–3.99 (m, 5H, H-5', H-6', 4-CH<sub>3</sub>OC<sub>6</sub>H<sub>4</sub>CH<sub>2</sub>), 3.77 (s, 3H, 4-CH<sub>3</sub>OC<sub>6</sub>H<sub>4</sub>CH<sub>2</sub>), 3.05 (s, 6H, 4-(CH<sub>3</sub>)<sub>2</sub>NC<sub>6</sub>H<sub>4</sub>CH = N), 2.07–1.93 (4s, 12H, CH<sub>3</sub>CO). Anal. Calcd. for C<sub>35</sub>H<sub>39</sub>N<sub>7</sub>O<sub>11</sub>S: C, 54.90; H, 5.13; N, 12.80. Found: C, 55.02; H, 4.94; N, 13.00.

8-(3,4-Dimethoxybenzylideneamino)-6-(4-methoxybenzyl)-2-(2,3,4,6-tetra-*O*-acetyl-β-D-glucopyranosyl)-2,8-dihydro-3-thioxo-1,2,4-triazolo[4,3-*b*][1,2,4]-triazin-7(*3H*)-one (8y). Using the general procedure, 7c gave 8y (75%); mp 218°C; IR (KBr) 1743 (C = O acetate), 1666 (C = O amide) cm<sup>-1</sup>; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 8.33 (s, 1H, N = CH), 7.84–7.00 (m, 7H, ArH), 6.82 (d, 1H, J = 9.0 Hz, H-1'), 5.49 (t, 1H, J = 9.3 Hz, H-2'), 5.43 (t, 1H, J = 9.3 Hz, H-3'), 5.02 (t, 1H, J = 9.3 Hz, H-4'), 4.24–3.92 (m, 5H, H-5', H-6', 4-CH<sub>3</sub>OC<sub>6</sub>H<sub>4</sub>CH<sub>2</sub>), 3.79, 3.82 (2s, 9H, 4-CH<sub>3</sub>OC<sub>6</sub>H<sub>4</sub>CH<sub>2</sub>, 3,4-(CH<sub>3</sub>O)<sub>2</sub>C<sub>6</sub>H<sub>3</sub>CH = N), 2.07–1.94 (4s, 12H, CH<sub>3</sub>CO). Anal. Calcd. for C<sub>35</sub>H<sub>38</sub>N<sub>6</sub>O<sub>13</sub>S: C, 53.70; H, 4.89; N, 10.74. Found: C, 53.50; H, 4.72; N, 10.85.

8-[(1-Naphthylmethylidene)amino]-6-(4-methoxybenzyl)-2-(2,3,4,6-tetra-*O*-acetyl-β-D-glucopyranosyl)-2,8-dihydro-3-thioxo-1,2,4-triazolo[4,3-*b*][1,2,4]-triazin-7(*3H*)-one (8z). Using the general procedure, 7c gave 8z (71%); mp 221–222°C; IR (KBr) 1751 (C = O acetate), 1674 (C = O amide) cm<sup>-1</sup>; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 8.33 (s, 1H, N = CH), 7.95–7.00 (m, 11H, ArH), 6.93 (d, 1H, J = 9.2 Hz, H-1'), 5.51 (t, 1H, J = 9.1 Hz, H-2'), 5.41 (t, 1H, J = 9.1 Hz, H-3'), 5.01 (t, 1H, J = 9.1 Hz, H-4'), 4.22–3.95 (m, 5H, H-5', H-6', 4-CH<sub>3</sub>OC<sub>6</sub>H<sub>4</sub>CH<sub>2</sub>), 3.77 (s, 3H, 4-CH<sub>3</sub>OC<sub>6</sub>H<sub>4</sub>CH<sub>2</sub>), 2.08–1.99 (4s, 12H, CH<sub>3</sub>CO). Anal. Calcd. for C<sub>37</sub>H<sub>36</sub>N<sub>6</sub>O<sub>11</sub>S: C, 57.51; H, 4.70; N, 10.87. Found: C, 57.64; H, 4.75; N, 10.94.

6-Benzyl(or substituted benzyl)-2-(2,3,4,6-tetra-*O*-acetyl-β-D-glucopyranosyl)-2,8-dihydro-3-thioxo-1,2,4-triazolo[4,3-b][1,2,4]-triazin-7(3*H*)-ones (9a–d). General procedure: To a cold stirred solution (at 0 C) of each of compounds 7a–d (10 mmol) in acetic acid (100 mL) was added dropwise a solution of sodium nitrite (6 g in 6 mL water) over a period of one hour and the reaction mixture was kept in the refrigerator overnight. The formed precipitate was collected by filtration, washed well with water, dried at room temperature and recrystallized from dichloromethane/petroleum ether (bp. 40–60°C) to give colorless crystals of compounds 9a–d.

**6-Benzyl-2-(2,3,4,6-tetra-***O***-acetyl-β-D-glucopyranosyl)-2,8-dihydro-3-thioxo-1,2,4-triazolo[4,3-b][1,2,4]-triazin-7(3***H***)-one (9a). Using the general procedure, <b>7a** gave **9a** (80%); mp 196°C; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 9.45 (s, 1H, NH-exchangeable), 7.38–7.21 (m, 5H, ArH), 6.62 (d, 1H, J = 9.4 Hz, H-1′), 5.64 (t, 1H, J = 9.5 Hz, H-2′), 5.42 (t, 1H, J = 9.5 Hz, H-3′), 5.02 (t, 1H, J = 9.5 Hz, H-4′), 4.21–3.99 (m, 5H, H-5′, H-6′, C<sub>6</sub>H<sub>5</sub>CH<sub>2</sub>), 2.12–1.92 (4s, 12H, CH<sub>3</sub>CO). Anal. Calcd. for C<sub>25</sub>H<sub>27</sub>N<sub>5</sub>O<sub>10</sub>S: C, 50.93; H, 4.62; N, 11.82. Found: C, 50.85; H, 4.71; N, 11.97.

**6-(4-Methylbenzyl)-2-(2,3,4,6-tetra**-*O*-acetyl-β-D-glucopyranosyl)-2,8-dihydro-3thioxo-1,2,4-triazolo[4,3-b][1,2,4]-triazin-7(3*H*)-one (9b). Using the general procedure, 7b gave 9b (74%); mp 212°C; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 9.57 (s, 1H, NHexchangeable), 7.30 (d, 2H, J = 7.7 Hz, ArH), 6.84 (d, 2H, J = 7.9 Hz, ArH), 6.64 (d, 1H, J = 9.4 Hz, H-1'), 5.63 (t, 1H, J = 9.5 Hz, H-2'), 5.42 (t, 1H, J = 9.5 Hz, H-3'), 5.04 (t, 1H, J = 9.5 Hz, H-4'), 4.21–3.99 (m, 5H, H-5', H-6', 4-CH<sub>3</sub>C<sub>6</sub>H<sub>4</sub>CH<sub>2</sub>), 2.26 (s, 3H, 4-CH<sub>3</sub>C<sub>6</sub>H<sub>4</sub>CH<sub>2</sub>), 2.11–1.92 (4s, 12H, CH<sub>3</sub>CO). Anal. Calcd. for C<sub>26</sub>H<sub>29</sub>N<sub>5</sub>O<sub>10</sub>S: C, 51.74; H, 4.84; N, 11.60. Found: C, 51.81; H, 4.80; N, 11.44.

**6-(4-Methoxybenzyl)-2-(2,3,4,6-tetra**-*O*-acetyl-β-D-glucopyranosyl)-2,8-dihydro-3-thioxo-1,2,4-triazolo[4,3-*b*][1,2,4]-triazin-7(*3H*)-one (9c). Using the general procedure, 7c gave 9c (70%); mp 228°C; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 9.41 (s, 1H, NHexchangeable), 7.26 (d, 2H, J = 7.9 Hz, ArH), 6.86 (d, 2H, J = 7.6 Hz, ArH), 6.60 (d, 1H, J = 9.4 Hz, H-1'), 5.61 (t, 1H, J = 9.5 Hz, H-2'), 5.41 (t, 1H, J = 9.5 Hz, H-3'), 5.01 (t, 1H, J = 9.5 Hz, H-4'), 4.23–4.07 (m, 5H, H-5', H-6', 4-CH<sub>3</sub>-OC<sub>6</sub>H<sub>4</sub>CH<sub>2</sub>), 3.72 (s, 3H, 4-CH<sub>3</sub>OC<sub>6</sub>H<sub>4</sub>CH<sub>2</sub>), 2.09–1.95 (4s, 12H, CH<sub>3</sub>CO). Anal. Calcd. for C<sub>26</sub>H<sub>29</sub>N<sub>5</sub>O<sub>11</sub>S: C, 50.40; H, 4.72; N, 11.30. Found: C, 50.41; H, 4.68; N, 11.24.

**6-(4-Chlorobenzyl)-2-(2,3,4,6-tetra**-*O*-acetyl-β-D-glucopyranosyl)-2,8-dihydro-**3-thioxo-1,2,4-triazolo[4,3-b][1,2,4]-triazin-7(3H)-one (9d).** Using the general procedure, **7d** gave **9d** (75%); mp 191°C; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 9.41 (s, 1H, NH-exchangeable), 7.24 (d, 2H, J = 7.8 Hz, ArH), 6.81 (d, 2H, J = 7.7 Hz, ArH), 6.63 (d, 1H, J = 9.4 Hz, H-1′), 5.64 (t, 1H, J = 9.5 Hz, H-2′), 5.40 (t, 1H, J = 9.5 Hz, H-3′), 5.04 (t, 1H, J = 9.5 Hz, H-4′), 4.22–4.03 (m, 5H, H-5′, H-6′, 4-ClC<sub>6</sub>H<sub>4</sub>CH<sub>2</sub>), 2.11–1.93 (4s, 12H, CH<sub>3</sub>CO). Anal. Calcd. for C<sub>25</sub>H<sub>26</sub>ClN<sub>5</sub>O<sub>10</sub>S: C, 48.12; H, 4.20; N, 11.22. Found: C, 48.00; H, 4.33; N, 11.26.

6-Benzyl(or substituted benzyl)-2-(2,3,4,6-tetra-*O*-acetyl-β-D-glucopyranosyl)-1,2,4-triazolo[4,3-*b*][1,2,4]-triazine-3,7(2*H*,8*H*)-dithiones (10a-d). General procedure: To a solution of each of compounds 9a-d (5 mmol) in dry pyridine (25 mL) was added phosphorus pentasulfide (10 mmol). The reaction mixture was then heated at reflux temperature for 6 hours. After cooling, the products were extracted from the oily materials with ethanol (50 mL). The supernatant solutions were decanted, acidified with acetic acid (5 mL), concentrated, and diluted with water. The precipitates were collected by filtration, dried at room temperature, dissolved in dichloromethane, treated with charcoal (2.5 g) and filtered. The filtrates were evaporated on a rotavap and the resulting solids were recrystallised from dichloromethane/petroleum ether (bp 40– 60°C) to give yellow crystals of compounds 10a-d.

**6-Benzyl-2-(2,3,4,6-tetra-***O***-acetyl-β-D-glucopyranosyl)-1,2,4-triazolo[4,3***b*][**1,2,4**]**-triazine-3,7(2***H***,8***H***)<b>-dithione (10a).** Using the general procedure, **9a** gave **10a** (50%); mp 184°C; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 9.36 (s, 1H, NH-exchangeable), 7.32–7.19 (m, 5H, ArH), 6.61 (d, 1H, J = 9.4 Hz, H-1′), 5.69 (t, 1H, J = 9.5 Hz, H-2′), 5.42 (t, 1H, J = 9.5 Hz, H-3′), 5.02 (t, 1H, J = 9.5 Hz, H-4′), 4.21–3.99 (m, 5H, H-5′, H-6′, C<sub>6</sub>H<sub>5</sub>CH<sub>2</sub>), 2.12–1.92 (4s, 12H, CH<sub>3</sub>CO). Anal. Calcd. for C<sub>25</sub>H<sub>27</sub>N<sub>5</sub>O<sub>9</sub>S<sub>2</sub>: C, 49.58; H, 4.49; N, 11.56. Found: C, 49.47; H, 4.31; N, 11.77. **6-(4-Methylbenzyl)-2-(2,3,4,6-tetra**-*O*-acetyl-β-D-glucopyranosyl)-1,2,4-triazolo[4,3-*b*][1,2,4]triazine-3,7(2*H*,8*H*)-dithione (10b). Using the general procedure, 9b gave 10b (51%); mp 204°C; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 9.37 (s, 1H, NH-exchangeable), 7.28(d, 2H, J = 7.5 Hz, ArH), 6.86 (d, 2H, J = 7.8 Hz, ArH), 6.59 (d, 1H, J = 9.4 Hz, H-1'), 5.65 (t, 1H, J = 9.5 Hz, H-2'), 5.42 (t, 1H, J = 9.5 Hz, H-3'), 5.04 (t, 1H, J = 9.5Hz, H-4'), 4.24–3.99 (m, 5H, H-5', H-6', 4-CH<sub>3</sub>C<sub>6</sub>H<sub>4</sub>CH<sub>2</sub>), 2.24 (4-CH<sub>3</sub>C<sub>6</sub>H<sub>4</sub>CH<sub>2</sub>), 2.13–1.92 (4s, 12H, CH<sub>3</sub>CO). Anal. Calcd. for C<sub>26</sub>H<sub>29</sub>N<sub>5</sub>O<sub>9</sub>S<sub>2</sub>: C, 50.40; H, 4.72; N, 11.30. Found: C, 50.42; H, 4.84; N, 11.24.

**6-(4-Methoxybenzyl)-2-(2,3,4,6-tetra-***O***-acetyl-β-D-glucopyranosyl)-1,2,4-tria**zolo[4,3-*b*][1,2,4]triazine-3,7(2*H*,8*H*)-dithione (10c). Using the general procedure, **9c** gave **10c** (48%); mp 198–200°C; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 9.75 (s, 1H, NHexchangeable), 7.32 (d, 2H, *J* = 7.6 Hz, ArH), 6.80 (d, 2H, *J* = 7.9 Hz, ArH), 6.62 (d, 1H, *J* = 9.4 Hz, H-1'), 5.62 (t, 1H, *J* = 9.5 Hz, H-2'), 5.41 (t, 1H, *J* = 9.5 Hz, H-3'), 5.04 (t, 1H, *J* = 9.5 Hz, H-4'), 4.24–4.02 (m, 5H, H-5', H-6', 4-CH<sub>3</sub>OC<sub>6</sub>H<sub>4</sub>CH<sub>2</sub>), 3.74 (s, 3H, 4-CH<sub>3</sub>OC<sub>6</sub>H<sub>4</sub>CH<sub>2</sub>), 2.07–1.95 (4s, 12H, CH<sub>3</sub>CO). Anal. Calcd. for C<sub>26</sub>H<sub>29</sub>N<sub>5</sub>O<sub>10</sub>S<sub>2</sub>: C, 49.13; H, 4.60; N, 11.02. Found: C, 49.22; H, 4.58; N, 11.12.

**6-(4-Chlorobenzyl)-2-(2,3,4,6-tetra**-*O*-acetyl-β-D-glucopyranosyl)-1,2,4-triazolo[4,3-*b*][1,2,4]triazine-3,7(2*H*,8*H*)-dithione (10d). Using the general procedure, 9d gave 10d (45%); mp 226°C; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 9.36 (s, 1H, NH-exchangeable), 7.23 (d, 2H, J = 7.6 Hz, ArH), 6.77 (d, 2H, J = 7.8 Hz, ArH), 6.60 (d, 1H, J = 9.4 Hz, H-1'), 5.66 (t, 1H, J = 9.5 Hz, H-2'), 5.42 (t, 1H, J = 9.5 Hz, H-3'), 5.01 (t, 1H, J = 9.5Hz, H-4'), 4.21–4.03 (m, 5H, H-5', H-6', 4-ClC<sub>6</sub>H<sub>4</sub>CH<sub>2</sub>), 2.07–1.93 (4s, 12H, CH<sub>3</sub>CO). Anal. Calcd. for C<sub>25</sub>H<sub>26</sub>ClN<sub>5</sub>O<sub>9</sub>S<sub>2</sub>: C, 46.91; H, 4.09; N, 10.94. Found: C, 47.09; H, 4.13; N, 11.12.

6-Benzyl(or substitutedbenzyl)-2-(2,3,4,6-tetra-*O*-acetyl-β-D-glucopyranosyl)-1,2,4-triazolo[4,3-*b*][1,2,4]-triazine-3,7(2*H*,8*H*)-diones (11a–d). General procedure: Each of compounds 9a–d (1 mmol) was dissolved in acetic acid (5 mL), then hydrogen peroxide (33%; 1 mL) was added. The reaction mixture was then boiled for 3 minutes, left to cool at room temperature, and diluted with water. The formed precipitate was collected by filtration, dried at room temperature, and recrystallised from ethanol to give colorless crystals of compounds 11a–d.

**6-Benzyl-2-(2,3,4,6-tetra-***O***-acetyl-β-D-glucopyranosyl)-1,2,4-triazolo[4,3***b*]**[1,2,4]-triazine-3,7(2H,8H)-dione (11a).** Using the general procedure, **9a** gave **11a** (60%); mp 178°C; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 9.32 (s, 1H, NH-exchangeable), 7.31–7.19 (m, 5H, ArH), 6.66 (d, 1H, J = 9.4 Hz, H-1'), 5.60 (t, 1H, J = 9.5 Hz, H-2'), 5.41 (t, 1H, J = 9.5 Hz, H-3'), 5.05 (t, 1H, J = 9.5 Hz, H-4'), 4.24–3.97 (m, 5H, H-5', H-6', C<sub>6</sub>H<sub>5</sub>CH<sub>2</sub>), 2.09–1.92 (4s, 12H, CH<sub>3</sub>CO). Anal. Calcd. for C<sub>25</sub>H<sub>27</sub>N<sub>5</sub>O<sub>11</sub>: C, 52.36; H, 4.75; N, 12.21. Found: C, 52.20; H, 4.79; N, 12.36.

**6-(4-Methylbenzyl)-2-(2,3,4,6-tetra**-*O*-acetyl-β-D-glucopyranosyl)-1,2,4-triazolo[4,3-*b*][1,2,4]triazine-3,7(2*H*,8*H*)-dione (11b). Using the general procedure, 9b gave 11b (65%); mp 185°C; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 9.47 (s, 1H, NH-exchangeable), 7.30 (d, 2H, J = 7.7 Hz, ArH), 6.84 (d, 2H, J = 7.9 Hz, ArH), 6.64 (d, 1H, J = 9.4 Hz, H-1'), 5.63 (t, 1H, J = 9.5 Hz, H-2'), 5.42 (t, 1H, J = 9.5 Hz, H-3'), 5.04 (t, 1H, J = 9.5 Hz, H-4'), 4.21–3.99 (m, 5H, H-5', H-6', 4-CH<sub>3</sub>C<sub>6</sub>H<sub>4</sub>CH<sub>2</sub>), 2.26 (s, 3H, 4-CH<sub>3</sub>C<sub>6</sub>H<sub>4</sub>CH<sub>2</sub>), 2.11–1.92 (4s, 12H, CH<sub>3</sub>CO). Anal. Calcd. for  $C_{26}H_{29}N_5O_{11}$ : C, 53.15; H, 4.98; N, 11.92. Found: C, 53.22; H, 4.88; N, 11.75.

**6-(4-Methoxybenzyl)-2-(2,3,4,6-tetra-***O***-acetyl-β-D-glucopyranosyl)-1,2,4-tria-zolo[4,3-***b***][<b>1,2,4**]**triazine-3,7(2***H***,8***H***)-<b>dione (11c).** Using the general procedure, **9c** gave **11c** (64%); mp 199°C; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 9.41 (s, 1H, NH-exchangeable), 7.26 (d, 2H, J = 7.9 Hz, ArH), 6.86 (d, 2H, J = 7.6 Hz, ArH), 6.60 (d, 1H, J = 9.4 Hz, H-1'), 5.61 (t, 1H, J = 9.5 Hz, H-2'), 5.41 (t, 1H, J = 9.5 Hz, H-3'), 5.01 (t, 1H, J = 9.5 Hz, H-4'), 4.23–4.07 (m, 5H, H-5', H-6', 4-CH<sub>3</sub>OC<sub>6</sub>H<sub>4</sub>CH<sub>2</sub>), 3.72 (s, 3H, 4-CH<sub>3</sub>OC<sub>6</sub>H<sub>4</sub>CH<sub>2</sub>), 2.09-1.95 (4s, 12H, CH<sub>3</sub>CO). Anal. Calcd. for C<sub>26</sub>H<sub>29</sub>N<sub>5</sub>O<sub>12</sub>: C, 51.74; H, 4.84; N, 11.60. Found: C, 51.89; H, 4.84; N, 11.67.

**6-(4-Chlorobenzyl)-2-(2,3,4,6-tetra-***O***-acetyl-β-D-glucopyranosyl)-1,2,4-triazolo** [**4,3-***b*][**1,2,4**]**triazine-3,7(2***H***,8***H***)-<b>dione** (**11d**). Using the general procedure, **9d** gave **11d** (71%); mp 179°C; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 9.41 (s, 1H, NH-exchangeable), 7.24 (d, 2H, *J* = 7.8 Hz, ArH), 6.81 (d, 2H, *J* = 7.7 Hz, ArH), 6.63 (d, 1H, *J* = 9.4 Hz, H-1'), 5.64 (t, 1H, *J* = 9.5 Hz, H-2'), 5.40 (t, 1H, *J* = 9.5 Hz, H-3'), 5.04 (t, 1H, *J* = 9.5 Hz, H-4'), 4.22–4.03 (m, 5H, H-5', H-6', 4-ClC<sub>6</sub>H<sub>4</sub>CH<sub>2</sub>), 2.11–1.93 (4s, 12H, CH<sub>3</sub>CO). Anal. Calcd. for C<sub>25</sub>H<sub>26</sub>ClN<sub>5</sub>O<sub>11</sub>: C, 49.39; H, 4.31; N, 11.52. Found: C, 49.29; H, 4.39; N, 11.57.

6-Benzyl(or substituted benzyl)-2-β-D-glucopyranosyl-2,8-dihydro-3-thioxo-1,2,4-triazolo[4,3-b][1,2,4]-triazin-7(3H)-ones (12a-d). General procedure: Dry gaseous ammonia was passed through a solution of each of compounds 9a-d (1 mmol) in dry methanol (10 mL) for about 1 hour while cooling and stirring. The reaction mixture was stirred at 0°C until complete as shown by TLC (9 to16 hours), using chloroform/ methanol (19 : 1) (R<sub>f</sub> = 0.70-0.75). The resulting mixture was then concentrated at reduced pressure to afford a solid residue which was boiled in chloroform (100 mL), collected by filtration, washed several times with boiled chloroform, dried at room temperature, and recrystallised from methanol/chloroform to give pale crystals of compounds 12a-d.

**6-Benzyl-2-β-D-glucopyranosyl-2,8-dihydro-3-thioxo-1,2,4-triazolo[4,3-b][1,2,4]-triazin-7(3H)-one (12a).** Using the general procedure, **9a** gave **12a** (60%); mp 262°C; IR (KBr) 3500–3200 (OH, NH), 1670 (C = O amide) cm<sup>-1</sup>; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 13.51 (s, 1H, NH-exchangeable), 7.62–7.12 (m, 5H, ArH), 6.37 (d, 1H, J = 9.0 Hz, H-1'), 5.22–4.63 (3d, 1t, 4H, 4OH-exchangeable), 4–3 (m, 8H, H-2', H-3', H-4', H-5', H-6', C<sub>6</sub>H<sub>5</sub>CH<sub>2</sub>). Anal. Calcd. for C<sub>17</sub>H<sub>19</sub>N<sub>5</sub>O<sub>6</sub>S: C, 48.45; H, 4.54; N, 16.62. Found: C, 48.33; H, 4.48; N, 16.49.

**6-(4-Methylbenzyl)-2-β-D-glucopyranosyl-2,8-dihydro-3-thioxo-1,2,4-tria-zolo[4,3-b][1,2,4]-triazin-7(3H)-one (12b).** Using the general procedure, **9b** gave **12b** (65%); mp 247°C; IR (KBr) 3500–3200 (OH, NH), 1669 (C = O amide) cm<sup>-1</sup>; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 13.47 (s, 1H, NH-exchangeable), 7.27 (d, 2H, J = 7.6 Hz, ArH), 6.85 (d, 2H, J = 7.9 Hz, ArH), 6.39 (d, 1H, J = 9.2 Hz, H-1'), 5.18–4.61 (3d, 1t, 4H, 4OH-exchangeable), 3.92–2.98 (m, 8H, H-2', H-3', H-4', H-5', H-6', 4-CH<sub>3</sub>C<sub>6</sub>H<sub>4</sub>CH<sub>2</sub>), 2.24 (s,3H, 4-CH<sub>3</sub>C<sub>6</sub>H<sub>4</sub>CH<sub>2</sub>). Anal. Calcd. for C<sub>18</sub>H<sub>21</sub>N<sub>5</sub>O<sub>6</sub>S: C, 49.65; H, 4.86; N, 16.08. Found: C, 49.58; H, 4.71; N, 16.25.

**6-(4-Methoxybenzyl)-2-β-D-glucopyranosyl-2,8-dihydro-3-thioxo-1,2,4-tria-zolo[4,3-***b***][<b>1,2,4]-triazin-7(3***H***)-one (12c).** Using the general procedure, **9c** gave **12c** (65%); mp 236°C; IR (KBr) 3500–3200 (OH, NH), 1671 (C = O amide) cm<sup>-1</sup>; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 13.54 (s, 1H, NH-exchangeable), 7.22 (d, 2H, J = 7.4 Hz, ArH), 6.81 (d, 2H, J = 7.5 Hz, ArH), 6.34 (d, 1H, J = 9.2 Hz, H-1'), 5.20–4.58 (3d, 1t, 4H, 4OH-exchangeable), 4.03–2.97 (m, 8H, H-2', H-3', H-4', H-5', H-6', 4-CH<sub>3</sub>OC<sub>6</sub>H<sub>4</sub>CH<sub>2</sub>), 3.81 (s,3H, 4-CH<sub>3</sub>OC<sub>6</sub>H<sub>4</sub>CH<sub>2</sub>). Anal. Calcd. for C<sub>18</sub>H<sub>21</sub>N<sub>5</sub>O<sub>7</sub>S: C, 47.89; H, 4.69; N, 15.51. Found: C, 47.94; H, 4.74; N, 15.33.

**6-(4-Chloroenzyl)-2-β-D-glucopyranosyl-2,8-dihydro-3-thioxo-1,2,4-triazolo [4,3-b][1,2,4]-triazin-7(3H)-one (12d).** Using the general procedure, **9d** gave **12d** (58%); mp 212°C; IR (KBr) 3500–3200 (OH, NH), 1670 (C = O amide) cm<sup>-1</sup>; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 13.52 (s, 1H, NH-exchangeable), 7.19(d, 2H, J = 7.3 Hz, ArH), 6.88 (d, 2H, J = 7.7 Hz, ArH), 6.37 (d, 1H, J = 9.0 Hz, H-1'), 5.22–4.63 (3d, 1t, 4H, 4OH-exchangeable), 4–3 (m, 8H, H-2', H-3', H-4', H-5', H-6', 4-ClC<sub>6</sub>H<sub>4</sub>CH<sub>2</sub>). Anal. Calcd. for C<sub>17</sub>H<sub>18</sub>ClN<sub>5</sub>O<sub>6</sub>S: C, 44.79; H, 3.98; N, 15.36. Found: C, 44.84; H, 4.14; N, 15.44.

#### REFERENCES

- Jones, D.H.; Slack, R.; Squires, S.; Wooldridge, K.R.H. Antiviral chemotherapy. I. The activity of pyridine and quinoline derivatives against neurovaccina in mice. J. Med. Chem. **1965**, *8* (5), 676–680.
- Goswami, B.N.; Kataky, J.C.S.; Baruah, J.N. Synthesis and antibacterial activity of 1-(2,4-dichlorobenzoyl)-4-substituted thiosemicarbazides, 1,2,4-triazoles and their methyl derivatives. J. Heterocycl. Chem. **1984**, *21* (4), 1225–1229.
- Holla, B.S.; Kalluraya, B.; Sridhar, K.R. Synthesis of 4-(5-nitro-2-furfurylidene)amino-3-mercapto-5-(substituted)-1,2,4-triazoles as possible antibacterial agents. Curr. Sci. 1987, 56 (5), 236–238.
- Abdou, N.A.; Amin, F.M. Synthesis and biological activity of certain 5-mercapto-3-(3-pyridyl)-4*H*-1,2,4-triazole derivatives. Mansoura J. Pharm. Sci. **1990**, *6* (2), 25– 29.
- Mishra, R.K.; Tewari, R.K.; Srivastava, S.K.; Bahel, S.C. Synthesis and antifungal activity of some 1,4-disubstitued-thiosemicarbazides, 2,5-disubstituted-1,3,4-thiadiazoles and 3,4-disubstituted-5-mercapto-1,2,4-triazoles. J. Indian Chem. Soc. 1991, 68 (2), 110–112.
- Mansour, A.K.; Eid, M.M.; Hassan, R.A.; Haemers, A.; Pattyn, S.R.; Vanden Berghe, D.A.; Van Hoof, L. Synthesis, reactions and biological evaluation of some 1,2,4-triazine derivatives. J. Heterocycl. Chem. **1988**, *25*, 279–283.
- Holỳ, A.; Bald, R. Nucleic acid components and their analogues. CXXXVIII. Synthesis of 2',3'-cyclic phosphates derived from some pyrimidine ribonucleosides and their behaviour towards pancreatic ribonuclease and ribonuclease T2. Collect. Czechoslov. Chem. Commun. 1971, 36, 2809–2823.
- Gerzon, K.; Kau, D. The adamantly group in medicinal agents. III. Nucleoside 5'adamantoates. The adamantoyl function as a protecting group. J. Med. Chem. 1967, 10, 189–199.
- 9. Witkowski, J.T.; Robins, R.K.; Sidwell, R.W.; Simon, L.N. Design, Synthesis, and

#### 1,2,4-Triazolo[4,3-b][1,2,4]Triazines

broad spectrum antiviral activity of  $1-\beta$ -D-ribofuranosyl-1,2,4-triazole-3-carboxamide and related nucleosides. J. Med. Chem. **1972**, *15* (11), 1150–1154.

- The Pharmacology of Ribavirin; Smith, R.A., Kirkpatrick, W., Eds.; Academic Press: London, 1980; 133–156.
- 11. In vitro and in vivo Inhibition of DNA Viruses by Ribavirin; Sidwell, R.W., Ed.; Academic Press: Orlando, FL, 1984; 19–32.
- 12. *Clinical Applications of Ribavirin*; Smith, R.A., Knight, V., Smith, J.A.D., Eds.; Academic Press: Orlando, FL, 1984; 222.
- 13. Niedballa, U.; Vorbrüggen, H. Pyrimidine Nucleosides. Ger. Offen. 1,919,307, Jan 14, 1971; Chem Abstr. **1971**, *74*, 88267d.
- Vorbrüggen, H.; Niedballa, U. Pyrimidine Nucleosides. S. Afr. Patent 70,02,144, Oct 26, 1970; Chem Abstr. 1971, 75, 20912a.
- Vorbrüggen, H.; Kolb, K.; Niedballa, U.; Strehlke, P. Therapeutical 2-Thio-6-Azauridines. Ger. Offen. 1,955,695, May 13, 1971; Chem Abstr. 1971, 75, 49513g.
- Niedballa, U.; Vorbrüggen, H. Pyrimidine Nucleosides. Ger. Offen. 1,943,428, Feb 25, 1971; Chem Abstr. 1971, 74, 100361q.
- Roy-Durman, P. Analogues of Nucleic Acid Componennts; Springer Verlag: Berlin-Heidelberg, New York, 1970; 42–45.
- Beranek, J.; Sorm, F. Acyl derivatives of 6-Azacytidine. Czech. Patent 110,829, May 15, 1964; Chem Abstr. 1964, 61, 13405f.
- Zabolotnyi, D.K. Institute of Microbiology. 6-Azacytidine. Fr. Patent 1,386,727, Jan 22, 1965, ; Chem Abstr. 1965, 62, 13222h.
- 20. Restivo, A.R. Azauridines. US Patent 3,412,083, Nov 19, 1968; Chem Abstr. **1969**, 70, 47807d.
- Mansour, A.K.; Eid, M.M.; Khalil, N.S.A.M. Selective synthesis and reactions of 6-substituted-2-β-galactosyl-1,2,4-triazines of potential anticancer activity. Nucleosides Nucleotides Nucleic Acids 2003, 22 (1), 21–44; and references cited therein.
- 22. Mansour, A.K.; Eid, M.M.; Khalil, N.S.A.M. Synthesis of some new  $2-\alpha$ -L-arabinopyranosyl-1,2,4-triazines of potential anticancer activity. Nucleosides Nucleotides Nucleic Acids **2003**, 22 (9), 1805–1823.
- 23. Eid, M.M.; Hassan, R.H. Synthesis and condensation reactions of some 6-substituted benzyl-3-hydrazino-2,5-dihydro-1,2,4-triazin-5-ones. Egypt. J. Pharm. Sci. **1990**, *31*, 337–342.
- Mansour, A.K.; Ibrahim, Y.A.; Eid, M.M.; Abdel-Hady, S.A.L. N-Glucosyl derivatives of some 1,2,4-triazines with tetra-*O*-acetyl-α-D-glucopyranosyl bromide. J. Carbohydr. Nucleosides Nucleotides 1981, 8 (2), 81–99.
- Mansour, A.K.; Ibrahim, Y.A.; Eid, M.M. Nucleoside derivatives of 1,2,4-triazine. Reaction of some derivatives of 1,2,4-triazine with tetra-O-acetyl-α-D-glucopyranosyl bromide. Z. Naturforsch. 1976, 31b, 505–508.
- 26. Eid, M.M.; Abdel-Hady, S.A.L.; Ali, H.A.W. Reaction of some 1,2,4-triazines with acetobromoglucose. Arch. Pharm. **1990**, *323* (4), 243–245.
- 27. Ibrahim, Y.A. Facile approach for the selective glycosidation of cyclic asymmetric amides and thioamides. Carbohydr. Lett. **1996**, *1*, 425–432.
- Mansour, A.K.; Ibrahim, Y.A.; Khalil, N.S.A.M. Selective synthesis and structure of 6-arylvinyl-2- and 4- glucosyl-1,2,4-triazines of expected interesting biological activity. Nucleosides Nucleotides 1999, 18 (10), 2256–2283.
- 29. Ibrahim, Y.A.; Abbas, A.A.; Elwahy, A.H.M. Selective synthesis and structure of

2-*N*- and 3-*S*-glucosyl-1,2,4-triazoles of potential biological interest. Carbohydr. Lett. **1999**, *3*, 331–338; and reference 12 cited therein.

- Ibrahim, Y.A. Synthesis and structure of 2-deoxyribosyl-6-azauracil derivatives. Carbohydr. Lett. 1996, 2, 189–195.
- 31. Hiari, K.; Sugimoto, H.; Mizushima, T. Preparation of Triazinone Derivatives as Antiulcer Agents. Jpn Patent 61,134,389, June 21, 1986; Chem Abstr. **1987**, *106*, 67353t.
- 32. Ahmed, R.; Ajaz, S. Synthesis of 6-(*p*-halobenzyl)-1,2,4-triazine-3,5(2*H*,4*H*)dithiones having potential antimicrobial activity. J. Chem. Soc. Pak. **1985**, 7 (4), 315–320.
- Hassan, R.A. Studies on 1,2,4-triazine derivatives. Ph.D. Thesis; Faculty of Science: Cairo University: Giza, Egypt, 1986; 73–123.

Received June 16, 2004 Accepted August 25, 2004