


ChemComm

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the RSC Publishing peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, which is prior to technical editing, formatting and proof reading. This free service from RSC Publishing allows authors to make their results available to the community, in citable form, before publication of the edited article. This *Accepted Manuscript* will be replaced by the edited and formatted *Advance Article* as soon as this is available.

To cite this manuscript please use its permanent Digital Object Identifier (DOI®), which is identical for all formats of publication.

More information about *Accepted Manuscripts* can be found in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics contained in the manuscript submitted by the author(s) which may alter content, and that the standard [Terms & Conditions](#) and the [ethical guidelines](#) that apply to the journal are still applicable. In no event shall the RSC be held responsible for any errors or omissions in these *Accepted Manuscript* manuscripts or any consequences arising from the use of any information contained in them.

Cite this: DOI: 10.1039/c0xx00000x

www.rsc.org/xxxxxx

ARTICLE TYPE

Transamidation of primary amides with amines catalyzed by zirconocene dichloride†

Benjamin N. Atkinson, A. Rosie Chhatwal, Helen V. Lomax, James W. Walton and Jonathan M. J. Williams*

Received (in XXX, XXX) Xth XXXXXXXXXX 20XX, Accepted Xth XXXXXXXXXX 20XX

DOI: 10.1039/b000000x

Zirconocene dichloride (Cp_2ZrCl_2) has been shown to be an effective catalyst for the transamidation of primary amides with amines in cyclohexane at 80 °C in 5-24 hours. For favourable substrates, the reaction can be performed at temperatures as low as 30 °C.












Amides are one of the most important functional groups, being widely found in natural products, pharmaceuticals and materials.¹ In order to avoid the use of stoichiometric activating agents, many catalytic methods for amide synthesis have been developed in recent years.² Amines can react directly with carboxylic acids, most often catalyzed by boronic acid derivatives,³ although metal catalysts, including our own recent work using zirconocene dichloride, can also be used.⁴ There have been several reports of the oxidative coupling of alcohols with amines to give amides,⁵ along with metal-catalyzed rearrangement reactions of oximes which usually only give primary amides.⁶

Amides are relatively inert in comparison with other acyl donors and therefore uncatalyzed transamidation reactions require forcing reaction conditions.⁷ The reaction has been catalyzed by imidazole,⁸ at least for formamide synthesis and by hydroxylamine hydrochloride.⁹ Other non-metal catalysts include boric acid¹⁰ and related compounds.¹¹ Stahl and co-workers have identified several catalysts,¹² including zirconium¹³ and aluminium-based¹⁴ catalysts, where the main focus has been on achieving equilibration between amides. Other catalysts include cerium oxide¹⁵ and Beller's recent report of the use of copper acetate.¹⁶ All of the reported catalysts have some drawback; the need for high temperatures, reversibility or high catalyst loading. Myers has reported that transamidation can be achieved catalytically under mild conditions, but the protocol requires derivatization of the amide with a stoichiometric activating agent.¹⁷

Further to our report of hydroxylamine hydrochloride acting as a catalyst for transamidation, we chose to investigate the use of metal catalysts for this process, with the reaction of primary

amides **1** with benzylamine **2** to give secondary amides **3**. Using toluene as a solvent, many Lewis acids showed at least some catalytic activity,[†] although Cp_2ZrCl_2 and Cp_2HfCl_2 stood out as catalysts capable of achieving close to full conversion in only 5 hours at reflux.

Table 1 Acylation of benzylamine with amides

				
Entry	Amide Product	Time (h)	Conversion ^a (%)	Isolated Yield (%)
1		5	100	83
2		5	100	88
3		8	96	84
4		18	100	93
5		18	89	76
6		18	100	84
7		18	100	86
8		24	87	73
9		18	91	84
10		5	100	92

^a Conversions were determined by analysis of the crude ¹H NMR spectra.

University of Bath, Claverton Down, Bath, BA2 7AY, UK. Email: j.m.j.williams@bath.ac.uk

†Electronic supplementary information (ESI) available: Full experimental details, product characterization and selected NMR spectra are provided. See DOI: 10.1039/b000000x/

A solvent screen[†] identified commercially available anhydrous cyclohexane as a particularly good solvent. The reagent grade solvent performed less well, indicating that the presence of

adventitious water is slightly detrimental to the reaction. Complete conversion was achieved when a small excess of amine **2** was used. The optimized reaction conditions allowed the reaction to be performed at only 80 °C in 5 hours.

Having established a viable catalytic transamidation under relatively mild conditions, we explored the reaction scope. We were pleased to find that benzylamine reacted with a range of primary amides at 80 °C (Table 1). Longer reaction times were required for less reactive amides such as benzamide and related aromatic amides (Table 1, entries 4-7). *N*-Boc prolinamide (Table 1, entry 8) was found to be less reactive, perhaps due to its chelating ability, but we were pleased to see that stereochemistry was fully retained in the secondary amide product.

Table 2 Further transamidation reactions

$\text{R}-\text{C}(=\text{O})\text{NH}_2 + \text{H}_2\text{N}-\text{R}' \xrightarrow[\text{Anhydrous Cyclohexane (1.0 M) 30-100 }^\circ\text{C}]{5 \text{ mol\% Cp}_2\text{ZrCl}_2}$ $\text{R}-\text{C}(=\text{O})\text{NHR}'$				
Entry	Amide product	Time (h)/Temp (°C)	Conversion ^a (%)	Isolated Yield (%)
1		5/80	100	78
2		5/80	100	81
3		5/80	100	69
4		18/80	95	86
5		5/80	100	91
6		5/80	100	89
7		8/80	100	90
8		18/100 ^b	81	54
9		18/100 ^b	80	68
10		5/30	100	88
11		5/30	100	95
12		5/30	100	90
13		5/30	100	89

^a Conversions were determined by analysis of the crude ¹H NMR spectra.

^b These reactions were performed in *n*-heptane.

A range of other amines was used successfully (Table 2). Most substrates underwent conversion into product at 80 °C

although the less nucleophilic amines, *p*-methoxyaniline and 2-aminothiazole (Table 2, entries 8 and 9) required a higher temperature. Remarkably, trifluoroacetamide underwent catalytic transamidation with (*S*)-(-)- α -methylbenzylamine at 30 °C and with retention of stereochemistry (Table 2, entry 10). Formamide

was also converted into the corresponding secondary and tertiary amides at the same temperature (Table 2, entries 11 – 13). Notable limitations[†] include the use of bulky amines (H₂N^tBu with butyramide) and amides (^tBuCONH₂ with benzylamine), which each gave less than 10% conversion into the desired secondary amides in their respective reactions. The transamidation of secondary amides and ureas, along with amidation of simple esters, were found to be less effective under the optimized reaction conditions.


Carbamates, however, were found to be good substrates for conversion into the corresponding mono-substituted ureas – the most convenient substrate was ethyl carbamate and the reaction was successful with primary amines, anilines and with piperidine (Table 3). Heptane was used as solvent in order to accommodate the higher temperature required.

Table 3 Reaction of amines with ethyl carbamate

$\text{H}_2\text{N}-\text{C}(=\text{O})\text{OEt} + \text{H}_2\text{N}-\text{R}' \xrightarrow[\text{Anhydrous Heptane (1.0 M) 100 }^\circ\text{C, 24 h}]{5 \text{ mol\% Cp}_2\text{ZrCl}_2}$ $\text{H}_2\text{N}-\text{C}(=\text{O})\text{NHR}'$			
Entry	Mono-substituted urea	Conversion ^a (%)	Isolated yield (%)
1		100	87
2		100	80
3		96	92
4		100	89
5		100	84

^a Conversions were determined by analysis of the crude ¹H NMR spectra.

Although it seemed unlikely that the amide was attacking the amine, we confirmed this with an isotope labeling study. The reaction of ¹⁵N-labeled benzamide with unlabeled benzylamine led to the exclusive formation of unlabeled secondary amide product (Scheme 1), indicating that the amine adds to the amide.




Scheme 1 Mechanistic studies

In a second experiment, a small amount of H/D exchange was observed when acetamide-d₃ was converted into the secondary amide. It is unlikely that a ketene mechanism plays any significant role in the reaction, as this would be expected to lead to a greater level of H/D exchange (D/H ~ 67/33).

Analysis of the ¹H NMR spectra of Cp₂ZrCl₂ in the presence of varying concentrations of either benzamide or benzylamine revealed that the cyclopentadienyl ligands were not displaced and that both substrates bound fairly weakly (*K*_a = 0.4 M⁻¹ for benzylamine and 1.5 M⁻¹ for benzamide).

A kinetic analysis of the reaction established that the reaction was first order in both Cp₂ZrCl₂ and benzamide. However, the reaction was found to be second order in benzylamine. One explanation for this might be that the amine is used as a base as well as a nucleophile in the reaction, but the reaction rate was found to be zero order with respect to tertiary amine when ⁱPr₃N⁺Et⁻ was added (in fact the rate decreased slightly).

One possible mechanism which is consistent with the data is presented in Scheme 2. Reaction of the zirconium catalyst with the amine leads to formation of a zirconium amide (i.e. Zr-NHR). Complexation of the carboxamide to the catalyst would be followed by addition of the amine to the carboxamide as shown in Scheme 2, intermediate **4**. We speculate that this addition may be assisted by the zirconium amide (Zr-NHR), which would rationalize why the reaction is second order in amine. After loss of ammonia from intermediate **6**, secondary amide would be exchanged from intermediate **7** to complete the catalytic cycle.



Scheme 2 Possible mechanism for transamidation

In summary, we have identified zirconocene dichloride as an effective catalyst for transamidation reactions of primary amides under milder conditions than previously reported.

We thank the EPSRC (BNA) and the Doctoral Training Centre in Sustainable Chemical Technology (HVL) for funding. We thank the University of Bath for further support.

Notes and references

- Many recent reviews including (a) E. Valeur and M. Bradley, *Chem. Soc. Rev.*, 2009, **38**, 606-631;
- C. L. Allen and J. M. J. Williams, *Chem. Soc. Rev.*, 2011, **40**, 3405-3415.
- (a) I. Georgiou, G. Ilyashenko and A. Whiting, *Acc. Chem. Res.*, 2009, **42**, 756-768; (b) K. Arnold, B. Davies, R. L. Giles, C. Grosjean, G. E. Smith and A. Whiting, *Adv. Synth. Catal.* 2006, **348**, 813-820; (c) T. Marcelli, *Angew. Chem. Int. Ed.*, 2010, **49**, 6840-6843.
- (a) C. L. Allen, A. R. Chhatwal and J. M. J. Williams, *Chem. Commun.*, 2012, **48**, 666-668; (b) M. Hosseini-Sarvari and H. Sharghi, *J. Org. Chem.*, 2006, **71**, 6652-6654; (c) T. H. T. Ogawa, T. Ikegami, N. Ono, H. Suzuki, *J. Chem. Soc. Perkin Trans. 1.*, 1994, 3473-3478; (d) A. C. Shekhar, A. R. Kumar, G. Sathaiyah, V. L. Paul, M. Sridhar and P. Shanthan Rao, *Tetrahedron Lett.*, 2009, **50**, 7099-7101; (e) H. Lundberg, F. Tinnis and H. Adolfsson, *Chem. Eur. J.*, 2012, **18**, 3822-3826.
- (a) C. Gunanathan, Y. Ben-David and D. Milstein, *Science*, 2007, **317**, 790-792; (b) L. U. Nordström, H. Vogt and R. Madsen, *J. Am. Chem. Soc.*, 2008, **130**, 17672-17673; (c) T. Zweifel, J.-V. Naubron and H. Grützmaier, *Angew. Chem. Int. Ed.*, 2009, **48**, 559-563; (d) A. J. A. Watson, A. C. Maxwell and J. M. J. Williams, *Org. Lett.*, 2009, **11**, 2667-2670;
- For recent examples see; (a) C. L. Allen, R. Lawrence, L. Emmett and J. M. J. Williams, *Adv. Synth. Catal.*, 2011, **353**, 3262-3268; (b) Gnanamgari and R. H. Crabtree, *Organometallics.*, 2009, **28**, 922-924; (c) M. A. Ali and T. Punniyamurthy, *Adv. Synth. Catal.*, 2010, **352**, 288-292; (d) R. N. S. Ramón, J. Bosson, S. Díez-González, N. Marion and S. P. Nolan, *J. Org. Chem.*, 2010, **75**, 1197-1202;
- (a) A. Galat and G. Elion, *J. Am. Chem. Soc.*, 1943, **65**, 1566-1567. For an example of a reactive amide, see; (b) M. Hutchby, C. E. Houlden, M. F. Haddow, S. N. G. Tyler, G. C. Lloyd-Jones and K. I. Booker-Milburn, *Angew. Chem. Int. Ed.*, 2012, **51**, 548-551.
- M. R. Suchý, A. A. H. Elmehrikki and R. H. E. Hudson, *Org. Lett.*, 2011, **13**, 3952-3955.
- C. L. Allen, B. N. Atkinson and J. M. J. Williams, *Angew. Chem. Int. Ed.*, 2012, **51**, 1383-1386.
- T. B. Nguyen, J. Sorres, M. Q. Tran, L. Ermolenko and A. Al-Mourabit, *Org. Lett.*, 2012, **14**, 3202-3205.
- (a) P. Starkov and T. D. Sheppard, *Org. Biomol. Chem.*, 2011, **9**, 1320-1323; (b) S. Çalimsiz and M. A. Lipton, *J. Org. Chem.*, 2005, **70**, 6218-6221; (c) A. Klapars, S. Parris, K. W. Anderson and S. L. Buchwald, *J. Am. Chem. Soc.*, 2004, **126**, 3529-3533; (d) A. Vasudevan, C. I. Villamil and S. W. Djuric, *Org. Lett.*, 2004, **6**, 3361-3364; (e) J. Lasri, M. E. González-Rosende and J. Sepúlveda-Arques, *Org. Lett.*, 2003, **5**, 3851-3853.
- (a) S. E. Eldred, D. A. Stone, S. H. Gellman and S. S. Stahl, *J. Am. Chem. Soc.*, 2003, **125**, 3422-3423; (b) D. A. Kissounko, I. A. Guzei, S. H. Gellman and S. S. Stahl, *Organometallics*, 2005, **24**, 5208-5210; (c) D. A. Kissounko, J. M. Hoerter, I. A. Guzei, Q. Cui, S. H. Gellman and S. S. Stahl, *J. Am. Chem. Soc.*, 2007, **129**, 1776-1783.
- N. A. Stephenson, J. Zhu, S. H. Gellman and S. S. Stahl, *J. Am. Chem. Soc.*, 2009, **131**, 10003-10008.
- (a) J. M. Hoerter, K. M. Cui, S. H. Gellman, Q. Cui and S. S. Stahl, *J. Am. Chem. Soc.*, 2007, **130**, 647-654; (b) J. M. Hoerter, K. M. Otte, S. H. Gellman and S. S. Stahl, *J. Am. Chem. Soc.*, 2006, **128**, 5177-5183. See also; (c) E. Bon, D. C. H. Bigg and G. Bertrand, *J. Org. Chem.*, 1994, **59**, 4035-4036.
- M. Tamura, T. Tonomura, K.-I. Shimizu and A. Satsuma, *Green Chem.*, 2012, **14**, 717-724.
- M. Zhang, S. Imm, S. Bähn, L. Neubert, H. Neumann and M. Beller, *Angew. Chem. Int. Ed.*, 2012, **51**, 3905-3910.
- T. A. Dineen, M. A. Zajac and A. G. Myers, *J. Am. Chem. Soc.*, 2006, **128**, 16406-16409.

[View Online](#)

Zirconocene dichloride (Cp_2ZrCl_2) has been shown to be an effective catalyst for the transamidation of primary amides with amines in cyclohexane at 80 °C in 5-24 hours. For favourable substrates, the reaction can be performed at temperatures as low as 30 °C.

